

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/23643868>

# Notes on Fermi–Dirac Integrals

Article · November 2008

Source: arXiv

---

CITATIONS

35

---

READS

928

2 authors:



Raseong Kim

Intel

67 PUBLICATIONS 648 CITATIONS

SEE PROFILE



M.s. Lundstrom

Purdue University

527 PUBLICATIONS 17,511 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Scientific Software Development for nanoHUB [View project](#)

All content following this page was uploaded by [M.s. Lundstrom](#) on 19 August 2014.

The user has requested enhancement of the downloaded file.

# Notes on Fermi-Dirac Integrals

## 2<sup>nd</sup> edition

Raseong Kim and Mark Lundstrom  
Network for Computational Nanotechnology  
Purdue University

June 27, 2008

### 1. Introduction

Fermi-Dirac integrals appear frequently in semiconductor problems, so an understanding of their properties is essential. The purpose of these notes is to collect in one place, some basic information about Fermi-Dirac integrals and their properties. To see how they arise, consider computing the equilibrium electron concentration per unit volume in a three-dimensional semiconductor with a parabolic conduction band from the expression,

$$n = \int_{E_C}^{\infty} g(E) f_0(E) dE = \int_{E_C}^{\infty} \frac{g(E) dE}{1 + e^{(E-E_F)/k_B T}}, \quad (1)$$

where  $g(E)$  is the density of states,  $f_0(E)$  is the Fermi function, and  $E_C$  is the conduction band edge. For three dimensional electrons,

$$g_{3D}(E) = \frac{(2m^*)^{3/2}}{2\pi^2 \hbar^3} \sqrt{E - E_C} \quad (2)$$

which can be used in (1) to write

$$n = \frac{(2m^*)^{3/2}}{2\pi^2 \hbar^3} \int_{E_C}^{\infty} \frac{\sqrt{E - E_C} dE}{1 + e^{(E-E_F)/k_B T}}. \quad (3)$$

By making the substitution,

$$\varepsilon = (E - E_C)/k_B T \quad (4)$$

eqn. (3) becomes

$$n = \frac{(2m^* k_B T)^{3/2}}{2\pi^2 \hbar^3} \int_0^{\infty} \frac{\varepsilon^{1/2} d\varepsilon}{1 + e^{\varepsilon - \eta_F}}, \quad (5)$$

where we have defined

$$\eta_F \equiv (E_F - E_C)/k_B T. \quad (6)$$

By collecting up parameters, we can express the electron concentration as

$$n_0 = N_{3D} \frac{2}{\sqrt{\pi}} F_{1/2}(\eta_F) \quad (7)$$

where

$$N_{3D} = 2 \left( \frac{2\pi m^* k_B T}{h^2} \right)^{3/2} \quad (8)$$

is the so-called effective density-of-states and

$$F_{1/2}(\eta_F) \equiv \int_0^\infty \frac{\varepsilon^{1/2} d\varepsilon}{1 + \exp(\varepsilon - \eta_F)} \quad (9)$$

is the Fermi-Dirac integral of order 1/2. This integral can only be done numerically. Note that its value depends on  $\eta_F$ , which measures the location of the Fermi level with respect to the conduction band edge. It is more convenient to define a related integral,

$$\mathcal{F}_{1/2}(\eta_F) \equiv \frac{2}{\sqrt{\pi}} \int_0^\infty \frac{\varepsilon^{1/2} d\varepsilon}{1 + \exp(\varepsilon - \eta_F)} \quad (10)$$

so that eqn. (7) can be written as

$$n = N_{3D} \mathcal{F}_{1/2}(\eta_F). \quad (11)$$

It is important to recognize whether you are dealing with the “Roman” Fermi-Dirac integral or the “script” Fermi-Dirac integral.

There are many kinds of Fermi-Dirac integrals. For example, in two dimensions, the density-of-states is

$$g_{2D}(E) = \frac{m^*}{\pi \hbar^2}, \quad (12)$$

and by following a procedure like that one we used in three dimensions, one can show that the electron density per unit area is

$$n_s = N_{2D} \mathcal{F}_0(\eta_F) \quad (13)$$

where

$$N_{2D} = \frac{m^* k_B T}{\pi \hbar^2}, \quad (14)$$

and

$$\mathcal{F}_0(\eta_F) = \int_0^\infty \frac{\varepsilon^0 d\varepsilon}{1 + e^{\varepsilon - \eta_F}} = \ln(1 + e^{\eta_F}) \quad (15)$$

is the Fermi-Dirac integral of order 0, which can be integrated analytically.

Finally, in one dimension, the density-of-states is

$$g_{1D}(E) = \frac{\sqrt{2m^*}}{\pi \hbar} \sqrt{\frac{1}{E - E_C}} \quad (16)$$

and the equilibrium electron density per unit length is

$$n_L = N_{1D} \mathcal{F}_{-1/2}(\eta_F) \quad (17)$$

where

$$N_{1D} = \frac{1}{\hbar} \sqrt{\frac{2m^* k_B T}{\pi}} \quad (18)$$

and

$$\mathcal{F}_{-1/2}(\eta_F) = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{\varepsilon^{-1/2} d\varepsilon}{1 + e^{\varepsilon - \eta_F}} \quad (19)$$

is the Fermi-Dirac integral of order  $-1/2$ , which must be integrated numerically.

## 2. General Definition

In the previous section, we saw three examples of Fermi-Dirac integrals. More generally, we define

$$\mathcal{F}_j(\eta_F) \equiv \frac{1}{\Gamma(j+1)} \int_0^\infty \frac{\varepsilon^j d\varepsilon}{1 + \exp(\varepsilon - \eta_F)}, \quad (20)$$

where  $\Gamma$  is the gamma function. The  $\Gamma$  function is just the factorial when its argument is a positive integer,

$$\Gamma(n) = (n-1)! \quad (\text{for } n \text{ a positive integer}). \quad (21a)$$

Also

$$\Gamma(1/2) = \sqrt{\pi} \quad (21b)$$

and

$$\Gamma(p+1) = p\Gamma(p) \quad (21c)$$

As an example, let's evaluate  $\mathcal{F}_{1/2}(\eta_F)$  from eqn. (20):

$$\mathcal{F}_{1/2}(\eta_F) \equiv \frac{1}{\Gamma(1/2+1)} \int_0^\infty \frac{\varepsilon^{1/2} d\varepsilon}{1 + e^{\varepsilon - \eta_F}}, \quad (22a)$$

so we need to evaluate  $\Gamma(3/2)$ . Using eqns. (21b) and (21c), we find,

$$\Gamma(3/2) = \Gamma(1/2+1) = \frac{1}{2} \Gamma(1/2) = \frac{\sqrt{\pi}}{2}, \quad (22b)$$

so  $\mathcal{F}_{1/2}(\eta_F)$  is evaluated as

$$\mathcal{F}_{1/2}(\eta_F) \equiv \frac{2}{\sqrt{\pi}} \int_0^\infty \frac{\varepsilon^{1/2} d\varepsilon}{1 + e^{\varepsilon - \eta_F}}, \quad (22c)$$

which agrees with eqn. (10). For more practice, use the general definition, eqn. (20) and eqns. (21a-c) to show that the results for  $\mathcal{F}_0(\eta_F)$  and  $\mathcal{F}_{-1/2}(\eta_F)$  agree with eqns. (15) and (19).

### 3. Derivatives of Fermi-Dirac Integrals

Fermi-Dirac integrals have the property that

$$\frac{d\mathcal{F}_j}{d\eta_F} = \mathcal{F}_{j-1}, \quad (23)$$

which often comes in useful. For example, we have an analytical expression for  $\mathcal{F}_0(\eta_F)$ , which means that we have an analytical expression for  $\mathcal{F}_{-1}(\eta_F)$ ,

$$\mathcal{F}_{-1} = \frac{d\mathcal{F}_0}{d\eta_F} = \frac{1}{1 + e^{-\eta_F}}. \quad (24)$$

Similarly, we can show that there is an analytic expression for any Fermi-Dirac integral of integer order,  $j$ , for  $j \leq -2$ ,

$$\mathcal{F}_j(\eta_F) = \frac{e^{\eta_F}}{(1 + e^{\eta_F})^{-j}} P_{-j-2}(e^{\eta_F}) \quad (25)$$

where  $P_k$  is a polynomial of degree  $k$ , and the coefficients  $p_{k,i}$  are generated from a recurrence relation [1]

$$p_{k,0} = 1 \quad (26a)$$

$$p_{k,i} = (1+i)p_{k-1,i} - (k+1-i)p_{k-1,i-1} \quad i = 1, \dots, k. \quad (26b)$$

#### 4. Asymptotic Expansions for Fermi-Dirac Integrals

It is useful to examine Fermi-Dirac integrals in the non-degenerate ( $\eta_F \ll 0$ ) and degenerate ( $\eta_F \gg 0$ ) limits. For the non-degenerate limit, the result is particularly simple,

$$\mathcal{F}_j(\eta_F) \rightarrow e^{\eta_F} \quad (27)$$

which means that for all orders,  $j$ , the function approaches the exponential in the non-degenerate limit. To examine Fermi-Dirac integrals in the degenerate limit, we consider the complete expansion for the Fermi-Dirac integral for  $j > -1$  and  $\eta_F > 0$  [2, 3]

$$\mathcal{F}_j(\eta_F) = 2\eta_F^{j+1} \sum_{n=0}^{\infty} \frac{t_{2n}}{\Gamma(j+2-2n)\eta_F^{2n}} + \cos(\pi j) \sum_{n=1}^{\infty} \frac{(-1)^{n-1} e^{-n\eta_F}}{n^{j+1}} \quad (28)$$

where  $t_0 = 1/2$ ,  $t_n = \sum_{\mu=1}^{\infty} (-1)^{\mu-1} / \mu^n = (1 - 2^{1-n}) \zeta(n)$ , and  $\zeta(n)$  is the Riemann zeta function. The expressions for the Fermi-Dirac integrals in the degenerate limit ( $\eta_F \gg 0$ ) come from (28) as  $\mathcal{F}_j(\eta_F) \rightarrow \eta_F^{j+1} / \Gamma(j+2)$  [4]. Specific results for several Fermi-Dirac integrals are shown below.

$$\mathcal{F}_{-1/2}(\eta_F) \rightarrow \frac{2\eta_F^{1/2}}{\sqrt{\pi}} \quad (29a)$$

$$\mathcal{F}_{1/2}(\eta_F) \rightarrow \frac{4\eta_F^{3/2}}{3\sqrt{\pi}} \quad (29b)$$

$$\mathcal{F}_1(\eta_F) \rightarrow \frac{1}{2}\eta_F^2 \quad (29c)$$

$$\mathcal{F}_{3/2}(\eta_F) \rightarrow \frac{8\eta_F^{5/2}}{15\sqrt{\pi}} \quad (29d)$$

$$\mathcal{F}_2(\eta_F) \rightarrow \frac{1}{6}\eta_F^3 \quad (29e)$$

Now we relate the complete expansion in (28) to the Sommerfeld expansion [5, 6]. The Sommerfeld expansion for a function  $H(\varepsilon)$  is expressed as

$$\int_0^\infty H(\varepsilon) f_0(\varepsilon) d\varepsilon = \int_0^{\eta_F} H(\varepsilon) d\varepsilon + \sum_{n=1}^\infty a_n \frac{d^{2n-1}}{d\varepsilon^{2n-1}} H(\varepsilon) \Big|_{\varepsilon=\eta_F} \quad (30)$$

where

$$a_n = 2 \left( 1 - \frac{1}{2^{2n}} + \frac{1}{3^{2n}} - \frac{1}{4^{2n}} + \dots \right), \quad (31)$$

and it is noted that  $a_n = 2t_{2n}$ . Then the Sommerfeld expansion for the Fermi-Dirac integral of order  $j$  can be evaluated by letting  $H(\varepsilon) = \varepsilon^j / \Gamma(j+1)$ , and the result is

$$\mathcal{F}_j(\eta_F) = 2\eta_F^{j+1} \sum_{n=0}^\infty \frac{t_{2n}}{\Gamma(j+2-2n)\eta_F^{2n}}. \quad (32)$$

Equation (32) is the same as the (28) except that the second term in (28) is omitted [3]. In the degenerate limit, however, the second term in (28) vanishes, so the (28) and (32) give the same results as (29a-e).

## 5. Approximate Expressions for Common Fermi-Dirac Integrals

The Fermi-Dirac integral can be quickly evaluated by tabulation [2, 4, 7, 8] or analytic approximation [9-11]. We briefly mention some of the analytic approximations and refer a Matlab script. Bednarczyk *et al.* [9] proposed a single analytic approximation which evaluates

the Fermi-Dirac integral of order  $j = 1/2$  with errors less than 0.4 % [12]. Aymerich-Humet *et al.* [10, 11] introduced an analytic approximation for a general  $j$ , and it gives an error of 1.2 % for  $-1/2 < j < 1/2$  and 0.7 % for  $1/2 < j < 5/2$ , and the error increases with larger  $j$ . The Matlab function, "FD\_int\_approx.m," calculates the Fermi-Dirac integral defined in (10) with orders  $j \geq -1/2$  using these analytic approximations.

If a better accuracy is required while keeping the calculation relatively simple, the approximations proposed by Halen and Pulfrey [13, 14] may be used. In this model, several approximate expressions are introduced based on the series expansion in (28), and the error is less than  $10^{-5}$  for  $-1/2 \leq j \leq 7/2$  [13]. The Matlab function, "FDjx.m," is the main function which calculates the Fermi-Dirac integrals using this model.

## 6. Numerical Evaluation of Fermi-Dirac Integrals

The Fermi-Dirac integrals can be evaluated accurately by numerical integration. Here we briefly review the approach by Press *et al.* for generalized Fermi-Dirac integrals with order  $j > -1$  [15]. In this approach, the composite trapezoidal rule with variable transformation  $\varepsilon = \exp(t - e^{-t})$  is used for  $\eta_F \leq 15$ , and the double exponential (DE) rule is used for larger  $\eta_F$ . The double precision (eps,  $\sim 2.2 \times 10^{-16}$ ) can be achieved after 60 to 500 iterations [15]. The Matlab function, "FD\_int\_num.m," evaluates the Fermi-Dirac integral numerically using the composite trapezoidal rule following the approach in [15].

## 7. References

- [1] G. Michele, "Algorithm 745: computation of the complete and incomplete Fermi-Dirac integral," *ACM Trans. Math. Softw.*, vol. 21, pp. 221-232, 1995.
- [2] R. Dingle, "The Fermi-Dirac integrals  $\mathcal{F}_p(\eta) = (p!)^{-1} \int_0^\infty \varepsilon^p (e^{\varepsilon - \eta} + 1)^{-1} d\varepsilon$ ," *Applied Scientific Research*, vol. 6, pp. 225-239, 1957.
- [3] R. B. Dingle, *Asymptotic Expansions: Their Derivation and Interpretation*. London: Academic Press, 1973.
- [4] J. McDougall and E. C. Stoner, "The computation of Fermi-Dirac functions," *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, vol. 237, pp. 67-104, 1938.
- [5] A. Sommerfeld, "Zur Elektronentheorie der Metalle auf Grund der Fermischen Statistik," *Zeitschrift für Physik A Hadrons and Nuclei*, vol. 47, pp. 1-32, 1928.
- [6] N. W. Ashcroft and N. D. Mermin, *Solid State Physics*. Philadelphia: Saunders College Publishing, 1976.



- [7] A. C. Beer, M. N. Chase, and P. F. Choquard, "Extension of McDougall-Stoner tables of the Fermi-Dirac functions," *Helvetica Physica Acta*, vol. 28, pp. 529-42, 1955.
- [8] P. Rhodes, "Fermi-Dirac functions of integral order," *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, vol. 204, pp. 396-405, 1950.
- [9] D. Bednarczyk and J. Bednarczyk, "The approximation of the Fermi-Dirac integral  $\mathcal{F}_{1/2}(\eta)$ ," *Physics Letters A*, vol. 64, pp. 409-410, 1978.
- [10] X. Aymerich-Humet, F. Serra-Mestres, and J. Millán, "An analytical approximation for the Fermi-Dirac integral  $F_{3/2}(\eta)$ ," *Solid-State Electronics*, vol. 24, pp. 981-982, 1981.
- [11] X. Aymerich-Humet, F. Serra-Mestres, and J. Millan, "A generalized approximation of the Fermi-Dirac integrals," *Journal of Applied Physics*, vol. 54, pp. 2850-2851, 1983.
- [12] J. S. Blakemore, "Approximations for Fermi-Dirac integrals, especially the function  $\mathcal{F}_{1/2}(\eta)$  used to describe electron density in a semiconductor," *Solid-State Electronics*, vol. 25, pp. 1067-1076, 1982.
- [13] P. V. Halen and D. L. Pulfrey, "Accurate, short series approximations to Fermi-Dirac integrals of order -1/2, 1/2, 1, 3/2, 2, 5/2, 3, and 7/2," *Journal of Applied Physics*, vol. 57, pp. 5271-5274, 1985.
- [14] P. Van Halen and D. L. Pulfrey, "Erratum: "Accurate, short series approximation to Fermi-Dirac integrals of order -1/2, 1/2, 1, 3/2, 2, 5/2, 3, and 7/2" [J. Appl. Phys. 57, 5271 (1985)]," *Journal of Applied Physics*, vol. 59, p. 2264, 1986.
- [15] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing*, 3rd ed. New York: Cambridge University Press, 2007.