

bounds is in order. While the lower bound would be particularly important, the improved upper bound would also be useful.

Applied Mathematics Laboratory  
David Taylor Model Basin  
Washington 7, D. C.

1. DANIEL SHANKS, "On the conjecture of Hardy & Littlewood concerning the number of primes of the form  $n^2 + a$ ," *Math. Comp.*, v. 14, 1960, p. 321-332.
2. DANIEL SHANKS & JOHN W. WRENCH, JR., "The calculation of certain Dirichlet series," *Math. Comp.*, v. 17, 1963, p. 136-154.
3. JOHN W. WRENCH, JR., "Evaluation of Artin's constant and the twin-prime constant," *Math. Comp.*, v. 15, 1961, p. 396-398.
4. PAUL T. BATEMAN & ROSEMARIE M. STEMMLER, "Waring's problem in algebraic number fields and primes of the form  $(p^r - 1)/(p^d - 1)$ ," *Illinois J. Math.*, v. 6, 1962, p. 142-156.

## An Approximation to the Fermi Integral $F_{1/2}(x)$

By H. Werner and G. Raymann

The Fermi Integral as defined, for instance, in the *Handbuch der Physik*, Bd. XX, S. 58 [1], is given by

$$(1) \quad F_p(x) = \int_0^\infty \frac{t^p}{e^{t-x} + 1} dt.$$

The function  $F_{1/2}(x)$  has for negative values of  $x$  an expansion of the form

$$(2) \quad F_{1/2}(x) = \frac{\sqrt{\pi}}{2} \sum_{v=1}^{\infty} (-1)^{v-1} \cdot \frac{e^{vx}}{v^{3/2}},$$

and for large positive  $x$  the asymptotic expansion

$$(3) \quad F_{1/2}(x) \sim x^{3/2} \left[ \frac{2}{3} + \frac{\pi^2}{12 \cdot x^2} + \left( \frac{1}{2} \right) \cdot \frac{7}{60} \cdot \frac{\pi^4}{x^4} + \dots \right. \\ \left. + \left( 2n - \frac{1}{2} \right) \frac{2^{2n-1} - 1}{n} |B_{2n}| \cdot \frac{\pi^{2n}}{x^{2n}} + \dots \right];$$

compare [2], formulas (10) and (12);

$B_{2n}$  are the Bernoulli numbers, given for example in [3], page 298. We obtained Chebyshev approximations to  $F_{1/2}(x)$ , based upon the table by McDougall and Stoner [4]. This table was subtabulated by interpolation with a fifth-degree polynomial. The approximations are

$$(4) \quad F_{1/2}^*(x) = e^x \sum_{v=0}^5 a_v e^{vx} \quad \text{for } -\infty < x \leq +1, \\ F_{1/2}^*(x) = x^{3/2} \left[ \frac{2}{3} + \sum_{v=0}^5 \frac{b_v}{x^{2v+2}} \right] \quad \text{for } +1 < x < +\infty,$$

---

Received March 27, 1961. This note was prepared at the AEG Forschungs-Institut, Frankfurt (Main).

the coefficients

$\nu$	$a_\nu$	$b_\nu$
0	+0.8860 7596	+0.8435 00
1	-0.3087 1705	+0.7108 09
2	+0.1463 8520	-3.7124 56
3	-0.0584 3877	+6.7056 28
4	+0.0143 1771	-5.5948 77
5	-0.0015 0176	+1.7777 87

With these approximations, the relative error  $|F_{1/2}(x) - F_{1/2}^*(x)|/F_{1/2}(x)$  is less than  $2 \cdot 10^{-4}$  and  $5 \cdot 10^{-4}$ , respectively.

Another intensive table of  $F_p(x)$  has been given by G. A. Chisnall [5] who also discusses in [6] a method for the interpolation of the existing tables of  $F_{1/2}(x)$ . It is not difficult to obtain analogous Chebyshev approximations to  $F_p(x)$  for any fixed values of  $p$  to a prescribed degree of accuracy if one is able to generate the function with this (or slightly more) accuracy.

Institut für Angewandte Mathematik  
der Universität Hamburg, Hamburg, and  
Deutsches Rechenzentrum  
Darmstadt, Germany

1. O. MADELUNG, Halbleiter in *Handbuch der Physik*, herausgegeben von S. Flügge, Bd. XX, Springer, Berlin, 1957, p. 58-59.
2. A. C. BEER, M. N. CHASE & P. F. CHOQUARD, "Extension of McDougall-Stoner tables of the Fermi-Dirac functions," *Phys. Acta Helv.* 28, 1955, p. 529-542.
3. E. JAHNKE & F. EMDE, unter Mitwirkung von F. Lösch, *Tafeln höherer Funktionen*, B. G. Teubner Verlagsgesellschaft, Stuttgart, 1960.
4. J. McDOUGALL & E. C. STONER, "The computation of Fermi-Dirac functions," *Trans. Roy. Soc., Ser. A* 237, 1939, p. 67-104.
5. G. A. CHISNALL, *Jodrell Bank Annals*, v. 1, 1956, p. 126-140.
6. G. A. CHISNALL, "A modified Chebyshev Everett interpolation formula," *MTAC* v. 10, 1956, p. 66-73.

## On the Congruences $(p-1)! \equiv -1$ and $2^{p-1} \equiv 1 \pmod{p^2}$

By Erna H. Pearson

The results of computations to determine primes  $p$  such that one of the relations

- (1)  $(p-1)! \equiv -1 \pmod{p^2}$ ,
- (2)  $2^{p-1} \equiv 1 \pmod{p^2}$

holds have been published previously [1-5]. The known Wilson primes (those satisfying (1)) are 5, 13, and 563, the last having been determined by Goldberg [3] in testing  $p < 10^4$ . Froberg [4] tested  $10^4 < p < 30,000$  without finding additional Wilson primes.

Froberg [4] determined  $p = 1093$  and  $p = 3511$  to be the only primes less than

---

Received April 6, 1962.