

Warsaw University of Life Sciences WULS – SGGW  
in Warsaw  
Faculty of Forestry

Eberswalde University for Sustainable Development – HNEE  
University of Applied Sciences  
Faculty of Forest and Environment

Tobias Seydewitz  
Album number SGGW: 178311  
Album number HNEE: 15210024

# Kompleksowa analiza wylesiania w krajach tropikalnych - bezpośrednie czynniki wylesiania, emisje dwutlenku węgla i równowaga wartości usług ekosystemów

A comprehensive study on deforestation in the tropics - direct deforestation drivers, carbon emissions and ecosystem service value balance

Master's Thesis  
on the course of - Forestry

Thesis written under the supervision of  
Dr. Prajal Pradhan  
Potsdam Institute of Climate Impact Research  
Research Domain II - Climate Climate Impacts & Vulnerabilities

Potsdam, 2018



### **Oświadczenie promotora pracy**

Oświadczam, że niniejsza praca została przygotowana pod moim kierunkiem i stwierdzam, że spełnia warunki do przedstawienia tej pracy w postępowaniu o nadanie tytułu zawodowego.

### **Declaration of the promoter**

I declare that this thesis was prepared under my supervision and I state that it meets the conditions for presenting such a body of work in the process of obtaining a professional title.

### **Erklärung des Betreuers**

Hiermit erkläre ich, dass die vorliegende Arbeit, unter meiner Leitung erstellt wurde und ich bestätige, dass sie die Bedingungen zur Verleihung des Abschlussdiploms erfüllt.

Data .....	Podpis promotora pracy .....
Date .....	Signature of the promoter .....
Datum .....	Unterschrift des Betreuers .....



## **Oświadczenie autora pracy**

Świadom odpowiedzialności prawnej, w tym odpowiedzialności karnej za złożenie fałszywego oświadczenia, oświadczam, że niniejsza praca dyplomowa została napisana przeze mnie samodzielnie i nie zawiera treści uzyskanych w sposób niezgodny z obowiązującymi przepisami prawa, w szczególności ustawą z dnia 4 lutego 1994 r.o prawie autorskim i prawach pokrewnych (Dz. U. Nr 90 poz. 631 z późn. zm.).

Oświadczam, że przedstawiona praca nie była wcześniej podstawą żadnej procedury związanej z nadaniem dyplomu lub uzyskaniem tytułu zawodowego test. Lorem ipsum dolores anno sacntum

Oświadczam, że niniejsza wersja pracy jest identyczna z załączoną wersją elektroniczną. Przyjmuję do wiadomości, że praca dyplomowa poddana zostanie procedurze antyplagiatoowej.

## **Declaration of the author**

Aware of the legal liability, including criminal liability for submitting a false statement, I declare that this thesis was written by myself alone and does not contain content obtained in a manner breaking applicable laws, in particular the Act of February 4, 1994 on copyright and related rights (Journal of Laws, no. 90, item 631, as amended).

I certify that the work has not previously been the basis for any procedure in connection with obtaining a diploma or professional title.

I declare that this version of the work is identical with the attached electronic version.

I acknowledge that the thesis is subject to anti-plagiarism procedures.

## **Erklärung des Autors**

Gesetzlicher Haftpflicht, besonders der strafrechtlichen Verantwortlichkeit für Abgabe der falschen Erklärung bewusst, erkläre ich hiermit, dass vorliegende Diplomarbeit selbständig angefertigt wurde und keinen Inhalt enthält, der widerrechtlich erworben wurde, insbesondere nicht mit dem Gesetz über Urheberrecht vom 4. Februar 1994 (GB. Nr. 90, Pos. 631 mit späteren Änderungen) übereinstimmend.

Ich erkläre auch, dass die Arbeit bisher keiner anderen Prüfungsbehörde vorgelegt wurde.

Der Durchführung einer elektronischen Plagiatsprüfung stimme ich hiermit zu. Die eingereichte elektronische Fassung der Arbeit entspricht der eingereichten schriftlichen Fassung exakt.

Data .....	Podpis autora pracy .....
Date .....	Signature of the author .....
Datum .....	Unterschrift des Autors .....







## **Streszczenie**

**Tytuł:** Text

Text

Słowa kluczowe: Text



## **Summary**

**Title:** Text

Text

Keywords: Text



## **Zusammenfassung**

**Titel:** Text

Text

Schlüsselwörter: Text








# Contents

1	Introduction	11
1.1	Tropical forest . . . . .	11
1.1.1	Current state . . . . .	11
1.1.2	Contribution to climate . . . . .	12
1.1.3	Forest definitions . . . . .	12
1.2	Deforestation . . . . .	13
1.2.1	Land use and land cover change . . . . .	13
1.2.2	Drivers of deforestation . . . . .	14
1.2.3	Emissions through deforestation . . . . .	15
1.2.4	Removal of AGB . . . . .	15
1.2.5	Soil organic carbon change and soil dynamics . . . . .	16
1.3	Ecosystem services . . . . .	16
1.3.1	Ecosystem service values . . . . .	16
1.3.2	Research objective and questions . . . . .	17
2	Data and methods	18
2.1	Data . . . . .	18
2.1.1	Spatial data . . . . .	18
2.1.1.1	Global Forest Change . . . . .	18
2.1.1.2	GlobeLand30 . . . . .	20
2.1.1.3	Intact Forest Landscapes . . . . .	21
2.1.1.4	Aboveground Woody Biomass . . . . .	22
2.1.1.5	Global Soil Organic Carbon . . . . .	23
2.1.1.6	Auxiliary . . . . .	24
2.1.2	Empirical data . . . . .	24
2.1.2.1	Soil Organic Carbon . . . . .	24
2.1.2.2	Ecosystem Service Values . . . . .	25
2.2	Methods . . . . .	25
2.2.1	Pre-processing . . . . .	25
2.2.2	Deforestation . . . . .	26
2.2.2.1	Forest definition . . . . .	26
2.2.2.2	Land use change driver . . . . .	27
2.2.2.3	Accuracy assessment . . . . .	27
2.2.3	Emissions . . . . .	28
2.2.3.1	Above ground biomass . . . . .	28
2.2.3.2	Soil organic carbon change . . . . .	28
2.2.4	Ecosystem service values . . . . .	29
2.2.4.1	Ecosystem service value loss . . . . .	29

2.2.4.2	Ecosystem service value gain . . . . .	29
2.2.5	Binning analysis . . . . .	30
<b>3</b>	<b>Results</b>	<b>34</b>
3.1	Forest definition and accuracy assessment . . . . .	34
3.2	Deforestation drivers . . . . .	35
3.2.1	Global . . . . .	35
3.2.2	Americas . . . . .	35
3.2.3	Asia . . . . .	35
3.2.4	Africa . . . . .	35
3.3	Deforestation emissions . . . . .	41
3.3.1	Global . . . . .	42
3.3.2	Americas . . . . .	42
3.3.3	Asia . . . . .	42
3.3.4	Africa . . . . .	42
3.4	Ecosystem service value balance . . . . .	42
3.4.1	Global . . . . .	42
3.4.2	Americas . . . . .	42
3.4.3	Asia . . . . .	42
3.4.4	Africa . . . . .	42
<b>4</b>	<b>Discussion</b>	<b>43</b>
<b>Acknowledgements</b>		<b>44</b>
<b>Bibliography</b>		<b>I</b>
<b>List of Figures</b>		<b>III</b>
<b>List of Tables</b>		<b>IV</b>
<b>List of Abbreviations</b>		<b>V</b>
<b>Appendix</b>		<b>VII</b>

# 1 Introduction

## 1.1 Tropical forest



**Figure 1.1.** Geographic tropical zone framed red and the tropical forest

### 1.1.1 Current state

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placemat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor

sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.1.2 Contribution to climate

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, plac-erat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.


### 1.1.3 Forest definitions

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, plac-erat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

## 1.2 Deforestation



**Figure 1.2.** Upper Brazil agriculture, middle Brazil urbanization, lower Indonesia large scale palm oil plantations

### 1.2.1 Land use and land cover change

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy

eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.2.2 Drivers of deforestation

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placemat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.2.3 Emissions trough deforestation

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placet ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.2.4 Removal of AGB

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placet ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.2.5 Soil organic carbon change and soil dynamics

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, plac-erat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

## 1.3 Ecosystem services

### 1.3.1 Ecosystem service values

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, plac-erat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum

sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 1.3.2 Research objective and questions

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placet ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

## 2 Data and methods

### 2.1 Data


#### 2.1.1 Spatial data

##### 2.1.1.1 Global Forest Change

Global Forest Change (GFC) 2000-2012 Version 1.0 is the first high-resolution dataset that provides a comprehensive view of the annual global forest cover change between 2000 and 2012 [Hansen et al. 2013; Li et al. 2017]. The initial GFC dataset released by Hansen et al. is extended by recent releases which encompass the annual forest cover changes between 2000-2013, 2000-2014, 2000-2015 and 2000-2016, respectively. All versions of this dataset have in common, that they are derived from growing season imagery captured by the remote sensing satellite Landsat 7 Enhanced Thematic Mapper Plus (ETM+) enhanced by band metrics of other sensors like Quickbird imagery, existing percent tree cover layers from Landsat data, and global Moderate Resolution Imaging Spectroradiometer (MODIS) percent tree cover [Hansen et al. 2013]. On the satellite imagery, a time-series spectral metrics analysis is applied to gather the global forest extent at 2000 as well as the annual forest loss and the accumulated gain for the period 2001 till 2012. Hence, GFC comprises three independent data layers tree cover, annually forest loss and forest gain divided into 10x10 degree tiles by the geodetic coordinate system World Geodetic System 1984 (WGS84) (EPSG:4326) with a spatial resolution of 1 arc-second per pixel. Furthermore, across the provided Geo-Tiff (GTiff) layers the pixel data is coded in unsigned 8-bit integers. Hansen et al. defined trees as all vegetation taller than 5 meters for their study. For each pixel covered by trees, a canopy density ranging from 0 to 100% is computed. Forest loss is defined as a stand displacement disturbance leading from a forest state to a non-forest state (e.g. canopy density >50% to 0%). Tree cover gain is defined as the inverse of loss where the canopy density must exceed 50% to get recognized.

Accuracy

The Hansen et al. GFC dataset is without any constraint publicly available for download. On the version 1.0 dataset homepage you may download "/\*.txt" files containing the download Uniform Resource Locator (URL) of the tiles for each sub dataset. The spatial location of an image tile can be determined with a tile identifier "{NAME}\_{LAT}[NS]\_{LNG}[WE]" were latitude (LAT) and longitude refer to the top left decimal corner coordinates in WGS84 and N or S as well W or E to the orientation on the hemisphere. For this project we needed the subdatasets: Treecover2000, lossyear and gain. The download process is fully automatized with an Python script by using the Standard Library (stdlib) modules urllib, re and threading. First this script downloads the required "/\*.txt" files and creates an list where each URL is a list element. Next it iterates over the list and extracts the corner coordinates from the file identifier with an Regular Expression (REGEX). This coordinate string is converted to an numeric identifier where latitude coordinates on the northern hemisphere are between [90, 0] and on the southern hemisphere between (0, -90]. Now a dataset tile is only downloaded if it is within the study extent between [30,-20] latitude. The acquired image tiles are shown in the top panel (green squares) in Figure 2.1. Because we have to download in total 678 images tile the entire download process is parallelized by means of multithreading.



**Figure 2.1. Map of downloaded dataset tiles:** This map shows the acquired image tiles for this study. From top to bottom in green Global Forest Change (GFC) dataset tiles (Treecover2000, lossyear and gain), the land cover dataset GlobeLand30 (GL30) image tiles in red, and in blue the Aboveground Biomass (AGB) dataset tiles. The orange filled shapes highlight countries within the tropical zone.

### 2.1.1.2 GlobeLand30

GlobeLand30 (GL30) is the first 30 meter per pixel spatial resolution global land cover dataset that provides a comprehensive view on the distribution of 10 different land cover classes (Table 2.1) over the earth [Chen et al. 2017]. Currently this dataset is for two different time frames available 2000 and 2010 [Chen et al. 2015]. The dataset is coded in unsigned 8 bit integers and as coordinates system it uses WGS84 in Universal Transverse Mercator (UTM) projection and is shipped as GTiff in a tiled manner where each tile covers 6x5 degrees [Chen et al. 2014]. For detection of the 10 land cover classes Chen et al. used a so called Pixel-Object-Knowledge (POK) oriented approach [Chen et al. 2015]. The assigned pixel values and the corresponding land cover class is listed in Table 2.1. They grouped mapping process in different stages where each land cover type is detected separately and deleted from the source satellite image the order is: water bodies, wetland, snow and ice, cultivated land and forest, shrub land, grass land and bare land synchronous. For the detection of a land cover type they used at pixel level one of the following classifiers: Decision Tree (DT), Support Vector Machine (SVM) or Maximum Likelihood Classifier (MLC). After pixel detection they grouped the classified pixel to an object and validated this object by expert knowledge. For their approach they used satellite imagery from Landsat and HJ1 and auxiliary data like MODIS NDVI.

Chen et al. estimates an overall accuracy of 80.33% for the product from 2010 and 78.6% for the product from 2000 (2000 only validated at Shaanxi province in China) [Chen et al. 2015]. Various scientists besides Chen et al. validated the mapping accuracy of GL30 at different regions and scales. Arsanjani et al. estimates an accuracy of 77.9% for Iran and an accuracy >80% for Germany [Arsanjani et al. 2016a,b]. Yang, Cao and Jacobson estimate an accuracy of 82.4%, 80.1% and 83.1% for China, Nepal and East Africa, respectively [reference](#). Unfortunately, there are no estimates for countries regions falling completely in the tropical zone.

Chen funded the GL30 land cover mapping to the UN but it is not barrier less public available. Restriction is registration on the dataset homepage but the author was not able to register at the platform. Fortunately the supervisor of this work had already an account for this page otherwise I would be fucked. To download tiles of this dataset a order application must be filled with the required image tiles identifiers and the the year must be selected. An tile identifier "[NS] {LAT}\_ {LNG}" consist of the hemisphere orientation followed by the top left corner latitude and logitude decimal coordinate (WGS84). The homepage provides an interface for selecting the required image tiles but it is broken and buggy, especially the selection of multiple tiles did not work. Fortunately they provide a vector file which contains the dataset tile polygons with assigned identifies. This file was used to select all required tiles within the tropical zone between approxiamtely 23°N and 23°S (WGS84), the selection

**Table 2.1. Classification schema of the GlobeLand30 product:** The code column is the assigned pixel value, type the corresponding land cover type and definition explains in broad terms which types of surfaces fall into the land cover type [Chen et al. 2017].

Code	Type	Definition
10	Cultivated land	used for agriculture, horticulture and gardens, including paddy fields, irrigated and dry farmland, vegetable and fruit gardens, etc.
20	Forest	covered by trees, vegetation covers over 30%, including deciduous and coniferous forest, and sparse woodland with cover 10-30%, etc.
30	Grassland	covered by natural grass with cover over 10%, etc.
40	Shrubland	covered by shrubs with cover over 30%, including deciduous and evergreen shrubs, and desert steppe with cover over 10%, etc.
50	Wetland	covered by wetland plants and water bodies, including inland marsh, lake marsh, river floodplain wetland, forest/shrub wetland, peat bogs, mangrove and salt marsh, etc.
60	Water bodies	in land area, including river, lake, reservoir, fish pond, etc.
70	Tundra	covered by lichen, moss, hardy perennial herb and shrubs in the polar regions, including shrub-, herbaceous-, wet- and barren-tundra, etc.
80	Artificial surfaces	modified by anthropogenic influence, including all kinds of habitation, industrial and mining area, transportation facilities, and interior urban green zones and water bodies, etc.
90	Bareland	with vegetation cover lower 10%, including desert, sandy fields, Gobi, bare rocks, saline and alkaline land, etc.
100	Snow and ice	covered by permanent snow, glacier and icecap

is shown in red in the middle panel at Figure 2.1. The corresponding identifiers were converted to a single line string and copied to application form. After submitting the form your order will be checked and approved. After one week we received 2 weeks limited access to an password protected ftp server where we downloaded the data. Due to restrictions this process of selecting data and download it could not be automatized with one pipeline only the selection and string conversion was automatized with a throw away script.[missing total number of tiles](#)

### 2.1.1.3 Intact Forest Landscapes

Intact Forest Landscapes (IFL) 2000 is a dataset comprising a mosaic of forest and naturally treeless ecosystems without signs of human activity and large enough to maintain all native biological diversity for the time period 2000 [Potapov et al. 2017]. Due to the fact that IFL comprises different intact natural landscape patterns like primary forests, non-forest ecosystems, temporary treeless areas after a natural disturbance, and water bodies the term is

not congruent to the term primary forest defined by the Food and Agriculture Organization of the United Nations (FAO) [FAO 2012]. But as mentioned IFLs includes large patches of primary forests with a minimum extent of 500 Km<sup>2</sup> therefore primary forests can be extracted from the layer. Still there are smaller fragments of primary forest outside of the IFLs. In regards of the extent a IFL has a minimum size of 500 Km<sup>2</sup>, a minimum width of 10 Km, and a minimum corridor/appendage width of 2 Km. Further a IFL should not contain any of the following: ecosystem alternation, fragmentation by infrastructure and disturbance, and areas altered or managed trough agriculture, logging, and mining. For mapping and detecting IFLs Potapov et al. used Landsat imagery and several auxiliary data sources like GFC, and national transportation maps. The dataset is may be downloaded as a Shapefile (SHP) with the coordinate reference system WGS84. Each polygon in the SHP represents an IFL patch at an certain location on our planet.

Data acquisition is pretty straight forward the IFL dataset is barrier-less public available for download. As mentioned it is an SHP so you must only download a single compressed archive. This download is realized with an Python script by using the stdlib module urllib.

#### 2.1.1.4 Aboveground Woody Biomass

The Aboveground live woddy Biomass density (AGB) raster dataset is prepared by Global Forest Watch (GFW) with an adapted approach of Baccini et al. [Baccini et al. 2012, 2015, 2017]. For the year 2000, this dataset comprises the aboveground biomass density in Mg C ha<sup>-1</sup> (mega gram carbon per hectare) per pixel, and a confidence estimate per pixel at a spatial resolution of approximately 1 arc-second. The dataset covering the global tropical zone as an mosaic of GTiff raster images where each tile has the Coordinate Reference System (CRS) WGS84 and is coded in float. For deriving biomass density GFW used canopy metrics from Geoscience Laser Altimeter System (GLAS) Light Detection and Ranging (LIDAR) footprint and several allometric equations. The resulting GLAS AGB estimates are used to train a Random Forest (RF) model based on Landsat 7 ETM+.

The AGB raster image tiles are public available for download on the homepage of GFW. As mentioned before this dataset covers only the tropical zone therefore no filtering is needed before download. To receive the URL of each image tile the GFW homepage provides an Geographic JavaScript Object Notation (GeoJSON) Application Programming Interface (API). The response of this API if requested is an GeoJSON feature collection containing the URL of the actual biomass layer, the URL of the uncertainty layer, and the rectangular bounds of each image tile. The data acquisition is automatized by a Python script by using the stdlib modules urllib, threading, and the open source library Geopandas. At first the GeoJSON is downloaded via an API call and eventually stored on disk. Next we

iterate over each feature of the GeoJSON feature collection and extract the URLs (biomass and confidence) of each tile. These URLs are piped to our multi-threaded downloader and eventually store on disk. Despite mentioning the confidence layers in the dataset description, the server answers with a 404 if the confidence URL is requested. Therefore the confidence layers are not available. In total we downloaded 105 different image tiles, their extent is shown in blue at the bottom panel of Figure 2.1.

#### 2.1.1.5 Global Soil Organic Carbon

The Global Soil Organic Carbon map (GSOCmap) is a joint project between Global Soil Partnership (GSP) and Intergovernmental Technical Panel on Soils (ITPS) to produce a global Soil Organic Carbon (SOC) content map by a country driven approach. This year 2018, the first iteration of this map in version 1.0 was released, and later followed by 1.1 (new country submission by Rwanda) and 1.2 (new country submissions by Chile and Colombia). The mapping project is intended as a long lived dataset which will improve over time and by new country submissions. Till now 67 (approximately 63 % of the global land mass) different countries submitted their SOC estimates. To empower national SOC mappings the International Soil Reference and Information Center (ISRIC) provides several covariate datasets like national Digital Elevation Map (DEM) maps, annual spectral remote sensing data or national soil type grids. Also the contributors can join a mapping training and use the GSOCmap cookbook as guidance for their mapping efforts. As exchange each country shares its own national GSOCmap meeting several criteria: reporting of the Meta-data of national SOC sampling (timeline, depth, bulk density and so on), uncertainty assessment, and the applied methods for estimating and interpolation of the SOC content. As interpolation method the guide organizations suggest the following approaches and more: simple geo-matching, class-matching, Multiple Linear Regression (MLR), RF or SVM. The national maps are aggregated to the final GSOCmap with a target resolution of 30 arc-seconds in the CRS WGS84. The dataset is one GTiff coded in float covering the entire globe where each pixel value is the SOC content in  $\text{Mg C ha}^{-1}$  at soil depth of 0-30cm [FAO and ITPS 2018].

issue border differences between countries caused by different methodologies and gap filling  
(if country doesn't use provided mask there is no overlap)

Is publicly available as geotiff, download at fao hp, download one huge gtiff, acquisition performed by python.

### 2.1.1.6 Auxiliary

As auxiliary source for country boundaries we used the Global Administrative Areas Map (GADM) [Hijmans et al. 2018].

## 2.1.2 Empirical data

### 2.1.2.1 Soil Organic Carbon

Don et al. performed the first study on tropical SOC change for soil depth between 0 and 30 cm. For the study a global meta-analysis is applied by using 358 (153 published and peer-reviewed) different studies to estimate SOC change for 12 major Land-use Change (LUC) types. The base date is derived from 39 different tropical countries covering all continents. Unfortunately Africa and East-Asia are under-sampled whereas South-America has the best data coverage. The meta-analysis is restricted to mineral soils therefore all wet soil types are excluded from analysis. Don et al. considered 5 different Land Use (LU) types for his study: primary forest, secondary forest, grassland, cropland, and perennial crops. Primary forest is defined as natural vegetation without human impacts which includes natural grassland and shrubland. Secondary forest are managed forests and regrown forest after partial destruction of the old stand. Grassland comprises pastures but excludes natural grasslands. Cropland is comprises annual crops like maize or beans and perennial crops could be coffee or sugar cane [Don et al. 2010]. For our study we used only the SOC change estimates for these LUC types which corresponds to the GL30 and IFL classification schema shown in Table 2.2.

**Table 2.2. Relative soil organic carbon change for certain land-use change types:** The Land-use change columns from and to defines the LUC type with corresponding relative Soil Organic Carbon (SOC) change and the Standard Error of the Mean (SEM) [Don et al. 2010].

LUC type		Relative SOC change	
From	To	[%]	SEM
Primary forest	Grassland	-12.1	±2.3
Primary forest	Cropland	-25.2	±3.3
Primary forest	Secondary forest	-8.6	±2.0
Secondary forest	Grassland	-6.4	±2.5
Secondary forest	Cropland	-21.3	±4.1

### 2.1.2.2 Ecosystem Service Values

## 2.2 Methods

### 2.2.1 Pre-processing

As mentioned in the data section, we have several datasets with different image properties like extent, coordinate reference system, coverage of the globe as single tile or multi-tile and so on. To perform further in detail analysis we need to create an similar image stack where each tile is aligned with the other images. In further descriptions we will refer to this stack as aligned image stack.

The first exercise for multi tile datasets is to find overlapping image tiles. At first we create a polygon mask as an shp-file of each dataset. To create this polygon mask we extract from each tile the bounding box and in case it is not in wgs84 we reproject it the bounds to this crs. During this process we recognized that the tiles of both gl30 (2000, 2010) datasets are corrupted located in utm zone 1 and 60. The bounds coordinates of the tiles in both utm zones exceed the maximum coordinates of the zone. As solution to this we excluded the gl30 tiles with the following identifiers: "n01\_00 s01\_00 s01\_10 s01\_15 s01\_20 s60\_00 s60\_05 s60\_10 s60\_15 n53\_00" Description: first we have our different datasets some are single tile data, some are multi tile data, some are vector data, we create a tile index mask for each multi tile dataset (GL30, GFC, AGB) in, extract the bounds of the tile, create a polygon from bounds, reproject polygon to wgs84, store polygon in a shapefile structure, assign to polygon as attribute the tile name file name as identifier, during the masking process i recognized that the gl30 dataset has coordinate system issues, the tiles at the edges of the earth , theses masks are than intersected to get overlapping tile stack, next step is to group the intersecting tiles by the key





Figure 2.2. Flowchart of tile alignment process:



**Figure 2.3. Map of aligned data tiles and sampling locations:** The map shows the location of the aligned multi-image stack tiles as black-framed square sized polygons, the sampling locations for accuracy assessment in red, and countries within the tropical zone in orange.

## 2.2.2 Deforestation

### 2.2.2.1 Forest definition

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

#### 2.2.2.2 Land use change driver

  Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

  Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

#### 2.2.2.3 Accuracy assessment

  Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

  Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

## 2.2.3 Emissions

### 2.2.3.1 Above ground biomass

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 2.2.3.2 Soil organic carbon change

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum

sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

## 2.2.4 Ecosystem service values

### 2.2.4.1 Ecosystem service value loss

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placemat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### 2.2.4.2 Ecosystem service value gain

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placemat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## 2.2.5 Binning analysis

The previous sections were focused on the generation of large scale spatial data. Now, a feasible method must be developed for analyzing, aggregating, interpreting and visualizing the output data. To develop a good approach we must formalize the problem domain. At first we are confronted with large N (many samples) which results in many variables (dimensionality) and complexity of relationships among this variables [Carr 1990]. From a visual/analytical perspective georeferenced raster maps can be interpreted as a multivariate scatter plot of large datasets where longitude and latitude represent the x and y coordinate of an data point and the pixel values (in this case nominal scaled) representing the third dimension as an group coloring. Therefore we have a large multidimensional dataset combined with a scatter plot visualization which leads commonly to over plotting issues and hidden point densities [Carr et al. 1987]. Due to the spatial nature of your data we are also confronted with not equal distributed data some regions show high data densities and other regions have sparse to no data. Also a severe problem domain is the frame size of our representation. Goal is to present data on a continental level which intensifies visual problems. Each pixel has a resolution of approximately 30x30m, the continental representation of americas spanning approximately 1200000x120000km<sup>2</sup>. Therefore small scale isolated changes are hidden and only large scale changes are visual detectable. Which results in hidden details and not perceivable patterns of change.

Goal should be to develop a process who solve this issues and generates satisfying output for our multivariate data. In case of raster data a re-sampling to coarser resolution could solve over plotting and resolution issues as well normalize the unequal distributed data. But the nature of re-sampling (for nominal data a nearest neighbor or majority wins [Reference](#)) would negate important spatial patterns as well frequency distributions. Another well known approach is to use binning of the spatial explicit data with a certain kind of regular polygon that is tessellating the plane [Carr et al. 1992]. Polygon tessellations provide numerous opportunities for presenting multivariate statistical summaries. The scaling of the polygon could be used to represent pixel densities within the polygon area, a polygon filling color gradient is applicable to show nominal or ordinal scaled data. Also it is imaginable to use the polygon interior for a pie chart. To use regular tessellation it is important to mention there are only three types of regular polygons tessellate the plane: squares, equilateral triangles and hexagons [Carr et al. 1992]. Square tessellation is the most common approach used for binning in spatial visualization. A raster image is a square tessellation. In a square mosaic each polygon shares 4 edge neighbors and 4 vertex neighbors [more explanation error distance disadvantages etc Hexagons properties, advantages disadvantages of both tessellations](#). Final goal is to show your analysis results of spatial explicit raster data in hexagonal binned form. For bivariate maps we choose a visual representation with scaled hexagons and colorization.

For multivariate details we choose a pie chart alike visualization. We split the hexagons horizontal in regards of the presented ratio. The ratios should be ordered descending so that the greatest ratio is south oriented. It is following a general description how we created the hexagon grids and how we tackled the polygon split problem.

To be flexible at hexagon construction we accept 4 different parameters as construction arguments:  $D$  long diagonal (Diameter of the circumscribing circle),  $d$  short diagonal (diameter of the inscribed circle),  $A$  area the hexagon should span and or  $e$  the edge length. One selected parameter of these is used to compute  $R$  the radius of the circumscribing circle with respect to input parameter as shown in equation 2.1.  $R$  is used to calculate the midpoint  $\langle c_x, c_y \rangle$  of the hexagon located in the first quadrant of the cartesian coordinate system Equation 2.2 and 2.3. Equation 2.4 shows the computation of the hexagon anti-clockwise vertex matrix. Whereas the two leftmost vertices (first and last row of the matrix  $\mathbf{H}$ ) are located at koordinatenursprung, will sagen auf deutsch korridanten at  $x=0$  und  $y$ =value of matrix. In summary equation 2.1 to 2.4 show the creation of an hexagon at the leftmost corner of first quadrant (Figure 2.4). The orientation is important for the subsequent mosaic creation.

$$R = \frac{\sqrt{2A}}{\sqrt[4]{27}} = \frac{D}{2} = \frac{d}{\sqrt{3}} = e \quad (2.1)$$

$$c_x = \frac{R\sqrt{3}}{2} \quad (2.2)$$

$$c_y = R \quad (2.3)$$


$$\mathbf{H} = \begin{bmatrix} 0 & c_x & 2c_x & 2c_x & c_x & 0 \\ R\sin\left(\frac{7\pi}{6}\right) + c_y & 0 & R\sin\left(\frac{11\pi}{6}\right) + c_y & R\sin\left(\frac{\pi}{6}\right) + c_y & 2R & R\sin\left(\frac{5\pi}{6}\right) + c_y \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \quad (2.4)$$

A polygon tessellation needs several polygons to create a grid in case of the creation of one hexagon with the presented algorithm needs approximately **benchmark** but the creation of **several N hexagons** needs approximately **benchmark**. Therefore it is much simpler to create only one hexagon with the presented algorithm and to create the grid polygons by copying the coordinates of the source polygon and translating them to their target position with a affine transformation matrix shown in equation 2.5. To create the grid we get the rectangular bounds of the area to tessellate as a matrix  $\mathbf{B} \in R^{2 \times 2}$  (equation 2.6), where the first column of the matrix contains the lower left corner and the second column the upper right corner of the image. Each subsequent translation in regards of  $x_{off}$  is  $x_1 + d$  for even rows and bla bla for odd rows.  $Y_{off}$  is computed by bla bla see figure 2.4.

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & x_{off} \\ 0 & 1 & y_{off} \\ 0 & 0 & 1 \end{bmatrix} \circ \mathbf{H} \quad (2.5)$$

$$\mathbf{B} = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} \quad (2.6)$$

Binning of raster data is easy we just have a point in polygon problem each points/pixels



**Figure 2.4. Hexagon tessellation:** Located at the left bottom corner in red a hexagon defined by its geometric properties the 6 vertex vectors  $\{\vec{v}_0, \dots, \vec{v}_5\}$  (black crosses), with center vector  $\vec{m}$ , edge length  $e$ ,  $R$  radius of the circumscribing circle,  $r$  radius of the inscribed circle and  $d$  the length of the short diagonal. Top right black dotted box are the bounds of an area which is tessellated by a hexagon grid in red. Each grid cell is translated from the origin hexagon at its position by computing the  $x_{off}$  and  $y_{off}$  offset with the presented equations at the left-hand side of the grid.

falling in hexagon are counted and aggregated through a function. In case of drivers of deforestation we count all driver classes per hexagon and compute ratios next we compute the sha **describe for each map how you build it** As mentioned before for the visualization of the drivers of deforestation map we want to segment the hexagons with horizontal lines and each segment should represent the share of the direct deforestation driver within the tessellated area. To compute the split line for a certain hexagon we need the hexagon R computeable from the area of the hexagon equation 2.7 and the rectangular bounds of the hexagon. We compute the relative share of an deforestation driver per hexagon this relative share can be used to compute the y-axis coordinate of an split line equation 2.8. A regular hexagon can not only be presented in it vertex form as shown above. We can also use functions to define the hexagon shape. A hexagon consist of 2 picewise functions where each function consist of 3 linear functions restricted to an intervall. If we invert these functions we can use these functions to compute the x-coordinate of the split line with the previous computed y-coordinate Equations 2.9 and 2.10. As a results we receive the solution matrix L which represents the horizontal line segment splitting the hexagon at the point where we want (driver ratio share) equation 2.11. The solution matrix can be plugged in to a polygon

split function which separates the hexagon polygon in a upper and lower part to do so we iterate over the hexagon vertices and decide if they are above or under the split line and append to a lower upper polygon. These list are our results [explain better split function](#).

$$R = \frac{\sqrt{2A}}{\sqrt[4]{27}} \quad (2.7)$$

$$y = \frac{P(y_2 - y_1)}{100} + y_1 \quad (2.8)$$


$$f^{-1}(y) = \begin{cases} -\frac{y-y_1}{\tan(\frac{\pi}{6})} + \frac{x_1+x_2}{2} & \text{if } y_1 \leq y < y_1 + R \sin(\frac{5\pi}{6}) \\ x_1 & \text{if } y_1 + R \sin(\frac{5\pi}{6}) \leq y < R(\sin(\frac{5\pi}{6}) + 1) \\ \frac{y-y_2}{\tan(\frac{\pi}{6})} + \frac{x_1+x_2}{2} & \text{if } R(\sin(\frac{5\pi}{6}) + 1) \leq y \leq y_2 \end{cases} \quad (2.9)$$

$$g^{-1}(y) = \begin{cases} \frac{y-y_1}{\tan(\frac{\pi}{6})} + \frac{x_1+x_2}{2} & \text{if } y_1 \leq y < y_1 + R \sin(\frac{5\pi}{6}) \\ x_2 & \text{if } y_1 + R \sin(\frac{5\pi}{6}) \leq y < R(\sin(\frac{5\pi}{6}) + 1) \\ -\frac{y-y_2}{\tan(\frac{\pi}{6})} + \frac{x_1+x_2}{2} & \text{if } R(\sin(\frac{5\pi}{6}) + 1) \leq y \leq y_2 \end{cases} \quad (2.10)$$

$$\mathbf{L} = \begin{bmatrix} f^{-1}(y) & g^{-1}(y) \\ y & y \end{bmatrix} \quad (2.11)$$

### 3 Results

#### 3.1 Forest definition and accuracy assessment



**Figure 3.1. Similarity distribution of GFC tree cover vs. GL30 tree cover at 2000:** Americas from JC<sub>0</sub> till JC<sub>3</sub>:  $\bar{x} \approx .71, .71, .70, .69$ ;  $\tilde{x} \approx .77, .77, .76, .75$ ;  $Q_1 \approx .64, .65, .64, .63$ ;  $Q_3 \approx .84, .84, .84, .83$ ; Asia from JC<sub>0</sub> till JC<sub>3</sub>:  $\bar{x} \approx .71, .71, .70, .69$ ;  $\tilde{x} \approx .77, .77, .76, .75$ ;  $Q_1 \approx .65, .66, .64, .63$ ;  $Q_3 \approx .84, .84, .84, .83$ ; Africa from JC<sub>0</sub> till JC<sub>3</sub>:  $\bar{x} \approx .39, .39, .34, .32$ ;  $\tilde{x} \approx .40, .40, .31, .29$ ;  $Q_1 \approx .14, .10, .00, .00$ ;  $Q_3 \approx .60, .61, .63, .56$

**Table 3.1. Two-sided Wilcoxon signed-rank test:**

$H_0: \tilde{x}_1 = \tilde{x}_2$												
Cls	Americas				Asia				Africa			
	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>
JC <sub>0</sub>	1.				1.				1.			
JC <sub>1</sub>	.00	1.			.72	1.			.22	1.		
JC <sub>2</sub>	.06	.36	1.		.00	.00	1.		.03	.03	1.	
JC <sub>3</sub>	.16	.50	.60	1.	.00	.00	.00	1.	.00	.00	.00	1.

**Table 3.2. One-sided Wilcoxon signed-rank test:**

$H_0: \tilde{x}_1 \leq \tilde{x}_2$													
		Americas				Asia				Africa			
	Cl	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>	JC <sub>0</sub>	JC <sub>1</sub>	JC <sub>2</sub>	JC <sub>3</sub>
$\text{H}^{\text{S}}$	JC <sub>0</sub>	1.	.00	.03	.08	1.	.64	1.	1.	1.	.11	.98	1.
$\wedge$	JC <sub>1</sub>	1.	1.	.18	.25	.36	1.	1.	1.	.89	1.	.99	1.
$\text{H}_0: \tilde{x}_1$	JC <sub>2</sub>	.97	.82	1.	.30	.00	.00	1.	1.	.02	.01	1.	1.
$\text{H}_0: \tilde{x}_2$	JC <sub>3</sub>	.92	.75	.70	1.	.00	.00	.00	1.	.00	.00	.00	1.

**Table 3.3.** Confusion matrix for accuracy assessment

		Reference												
		Cl	10	20	25	30	40	50	60	80	90	Tot	UAc	Om
Prediction		10	732	38	62	15	16	2	3	5	0	873	.84	.16
		20	42	751	57	189	31	12	0	17	4	1103	.68	.32
		25	29	202	1155	173	22	10	5	11	4	1611	.72	.28
		30	36	187	32	1466	73	21	0	17	0	1832	.80	.20
		40	14	21	4	41	352	1	1	2	1	437	.81	.19
		50	0	5	3	10	4	50	0	1	0	73	.68	.32
		60	2	1	0	3	0	2	18	2	0	28	.64	.36
		80	3	4	0	1	1	1	0	50	0	60	.83	.17
		90	0	0	0	1	0	0	0	3	5	9	.56	.44
		Tot	858	1209	1313	1899	499	99	27	108	14	6026		
		PAC	.85	.62	.88	.77	.71	.51	.67	.46	.36			OvAc
		Com	.15	.38	.12	.23	.29	.49	.33	.54	.64			.75

## 3.2 Deforestation drivers

### 3.2.1 Global


### 3.2.2 Americas

### 3.2.3 Asia


### 3.2.4 Africa

**Table 3.4.** Absolute in km<sup>2</sup>


Type	Class	Americas	Asia	Africa
Agriculture	Cropland	rel. abs.	24.37 95908	18.37 38719
	Grassland	rel. abs.	46.19 181781	8.41 17726
Forestry/Plantations	Regrowth	rel. abs.	14.40 56671	70.27 148111
	Shrubland	rel. abs.	12.69 49941	1.11 2340
Urban/Mining	Artificial	rel. abs.	0.41 1614	0.46 970
	Bareland	rel. abs.	0.10 394	0.03 63
Natural	Wetland	rel. abs.	1.50 5903	0.97 2045
	Water	rel. abs.	0.32 1259	0.38 801
Forest loss		rel. abs.	3.87 393550	4.68 210774
Forest cover		abs.	10223187	177400
		abs.	4457940	10496591




**Figure 3.2.**




**Figure 3.3.**




**Figure 3.4.**




**Figure 3.5.**




**Figure 3.6.**




**Figure 3.7.**



**Figure 3.8.**




**Figure 3.9.**



**Figure 3.10.**

### 3.3 Deforestation



**Figure 3.12.**

**Table 3.5.** Soil organic carbon emissions

Region	SC1 [Gt CO <sub>2</sub> ]			SC2 [Gt CO <sub>2</sub> ]			SC3 [Gt CO <sub>2</sub> ]		
	min	mean	max	min	mean	max	min	mean	max
Americas	0.80	0.96	1.12	0.45	0.61	0.77	0.43	0.59	0.76
Asia	0.66	0.81	0.97	0.22	0.28	0.34	0.22	0.28	0.33
Africa	0.32	0.39	0.45	0.17	0.23	0.29	0.16	0.23	0.29

3.3.1 Global

3.3.2 Americas

3.3.3 Asia

3.3.4 Africa

## 3.4 Ecosystem service value balance




Figure 3.13.

3.4.1 Global

3.4.2 Americas

3.4.3 Asia

3.4.4 Africa

## 4 Discussion

# Acknowledgements

# Bibliography

- Arsanjani J. J., See L., and Tayyebi A. Assessing the suitability of GlobeLand30 for mapping land cover in Germany. *International Journal of Digital Earth*, 2016a. doi: 10.1080/17538947.2016.1151956.
- Arsanjani J. J., Tayyebi A., and Vaz E. GlobeLand30 as an alternative fine-scale global land cover map: Challenges, possibilities, and implications for developing countries. *Habitat International* xxx, pages 1–7, 2016b. doi: 10.1016/j.habitint.2016.02.003.
- Baccini A., Goetz S., Walker W., Laporte N., Sun M., Sulla-Menashe D., Hackler J., Beck P., Friedl M., Samanta S., and Houghton R. Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature Climate Change*, 2, March 2012. doi: 10.1038/nclimate1354.
- Baccini A., Walker W., Carvahlo L., Farina M., Sulla-Menashe D., and Houghton R. Tropical forests are a net carbon source based on new measurements of gain and loss. Online accessed through Global Forest Watch, 2015. URL <https://www.globalforestwatch.org>.
- Baccini A., Walker W., Carvalho L., Farina M., and D. Sulla-Menashe R. A. H. Tropical forests are a net carbon source based on aboveground measuremeasure of gain and loss. *Science*, September 2017. doi: 10.1126/science.aam5962.
- Carr D. B. Looking at large data sets using binned data plots. resreport, U.S. Department of Energy, 1990.
- Carr D. B., Littlefield R. J., Nicholson W. L., and Littlefield J. S. Scatterplot matrix techniques for large N. *Journal of the American Statistical Association*, 82(398), 1987.
- Carr D. B., Olsen A. R., and White D. Hexagon mosaic maps for display of univariate and bivariate geographical data. *Cartography and Geographic Information Systems*, 19(4), 1992.
- Chen J., Chen J., Liao A., Cao X., Chen L., Chen X., He C., Han G., Peng S., Lu M., Zhang W., Tong X., and Mills J. *30-meter Global Land Cover Dataset - Product Description*. National Geomatics Center of China, May 2014.
- Chen J., Chen J., Liao A., Cao X., Chen L., Chen X., He C., Han G., Peng S., Lu M., Zhang W., Tong X., and Mills J. Global land cover mapping at 30 m resolution: A POK-based operational approach. *ISPRS Journal of Photogrammetry and Remote Sensing*, 103:7–27, 2015. doi: 10.1016/j.isprsjprs.2014.09.002. URL <http://www.globallandcover.com>.
- Chen J., Cao X., Peng S., and Ren H. Analysis and Applications of GlobeLand30: A Review. *International Journal of Geo-Information*, 6(230), July 2017. doi: 10.3390/ijgi6080230.
- Don A., Schumacher J., and Freibauer A. Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis. *Global Change Biology*, 17(4):1658–1670, November 2010. doi: 10.1111/j.1365-2486.2010.02336.x.

FAO. FRA 2015 terms and definitions. resreport, Food and Agriculture Organization of the United Nations, 2012.

FAO and ITPS. Global Soil Organic Carbon Map. resreport, FAO and ITPS, 2018.

Hansen M. C., Potapov P. V., Moore R., Hancher M., Turubanova S. A., Tyukavina A., Thau D., Stehman S. V., Goetz S. J., Loveland T. R., Kommareddy A., Egorov A., Chini L., Justice C. O., and Townshend J. R. G. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science*, 342(6160):850–853, November 2013. doi: 10.1126/science.1244693. URL [https://earthenginepartners.appspot.com/science-2013-global-forest/download\\_v1.0.html](https://earthenginepartners.appspot.com/science-2013-global-forest/download_v1.0.html).

Hijmans R., Garcia N., Kapoor J., Rala A., Maunahan A., and Wieczorek J. GADM database of Global Administrative Areas. Online, May 2018. URL <https://www.gadm.org>. Version 3.6.

Li Y., Sulla-Menashe D., Motesharrei S., Kalnay X.-P. S., Ying Q., Li S., and Ma Z. Inconsistent estimates of forest cover change in China between 2000 and 2013 from multiple datasets: differences in parameters, spatial resolution, and definitions. *Scientific Reports*, 7(8748), August 2017. doi: 10.1038/s41598-017-07732-5.

Potapov P., Hansen M. C., Laestadius L., Turubanova S., Yaroshenko A., Thies C., Smith W., Zhuravleva I., Komarova A., Minnemeyer S., and Esipova E. The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Science Advances*, 3, 2017. doi: 10.1126/sciadv.1600821. URL <http://www.intactforests.org/>.

# List of Figures

1.1	Tropical zone . . . . .	11
1.2	Deforestation examples . . . . .	13
2.1	Map of downloaded dataset tiles . . . . .	19
2.2	Flowchart of tile alignment process . . . . .	25
2.3	Map of aligned data tiles and sampling locations . . . . .	26
2.4	Hexagon tessellation . . . . .	32
3.1	Similarity distribution of GFC tree cover vs. GL30 tree cover at 200 . . . . .	34
3.2	Ecosystem service values . . . . .	36
3.3	Ecosystem service values . . . . .	37
3.4	Ecosystem service values . . . . .	37
3.5	Ecosystem service values . . . . .	38
3.6	Ecosystem service values . . . . .	38
3.7	Ecosystem service values . . . . .	39
3.8	Ecosystem service values . . . . .	39
3.9	Ecosystem service values . . . . .	40
3.10	Ecosystem service values . . . . .	40
3.11	Ecosystem service values . . . . .	41
3.12	Ecosystem service values . . . . .	41
3.13	Ecosystem service values . . . . .	42

# List of Tables

2.1	Classification schema of the GlobeLand30 product . . . . .	21
2.2	Relative soil organic carbon change for certain land-use change types . . . . .	24
3.1	Two-sided Wilcoxon signed-rank test . . . . .	34
3.2	One-sided Wilcoxon signed-rank test . . . . .	35
3.3	Accuracy assessment . . . . .	35
3.4	Deforestation driver . . . . .	36
3.5	Soil organic carbon emissions . . . . .	41

# List of Abbreviations

<b>AGB</b>	Aboveground live woddy Biomass density
<b>API</b>	Application Programming Interface
<b>CRS</b>	Coordinate Reference System
<b>DEM</b>	Digital Elevation Map
<b>DT</b>	Decision Tree
<b>ETM+</b>	Enhanced Thematic Mapper Plus
<b>FAO</b>	Food and Agriculture Organization of the United Nations
<b>GADM</b>	Global Administrative Areas Map
<b>GFC</b>	Global Forest Change
<b>GFW</b>	Global Forest Watch
<b>GIS</b>	Geographic Information System
<b>GL30</b>	GlobeLand30
<b>GLAS</b>	Geoscience Laser Altimeter System
<b>GSOCmap</b>	Global Soil Organic Carbon map
<b>GSP</b>	Global Soil Partnership
<b>GTiff</b>	Geo-Tiff
<b>GeoJSON</b>	Geographic JavaScript Object Notation
<b>IFL</b>	Intact Forest Landscapes
<b>IPCC</b>	Intergovernmental Panel on Climate Change
<b>ISRIC</b>	International Soil Reference and Information Center
<b>ITPS</b>	Intergovernmental Technical Panel on Soils
<b>LIDAR</b>	Light Detection and Ranging
<b>LU</b>	Land Use
<b>LUC</b>	Land-use Change
<b>LULC</b>	Land Use/Land Cover
<b>MLC</b>	Maximum Likelihood Classifier
<b>MODIS</b>	Moderate Resolution Imaging Spectroradiometer
<b>POK</b>	Pixel-Object-Knowledge
<b>REGEX</b>	Regular Expression
<b>RF</b>	Random Forest
<b>SHP</b>	Shapefile
<b>SOC</b>	Soil Organic Carbon
<b>SVM</b>	Support Vector Machine
<b>URL</b>	Uniform Resource Locator
<b>UTM</b>	Universal Transverse Mercator
<b>WGS84</b>	World Geodetic System 1984
<b>stdlib</b>	Standard Library

**MLR**

Multiple Linear Regression

# Appendix

Wyrażam zgodę na udostępnienie mojej pracy w czytelniach Biblioteki SGGW w tym w Archiwum Prac Dyplomowych SGGW.

I agree to share my work in the reading rooms of the SGGW Library, including the SGGW Theses Archive.

Ich erteile meine Zustimmung zur Veröffentlichung meiner Arbeit in der Bibliothek der SGGW (Warschauer Naturwissenschaftliche Universität), einschließlich des Archivs der Diplomarbeiten.

.....  
*(czytelny podpis autora pracy)  
(legible signature of the author)  
(lesbare Unterschrift des Autors der Arbeit)*