

Lab 2 – Homework Solutions

September 23, 2022

A Homework

Homework 2.1 (Bisection method). Consider the following equation

$$\cot(x) = \frac{x^2 - 1}{2x}.$$

- Define three intervals containing the three smallest positive roots.
- Find the biggest root among the ones at item a), using bisection method with tolerance equal to 10^{-6} .
- Plot the evolution of the error as a function of the number of the iterations.

Solution Homework 2.1.

```
function rootfinding_function_plot(f, a, b, new_figure)

    % Check if a new graphic window is requested
    if ((nargin < 4) || new_figure)
        figure
    end

    hold on, box on
    x_plot = linspace(a, b, 1000);

    % Plot the function
    plot(x_plot, f(x_plot), 'LineWidth', 2)

    % Plot a zero-line
    plot(x_plot, 0*x_plot, 'k-', 'LineWidth', 1)

    xlabel('x','FontSize', 16)
    ylabel('f(x)','FontSize', 16)
    set(gca,'FontSize', 16)
    set(gca,'LineWidth', 1.5)

end

% HOMEWORK 2.1

f = @(x) cot(x);
g = @(x) (x.^2 - 1)./(2*x);

epsilon = 0.01;
rootfinding_function_plot(f, 0+epsilon, pi-epsilon, true);
rootfinding_function_plot(f, pi+epsilon, 2*pi-epsilon, false);
rootfinding_function_plot(f, 2*pi+epsilon, 3*pi-epsilon, false);
rootfinding_function_plot(g, 0, 3*pi, false);

% Three intervals for the roots are then [1,2], [3,4] and [6,7].
```

```

tol = 1e-6;
h = @(x) cot(x) - (x.^2 - 1)./(2*x);

rootfinding_function_plot(h, 0+epsilon, pi-epsilon, true);
rootfinding_function_plot(h, pi+epsilon, 2*pi-epsilon, false);
rootfinding_function_plot(h, 2*pi+epsilon, 3*pi-epsilon, false);

I = [6.5 7] % cannot use [6, 7] since h is discontinuous there
[xi, x] = bisection(h, I(1), I(2), tol);
xi
iter = numel(x)

% Since the exact value of the roots is NOT known, we perform
% again the bisection method with a very small tolerance, and consider
% the final approximation as the exact root of the equation.

figure

% First root
[xiex, xex] = bisection(h, I(1), I(2), 1e-10*tol);
xiex
err = abs(x - xiex);

subplot(1,2,1)
semilogy(err, 'LineWidth',2)
box on, hold on
semilogy(tol*ones(iter,1), 'r--', 'LineWidth',2)
xlim([0 iter+1])
set(gca,'FontSize',16)
xlabel('Iteration','FontSize',16)
ylabel('Error','FontSize',16)

```

Homework 2.2 (Bisection method). Consider the following equation

$$e^{-(x-2)^2} = 1 - e^{x-4}$$

- Plot function $f(x) = 0$ obtained from the above equation by writing all the terms in the first member and determine the number of real solutions.
- Apply the bisection method by choosing a tolerance equal to 10^{-3} and initial interval $I_1 = [0, 5]$, $I_2 = [1, 6]$, $I_3 = [1, 5]$ and $I_4 = [-1, 5]$. Can you find all the roots?

Solution Homework 2.2.

```

% HOMEWORK 2.2

f = @(x) exp(-(x-2).^2) + exp(x-4) - 1;
rootfinding_function_plot(f, 0, 5);

% There are three solutions in the interval [1, 5].

tol = 1e-6;

I1 = [0 5];
[xi1, x_iter1] = bisection(f, I1(1), I1(2), tol);
xi1

I2 = [1 6];
[xi2, x_iter2] = bisection(f, I2(1), I2(2), tol);
xi2

I3 = [1 5];
[xi3, x_iter3] = bisection(f, I3(1), I3(2), tol);
xi3

I4 = [-1 5];

```

```
[xi4, x_iter4] = bisection(f, I4(1), I4(2), tol);
xi4
```

```
% In the first and fourth interval the bisection method converges to the root with
% minimum magnitude; similarly, in the second and third interval the bisection method
% converges to the root with maximum magnitude. The second root cannot be reached.
%
% This is caused by the iterative procedure that halves the interval: at the
% very first iteration the half-interval that contains the second root is discarded.
```

```
bisection_plot(f, I2(1), I2(2), tol);
bisection_plot(f, I3(1), I3(2), tol);
```

```
% This example shows that the choice of the initial interval is important.
```

Homework 2.3 (Newton method). a. Approximate the root x^* of $\tan(x) = 2x$ in the interval $(0, \frac{\pi}{2})$ up to a tolerance equal to 10^{-15} .

b. Approximate the zero of $\sin(x)$ in the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ using Newton method and starting from the value $x^{(0)} = x^*$. What do you expect? What do you observe?

Solution Homework 2.3.

```
function [xi, x_iter] = newton25(f, df, x0, tol, maxit, multiplicity)
%NEWTON Find a root of the equation f(x) = 0 using the Newton method, starting from the
%initial guess x0.
%
% [xi, x_iter] = NEWTON(f, df, x0, tol, maxit, multiplicity)
%
% Inputs : f = function handle to the function f(x)
%          df = function handle to the derivative of the function f(x)
%          x0 = initial guess
%          tol = requested tolerance
%          maxit = maximum number of iterations
%          multiplicity = multiplicity of the root
% Output :
%          xi = approximation of the root
%          x_iter = vector of the approximations of the root at each step

if (nargin < 6)
    multiplicity = 1;
end

x_iter(1) = x0;

for (iter = 1:maxit)
    newton_method = @(x) x - multiplicity*f(x)/df(x);
    x_iter(iter+1) = newton_method(x_iter(iter));

    if (abs (x_iter(iter+1) - x_iter(iter)) < tol)
        break;
    end

end

xi = x_iter(end);

end

% HOMEWORK 2.3

f = @(x) tan(x) - 2*x;
epsilon = 0.01;
rootfinding_function_plot(f, 0+epsilon, pi/2-epsilon, true);

% From the plot we choose the interval I = [1, 1.5] that ensures the validity of the Bolzano's
% theorem, and we can use e.g. the bisection method to find x*.
```

```

xstar = bisection(f, 1, 1.5, 1e-15);

% The Newton method for the approximation of  $\sin(x) = 0$  is
%
%  $x\{k+1\} = x\{k\} - \{\sin(x\{k\})\}/\{\cos(x\{k\})\} = x\{k\} - \tan(x\{k\})$ .
%
% Therefore, if  $x\{0\} = x^*$ 
%
%  $x\{1\} = x\{0\} - \tan(x\{0\}) = x^* - \tan(x^*) = x^* - 2x^* = -x^* = -x\{0\}$ 
%  $x\{2\} = x\{1\} - \tan(x\{1\}) = -x^* + \tan(x^*) = -x^* + 2x^* = x^* = x\{0\}$ 
%
% The method is describing the orbit  $\{x^*, -x^*\}$ , and thus cannot converge!

g = @(x) sin(x);
dg = @(x) cos(x);

[xi, x_iter] = newton25(g, dg, xstar, 1e-6, 25);
x_iter

% The first 10 iterations of the Newton method seem to describe the orbit  $\{x^*, -x^*\}$ , but
% after the first few iterations some numerical error arises and this makes the algorithm move
% from the orbit and then to converge to the solution.

```

Homework 2.4 (Newton method). Consider the following function

$$f(x) = e^{ax} - 1 = 0, \quad a \neq 0.$$

For any value of the parameter a the unique solution is clearly $\xi = 0$.

- Consider the case of $a = 200$ and apply Newton method starting from $x^{(0)} = 1$ with tolerance equal to 10^{-3} and 10^{-12} .
- Repeat item a) for $a = 10^{-3}$.
- For both items b) and c), compare the absolute value of the residual with the error. Justify the obtained results.

Solution Homework 2.4.

```

% HOMEWORK 2.4

a = 200;
f = @(x) exp(a*x) - 1;
df = @(x) a*exp(a*x);

maxit = 1000;
x0 = 1;

[xi, x] = newton25(f, df, x0, 1e-3, maxit);
err_1_3 = abs(xi)
absres_1_3 = abs( f(xi) )

[xi, x] = newton25(f, df, x0, 1e-12, maxit);
err_1_12 = abs(xi)
absres_1_12 = abs( f(xi) )

%%
a = 1e-3;
f = @(x) exp(a*x) - 1;
df = @(x) a*exp(a*x);

maxit = 1000;
x0 = 1;

[xi, x] = newton25(f, df, x0, 1e-3, maxit);

```

```

err_2_3 = abs(xi)
absres_2_3 = abs( f(xi) )

[xi, x] = newton25(f, df, x0, 1e-12, maxit);
err_2_12 = abs(xi)
absres_2_12 = abs( f(xi) )

disp('    tol      absres      error')
disp([1e-3 absres_1_3 err_1_3;
      1e-12 absres_1_12 err_1_12])

disp('    tol      absres      error')
disp([1e-3 absres_2_3 err_2_3;
      1e-12 absres_2_12 err_2_12])

% We know that abs{f(x)} < abs{f'(xi)}abs{x - xi}
% Hence if the derivative of f at xi is big the residual can be big
% while the solution is already well approximated.
% Conversely if the derivative is small, the residual decreases fast and
% the convergence criterion can result to be too weak.

```

Homework 2.5 (Newton method). The equation $x^3 - x^2 - x + 1 = 0$ has a double root at $\xi = 1$. Given $x^{(0)} = 2$, does Newton method converge to ξ ? How can you improve the method for approximating ξ ?

Solution Homework 2.5.

```

clear all
close all
clc
% HOMEWORK 2.5

f = @(x) x.^3 - x.^2 - x + 1;
df = @(x) 3*x.^2 - 2*x - 1;
rootfinding_function_plot(f, -2, 2, true);

tol = 1e-6
nmax = 100;
[xin, xn] = newton25(f, df, 2, tol, nmax);
xin
itern = numel(xn)

[xinmod, xnmod] = newton25(f, df, 2, tol, nmax, 2);
xinmod
iternmod = numel(xnmod)

err_newton = abs(xn - 1);
err_newton2 = abs(xnmod - 1);

figure
semilogy(err_newton, 'bs--','LineWidth',2)
hold on, box on
semilogy(err_newton2, 'ro-','LineWidth',2)
axis([0 itern+1 1e-12 10])
set(gca,'FontSize',16)
set(gca,'LineWidth',1.5)
xlabel('iterations','FontSize',16)
ylabel('error','FontSize',16)
h = legend('Newton', 'Modified Newton');
set(h,'FontSize',16)

```