

3.24pt

Numerical Analysis

Lab 8

Luca Liverotti

A.Y. 2022-2023

December 2, 2022

Polynomial interpolation

Given a function $f(x)$, an **interpolating function** is generally defined as

$$g(x) = \sum_{k=0}^n \alpha_k \phi_k(x) \quad \text{s.t.} \quad g(x_i) = f(x_i) \quad i = 0, 1, \dots, n$$

where $\{x_i\}_{i=0}^n$ and $\{\phi_i(x)\}_{i=0}^n$ are respectively sets of $n+1$ **nodes** and **basis functions**, and the coefficients α_k are called **weights**.

Definition 8.1 (Lagrange interpolating polynomial)

Characteristic polynomials are defined as

$$\varphi_k(x) = \frac{\prod_{i \neq k} (x - x_i)}{\prod_{i \neq k} (x_k - x_i)}$$

and they are s.t. $\varphi_k(x_i) = \delta_{ik}$. Using them as basis functions, it holds $\alpha_i = f(x_i)$ so that we obtain the Lagrange

$$\Pi_n f(x) = \sum_{k=0}^n f(x_k) \varphi_k(x).$$

Polynomial interpolation

Theorem 8.1 (Interpolation error)

Let $\Pi_n f(x)$ be the interpolating polynomial of order n at the nodes $x_i \in [a, b]$. If $f(x) \in \mathcal{C}^{n+1}([a, b])$ then

$$E_n f(x) = f(x) - \Pi_n f(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} \prod_{i=0}^n (x - x_i) \quad \xi \in [a, b], \forall x \in [a, b].$$

If the nodes are equally spaced with step h , it holds

$$\|E_n f(x)\|_{L^\infty([a, b])} \leq \frac{\|f^{(n+1)}(x)\|_{L^\infty([a, b])}}{4(n+1)} h^{n+1}$$

This does not ensure a priori the uniform convergence of $\Pi_n f(x)$ to $f(x)$ for $n \rightarrow \infty$.

Polynomial interpolation

Definition 8.2 (Piecewise linear interpolation)

Given of nodes: $x_0 < x_1 < \dots < x_n$, we denote by I_i the interval $[x_i, x_{i+1}]$ and by H the maximum length of these intervals.

We denote by $\Pi_1^H f$ the piecewise linear interpolating polynomial of f given by:

$$\Pi_1^H f(x) = f(x_i) + \frac{f(x_{i+1}) - f(x_i)}{x_{i+1} - x_i}(x - x_i) \quad \forall x \in I_i.$$

Theorem 8.2 (Interpolation error)

If $f \in C^2(I)$, where $I = [x_0, x_n]$, then

$$\|f(x) - \Pi_1^H f(x)\|_{L^\infty(I)} \leq \frac{H^2}{8} \|f''(x)\|_{L^\infty(I)}$$

This bound allows us to ensure the uniform convergence of the interpolating polynomial to f .

Polynomial interpolation

MATLAB commands

Lagrange interpolation:

```
>> help polyfit
```

```
>> help polyval
```

Piecewise polynomial interpolation:

```
>> help interp1
```

Cubic spline:

```
>> help spline
```

Polynomial interpolation

Notice that the commands `polyfit` and `polyval` can be used to compute both the interpolation and a least-square approximation

`polyfit(x_nodes, y_nodes, N)`

$N + 1 = \#x_nodes \implies$ Lagrange interpolation;
 $N + 1 \neq \#x_nodes \implies$ least square interpolation.

Polynomial interpolation

Exercise 8.1

Approximate the function $f(x) = x \sin(x)$ in the interval $[-2, 6]$.

- a Consider a set of equally spaced nodes: find the interpolating polynomial using Lagrange basis polynomials (`polyfit`, `polyval` commands) of degree 4, 6, 8. Plot the function $f(x)$ and the associate interpolating polynomials.
- b Estimate the interpolation error when resorting to a quadratic interpolation ($n=2$), using the theoretical results. Then evaluate the interpolating polynomial in 1000 equally spaced points and plot the interpolation error. Motivate the result.
- c Compute with Matlab the piecewise linear interpolant (`interp1` command) and the cubic spline (`spline` command) associated with $n + 1 = 6, 11, 21$ equally spaced nodes.

Polynomial interpolation

Exercise 8.2 (Runge's counterexample)

Consider the function $f(x) = \frac{1}{1+x^2} \quad -5 \leq x \leq 5$.

- a** Verify with Matlab that the interpolating polynomials $\Pi_n f(x)$ using equally spaced nodes are such that

$$\lim_{n \rightarrow \infty} |f(x) - \Pi_n f(x)| \neq 0.$$

Check this statement graphically and by computing the infinity norm

- b** Compute the piecewise linear interpolation (`interp1` command) with $n = 1, 2, 4, 8, 16, 32$ subintervals and plot the result. Find the error with the respect to the infinity norm and verify that it converges quadratically as a function of the distance between two nodes.
- c** Interpolate the Runge function using a piecewise cubic spline (`spline` command) with $n = 1, 2, 4, 8, 16, 32$ subintervals. Plot the result.
- d** Compute $\Pi_n f(x)$ using the Chebyshev nodes:

$$x_i = \frac{a+b}{2} + \frac{b-a}{2} \hat{x}_i, \quad \hat{x}_i = -\cos(\pi i/n), i = 0, \dots, n$$

Exercise 8.3

Consider the function

$$f(x) = \left| x - \frac{\pi}{12} \right| \quad -1 \leq x \leq 1.$$

- a** Verify that the interpolating polynomials $\Pi_n f(x)$ based on equally spaced nodes exhibit the Runge's phenomenon.
- b** Compute the piecewise linear interpolation (`interp1` command) based on $n = 1, 2, 4, 8, 16, 32$ intervals and compute the associated error with the respect to infinity norm. Is the quadratic convergence ensured in such a case?
- c** Interpolate the function using a piecewise cubic spline (`spline` command) with $n = 1, 2, 4, 8, 16, 32$ subintervals. Plot the result.
- d** Interpolate the function using the least square approach by setting the number of nodes $n = 10$ and varying the degree of the interpolant. Plot the results and check when the least-square approximation exactly matches the values `y_nodes` in correspondence with the `x_nodes`.

Table of contents

3.24pt

Homework 8.1

Consider the function $f(x) = e^x$ in the interval $[-1, 1]$ and the associated piecewise linear interpolating polynomial on a set of nodes $\{x_i\}$.

- a** Determine how many equally spaced nodes x_i are needed in order to ensure an interpolation error with infinity norm smaller than 10^{-3} .
- b** Construct the interpolating polynomial based on such a number of nodes, compute the associated error and check if the request on accuracy is satisfied.

Homework 8.2

Consider the function

$$f(x) = \frac{1}{(x - 0.3)^2 + 0.01} + \frac{1}{x^2 + 0.04} - 6.$$

- a** Sample $f(x)$ at 10 equally spaced nodes in the interval $[-1, 3]$ and construct the minimum degree interpolating polynomial.
- b** Numerically compute the error. Provide a method to improve the result.

Homework 8.3

Let $f(x) = x^5 + 1$ and $x_i = 0.2 \cdot i$, with $i = 0, 1, 2, 3$.

- a Let $\Pi_n f(x)$ be the interpolating polynomial of $f(x)$ at the nodes x_i . Compute $\Pi_n f(x)$ and provide an upper bound for $|E_n(x)| = |\Pi_n f(x) - f(x)|$.
- b Let $\Pi_1^H f(x)$ be the piecewise linear interpolation of $f(x)$ at the nodes x_i . Compute an upper bound for $|E_1^H(x)| = |\Pi_1^H f(x) - f(x)|$.

Homework 8.4

Consider the following function

$$f(x) = \begin{cases} 0 & x \in [0, 1) \\ 1 & x \in [1, 2]. \end{cases}$$

First interpolate the function using the Lagrange polynomial with an increasing number of nodes. Then repeat the procedure using the piecewise linear and a least-squares approximation. What do you observe? Justify the answer.

Temporary page!

\LaTeX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it. If you rerun the document (without altering it) this surplus page will go away, because \LaTeX now knows how many pages to expect for this document.