

A
constant
Events per division
1
5
3
10
87.4%
65.4%
48.7%
19.0%
3
2
1
Cumulative occurrence (proportion of runs)
Age (years)
B
increasing
Events per division
5
14
10
20
83.7%
74.7%
63.5%
44.0%
C
multistage
Events per division
2
4
3
6
99.1%
61.4%
21.0%
3.6%
D
constant + aging
Events per division
1
5
3
10
80.2%
56.4%
38.8%
15.0%
E
increasing + aging
Events per division
5
14
10
20
67.9%
57.8%
46.6%
25.6%
CTOR = 0.01
F
step 1
step 2
step 3
Observed − PH-predicted MEDA (years)
A
B
C
D
G
increasing + aging
Events per division
5
14
10
20
82.9%
74.9%
64.9%
44.5%
CTOR = 0.02
H
increasing + aging
79.9%
72.6%
65.7%
53.9%
Events per division
5
14
10
20
CTOR = 0.2
I
step 1
step 2
step 3
Observed − PH-predicted MEDA (years)
E
0.01
G
0.02
H
0.2
J
Inherited first hit (multistage)
non-carrier (3 events)
carrier (3 events)
non-carrier (4 events)
67.1%
61.4%
21.5%
Cumulative occurrence (proportion of runs)
Age (years)
K
carrier / non-carrier: ×0.74 per decade (0.65–0.84)
4 / 3 events per division: ×0.83 per decade (0.72–0.97)
Hazard ratio
Age (years)