

Homework 5

程昊一


*All of my homework is available at <https://github.com/lacampanella233/Homework.git>.

1 Problems from Artin

Problem 1 (I.2.6). Sketch the effect on the plane $\mathbb{R}^2$ of multiplication by the matrix

$$A = \begin{bmatrix} 2 & -1 \\ 2 & 3 \end{bmatrix}.$$

Solution.



Problem 2 (I.2.10). Let A be a square matrix. Prove that there exist elementary matrices $E_1, \dots, E_k$

such that

$$E_k \cdots E_1 A$$

is either the identity matrix or has its bottom row equal to zero.

Proof. If A is invertible, then one can transform A to I by elementary row transformation, thus there exists elementary matrices $E_1, \dots, E_k$ such that $E_k \cdots E_1 A = I$.

If A is not invertible, then by elementary row transformation, we can transform A to a row-echelon form, which has its bottom row equal to zero. Thus there exists elementary matrices $E_1, \dots, E_k$ such that $E_k \cdots E_1 A$ has its bottom row equal to zero. $\square$

Problem 3 (I.4.3). Prove that a matrix in which each row contains exactly one entry equal to 1, each column also contains exactly one entry equal to 1, and all other entries are 0, is a permutation matrix.

Proof. Denote the matrix by A . By the definition of A , we can write

$$A = \sum_{i=1}^n E_{i, \sigma(i)}, \quad \sigma(i) \in \{1, 2, \dots, n\}.$$

Moreover, since each column of A contains exactly one entry equal to 1, we have $\sigma(i) \neq \sigma(j)$ for $i \neq j$. Thus σ is a permutation of $\{1, 2, \dots, n\}$, i.e. $\sigma \in S_n$. Therefore A is a permutation matrix. $\square$

Problem 4 (I.4.6). What is the permutation matrix corresponding to the permutation $i \mapsto n - i$?

Solution.

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 1 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix}.$$

Problem 5 (I.5.3). Let $A = (a_{ij})$ be an $n \times n$ matrix with integer entries. Prove that A^{-1} has integer entries if and only if

$$\det(A) = \pm 1.$$

Proof. If A^{-1} has integer entries, then $\det A^{-1} \in \mathbb{Z}$. Since $\det(A^{-1}) = (\det A)^{-1}$, we have $(\det A)^{-1} \in \mathbb{Z}$, which implies $\det A = \pm 1$.

Conversely, if $\det A = \pm 1$, then $A^{-1} = (\text{adj } A)/(\det A)$ has integer entries since $\text{adj } A$ has integer entries and $\det A = \pm 1$. $\square$

2 Problems from LADR

Problem 6 (4-10). For $p \in \mathcal{P}(\mathbb{R})$, define $Tp : \mathbb{R} \rightarrow \mathbb{R}$ by

$$(Tp)(x) = \begin{cases} \frac{p(x) - p(3)}{x - 3}, & \text{if } x \neq 3, \\ p'(3), & \text{if } x = 3. \end{cases}$$

for each $x \in \mathbb{R}$. Show that $Tp \in \mathcal{P}(\mathbb{R})$ for every polynomial $p \in \mathcal{P}(\mathbb{R})$ and also show that $T : \mathcal{P}(\mathbb{R}) \rightarrow \mathcal{P}(\mathbb{R})$ is a linear map.

Proof. Since the polynomial $p(x) - p(3)$ vanishes at $x = 3$, it has a root $x = 3$, hence a factor $x - 3$. Write

$$p(x) - p(3) = q(x)(x - 3).$$

Take derivative of the equation above and set $x = 3$, we can get $p'(3) = q(3)$. Then $T(p) = q \in \mathcal{P}(\mathbb{R})$.

Let's prove that T is a linear map.

- for $p, q \in \mathcal{P}(\mathbb{R})$,

$$T(p + q) = \frac{p(x) + q(x) - p(3) - q(3)}{x - 3} = \frac{p(x) - p(3)}{x - 3} + \frac{q(x) - q(3)}{x - 3} = T(p) + T(q).$$

- for $p \in \mathcal{P}(\mathbb{R})$ and $c \in \mathbb{R}$,

$$T(cp) = \frac{cp(x) - cp(3)}{x - 3} = c \frac{p(x) - p(3)}{x - 3} = cT(p).$$

Therefore T is a linear map. □

Problem 7 (4-12). Suppose m is a nonnegative integer and $p \in \mathcal{P}_m(\mathbb{C})$ is such that there are distinct real numbers $x_0, x_1, \dots, x_m$ with $p(x_k) \in \mathbb{R}$ for each $k = 0, 1, \dots, m$. Prove that all coefficients of p are real.

Proof. By the Lagrange interpolation formula, we can write

$$p(x) = \sum_{k=0}^m p(x_k) \prod_{j \neq k} \frac{x - x_j}{x_k - x_j}.$$

Since $p(x_k) \in \mathbb{R}$ for each $k = 0, 1, \dots, m$, the coefficients of p are all real. □

Problem 8 (4-13). Suppose $p \in \mathcal{P}(\mathbf{F})$ with $p \neq 0$. Let

$$U = \{pq : q \in \mathcal{P}(\mathbf{F})\}.$$

1. Show that $\dim(\mathcal{P}(\mathbf{F})/U) = \deg p$.
2. Find a basis of $\mathcal{P}(\mathbf{F})/U$.

Solution. Let $\deg p = m$. We will show that $\{1 + U, x + U, \dots, x^{m-1} + U\}$ is a basis of $\mathcal{P}(\mathbf{F})/U$.

First, we show that $\{1 + U, x + U, \dots, x^{m-1} + U\}$ is linearly independent. Suppose $c_0, c_1, \dots, c_{m-1} \in \mathbf{F}$ such that

$$c_0(1 + U) + c_1(x + U) + \dots + c_{m-1}(x^{m-1} + U) = 0 + U.$$

Then $c_0 + c_1x + \dots + c_{m-1}x^{m-1} \in U$, which implies that there exists $q \in \mathcal{P}(\mathbf{F})$ such that

$$c_0 + c_1x + \dots + c_{m-1}x^{m-1} = pq.$$

Since $\deg p = m$ and $\deg(q) < m$, this is impossible unless $c_0 = c_1 = \dots = c_{m-1} = 0$. Therefore, $\{1 + U, x + U, \dots, x^{m-1} + U\}$ is linearly independent.

Next, we show that $\{1 + U, x + U, \dots, x^{m-1} + U\}$ spans $\mathcal{P}(\mathbf{F})/U$. Let $r \in \mathcal{P}(\mathbf{F})$. By the division algorithm, there exist unique polynomials $q, r' \in \mathcal{P}(\mathbf{F})$ such that

$$r = pq + r', \quad \deg(r') < \deg(p).$$

Then $r + U = pq + r' + U = r' + U$, and since $r' = \sum_{k=0}^{m-1} a_k x^k$ for some $a_k \in \mathbf{F}$, we have

$$r' + U = \sum_{k=0}^{m-1} a_k (x^k + U).$$

Therefore, $\{1 + U, x + U, \dots, x^{m-1} + U\}$ spans $\mathcal{P}(\mathbf{F})/U$. Hence, it is a basis.

Problem 9 (4-14). Suppose $p, q \in \mathcal{P}(\mathbb{C})$ are nonconstant polynomials with no zeros in common. Let $m = \deg p$ and $n = \deg q$. Use linear algebra as outlined below in (a)–(c) to prove that there exist $r \in \mathcal{P}_{n-1}(\mathbb{C})$ and $s \in \mathcal{P}_{m-1}(\mathbb{C})$ such that

$$rp + sq = 1.$$

(a) Define

$$T : \mathcal{P}_{n-1}(\mathbb{C}) \times \mathcal{P}_{m-1}(\mathbb{C}) \rightarrow \mathcal{P}_{m+n-1}(\mathbb{C})$$

by

$$T(r, s) = rp + sq.$$

Show that the linear map T is injective.

(b) Show that the linear map T in (a) is surjective.

(c) Use (b) to conclude that there exist $r \in \mathcal{P}_{n-1}(\mathbb{C})$ and $s \in \mathcal{P}_{m-1}(\mathbb{C})$ such that $rp + sq = 1$.

Proof. WLOG, we can assume that $p = (x - x_1)(x - x_2) \cdots (x - x_m)$ and $q = (x - y_1)(x - y_2) \cdots (x - y_n)$ are monic.

(a) Suppose that $T(r, s) = 0$ for some $r \in \mathcal{P}_{n-1}(\mathbb{C})$ and $s \in \mathcal{P}_{m-1}(\mathbb{C})$. Then $rp + sq = 0$. Since $p(x_i) = 0 \neq q(x_i)$ for each $i = 1, 2, \dots, m$, we have $s(x_i) = 0$ for each $i = 1, 2, \dots, m$. Since $s \in \mathcal{P}_{m-1}(\mathbb{C})$, this implies that $s = 0$. Similarly, since $q(y_j) = 0$ for each $j = 1, 2, \dots, n$, we have $r(y_j) = 0$ for each $j = 1, 2, \dots, n$. Since $r \in \mathcal{P}_{n-1}(\mathbb{C})$, this implies that $r = 0$. Therefore, T is injective.

(b) Since $\dim(\mathcal{P}_{n-1}(\mathbb{C}) \times \mathcal{P}_{m-1}(\mathbb{C})) = n + m = \dim(\mathcal{P}_{m+n-1}(\mathbb{C}))$, and T is injective, we can conclude that T is surjective.

(c) Since T is surjective, there exist $r \in \mathcal{P}_{n-1}(\mathbb{C})$ and $s \in \mathcal{P}_{m-1}(\mathbb{C})$ such that $T(r, s) = rp + sq = 1$.

□