

Homework 4

程昊一

*All of my homework is available at <https://github.com/lacampanella233/Homework.git>.

Problem 1 (3E-1). Suppose T is a function from V to W . The graph of T is the subset of $V \times W$ defined by

$$\text{graph of } T = \{(v, Tv) \in V \times W : v \in V\}.$$

Prove that T is a linear map if and only if the graph of T is a subspace of $V \times W$.

Proof. Denote the graph of T by $G \subset V \times W$. On the one hand, if T is a linear map, then for and $(v_1, Tv_1) \in G$ and $(v_2, Tv_2) \in G$, and $\lambda \in \mathbb{F}$, we have

$$(v_1, Tv_1) + (v_2, Tv_2) = (v_1 + v_2, T(v_1 + v_2)) \in G, \quad \lambda(v_1, Tv_1) = (\lambda v_1, T(\lambda v_1)) \in G.$$

Hence, G is a subspace of $V \times W$.

On the other hand, if G is a subspace of $V \times W$, then for any $v_1, v_2 \in V$ and $\lambda \in \mathbb{F}$, we have

$$(v_1, Tv_1) + (v_2, Tv_2) = (v_1 + v_2, Tv_1 + Tv_2) \in G, \quad \lambda(v_1, Tv_1) = (\lambda v_1, \lambda Tv_1) \in G,$$

which implies that $T(v_1 + v_2) = Tv_1 + Tv_2$ and $T(\lambda v_1) = \lambda Tv_1$. Hence, T is a linear map. $\square$

Problem 2 (3E-9). Prove that a nonempty subset A of V is a translate of some subspace of V if and only if

$$\lambda v + (1 - \lambda)w \in A$$

for all $v, w \in A$ and all $\lambda \in \mathbb{F}$.

Proof. On the one hand, if A is a translate of U , then there exists $t \in V$ such that $A = t + U$. For any $v, w \in A$ and $\lambda \in \mathbb{F}$, we have $v = t + u_1$ and $w = t + u_2$ for some $u_1, u_2 \in U$. Then

$$\lambda v + (1 - \lambda)w = \lambda(t + u_1) + (1 - \lambda)(t + u_2) = t + (\lambda u_1 + (1 - \lambda)u_2) \in t + U.$$

On the other hand, if $\lambda v + (1 - \lambda)w \in A$ for all $v, w \in A$ and all $\lambda \in \mathbb{F}$, We will show that there exists $t \in V$ and $U \subset V$ such that $A = t + U$.

Pick $a_0 \in A$ and $a_0 = t + u_0$. Denote

$$U = \{a - a_0 : a \in A\}.$$

It suffices to prove U is a subspace of V .

- (a) U is closed under addition. That's because for u_1, u_2 in U , set $a_1 = a_0 + u_1, a_2 = a_0 + u_2$. Then $a_1/2 + a_2/2 = a_0 + (u_1 + u_2)/2 \in A$, hence $2(a_1/2 + a_2/2) + (1-2)a_0 = a_0 + (u_1 + u_2) \in A$, which implies $u_1 + u_2 \in U$.
- (b) U is closed under scalar multiplication. That's because for $u \in U$ and $\lambda \in \mathbb{F}$, set $a = a_0 + u$. Then $\lambda a + (1-\lambda)a_0 = a_0 + \lambda u \in A$, which implies $\lambda u \in U$.

□

Problem 3 (3E-14). Suppose U and W are subspaces of V and $V = U \oplus W$. Suppose $w_1, \dots, w_m$ is a basis of W . Prove that $w_1 + U, \dots, w_m + U$ is a basis of V/U .

Proof. First we will show that $w_1 + U, \dots, w_m + U$ is linearly independent. Suppose $\lambda_1(w_1 + U) + \dots + \lambda_m(w_m + U) = 0 + U$ for some $\lambda_1, \dots, \lambda_m \in \mathbb{F}$. Then $\lambda_1 w_1 + \dots + \lambda_m w_m \in U$. Since $U \cap W = \{0\}$, we have $\lambda_1 w_1 + \dots + \lambda_m w_m = 0$, which implies $\lambda_1 = \dots = \lambda_m = 0$. Hence, $w_1 + U, \dots, w_m + U$ is linearly independent.

Next we will show that $w_1 + U, \dots, w_m + U$ spans V/U . For any $v + U \in V/U$, we can write $v = u + w$ for some $u \in U$ and $w \in W$. Since $w_1, \dots, w_m$ is a basis of W , we can write $w = \lambda_1 w_1 + \dots + \lambda_m w_m$ for some $\lambda_1, \dots, \lambda_m \in \mathbb{F}$. Then

$$v + U = u + w + U = \lambda_1(w_1 + U) + \dots + \lambda_m(w_m + U).$$

Hence, $w_1 + U, \dots, w_m + U$ spans V/U .

Combining the above two paragraphs, we conclude that $w_1 + U, \dots, w_m + U$ is a basis of V/U . □

Problem 4 (3E-15). Suppose U is a subspace of V and $v_1 + U, \dots, v_m + U$ is a basis of V/U and $u_1, \dots, u_n$ is a basis of U . Prove that $v_1, \dots, v_m, u_1, \dots, u_n$ is a basis of V .

Proof. First we will show that $v_1, \dots, v_m, u_1, \dots, u_n$ is linearly independent. Suppose $\lambda_1 v_1 + \dots + \lambda_m v_m + \mu_1 u_1 + \dots + \mu_n u_n = 0$ for some $\lambda_1, \dots, \lambda_m, \mu_1, \dots, \mu_n \in \mathbb{F}$. Then $\lambda_1(v_1 + U) + \dots + \lambda_m(v_m + U) = 0 + U$, which implies $\lambda_1 = \dots = \lambda_m = 0$. Hence, $\mu_1 u_1 + \dots + \mu_n u_n = 0$, which implies $\mu_1 = \dots = \mu_n = 0$. Therefore, $v_1, \dots, v_m, u_1, \dots, u_n$ is linearly independent.

Next we will show that $v_1, \dots, v_m, u_1, \dots, u_n$ spans V . Since $v_1 + U, \dots, v_m + U$ is a basis of V/U , we can write $v + U = \lambda_1(v_1 + U) + \dots + \lambda_m(v_m + U)$ for some $\lambda_1, \dots, \lambda_m \in \mathbb{F}$. Then

$$v = \lambda_1 v_1 + \dots + \lambda_m v_m + u$$

for some $u \in U$. Since $u_1, \dots, u_n$ is a basis of U , we can write $u = \mu_1 u_1 + \dots + \mu_n u_n$ for some $\mu_1, \dots, \mu_n \in \mathbb{F}$. Then

$$v = \lambda_1 v_1 + \dots + \lambda_m v_m + \mu_1 u_1 + \dots + \mu_n u_n.$$

Hence, $v_1, \dots, v_m, u_1, \dots, u_n$ spans V .

Combining the above two paragraphs, we conclude that $v_1, \dots, v_m, u_1, \dots, u_n$ is a basis of V . $\square$

Problem 5 (3E-18). Suppose that U is a subspace of V such that V/U is finite-dimensional.

(a) Show that if W is a finite-dimensional subspace of V and $V = U + W$, then $\dim W \geq \dim V/U$.

(b) Prove that there exists a finite-dimensional subspace W of V such that $\dim W = \dim V/U$ and $V = U \oplus W$.

Proof. (a) Suppose W is a finite-dimensional subspace of V and $V = U + W$. Then the map $T: W \rightarrow V/U$ defined by $Tw = w + U$ is a surjective linear map. Hence, we have

$$\dim W = \dim \ker T + \dim V/U \geq \dim V/U.$$

(b) Since V/U is finite-dimensional, we can pick a basis $v_1 + U, \dots, v_m + U$ of V/U . Denote $W = \text{span}\{v_1, \dots, v_m\}$. We will show that $V = U \oplus W$ and $\dim W = \dim V/U$.

First we will show that $V = U + W$. For any $v \in V$, we can write $v + U = \lambda_1(v_1 + U) + \dots + \lambda_m(v_m + U)$ for some $\lambda_1, \dots, \lambda_m \in \mathbb{F}$. Then

$$v = \lambda_1 v_1 + \dots + \lambda_m v_m + u$$

for some $u \in U$. Hence, $V = U + W$.

Next we will show that $U \cap W = \{0\}$. Suppose $u \in U \cap W$. Then $u = \lambda_1 v_1 + \dots + \lambda_m v_m$ for some $\lambda_1, \dots, \lambda_m \in \mathbb{F}$. Hence, we have

$$\lambda_1(v_1 + U) + \dots + \lambda_m(v_m + U) = 0 + U,$$

which implies $\lambda_1 = \dots = \lambda_m = 0$. Therefore, $u = 0$ and $U \cap W = \{0\}$.

Combining the above two paragraphs, we conclude that $V = U \oplus W$.

Finally, since $v_1 + U, \dots, v_m + U$ is a basis of V/U , we have $\dim W = \dim \text{span}\{v_1, \dots, v_m\} = m = \dim V/U$. $\square$

Problem 6 (3F-5). Suppose $T \in \mathcal{L}(V, W)$ and $w_1, \dots, w_m$ is a basis of $\text{im } T$. Hence for each $v \in V$, there exist unique numbers $\varphi_1(v), \dots, \varphi_m(v)$ such that

$$Tv = \varphi_1(v)w_1 + \dots + \varphi_m(v)w_m,$$

thus defining functions $\varphi_1, \dots, \varphi_m$ from V to $\mathbb{F}$. Show that each of the functions $\varphi_1, \dots, \varphi_m$ is a linear functional on V .

Proof. Without loss of generality, we only need to show that φ_1 is a linear functional on V . For any $v_1, v_2 \in V$ and $\lambda \in \mathbb{F}$, we have

$$\varphi_1(v_1 + v_2)w_1 + \dots = v_1 + v_2 = (\varphi_1(v_1) + \varphi_1(v_2))w_1 + \dots \implies \varphi_1(v_1 + v_2) = \varphi_1(v_1) + \varphi_1(v_2),$$

$$\lambda\varphi_1(v_1)w_1 + \dots = \lambda v_1 = \varphi_1(\lambda v_1)w_1 + \dots \implies \varphi_1(\lambda v_1) = \lambda\varphi_1(v_1).$$

$\square$

Problem 7 (3F-8). Suppose $v_1, \dots, v_n$ is a basis of V and $\varphi_1, \dots, \varphi_n$ is the dual basis of V' . Define $\Gamma : V \rightarrow \mathbb{F}^n$ and $\Lambda : \mathbb{F}^n \rightarrow V$ by

$$\Gamma(v) = (\varphi_1(v), \dots, \varphi_n(v))$$

and

$$\Lambda(a_1, \dots, a_n) = a_1 v_1 + \dots + a_n v_n.$$

Explain why Γ and Λ are inverses of each other.

Proof.

$$\Lambda\Gamma(v) = \Lambda(\varphi_1(v), \dots, \varphi_n(v)) = \varphi_1(v)v_1 + \dots + \varphi_n(v)v_n = v,$$

$$\Gamma\Lambda(a_1, \dots, a_n) = \Gamma(a_1 v_1 + \dots + a_n v_n) = (\varphi_1(a_1 v_1 + \dots + a_n v_n), \dots, \varphi_n(a_1 v_1 + \dots + a_n v_n)) = (a_1, \dots, a_n).$$

□

Problem 8 (3F-10). Suppose m is a positive integer.

(a) Show that $1, x-5, \dots, (x-5)^m$ is a basis of $\mathcal{P}_m(\mathbb{R})$.

(b) What is the dual basis of the basis in (a)?

Solution. (a) For any $p \in \mathcal{P}_m(\mathbb{R})$, we can write uniquely

$$p(x) = \sum_{k=0}^m \frac{p^{(k)}(5)}{k!} (x-5)^k,$$

which is not hard to verify.

(b) From the formula in (a), we can see that the dual basis of $1, x-5, \dots, (x-5)^m$ is

$$\varphi_k(p) = \frac{p^{(k)}(5)}{k!}, \quad 0 \leq k \leq m.$$

Problem 9 (3F-11). Suppose $v_1, \dots, v_n$ is a basis of V and $\varphi_1, \dots, \varphi_n$ is the corresponding dual basis of V' . Suppose $\psi \in V'$. Prove that

$$\psi = \psi(v_1)\varphi_1 + \dots + \psi(v_n)\varphi_n.$$

Proof. For any $v \in V$, we can write $v = \varphi_1(v)v_1 + \dots + \varphi_n(v)v_n$. Applying ψ to both sides, we get

$$\psi(v) = \psi(\varphi_1(v)v_1 + \dots + \varphi_n(v)v_n) = \varphi_1(v)\psi(v_1) + \dots + \varphi_n(v)\psi(v_n),$$

which is exactly we want to prove. □