


GROUP PROJECT 2.1, FLAVOR A

Some Group

Jane Doe  
12345678

JaneJohn Smith  
10010010

Alex kwikwōł  
9999<sup>9999</sup>



12 km <sup>wut</sup> lim<sub>x→0</sub>

1. (3 points) one

("1": (a) (Extra, no) two  
1)  
("1": i. (1 point) three  
1'  
"1{"1":  
a": 1,  
1) "1-  
a":  
1,  
"1-  
a-  
i":  
1)

what

User Manual  
**Math**  
Formatting math equations is probably the reason you are here. Unlike LaTeX, math in Typst is simple.

$E = m c^2$   
 $e^{i \pi} = -1$   
 $x = \frac{-b \pm \sqrt{b^2 - 4 a c}}{2 a}$

$E = mc^2$   
 $e^{i\pi} = -1$   
 $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

For “block” or “display” math, leave a space or newline between the dollar sign and the equations.

$E = m c^2$ 

$E = mc^2$ 

(1)

Documented are built-in **math functions** and **symbols**

## Numbering and Referencing Equations

Note that you must enable equation numbering to reference equations, which is set by this template.

```
$
e^(i pi) = -1 #<euler>
$
@euler is Euler's identity. \
#link(<euler>)[This] is the same thing.
```

$$e^{i\pi} = -1 \tag{2}$$

**Equation 2** is Euler's identity.  
**This** is the same thing.

## Extra Math Symbols and Functions

The `physica` package provides additional math symbols and functions.

```
$A^T, curl vb(E) = - pdv(vb(B), t)$
$tensor(Lambda,+mu,-nu) = dmat(1,RR)$
$f(x,y) dd(x,y)$
```

$$\left. \begin{array}{l} A^T, \nabla \times \boldsymbol{E} = -\frac{\partial \boldsymbol{B}}{\partial t} \\ \Lambda^\mu{}_\nu = \begin{pmatrix} 1 & \\ & \mathbb{R} \end{pmatrix} \\ f(x,y) \, dx \, dy \end{array} \right|$$

It is imported in this template.

## Units and Quantities

Although no as common as in physics, we do sometimes need to use units and quantities. This template uses the `metro` package for this purpose. If you prefer, you can also use the `unify` package.

---

Other helps: `introduction`, `getting-started`, `setup`, `author`, `drawing`, `question`, `solution`, `caveats`.