

THE PHYSICS COMPENDIUM

Matthew Lach

The Method 1.6 — a reading copy of the seated volume,
pressed 5 September 2026

Contents

Λ — the elemental index	3
Λ_{spectra} — the index of Rydberg channels	3
The point of observability — what the equation actually delivers	5
Kinematic and stateful — an interface the register found in its own objects	5
Placing a cell and valuing it are different operations	6
Where relativity enters, and where it does not	6
The bracket — the transition from a channel to an interval	7
What the interface still does not state	8
THE LÖWDIN-SOLUTION INDEXES — Λ_{chain}, Λ_{cinf}, Λ_{V5}	9
Λ_{chain} — the derived filling index	9
Λ_{cinf} — the twin index at $c \rightarrow \infty$	9
Λ_{V5} — the correlation closure at the contested rows	10
The corridor law — interface only	10
The collapse condition — interface only	10
The defect — interface only	11
Λ_3 — THE THREE-BODY INDEX	12
Λ_{phys} — THE INDEX OF PHYSICAL PARAMETERS	14
What the index shows about the work's dependence	14
The parameters	15
The three failures that are measured rather than anticipated	21
Λ_{chem} — THE CHEMICAL PROPERTIES INDEX	22
Why the compendium needed it	22
The coordinates	22
The fifth seat	22
The closure, and the boundary it declares	22
The last cell, and what filled it	23
The routing, by residual	23
Λ_{PCA} — PHYSICS $\oplus$ CHARGE $\oplus$ AMPLITUDE, AS ONE	24
The merge	24
What it gives that neither parent does	24
Λ_{amp} — the amplitude's own index	24
A CORRECTION TO SEATON'S RATIO	25
THE CHANNEL CONSTANTS IN Λ_{phys}	26
Failure modes	26
THE PRIOR-ART CHAIN — the seed's forced structure	26

Compiled 2026-08-12. **248 registered objects, 17 of them physical mechanisms.**

This compendium states the **interface** between a physical quantity and the index that holds it. It is not a list of mechanisms — those are in the Mathematical Compendium, and a third copy would drift as register 778’s list did. It states, for each index, what the cells stand for, what number is attached to them, the rule taking one to the other, which constants that rule requires, **what must be measured rather than computed**, and what physics does and does not do.

Four faults in a single session were transitions performed without their rule written down: register 868 used R_∞ for every species where the reduced-mass constant is correct; register 879 wrote Li III’s ionisation limit as a formula where a measurement was required; register 766 passed atomic numbers where the core’s charge is needed; register 782 set a verification column equal to its own denominator. **This document exists because those four share one cause.**

Λ — the elemental index

What it indexes. Electron configurations of the ground states of the elements, on eight coordinates (n, ℓ , k, q, e, f, g, 2S). **976 cells.**

The quantity. None. A cell of Λ carries no physical number: it is an *arrangement*, not a measurement.

The transition. A configuration is read off the aufbau order and its subshells mapped to coordinates. the index against the real subshells establishes that Λ ’s eight constraints hold on 247 real subshells across all 118 elements — **18,288 tests, no failures.**

Constants required. None.

Must be measured. Nothing. Λ is closed — $|\Lambda| = |\mathcal{R}(\Lambda)| = 976$, $\mathbf{E}(\Lambda) = \mathbf{0}$ — and the partition identity holds exactly: $6,912 = 976 + 0 + 5,936$ (register 833).

What physics does. It supplies the aufbau ordering and the shell capacities, and nothing else. **What it does NOT do:** confer predictive power. §18 (and the preface) states that Λ “*contains no physics that was not already in the quantum numbers; it is a recoding ... a recoding adds nothing.*” Register 834 measured the consequence: on the same closed index, the T-bracket passes 789 of 789 and the δ -bracket passes 546 of 789. **Closure coexists with a bracket that holds and one that fails, so it cannot be what makes either true.**

Λ_{spectra} — the index of Rydberg channels

What it indexes. Rydberg series: a fixed parent core, a fixed ℓ , n running. **596 channels across 70 spectra, 2,269 interior cells.** The complete index over the same alphabets is closed: $|X| = |\mathcal{R}(X)| = 624$, $\mathbf{E} = \mathbf{0}$ (register 853).

The quantity. The **quantum defect** δ — a continuous real number measuring how far the Rydberg orbital penetrates the ionic core. It is not an integer and has no factorization (register 896).

The transition. For a level at energy E below a limit I,

$$n^* = Z_c \sqrt{R_M / (I - E)} \quad \delta = n - n^*$$

Constants required, and which value:

- **Z_c — the charge of the CORE, not the atom.** NIST states the formula as $E_{nl} = -Z_c^2/(n-\delta)^2$, “where Z_c is the charge of the core”. Be III and B III are third spectra and take $Z_c = 3$, not 4 and 5. Passing atomic numbers gave defects of -2 to -5 with spreads above 1.0 (registers 766, 883).
- **R_M — the reduced-mass Rydberg, $R_\infty/(1 + m_e/M)$, not R_∞ .** For helium 109,722.386 against 109,737.316. Using R_∞ understates every defect by $n^*(\sqrt{R_\infty/R_M} - 1)$ — 0.00240 for helium at $n^* = 35$ — which drove He I’s high- ℓ defects **negative**. Precision spectroscopy writes its constant this way as a matter of course (registers 868, 883).
- **The exception.** Where a level table is itself a theoretical hydrogenic one computed from R_∞ , R_∞ is what belongs in the formula: the constant must match the **data’s** convention, not the physics in the abstract. Applying a reduced mass to Li III’s table drove its spreads from 0.0000 to 0.0125 (register 872).

Must be measured, not computed. The ionisation limit. Writing it as a formula is the single most damaging thing that can be done to a channel. Li III’s limit was recorded as “ $Z^2R = 9 \times 109737.31568$ (hydrogenic, $Z=3$)” — a bare Coulomb expression that omits the QED and relativistic terms. Fitting the limit from the series gives **987,662.29 $\pm$ 0.36, higher by 26.45 cm^{-1}** , and every Li III channel moves from $-0.0038\dots-0.0082$ to a uniform **+0.0003** (register 879). The same deficit appears at B V scaled by Z^4 : **678 cm^{-1} above $25R_\infty$** (register 895).

How a wrong limit announces itself. A defect that should be constant instead deviates, and **the power of n^* names the cause:**

n^{*3}	a shift in the ionisation limit	
n^{*7}	a magnetic field	— needs tens of tesla
n^{*10}	an electric field	— begins at $n > 60$

Two of the three cannot fire on this compendium. Its highest member anywhere is **$n = 56$** (Si I’s nd (3/2,5/2)); eight channels exceed $n = 40$ and **none exceeds 60**. So every residual chased here was a limit, and none was a field — not by luck but because ASD’s tables stop below where field effects begin (register 915).

And the limit term’s tolerance is published: a shift of even 0.2 cm^{-1} “causes totally different behavior of quantum defect versus n ”. Li III’s was **26.45** — a hundred and thirty times that (register 914).

Li III’s residual went as n^3 and the limit was the fault. This is published and was derived independently here (registers 877, 885).

Where the index STOPS, and why that is a decision. Λ is finite because all 118 ground configurations are known within its caps. **The spectra index is not:** Rydberg series are unbounded in n and ℓ runs to $n-1$, so it is **infinite unless capped**, and every cap is a choice. Three defensible ones give:

the measured alphabet	1,312 cells	$E = 1,084$
charge 1.. $Z-1$, elements held	4,376 cells	$E = 4,148$
all 118 elements, ℓ 0-7	55,224 cells	$E = 54,996$

A factor of fifty between them, and all three close when complete. Closure is a result about $\mathcal{R}$; the E that accompanies it is a result about the cap. At the physical cap the compendium holds **0.413%** — Λ holds 100% within its own (registers 959–962).

What physics does. It sets δ through core penetration, and the sixteen mechanisms of the Mathematical Compendium’s § P. The spectral mechanisms describe how. **What it does NOT do:** make δ an integer, or give the index predictive power over cells it does not contain. Of 624 cells, 160 are measured and **193 are fully isolated** — no measured channel adjacent in ℓ or in an isoelectronic sequence, so no bound reaches them (register 890).

The point of observability — what the equation actually delivers

The quantum defect is a phase shift. Outside the ionic core, where the potential is purely Coulombic, the radial Schrödinger solution is a superposition of the regular and irregular Coulomb functions:

$$\psi = \cos(\pi\delta) \cdot f(E, \ell, r) - \sin(\pi\delta) \cdot g(E, \ell, r)$$

and $\pi\delta$ is the phase shift with respect to the pure hydrogenic solution, due to all short-range non-Coulombic interactions at $r < r_c$. Seaton's theorem continues it across threshold, $\delta_\ell(0) = \pi\mu_\ell(0)$, so the same number describes bound levels and scattering.

The consequence, and it is sharp. The equation depends on δ only through $\cos \pi\delta$ and $\sin \pi\delta$. **What it determines is δ modulo 1. The integer part is not in the equation** — it is supplied afterwards by counting nodes inside the core, and it is a labelling, not a measurement.

What this costs the compendium. Register 891 tested the rule $\text{floor}(\delta) = (\text{core orbitals of that } \ell) - 1$ and found it holding for 318 of 407 channels — 78%, with Al I's ns at 1.767 where the rule requires 2.x. **It was testing a convention.** And testing the l-collapse (register 693) both ways:

on the full defect	150 correct, 8 inverted	94.9%
on $\delta \bmod 1$ (observable)	127 correct, 31 inverted	80.4%

The ordering survives on the assigned value and degrades on the observable one. So the l-collapse is a statement about node-counting as much as about penetration. Only **96 of 431** channels have $|\delta| > 1$ at all — the rest have no integer part to assign, and for them the distinction does not arise (registers 910–913).

The general rule this gives. *A mechanism that holds on the assigned value and fails on the observable one is a statement about the index, not about the atom.* That is §18's distinction — a recoding adds nothing — reappearing at the level of a single number rather than a whole lattice.

Kinematic and stateful — an interface the register found in its own objects

A geometric hypothesis cannot do a state's work, and the register's one OPEN object is what happens when it is asked to.

the presymplectic potential assumes $\Theta = 0$ — the non-expanding condition, geometry and nothing else — and it succeeds: the presymplectic form is block diagonal in y and the commutator is $(1/4\sqrt{q}) \operatorname{sgn}(u-u') \delta^{(d-2)}(y-y')$.

the open object — half-sided modular inclusion on a non-expanding horizon adds exactly one hypothesis, ω Hadamard, and that is a condition on a STATE.

Every obstruction encountered sits on the stateful side:

- a geometric modular flow needs a Killing vector — the flow is stateful and the geometry must match it
- half-sided modular inclusion $\iff$ a positive generator, and positivity is a property of a state
- $\Theta = 0$ is invariant under $\ell \rightarrow a(y)\ell$, so **the geometry does not fix the state**
- smoothness supplies a *local* Hartle–Hawking state, and local states do not glue
- Hadamard is microlocal and therefore too weak to select one
- and uniqueness, where it is available at all, needs invariance under a Killing flow

And the split is NOT geometry against state, which was the first reading and was wrong. The standard definition of a non-expanding horizon has three conditions and the third is “*Einstein field equations hold on Δ , and the stress-energy tensor T_{ab} is such that $-T^a{}_b \ell^b$ is future causal*” — **an energy condition, hence a condition**

on the state. Through the kk -component of the Einstein equation it forces the null-null flux $T_{ab} \ell^a \ell^b$ and the shear σ_{ab} to VANISH, giving $\mathcal{E}_\ell q_{ab} = 0$. *So the presymplectic potential's hypothesis is not an assumption but a consequence of the horizon's own definition, which is why it succeeds.*

What the condition does not do is reach the right order. It fixes the BACKGROUND — vanishing shear, vanishing null flux — while the linearised Raychaudhuri $\partial_u \delta\Theta = -8\pi G_N T_{uu}$ carries $T_{uu} \neq 0$ on the perturbations, which is what the theory is about. **Half-sidedness is a spectral statement about those perturbations, and future-causality of $-T^a_b \ell^a \ell^b$ is not one** (registers 1038–1042).

The dimensional ledger states the split arithmetically: $d - 1 = 1 + (d - 2)$, one HSMI per generator supplying the affine line and the transverse direct integral coming from the kinematic side. *The stateful object supplies one dimension; the kinematic one supplies $d - 2$.*

And it explains one result. Computing the free half on one generator gives a positive translation generator immediately — negative to positive spectral weight 5×10^{-5} — **because a state was supplied by hand, the vacuum on a null line.** *The moment a state is given the inclusion follows; nothing on a non-expanding horizon gives one.* Registers 1033–1037.

Placing a cell and valuing it are different operations

$\mathcal{R}$ places. Mechanisms value. The distinction is invisible in Λ , where a cell is an arrangement and carries no number, so recovering the cell recovers everything. It is the whole difficulty in Λ_{spectra} , where the cell carries a quantum defect and that is its entire content.

placed by $\mathcal{R}$	1,762 of 1,775 cells	99.3%
valued	525	32%

A relation is usable as a STEP — carrying a value from one cell to another — only where the physics has collapsed to ONE PARAMETER. Measured on pairs where both cells are known, as a geometric scatter:

N_e at fixed charge $\div$ Thomas-Fermi, s+p, $N_e \geq 9$	1.07
elem, s+p, $N_e \geq 9$, raw	1.10
ℓ at $\ell \geq 4$, Seaton's ratio, filtered	1.12
charge at fixed N_e , s	1.23
charge at fixed N_e , p	1.38
ℓ at p, d, f	3.31
elem, all ℓ and N_e	3.85
N_e at fixed charge, raw	4.12

Above $\ell = 4$ the electron never enters the core and only the polarisability matters — one number, and Seaton's formula follows. At $\ell \leq 1$ with $N_e \geq 9$ penetration is statistical and Thomas-Fermi governs. At $\ell = 2, 3$ or $N_e < 9$ neither dominates, and four separate corrections — Seaton's ratio, core-orbital counts, the defect's own magnitude, a free Thomas-Fermi exponent — each left the scatter where it was.

The transition zone is not a missing systematic. It is the absence of one, and those cells require measurement rather than propagation (registers 1102–1118).

Where relativity enters, and where it does not

One place: the ionisation limit of a hydrogenic ion. A Coulomb expression $Z^2 R$ gives the limit only to order $(Z\alpha)^2$. The Sommerfeld correction — special relativity applied to the atom in 1916, and shown by Gordon and Darwin to follow exactly from the Dirac equation — adds

$$\Delta E = Z^4 \alpha^2 R_M / 4 \quad \text{for the } 1s \text{ state}$$

Measured against the reduced-mass baseline $Z^2 R_M$, three hydrogenic species give:

Li III	deficit 103.94	Dirac 118.32	ratio 0.878
Be IV	deficit 329.80	Dirac 373.97	ratio 0.882
B V	deficit 816.68	Dirac 913.03	ratio 0.894

0.8849 ± 0.0069 across $Z = 3, 4, 5$ (register 906). The missing 11.5% is the **1s Lamb shift**, which raises the level and reduces the binding; it goes as $\alpha^3 Z^4$ against Dirac's $\alpha^2 Z^4$, so their ratio is order α times a slowly varying function — which is why three elements agree to under one per cent (register 908).

The practical consequence. Register 879's Li III fault was a Coulomb expression used where a relativistic one was required: the limit was written as $9R_\infty$ and was low by 26.45 cm^{-1} , showing as a spurious defect of -0.0048 growing as n^3 across every ℓ . **A hydrogenic limit computed rather than measured will always be low by this term.**

Where relativity does NOT enter. General relativity has no measurable term at these energies, and the compendium makes no use of it. The relativistic content here is entirely the special-relativistic kinetic correction, spin-orbit coupling and the Darwin term — the three that make up fine structure — plus the QED Lamb shift, which is not relativity at all (register 909).

The bracket — the transition from a channel to an interval

What it indexes. An interior cell: a series member with a measured neighbour on each side.

The quantity. Not a value but an **interval**, and the book is explicit that this is the point: §22 says it “*fits nothing, assumes no functional form, and returns an interval rather than a value. It is a deduction, not a prediction.*”

The transition — and there are TWO, which the book does not distinguish.

- **§22.1, on binding energy.** $T = I - E$ is monotone in n , so $T(n)$ lies between $T(n-1)$ and $T(n+1)$. **Passes 789 of 789.** A Rydberg series rises toward its limit by construction, so this form **cannot fail** (registers 797, 800, 830).
- **§25.6.1, on the defect.** $\delta(n)$ lies between $\delta(n-1)$ and $\delta(n+1)$, resting on “*within a channel, δ falls monotonically with n* ”. **Passes 546 of 789.**

The trade-off, stated. §22.1's bracket is a genuine deduction and carries no information. §25.6.1's carries information and is not a deduction. **Chapter 22's modesty is exactly what buys the 100%, and the price is that the 100% was never evidence** (register 803).

And the monotonicity claim is the wrong form. The extended Ritz expansion is

$$\delta(n) = \delta_0 + \delta_2 / (n - \delta_0)^2 + \delta_4 / (n - \delta_0)^4 + \dots$$

and caesium's nF series is measured with $\delta_2 = -0.2014(16)$ — **negative** — its defects **RISING** with n . So the law is that **δ approaches δ_0 monotonically**; the direction is $\text{sign}(\delta_2)$, a property of the channel, and falling is only the common case (register 858).

What physics does. It guarantees the monotonicity of T . **What it does NOT do:** guarantee the monotonicity of δ , which holds in 113 of 130 resolved steps — 87%, interval 80–92% — and only where a step exceeds its own uncertainty. With error estimated from quoted decimals instead, the same test gives 55%, and the intervals do not overlap (register 822).

What the interface still does not state

Uncertainties. Five of 49 species files carry a quoted uncertainty column. The monotonicity law is testable only where a step can be distinguished from its own error, and **44 of 49 species do not supply what that needs** (register 807).

The bracket's own inputs. The bracket reads untested for every channel built here, because running the bracketing method needs measured neighbours and a tolerance and neither was supplied. **285 of 431 channels are unverified on the compendium's central claim**, and the column says so rather than asserting a verification that never ran (register 782).

Appendix B's inputs. 133 channels were lifted with their values and without their levels. `spectra_raw/` holds no He I file, so when register 868 corrected the Rydberg constant those channels **could not be recomputed**. *A result kept without its inputs cannot be corrected when its method is* (register 871).

THE LÖWDIN-SOLUTION INDEXES — Λ_{chain} , Λ_{cinf} , Λ_{V5}

Interface entries for the indexes of Chapter 35, in this compendium's own template: what the cells stand for, what number is attached, the rule taking one to the other, the constants that rule requires, what must be measured rather than computed, and what physics does and does not do. Register 1701–1712. The mechanisms are stated once, in the Mathematical Compendium (family LS); a second copy would drift.

Λ_{chain} — the derived filling index

What it indexes. One row per element, $Z = 2\text{--}120$: the entrant channel, its depth, its margin, and the full candidate spectrum of the $V^{\{N-1\}}$ walk. **119 rows; 107 scored, 12 unwitnessed.**

The quantity. Per row: D_{ent} and the margin, in hartree; per candidate channel, its converged depth in the frozen field of $(Z, \text{cfg}(Z-1))$.

The transition. The entrant operator: converge the field of the ion carrying $\text{cfg}(Z-1)$; solve each frontier channel; deepest wins; chain forward. Nothing in the rule is adjustable.

Constants required. One: $c = 137.035999$ (Koelling–Harmon scalar-relativistic kernel). *No screening constant, no fitted parameter, no observed energy.*

Must be measured. Nothing upstream. The measured ground configurations enter once, downstream, as the score target: **107 of 107** (register 1701–1702). Rows 109–120 have no measurement to meet and are held **unwitnessed** — the book's own grade: not undefected for want of trying.

What physics does. It supplies the field and the depths; the ordering law is then read off, with its three derived exceptions (collapse) and its derived absence (no g block). **What it does NOT do:** it does not decide total-energy rearrangements inside an already-open block — the chromium class is outside this index's claim, and stating that is scope, not weakness (register 1701).

Λ_{cinf} — the twin index at $c \rightarrow \infty$

What it indexes. The identical walk with the one constant removed. 107 rows.

The quantity. The same depths and entrants, non-relativistically.

The transition. Identical operator, $c \rightarrow \infty$.

Constants required. None.

Must be measured. Nothing; it is compared, not scored: **eleven entrants differ from Λ_{chain}** (Mn, Zn, Ag, Cd, Nd, Pm, Sm, Lu, Hg, Lr, Rf), and every difference is an error against nature, since the relativistic walk scores 107/107 (register 1706).

What physics does. It states that the observed periodic table is not a solution of the non-relativistic equation.

What it does NOT do: serve as a fallback ordering anywhere; its role is the counterfactual, and it must never be consulted as data.

Λ_V5 — the correlation closure at the contested rows

What it indexes. The five elements where the mean-field competition is close: $Z = 38, 56, 72, 89, 105$. Per row: the second-order correlation differential between entrant and runner-up, complete (entrant pairs, core–core pairs, the near-degenerate block resummed), expressed over the mean-field margin.

The quantity. $\text{dm}2/\text{margin} = +1.33, +1.24, +2.6 [2.57\text{--}2.73], +2.2 [1.92\text{--}3.32], +2.16$.

The transition. Second-order many-body perturbation on the sealed field; brackets are declared estimate envelopes, points where single-valued.

Constants required. The same single c , inherited.

Must be measured. Nothing. The clause is a computation whose only external contact is the field it inherits.

What physics does. It closes the one silent failure mode of a mean-field derivation: **every contested competition widens; none reverses; an envelope entirely on the positive side cannot flip a positive sign** (register 1705).

What it does NOT do: bound the uncontested rows — there the mean-field margin needs no help, and claiming otherwise would be a pooled assertion of exactly the kind the domain protocol blocks.

The corridor law — interface only

Chapter 34's law, as physics. The mechanism — the corridor as a system of linear inequalities, and the hull form beneath it — is stated once, in the Mathematical Compendium addition; this entry records what the law claims about nature and where it could have failed.

derived here · from this work · valid $Z = 3\text{--}108$ · four objects state it; the transitive count is not computable at this build (register 1814)

What it states. The subshell the differentiating electron enters is always one the v form can select at all: a vertex of the lower convex hull of the points $(\sqrt{r}, n)$, $r = p + q/2(2\ell+1)$, over the Pauli-admissible subshells — equivalently, its corridor of admissible slopes a is non-empty. It holds at all 106 steps, $Z = 3$ to 108, in the node-only form and the finished form alike, and it could have failed at any of them (§34.5; registers 1445, 1460, 1463).

The transition. From the observed ground configurations (NIST ASD 5.12, register 1306) to a corridor per step: the two hull-edge slopes flanking the entrant. The identity of the corridor with the hull-edge slopes is asserted at 0 mismatches over 106 steps, in both forms.

Must be measured. Every step. The law is a statement about which configurations nature realises, so it is *verified* and never *proved*; the theorem beside it — the necessity of state — is the mathematics and needs no observation, and the law is what observation adds.

What physics does. It supplies the entrant; the law says the entrant is never outside the form's reach. **What it does NOT do:** select the entrant — every step admits two to six hull vertices and one carried number decides among them (§34.6); and non-emptiness refutes nothing: an empty corridor refutes the form and a non-empty one does not confirm it (register 1460). **Where it fails:** not known past $Z = 108$, where NIST lists no neutral ground configuration.

The collapse condition — interface only

Per this compendium's charter, the mechanism is stated once, in the Mathematical Compendium addition; a third copy would drift. This entry records the interface.

What it decides for these indexes. Exactly the tie-break exception set {La, Ac, Th} in Λ_{chain} ; and the $Z = 91$ sign held at SCF tolerance, entered as branch content rather than resolved by force (register 1703). **It occupies precisely the domain where Chapter 34’s corridor is silent.**

Must be measured. Nothing; the condition is computed from the same field as the walk.

What it does NOT do. Perturb Clause 1 anywhere: the ordering clause is exception-free with the collapse rows included.

The defect — interface only

Mechanism and decomposition are stated once, in the Mathematical Compendium addition (“The defect, closed”); this entry records the interface.

What it touches in these indexes. Nothing in any scored value: the defect is internal to the walk’s energy bookkeeping, and its closure (register 1707–1711) is the audit that no unexplained numerical content sits under the indexes above.

Must be measured. Nothing — and that is the entry’s point: a discrepancy that data cannot violate was defending the derivation, and once named it became mathematics. **The fault ledger’s two reclassifications live here.**


Λ_3 — THE THREE-BODY INDEX

Chapter 36; register 1713–1724. Objects in the Mathematical Compendium, family 3B.

What it indexes. Motions of three point masses under Newtonian gravity, at fixed E and L , by asymptotic class: five cells. It is the second index in the compendium built from physics rather than from Λ (the first is Λ_{spectra}), and the first whose subject is classical.

The coordinate supply. The masses enter through exactly six numbers — three c_{ij} and three unit vectors b_{ij} — and nowhere else. Everything structural (S^2 , the metric, N_8 , K_3) is mass-free.


Fig. 1 The shape sphere: 3 Euler (E), 2 Lagrange (L), 3 binary-collision rays (B). Masses move the points; they never change the count.



S^2 with the Euler, Lagrange and binary points; two mass cases.

Kinematic and stateful. Λ_3 's cells are configurations; by §12.11.1.3 the index has no time column, and the flow is the geodesic flow of the Jacobi–Maupertuis metric. Time is a quadrature and carries the transcendental part (Sundman series; Painlevé transcendents on the natural boundary). This is the interface *Kinematic and stateful* (above) found in the register's own objects, now on a classical subject: *placing* a cell (which stratum) and *valuing* it (the geodesic) are different operations.

Where the physics is exact and where it is envelope. Exact: the five fixed points, the zero-velocity surface $\{E + U \geq 0\}$ (Hill 1878), Hill stability of a hierarchy (Marchal–Bozis 1982). Envelope: every stratum boundary, by the triangle form on K_3 (§36).



U on the collinear circle, three mass cases.

The point of observability. What Λ_3 delivers for a given state: the stratum, the family within it (torus / braid word / symbolic sequence / distribution $P(\epsilon)$), and the geodesic as far as the natural boundary. What it never delivers: $r_i(t)$. §25.6 is why.

Failure modes, measured rather than anticipated. 1. A factor of the hyper-radius carried into the potential on shape space (§11) from an inherited convention: 13/13 failures, uniform, hence in the audit not the object. Corrected. 2. An inherited degree-8 polynomial with two wrong coefficients: caught by the second route (N_8). Withdrawn. 3. Threshold silence: L4/L5 stability at $\mu < 0.0385209$ is a number; the method is silent (§31.1.1). Not a failure, a boundary.

Λ_{phys} — THE INDEX OF PHYSICAL PARAMETERS

Every number the work takes from physics, with what it is, where it comes from, and where it stops being right. A parameter with no stated failure mode is a parameter being used outside a domain nobody has checked.

27 parameters. Coordinates: **kind** (exact $\rightarrow$ fitted), **source** (mathematics $\rightarrow$ this work), **domain** (universal $\rightarrow$ one species, and the three-body problem’s own range), and how many registered objects rest on it.

What the index shows about the work’s dependence

kind	universal	all elements	a region	one species	the three-body problem
exact by definition	2	0	0	0	0
measured constant	3	0	0	1	0
read from a table	0	3	2	1	0
derived here	0	2	1	0	0
fitted here	0	1	6	1	0
free parameter (three-body: $m_1 m_2 m_3$, E, L)	0	0	0	0	3
exact by scaling (G)	1	0	0	0	0

The speed of light c (measured constant, universal) joins the second row with Chapter 35; the four three-body inputs are the problem’s, not this work’s, and are entered as the last two rows.

“Objects rest on it” is counted from the Mathematical Compendium at this build, on one rule for all twenty-seven (register 1740): the seeds are the objects whose statement or title names the parameter — its symbol, its value or its name — and the count is the seeds plus every object that depends on one of them, transitively, in the compendium’s own dependency graph. Seeds and counts: subshell capacity: Pauli, $k \leq 4\ell + 2$ (§7.1), the generating function (§11.3), the membership function (§11.1), the first factor (§12.7), the second factor (§12.7), parastatistics $\rightarrow$ 66; angular momentum bound: node counting, $\ell \leq n - 1$ (§7.1), the generating function, the membership function, the first factor $\rightarrow$ 72; Rydberg constant: the five-sigma admissibility threshold — where the bracket may be applied at all (§22.5), when the bracket fails (§23.3), what silence implies (§23.5), the value-one crossing — where the bracket’s price V passes unity (§23.15) $\rightarrow$ 5; proton-electron mass ratio: none $\rightarrow$ 0; Cs I np defect: none $\rightarrow$ 0; ionisation limit: the refusal map (§6.2), constructed limit (register 711), self-determined limit (register 684), the precision lens (register 749) $\rightarrow$ 4; aufbau ordering: the Pauli bound (register 1141), the ordering law, five clauses (§35) $\rightarrow$ 18; Janet block boundary: orbital collapse (register 719), the Janet collapse (register 1187), the collapse condition (§35) $\rightarrow$ 13; Hund’s first rule: none $\rightarrow$ 0; core dipole polarisability: the polarisation formula, core polarisation (register 731), the polarisability index (register 1165), Seaton’s term (register 1165) $\rightarrow$ 18; actinide defects: none $\rightarrow$ 0; Seaton polarisation constant: the polarisation formula, Seaton’s term $\rightarrow$ 16; reduced-mass Rydberg: none $\rightarrow$ 0; bracket yardstick: the five-sigma admissibility threshold — where the bracket may be applied at all, when the bracket fails, what silence implies $\rightarrow$ 4; channel equation, a: the channel equation, final form (register 1205) $\rightarrow$ 9; channel equation, e0: the channel equation, final form $\rightarrow$ 9; channel equation, e1: the channel equation, final form $\rightarrow$ 9; channel equation, k: the channel equation, final form $\rightarrow$ 9; channel equation, h: the channel equation, final form $\rightarrow$ 9; isoelectronic ladder, B: the isoelectronic ladder (register 708), the channel equation, final form $\rightarrow$ 11; five-sigma admissibility: the five-sigma admissibility threshold — where the bracket may be applied at all $\rightarrow$ 2; exchange coefficient: the exchange factor (register

987), the channel equation, first form (register 1145) $\rightarrow 15$; speed of light: the entrant operator (§35), the ordering law, five clauses, the twin operator, $c \rightarrow \infty$ (§35) $\rightarrow 8$; three-body masses: the potential on shape space (§11), the algebraic variety, the five fixed points (§D.5.9) $\rightarrow 5$; three-body energy: the Jacobi–Maupertuis metric, the index Λ_3 (§36) $\rightarrow 2$; three-body angular momentum: the index $\Lambda_3 \rightarrow 1$; gravitational constant: none $\rightarrow 0$. A parameter no statement names — the proton–electron mass ratio, the caesium defect, the actinide defects, the reduced-mass Rydberg, Hund’s first rule — counts 0 on this rule: it enters through a value the objects use, not through a statement, and the rule says so rather than guessing. Prior counts, generated on a rule this volume did not print and standing to register 1739 where they differ: subshell capacity 12, angular momentum bound 12, Rydberg constant 9, proton-electron mass ratio 9, Cs I np defect 2, ionisation limit 25, aufbau ordering 19, Janet block boundary 15, Hund’s first rule 12, core dipole polarisability 10, actinide defects 2, Seaton polarisation constant 10, reduced-mass Rydberg 9, channel equation, a 8, channel equation, e0 8, channel equation, e1 8, channel equation, k 8, channel equation, h 8, isoelectronic ladder, B 3, five-sigma admissibility 3, exchange coefficient 1.

8 of the 27 parameters are this work’s own. Of those, **0 are universal**, **1 hold across all elements**, and **7 hold only in a region or for one species**. No number this work fitted claims wide validity.

And the dependence concentrates on the LATTICE’S OWN BOUNDS, then on tables. On the printed rule the two exact-by-definition constraints carry the most objects, because the index of Part III is built from them; the tables come next (register 1740; before it the three heaviest were the ionisation limit, the aufbau ordering and the Janet boundary — the prior counts stand in the paragraph above):

parameter	objects resting on it	kind
angular momentum bound	72	exact by definition
subshell capacity	66	exact by definition
aufbau ordering	18	read from a table
core dipole polarisability	18	read from a table
Seaton polarisation constant	16	derived here
exchange coefficient	15	fitted here

The parameters

subshell capacity — $4\ell+2$, an exact integer

exact by definition · from published literature · valid universal · 66 objects rest on it

What it is. the number of electrons a subshell of angular momentum ℓ can hold: 2 spin states times $2\ell+1$ orbital states.

Where it comes from. Stoner, The distribution of electrons among atomic levels, Phil. Mag. 48 (1924) 719-736; made exclusive by Pauli, Z. Phys. 31 (1925) 765-783

Where it fails. for parastatistics of order m the capacity is $m(4\ell+2)$ — parastatistics shows the index still closes, so the constraint is a cap and not a magic number.

angular momentum bound — $\ell \leq n-1$, an exact integer

exact by definition · from published literature · valid universal · 72 objects rest on it

What it is. the orbital angular momentum of a bound state cannot reach the principal quantum number; it is the condition for the radial function to have a node structure.

Where it comes from. Bohr, Phil. Mag. 26 (1913) 1-25; Schrödinger, Ann. Phys. 79 (1926) 361-376

| **Where it fails.** never within non-relativistic quantum mechanics.

Rydberg constant — $R_{\infty} = 109737.31568 \text{ cm}^{-1}$

measured constant · from international standard · valid universal · 5 objects rest on it

What it is. the binding energy of a hypothetical one-electron atom with an infinitely heavy nucleus, expressed as a wavenumber. It sets the scale of every atomic term value.

Where it comes from. Rydberg, K. Sven. Vetensk. Akad. Handl. 23 (1890) for the empirical constant; value CODATA 2018, uncertainty $1.9\text{e-}6 \text{ cm}^{-1}$ — eleven significant figures

| **Where it fails.** never as a constant; but using it in place of the reduced-mass R_M is wrong for every species (register 868). The error is 1 part in 1836 A, largest at hydrogen.

proton-electron mass ratio — $m_p/m_e = 1836.15267343$

measured constant · from international standard · valid universal · 0 objects rest on it

What it is. the ratio of the proton rest mass to the electron rest mass; the quantity that makes the reduced-mass correction small but not negligible.

Where it comes from. CODATA 2018, uncertainty $1.1\text{e-}10$

| **Where it fails.** never; it enters only through R_M , so an error here is an error there.

Cs I np defect — 3.5667

measured constant · from published literature · valid one species · 0 objects rest on it

What it is. the limiting quantum defect of the caesium np series, the compendium's only MEASURED far anchor.

Where it comes from. arXiv:1706.06237 (2017), $n = 70\text{-}100$; earlier 3.55925 at $n = 9\text{-}50$ (Lorenzen & Niemax, Z. Phys. A 315, 1984)

| **Where it fails.** the two values differ by 0.007 because of stray-field shifts; the compendium uses the field-free one.

ionisation limit — I , per species, cm^{-1}

read from a table · from published literature · valid one species · 4 objects rest on it

What it is. the energy at which a Rydberg series converges — the term value of the ion's ground state relative to the neutral. Every defect is measured against it.

Where it comes from. the extrapolation method is Rydberg 1890 and Ritz, Ann. Phys. 12 (1903) 264-310; values from NIST ASD where published, built by summing two spectra where not (constructed limit (register 711)), joint-fit where neither

| **Where it fails.** wherever it was CONSTRUCTED rather than published — such a channel is not a measurement against an independent standard, and the compendium marks the column.

aufbau ordering — $n+1$, then n , a permutation of subshells

read from a table · from published literature · valid all elements · 18 objects rest on it

What it is. the order in which subshells fill as Z increases: lowest $n+l$ first, and within one $n+l$ value, lowest n first.

Where it comes from. Janet 1928 (Considerations sur la structure du noyau de l'atome, Beauvais 1929), who recognised it before Madelung 1936; shell lengths by the Klechkovski-Hakala formulas

Where it fails. at about twenty known exceptions among the elements — Cr, Cu, Nb, Mo, Ru, Rh, Pd, Ag, La, Ce, Gd, Pt, Au, Ac, Th, Pa, U, Np, Cm, Lr. The compendium reads the OBSERVED configuration, so the exceptions are data and not failures. The rule's ORIGIN was unresolved when this entry was written (Löwdin's challenge, Allen & Knight, Int. J. Quantum Chem. 90 (2003) 80-88); it is now derived — Λ_{chain} above, Chapter 35, register 1701 — and the three tie-break exceptions La, Ac, Th are derived with it. The remaining exceptions of the Cr class are total-energy rearrangements inside an open block, outside Λ_{chain} 's claim.

Janet block boundary — $Z = 21, 57, 89$, an exact integer

read from a table · from published literature · valid all elements · 13 objects rest on it

What it is. the atomic number at which a new $n+l$ block opens — $n+l = 5$ at Sc, 7 at La, 8 at Ac — which is where the corresponding orbital contracts into the core.

Where it comes from. the block structure is Janet 1928; the CONTRACTION is Goeppert-Mayer, Phys. Rev. 60 (1941) 184-187, and Griffin, Andrew & Cowan, Phys. Rev. 177 (1969) 62-71

Where it fails. as a SHARP threshold: collapse is rapid, not instantaneous. Ca I at $Z = 20$ has $nd = 0.908$ against a threshold of 21, and Ba II at 56 has $nf = 0.756$ against 57.

Hund's first rule — max S for the core, a term selection

read from a table · from published literature · valid all elements · 0 objects rest on it

What it is. of the terms a configuration allows, the one of highest total spin lies lowest in energy; it fixes which multiplicities a core can present.

Where it comes from. Hund, Z. Phys. 33 (1925) 345-371

Where it fails. for a core in an excited term rather than its ground term. The index takes the ground term, so an excited-core series is outside it.

core dipole polarisability — α_d , per core, a_0^3

read from a table · from published literature · valid a region of the index · 18 objects rest on it

What it is. the induced dipole moment of the ionic core per unit applied field; the leading term in the potential a distant electron feels beyond the Coulomb tail.

Where it comes from. Born & Heisenberg, Z. Phys. 23 (1924) 388-410; systematic values Mayer & Mayer, Phys. Rev. 43 (1933) 605-611; modern e.g. Cs+ 15.696(16) a_0^3 , arXiv:2502.20961

Where it fails. below $l = 4$, where penetration dominates and Seaton's formula does not apply. AND for nf treated as non-penetrating: arXiv:2502.20961 reports α_d and α_q that then disagree with the ng energies.

actinide defects — 5.2, 4.75, 3.8, 2.0, theoretical

read from a table · from published literature · valid a region of the index · 0 objects rest on it

What it is. the asymptotic ns, np, nd and nf quantum defects predicted for the actinides, $Z = 89-103$; the compendium's only high- Z anchors.

Where it comes from. arXiv:2508.06733 (2025), theoretical

Where it fails. as MEASUREMENTS — they are calculated. Four of the seven far anchors are these, and the equation's high- Z arm moves if they are revised (register 1202).

Seaton polarisation constant — $3 \alpha c^2 / K(l)$

derived here · from published literature · valid a region of the index · 16 objects rest on it

What it is. the limiting quantum defect of a non-penetrating series, set by the core's polarisability and the centrifugal factor $K(l) = l(l+1)(2l-1)(2l+1)(2l+3)$.

Where it comes from. Seaton, MNRAS 118 (1958) 504-518; Drake & Swainson, Phys. Rev. A 44 (1991) 5448

Where it fails. at $l < 4$, and wherever the quadrupole term matters. The compendium measures the ratio δ_2/δ_0 against $-l(l+1)/3$ at 1.25 rather than 1.00.

reduced-mass Rydberg — $R_M = R_{\text{inf}} / (1 + 1/(A m_p))$

derived here · from international standard · valid all elements · 0 objects rest on it

What it is. the Rydberg constant corrected for a nucleus of finite mass A , by replacing the electron mass with the electron-nucleus reduced mass.

Where it comes from. Bohr 1913: after Fowler objected that the Pickering series did not fit, Bohr replaced the electron mass with the reduced mass and obtained five-digit agreement, which identified the series as ionised helium. Bohr, Nature 95 (1915) 6-7 for his later comment.

Where it fails. for an ion of unknown isotopic composition: A is the dominant isotope, and a mixed sample shifts R_M by roughly the isotope shift.

bracket yardstick — $2 Z^2 R / \nu^3$

derived here · from published literature · valid all elements · 4 objects rest on it

What it is. the local spacing between adjacent Rydberg levels — the derivative of the term formula. A perturbation larger than half of it reorders the levels.

Where it comes from. derived from Rydberg's term formula (1890)

Where it fails. for a PERTURBED series, where a level of another channel crosses in. Fano, Phys. Rev. 124 (1961) 1866-1878; Lu & Fano, Phys. Rev. A 2 (1970) 81-86.

channel equation, a — 0.3772

fitted here · from this work · valid a region of the index · 9 objects rest on it

What it is. the overall amplitude of the penetrating branch: how much defect one core orbital of the same l produces at unit electron count and unit charge.

Where it comes from. fitted here by least squares on 277 in-region channels plus 7 far anchors; the FORM it scales is the penetration picture of Hartree, Proc. Camb. Phil. Soc. 24 (1928) 89-110

Where it fails. outside $Z \leq 92$, charge ≤ 10 , $l \leq 4$, single-parent core.

channel equation, e_0 — 0.8297

fitted here · from this work · valid a region of the index · 9 objects rest on it

What it is. the intercept of the exponent on the core-orbital count p — how sharply the defect grows with the number of same- l orbitals in the core, at small Ne .

Where it comes from. fitted here; the p -count itself is the Pauli occupancy (Pauli 1925)

Where it fails. as above. Its value moved from 0.583 to 0.830 when the far anchors were added, so it is sensitive to the high- Z end.

channel equation, e_1 — -0.0900

fitted here · from this work · valid a region of the index · 9 objects rest on it

What it is. the log- Ne slope of that exponent: the rate at which additional core orbitals stop mattering as the atom grows.

Where it comes from. fitted here; that screening saturates with electron count is the Thomas-Fermi picture of Fermi, Z. Phys. 48 (1928) 73-79

Where it fails. as above; it was -0.027 before the anchors, a factor of three.

channel equation, k — 0.4942

fitted here · from this work · valid a region of the index · 9 objects rest on it

What it is. the Thomas-Fermi exponent: how the defect scales with the number of electrons in the core, through the screening the statistical model predicts.

Where it comes from. fitted here; the scaling is Fermi, Z. Phys. 48 (1928) 73-79, who applied the statistical model to Rydberg corrections directly

Where it fails. MEASURABLY: k rises with charge, from 0.84 at charge 1 to 1.52 at charge 4 (register 1214). A single value is a simplification and the register says so.

channel equation, h — 0.5415

fitted here · from this work · valid a region of the index · 9 objects rest on it

What it is. the amplitude of the collapse branch: how much defect a collapsed orbital produces where the core has no orbitals of its own l .

Where it comes from. fitted here; the collapse is Goeppert-Mayer 1941 and Griffin, Andrew & Cowan 1969

Where it fails. below the Janet boundary, where the polarisation floor is absent.

isoelectronic ladder, B — per sequence

fitted here · from this work · valid a region of the index · 11 objects rest on it

What it is. the single free coefficient of $\delta(c) = B \ln(c+1)/c$ along a sequence of fixed electron count, with the constant term fixed at zero by the hydrogenic edge.

Where it comes from. fitted here per sequence on the 52 with three or more charge states; the isoelectronic method is Edlen, Handbuch der Physik XXVII (1964) 80-220

Where it fails. as a CLOSED FORM: no expression of B in the coordinates beats the per-sequence fit by less than a factor of twenty, and PMC3671562 reports the same for ionisation energies.

five-sigma admissibility — $r \geq 5$, a threshold

fitted here · from this work · valid all elements · 2 objects rest on it

What it is. the bracket is applied only where the signal exceeds five times the measurement uncertainty, so that a bracket failure is a fact about the levels and not noise.

Where it comes from. the convention is particle physics's; its history and its critics are in Lyons, Discovering the significance of 5 sigma, arXiv:1310.1284 (2013)

Where it fails. it is a CHOICE. At three sigma more cells enter and more brackets fail; the compendium states the threshold rather than tuning it.

exchange coefficient — $s = -0.0782$, WITHDRAWN

fitted here · from this work · valid one species · 15 objects rest on it

What it is. the fractional difference between a triplet channel's defect and its singlet partner's, from the exchange interaction between the Rydberg electron and the core.

Where it comes from. fitted at register 987, WITHDRAWN at 1168; the physics is Heisenberg, Z. Phys. 38 (1926) 411-426, with Hund's first rule (1925)

Where it fails. everywhere: the fitted sign is opposite to the measured one, and a uniform s gives 0 of 66 pairs or 66 of 66.

The fifteen, read from the graph (register 1749): six were built on the superseded channel equation that carried s — the exchange factor (register 987), the channel equation, first form (register 1145), the bridge equation (register 1169), the validated domain (register 1193), the two-ended anchor (register 1201), the three blindnesses (register 1169) — and nine are the standing equation the channel equation, final form (register 1205) and the Löwdin objects beneath it, which reach s only by provenance: the channel equation, final form descends from the channel equation, first form through the validated domain and the two-ended anchor, and its statement carries no exchange factor. The count stands on the printed rule; the split says what it measures.

speed of light — $c = 137.035999$ a.u.

measured constant · from international standard · valid universal · 8 objects rest on it

What it is. the one entered number of the Löwdin solution: the constant in the Koelling–Harmon scalar-relativistic kernel of the entrant operator. No screening constant, no fitted parameter, no observed energy enters beside it.

Where it comes from. CODATA; the inverse fine-structure constant in Hartree atomic units.

Where it fails. at $c \rightarrow \infty$ — and the failure is measured, not anticipated: the twin index Λ_{cinf} disagrees with Λ_{chain} at eleven elements, each an error against nature (register 1706).

three-body masses — m_1, m_2, m_3 , three positive reals

free parameter · from the problem statement · valid all mass triples · 5 objects rest on it

What it is. the masses of the three bodies, unspecified; they enter Λ_3 through exactly six numbers — three c_{ij} and three unit vectors b_{ij} — and nowhere else.

Where it comes from. the problem as posed; the six-number coordinate supply is Montgomery 2014 §11.

Where it fails. nowhere structural — S^2 , the metric, N_8 and K_3 are mass-free. The masses move the five fixed points and nothing else.

three-body energy — E , a real

free parameter · from the problem statement · valid $E < 0$ for bound strata · 2 objects rest on it

What it is. the conserved energy, entering as the conformal factor $E + U$ of the Jacobi–Maupertuis metric.

Where it comes from. Maupertuis 1744; Jacobi 1837.

Where it fails. at the zero-velocity surface $\{E + U = 0\}$ (Hill 1878), where the metric degenerates and the geodesic flow stops.

three-body angular momentum — L , a real

free parameter · from the problem statement · valid all L · 1 object rests on it

What it is. the conserved angular momentum, carrying the reduction $6 \rightarrow 4$ of the phase space.

Where it comes from. standard; the reduction is Jacobi 1842.

Where it fails. at the angular-momentum stratum boundary, which is envelope, not exact (the triangle form on K_3 (§36)).

gravitational constant — $G = 1$

exact by scaling · from convention · valid universal · 0 objects rest on it

What it is. the coupling of Newton’s potential, set to 1 by choice of units.

Where it comes from. scaling; carries no content once units are fixed.

Where it fails. nowhere.

The three failures that are measured rather than anticipated

Most of the failure modes above are domain statements: the parameter is right inside its range and untested outside. Three are different — the compendium has MEASURED the failure.

Seaton’s polarisation constant. The ratio δ_2/δ_0 should be $-\ell(\ell+1)/3$ with the polarisability cancelling. On fourteen channels with $\delta_0 > 0.01$ the observed ratio is **1.25 times** the predicted one, and the departure is a per-core constant reproducible within a species to two decimals — Si III g gives 1.77 four times, Si III h gives 1.12 twice.

The Thomas–Fermi exponent k . Fitted per charge state, it reads **0.84, 0.91, 1.03, 1.40** at s and **1.01, 1.13, 1.28, 1.52** at p, for charges 1 to 4, with trend R^2 between 0.87 and 0.98. **The equation carries a single $k = 0.494$.** Register 1214.

The exchange coefficient. The fitted sign is opposite to the measured one, and no uniform coefficient can work: exchange acts through an overlap that falls sharply with ℓ , so one constant gives 0 of 66 triplet-above-singlet pairs or 66 of 66. Withdrawn at register 1168.

These three are why the compendium states failure modes at all. A parameter whose failure has been measured is more useful than one whose validity has been assumed, and the difference is not visible unless both are written down.

Λ_{chem} — THE CHEMICAL PROPERTIES INDEX

Registers 1339, 1343–1346. Forty-two chemical properties of a species, indexed on what each one IS and on which of physics, charge or amplitude it supplies.

Why the compendium needed it

Repeatedly in the Löwdin work a residual was left unexplained because a universal account of it was sought. **Each time the answer was a property of the individual species.** The record: the 1.029 factor on $t(\ell)$; the occupancy slope varying from 0.021 to 0.243; no f corridor, hence $\sqrt{6}$ untested; the crossing charge set by n_f ; Seaton's ratio valid only at $p = 0$.

Λ_{phys} had been saying this all along — no parameter of this work sits in the universal column. Λ_{chem} is the routing table that makes the statement usable: before calling a residual unexplained, look up which chemical property supplies it and which class it holds on.

The coordinates

KIND — what the property measures: *count · symmetry · size · energy · rate*

SEAT — which part of the species it belongs to: *the nucleus · the core · the subvalence shell · the valence shell · the aggregate*

PCA — which of *physics · charge · amplitude* it supplies

The fifth seat

Twelve of the forty-two are properties of **MATTER IN BULK** and not of an isolated atom at all: density, melting point, boiling point, hardness, crystal structure, electrical and thermal conductivity, colour of the metal, smell, taste, metallic character, reactivity.

With eleven hand-picked properties the index could not say this. It appeared the moment every chemical property of a species was demanded, and it is the reason the compendium can now distinguish an atomic property from a bulk one at all.

The closure, and the boundary it declares

E = 0 on fourteen cells over the subvalence and valence shells. And the defect climbs as foreign seats are added:

seats included	cells	E
subvalence + valence	14	0
+ the core	15	3
+ the nucleus	20	8
+ the aggregate	21	11
all five	26	19

PCA's domain is the subvalence and valence shells, and the index declares it by degrading monotonically outside it. *That is not a failure to close. It is a boundary stated — as Seaton's ratio is valid at $p = 0$ and undefined beyond.*

The last cell, and what filled it

All 1,440 orderings of the three axes give minimum $E = 1$, so the defect is structural rather than a labelling artefact. **The cell is symmetry $\times$ the valence shell $\times$ a charge role.**

The fill is the ground TERM, which changes with charge:

N_e	$c=1$	$c=2$	$c=3$	$c=4$
20	1S_0	3D_1	3F_2	3F_2
38	1S_0	3D_1	3F_2	3F_2
56	1S_0	3F_2	3H_4	—
88	1S_0	1S_0	$^3H_4^\circ$	—

The sequence $^1S_0 \rightarrow ^3D_1 \rightarrow ^3F_2$ is identical at two electron counts. Every ladder table carried the term symbol; the configurations were recorded and the terms were not.

The routing, by residual

residual	property	class
the 1.029 factor on $t(\ell)$	subshell radius	one subshell
no f corridor	centrifugal barrier	d and f only
the occupancy slope	subshell radius	one subshell
the crossing charge 2, 3, 5	closed f shell n_f	period 6, 7

Both of the first two were routed and both dissolved. *The f corridor does not exist because $p = 0$ is the node floor, so no rival lies below and $L = -\infty$ — the missing test is forbidden by the node count, not by missing data. And the 1.029 was arithmetic: the excess above U is 0.028 of the spread at p and 0.785 at d , a ratio of 27.7, and the apparent 2.9% agreement came from comparing a ratio at p , where $t \approx 1$, with a ratio at d , where $t \approx 1.78$.*

Λ_{PCA} — PHYSICS $\oplus$ CHARGE $\oplus$ AMPLITUDE, AS ONE

Registers 1297, 1323–1324, 1345. The merged index that correlates every chemical property to the physics that requires it.

The merge

Λ_{phys} closes on (source, domain). Λ_{charge} closes on (role, carrier, sign, regime). The shared axis is domain $\equiv$ regime — both say where a statement holds, and the charge regime is the physics domain at finer resolution.

Merged on that axis alone: $E = 0$ on nine cells.

	universal	all elements	low	neutral	hydrogenic	one species
standard	5	.	.	.	.	.
mathematics	3	1	.	.	.	.
literature	.	7	1	.	.	.
this work	.	2	5	2	4	.

A staircase, and the diagonal is the whole content: nothing standard is restricted to a region, and nothing this work produced claims universal validity.

What it gives that neither parent does

The per-atom calibration. A parameter's domain now reads as a SET OF ATOMS — *universal* is every atom and ion, *all elements* every neutral, *neutral* $c = 1$, *low* $c = 2$, *hydrogenic* $c \geq 3$, *one species* a single (Z, c) . For any given species the index states which parameters apply to it.

And the closing order puts **low** before **neutral** — charge 2 is the most particular domain. The isoelectronic ladders confirm it independently: $c = 2$ is the only charge with two-sided brackets.

Λ_{amp} — the amplitude's own index

The amplitude is the electron–electron term, and the multipole expansion of $1/r_{ij}$ gives, for each subshell pair, an exact finite list: F^k for k even up to $2\min(\ell, \ell')$, G^k for $k \equiv \ell + \ell' \pmod{2}$ from $|\ell - \ell'|$ to $\ell + \ell'$.

Twenty cells, $E = 0$, a triangle $0 \leq i \leq \ell$ — once indexed on position within sequence rather than multipole rank. Indexed on the raw rank the single defect is F^1 , a term parity forbids.

And it explains the Slater result exactly. At s/s the exchange G^0 IS the direct F^0 , so dropping exchange costs nothing; at f/f it discards four independent quantities. The count of lost integrals is the count of failures: Slater screening threads 5 of 5 s-block brackets and 0 of 1 f.

The angular factor of the sole exchange integral for an s electron against an ℓ electron is exactly $1/(2\ell+1)$ — 1, 1/3, 1/5, 1/7 at s, p, d, f, to machine precision. The reciprocal of the subshell's orbital degeneracy, a count Λ already holds, and it agrees independently with $h(\ell) = h_0\sqrt{(2\ell+1)}$ measured from the collapse data.

A CORRECTION TO SEATON'S RATIO

Previously recorded: the ratio δ_2/δ_0 comes out at 1.25 where the dipole term gives 1.00, and the departure is a per-core constant reproducible to two decimals.

That measurement was made on penetrating channels, where the relation is undefined. Seaton's derivation assumes the outer electron is non-penetrating, so the core polarisability is the whole interaction.

Stated properly:

	series	ratio to Seaton	sd
$p = 0$	3	1.150	0.206
$p \geq 1$	10	-0.015	0.177

On the non-penetrating branch the relation holds to 15%. On the penetrating branch the ratio is zero — the relation carries no information at all.

Not “broken at d and sound at f ”. Valid at $p = 0$ and undefined at $p \geq 1$, which is a domain statement and exactly what Λ_{phys} 's domain axis is for.

And the sign of δ_2 is penetration, not ℓ : 0 of 3 positive at $p = 0$, 2 of 2 at $p = 5$, monotone in p between.

THE CHANNEL CONSTANTS IN Λ_{phys}

Owed since register 1296; written. The failure modes are the point — a constant listed without the conditions under which it stops meaning anything is a number pretending to be a law.

standing	count	what it means
attributed / derived	5	traceable to a published result or a proof
measured	6	read from data, with a stated spread
fitted	3	set to make the form agree; carries no physics

The three fitted constants are the amplitude law's, and they are why nothing in the *amplitude* work answers Löwdin's challenge. Written; the challenge was closed at register 1701 by a different instrument, Λ_{chain} , which carries no fitted constant. The sentence is kept because it was true of what it described.

Failure modes

constant	fails when	how it shows
the amplitude a	across a language boundary	six placement rules dead; a seventh prohibited
$\beta = 2/3$	-	no failure mode found; attributed, arity three
the gate $\ell(\ell+1)$	never as a gate	but SIX appearances under six discipline names
$u_0 = 4$	outside the collapse region	the switch is local to the d-wave entry
the nu form	as a MEASURE	bounded delta against diverging $a\sqrt{p}$ — impossible
the three fitted	held out	the walk scores 90 against Madelung's 96

The last row is the one to read. A constant fitted on the data it explains scores 99; held out it scores below the rule it was built to improve on. A failure mode is not a caveat — it is the boundary of the domain, and Λ_{phys} exists to carry boundaries.

THE PRIOR-ART CHAIN — the seed's forced structure

Every piece of machinery in the seed results is attributed; no novelty is claimed for any of it. The application — measuring the forced content of an index's minimum seeds across caps, and locating its non-uniqueness as a Krein–Milman failure — is the contribution.

- **Moore (1910)** — closure operators; the root of $\mathcal{R}$ itself.
- **Baker & Pixley, Math. Z. 143 (1975) 165–174** — binary determination (2,d-interpolation): why pairwise envelopes decide membership, the generation criterion's licence.
- **Deville, Barták & Van Hentenryck (1999)** — the staircase/ (α, β) -monotone constraint class; $\mathcal{R}$ is its $(\leq, \leq)$ corner.
- **Karp (1972); Chvátal (1979)** — set cover, and the reduction rule: a set present in every minimum cover is forced by an element it uniquely covers. Carries the forced-set half of the non-uniqueness result.
- **Shannon, Trans. AIEE 57 (1938) 713–723** — reading structure as binary words; retained for the bit-encoding itself.
- **Edelman, Alg. Univ. 10 (1980) 290–299; Edelman & Jamison, Geom. Dedicata 19 (1985) 247–270** — anti-exchange closures / convex geometries and the combinatorial Krein–Milman theorem. *The load-bear-*

ing new citation: Λ_8 's empty extreme-point set with seed 7 is a maximal failure of the Krein–Milman property, which is what permits 24,585 minimum generating sets.

- **Dilworth (1940)** — lattices with unique irredundant decompositions: uniqueness under conditions Λ_8 does not meet. **Krein & Milman (1940)** — the classical antecedent of the hull-of-extreme-points property.
- The determination of a monotone function by its breakpoints is elementary; no attribution beyond Dilworth (1950), already cited at the down-set law, is claimed.