

The Method 1.6 — Mathematical Compendium

Matthew Lach

2026-08-26

Contents

I · THE CYCLE	8
The eighteen roots	9
II · THE OPERATORS	9
III · THE LANGUAGES, AND WHY EACH IS NECESSARY	10
Why none is redundant	11
IV · THE OBJECTS, BY FAMILY	12
A. The operators — closure, envelope, orientation — 43 objects	12
A.E — closure defect	12
A.EW — the metric defect	13
A.R — the recovery operator	13
A.S — the step operator	13
A.W — the walk	14
A.alph — projection onto coordinate i	14
A.anchor — the anchor	14
A.blind — the three blindnesses	15
A.bound — the defect bounds	15
A.bpc — binary path consistency / 2-decomposability (Kimura et al. 2024)	15
A.cert — the certificate condition	15
A.clos — closure operator (Moore family)	16
A.dechter — Dechter 1992	16
A.define — the defining letter	16
A.dens — density and informativeness	16
A.derived — adjunction never repairs (M Thm 10.1)	17
A.ebits — description length	17
A.env — monotone upper envelope / staircase bound	17
A.erel — E is coordinate-relative	17
A.expr — the expression always exists	18
A.ext — extensivity	18
A.fix — Bergman double-projection; Baker-Pixley 1975 (majority term)	18
A.freuder — Freuder 1982	19
A.gc — global consistency (CSP)	19
A.intext — interior and exterior	19
A.logic — the logic level	20
A.modeA — failure mode A — ordering	20
A.modeB — failure mode B — arity	20
A.montanari — Montanari 1974	20
A.morph — the cap as a morphism	21
A.orient — orientation cost	21
A.prod — the product rule for closure	21
A.prodE — the product rule for the defect	21
A.r4 — the four-orientation operator	22
A.relax — $\mathcal{R}$ as a relaxation	22
A.rule — the tightening rule	22
A.seaton — the polarisation formula	23
A.slack — slack	23
A.stair — staircase / connected row-convex	23
A.staircls — the one-corner characterisation	23
A.stat2 — the statistics operator	24

A.three — the three populations	24
A.two — the binary-projection criterion	24
S. The seed — generation, covering, structure — 18 objects	25
S.bits — the seed as binary	25
S.bounds — the bounds are recoverable	25
S.box — the box seed law, $d + c - 2$	25
S.car — the Carathéodory lower bound	26
S.channel — the channel conditions	26
S.core — the seed's necessary core	26
S.cover — the seed is a set cover	26
S.down — the down-set seed law, $d + c - 1$	27
S.erasure — the seed is an erasure structure	27
S.ground — the ground-state derivation	27
S.lam — Λ 's seed — seven cells	27
S.open — the open-index form, seed + E	28
S.parent — the parent-term wall	28
S.regime — the regime coordinate	29
S.ritz — the channel curve	29
S.seed — a seed	29
S.status — the existence partition	30
S.unit — the unit cell	30
F. The family of closed indexes — 4 objects	31
F.exact — the family's defect, exactly	31
F.moore — the family of closed sets	31
F.open — the family theorem	31
F.unstable — the unstable family	31
G. Graphs, constraints and languages — 12 objects	32
G.allcons — the index of all constraints	32
G.book — the book as an index	32
G.cons — the constraint index	32
G.graph — the constraint graph	32
G.near — the seven near-misses	33
G.prot — the protocol index	33
G.ref — the reference index	33
G.reg — the register as an index	33
G.shape — shape decides the seed	34
G.stat — statistics as a language	34
G.tower — the cycle rank of the tower	34
G.trip — independence is a triple property	34
L. Λ 's own constraints — 41 objects	35
L.E0 — the closure theorem	35
L.F — the generating function	35
L.F1 — the cardinality	35
L.Fm1 — the alternating specialisation	35
L.alpha — the join-irreducible count	36
L.amp — the amplification	36
L.arith — divisor lattice embedding	36
L.birk — Birkhoff representation 1937	36
L.bits — the Boolean representation	37
L.box — the box factorises	37

L.c1 — node counting	37
L.c2 — Pauli	37
L.c3 — the transfer bound	38
L.c4 — node counting	38
L.c5 — Pauli	38
L.c6 — the second transfer bound	38
L.c7 — vector coupling	38
L.c8 — the occupancy floor	39
L.chains — the linear extensions	39
L.chi — the membership function	39
L.circuit — the implication circuit	39
L.closed — sublattice of a product	39
L.def — the definition of Λ	40
L.dim — order dimension (Dushnik-Miller)	40
L.dist — Birkhoff / product of chains	40
L.metric — the lattice metric	40
L.mobius — Möbius function of a distributive lattice	41
L.modular — modularity (equality, not submodularity)	41
L.occ — the interval measure	41
L.omega — distinct-prime-counting function	41
L.pal — palindromic rank polynomial $\iff$ self-dual	41
L.pushback — join-prime and meet-prime	42
L.rankpoly — the rank polynomial	42
L.real — the index against the real subshells	42
L.skew — the rank skew	42
L.sperner — Sperner property / Dilworth	43
L.step — the interval-removal step	43
L.total — the membership function is total	43
L.tree — the constraint tree	43
L.void — the void	44
L.voidfrac — the void-free fraction	44
T. The tower and its couplings — 16 objects	44
T.a10 — Racah 1943 seniority	44
T.a11 — the parent bound	44
T.a12 — the recoupling bound	45
T.a13 — the spin-half bound	45
T.a9 — the seniority floor	45
T.a9p — the admissible Pauli cut	45
T.dens — the axis density	46
T.dich — the counting-coupling dichotomy	46
T.excl — the three excluded forms	46
T.invariant — basis independence of the level set	46
T.para — parastatistics	47
T.real — the bounds against real terms	47
T.scheme — the four coupling schemes	47
T.tight — constraint tightness as a count	48
T.tower — the tower	48
T.trad — the tree or the tightness	48
E. Empirical indexes — table, nuclide, layout — 4 objects	49
E.ioniz — the refusal map	49

E.layout — the price of a layout	49
E.nuclide — the nuclide chart's defect	49
E.table — the thirty-six decomposed	50
B. The bracket — 21 objects	50
B.V — the price V — the bracket's width against its error	50
B.V43 — the four-thirds law	50
B.Vexact — the exact price	50
B.adm — the five-sigma admissibility threshold — where the bracket may be applied at all	5
B.aitken — Aitken Δ^2 / Seki Kōwa	51
B.brk — monotone interpolation bracket	51
B.coll — the assembly rule — a composite observable's scaling exponent is $2a + 3b$	5
B.fail — when the bracket fails	52
B.floor2 — the floor at two	52
B.floor32 — the Rydberg floor	52
B.frac — the fractional widths	52
B.hstar — the optimal step	53
B.newton — Newton decrement, Nesterov-Nemirovskii 1994	53
B.nuV — the value-one crossing — where the bracket's price V passes unity	53
B.ordbr — the ordered bracket	54
B.ordk — the admissible order	54
B.pareto — the width-price front	54
B.pole — the linear pole	54
B.rank1 — the rank-one factorisation	55
B.selfconc — self-concordance	55
B.silence — what silence implies	55
K. Transit and information — 24 objects	55
K.arrow — the arrow index	55
K.axis — the axis test	56
K.cat — the tower is a category	56
K.clock — the composition clock	56
K.clockfail — the clock that does not tick	56
K.comp — composition	57
K.corner — corners and faces	57
K.coupling — the refuted conjecture	57
K.deadend — the dead ends, defined	58
K.decay — data-processing inequality + Pinsker	58
K.girth — the girth is four	58
K.helly — Helly number	58
K.jump — the categorical jump	59
K.langclose — closure as agreement	59
K.markov — the Markov property	59
K.peak — the composability peak	59
K.produce — the production rule	60
K.quiver — the composition quiver	60
K.redun — the projection ladder	60
K.three — the three-coordinate rule	61
K.transit — the transit profile	61
K.twocol — space is what an index holds, time is what it composes	61

K.window — the composable window	62
K.zcross — the periodic table as a coordinate	62
M. The modular chain — 9 objects	62
M.C1 — the presymplectic potential	62
M.C2 — the open object — half-sided modular inclusion on a non-expanding horizon ·	
GRADED OPEN	63
M.hsml — Borchers 1992 / Wiesbrock 1993	64
M.ledger — the modular ledger	64
M.rs — Reeh-Schlieder	65
M.semi — semifinite carries a trace	65
M.sorce — Sorce 2024	65
M.takesaki — Takesaki duality 1973	65
M.tt — Tomita-Takesaki	66
W. The violation index — 9 objects	66
W.core15 — the core at fifteen letters	66
W.core9 — the core at nine letters	66
W.face — the null surface is a face	67
W.frontier — the frontier metric — distance and status on the fifteen-letter alphabet	
.....	6
W.jur — jurisdiction does not change the defect	67
W.onehot — the one-hot partition	68
W.rel — Brunetti-Fredenhagen-Verch 2003	68
W.scale — the defect does not scale	68
W.sup — the core is minimal	68
EM. The electromagnetic quotient — 7 objects	69
EM.cross — the selection-rule crossing	69
EM.image — the complete rectangle	69
EM.map — electric/magnetic multipole selection rules	69
EM.notcomp — selection and composition are unrelated	70
EM.parity — the parity hole	70
EM.quotient — the electromagnetic quotient	70
EM.spin — the spin diagonal	70
C. Protocol and audit objects — 9 objects	71
C.Aq — the first factor	71
C.Bq — the second factor	71
C.box — the box generating function	71
C.compare — the comparison audit	71
C.cut — every cut closes	72
C.fib — the fibred count	72
C.local — local closure is not implied	72
C.pareto — the fibre trade	72
C.qmean — the transfer distribution	73
I. The intake — interval maps and convexity — 3 objects	73
I.convex — the convexity criterion	73
I.interval — the interval map	73
I.shape — the shape of an interval	73
P. The spectral mechanisms — what moves a defect and by how much — 17 objects	
.....	6
P.buildlimit — constructed limit	74
P.charge — the same-element ladder	74
P.converge — self-determined limit	74

P.coreblind — parent-term independence	75
P.dcollapse — orbital collapse	75
P.iso — the isoelectronic ladder	75
P.jj — the coupling-scheme marker	75
P.jsplit — the j-splitting	76
P.lcollapse — the l-collapse	76
P.lens — the precision lens	76
P.mono — the monotonicity of the defect	77
P.perturb — series perturbation	77
P.polar — core polarisation	77
P.qdt — core penetration	78
P.selfsame — series self-consistency	78
P.termsplit — core angular structure	78
P.trunc — truncation loss	78
Q. The channel equation — the closed form and its terms — 11 objects	79
Q.alpha — the polarisability index	79
Q.anchor — the two-ended anchor	79
Q.bound — the Pauli bound	80
Q.bridge — the bridge equation	80
Q.collapse — the Janet collapse	81
Q.delta — the channel equation	81
Q.exch — the exchange factor · GRADED OPEN	81
Q.final — the channel equation	82
Q.pen — the deficit term	82
Q.pol — Seaton's term	83
Q.region — the validated domain	83
LS. The Löwdin solution — the order derived from the equation — 8 objects	83
LS.ent — the entrant operator	84
LS.law — the ordering law, five clauses	84
LS.coll — the collapse condition	84
LS.pin — the pinned-channel theorem (no g block)	84
LS.asym — the state-dependent multiplier identity	85
LS.quart — the exact-quartic decomposition	85
LS.chord — the defect closed: chord = rot + perp	85
LS.twin — the twin operator, $c \rightarrow \infty$	86
3B. Three bodies — the index of families — 9 objects	86
3B.shape — shape space and the shape sphere	86
3B.metric — the shape metric	86
3B.JM — the Jacobi-Maupertuis metric	86
3B.pot — the potential on shape space	86
3B.norm — the algebraic variety	87
3B.five — the five fixed points	87
3B.tri — the triangle form on K_3	87
3B.def — the deficit	88
3B.index — the index Λ_3	89
V · THE CHAINS	89
VI · WHAT IS UNFINISHED	90
M.C2 — half-sided modular inclusion on a non-expanding horizon	90
Q.exch — the exchange factor	91
VII · THE BIBLIOGRAPHY	92

The works this compendium leans on most	97
THE MATHEMATICS OF THE LÖWDIN WORK	98
The corridor — a system of linear inequalities	98
The staircase algebra of a closed index	98
The Slater triangle, and a parity defect resolved	99
The falsification of a selection rule	99
The necessity of state	99
The observability boundary	99
Λ_{spectra} 's closure, and what the limit leaves	100
Attributions	101
What this work introduces, in one list	101
The Löwdin indexes — Λ_{law} and Λ_{const}	102
Λ_{law} — seven laws on (law, carrier). $E = 0$	102
Λ_{const} — fourteen constants on (role, carrier). $E = 0$ at 2 of 576.	102
The refusal, which is the more useful half	102

THE METHOD 1.6 — MATHEMATICAL COMPENDIUM

Compiled 2026-08-12; the counts below were recounted from this file on 2026-08-24 (register 1739). **265 objects · 18 roots · 263 settled · 2 unfinished.**

Verification coverage, stated rather than implied. 103 assertions name 46 of these objects and recompute each stated value against the register; **no verifier touches the other 167**. Of the check fields, **85 state a finding, 91 repeat the object's own key, 19 name a function, and 18 are empty** — so a check field present is not a verification performed. The check fields are classified at register 849.

This volume is drawn from the register, so it cannot drift from what the book actually holds.

I · THE CYCLE

The whole of this work is one recursion, stated at §14.5.2:

$\mathbf{X}_{(n+1)} = \mathcal{R}(\mathbf{X}_n \cup \Delta_n)$, halting when $\Delta_n \subseteq \mathcal{R}(\mathbf{X}_n)$ — when the increment is already implied.

It has never halted. Its falsifier is the ratio of closure to seed:

closure — objects the register holds	265
seed — objects nothing derives	17
ratio	15.6 : 1

The seed has not moved across every cycle of this project. If a genuinely new mathematical object entered that nothing existing derived, it would move — and that is the test.

The eighteen roots

object	statement
3B.shape	$\mathbb{R}^3 = \mathbb{C}^3 / (\text{translations} \times \text{rotations})$; $S^2 =$ that / scale — shape space (Montgomery 2014; Hopf 1931)
A.alph	$\hat{A}_i(X) = \{ x_i : x \in X \}$
A.seaton	$\delta(n) = (\alpha/K(l))(3 - l(l+1)/n^2)$ for a non-penetrating series, so $\delta_0 = 3\alpha/K$ and $\delta_2 = -l(\alpha/K)$
B.adm	$r = 2Z^2R/(\nu^3\sigma) \geq 5$
B.brk	$T(n)$ lies between $T(n-1)$ and $T(n+1)$
B.coll	an observable $\langle r \rangle^{a/(\Delta E)} b$ scales as $\nu^{\{2a+3b\}}$
B.nuV	$\nu_V = (3Z^2R/5q)^{\{1/4\}}$
L.c1	$\ell \leq n - 1$
L.c2	$k \leq 2(2\ell+1) = 4\ell+2$
L.c3	$q \leq k$
L.c4	$f \leq e - 1$
L.c5	$g \leq 4f+2$
L.c6	$g \leq q$
L.c7	$2S \leq k$
L.c8	$k \geq 1$ (definitional restriction, not a bound)
M.rs	the vacuum is cyclic and separating for local algebras
P.qdt	the quantum defect measures how far a Rydberg orbital reaches into the ionic core
P.trunc	truncation removes most channels and degrades those it leaves

II • THE OPERATORS

Every derivation in the book runs through these. Ten are this work's own; two (π , N_8) are cited and used as found.

symbol	definition	scope	origin
$\wedge \vee$	meet and join — greatest lower and least upper bound	cells	Birkhoff, standard
$\hat{\phi}$	the envelope: $\max x_i$ over cells with $x_j \leq v$	sets of cells	§14.5

symbol	definition	scope	origin
$\mathcal{R}$	closure at the ($\leq, \leq$) corner of Deville's staircase class	sets of cells	§14.5.4
$\mathcal{R}_4$	closure over all four orientations	sets of cells	§14.5.5
E	$E(X) = \mathcal{R}(X) - X $ — the defect	indexes	§6.1
E₄	the defect under $\mathcal{R}_4$; $E - E_4$ is the orientation cost	indexes	§14.5.5
d	$d(x,y) = \tau(\text{lcm/gcd}) = \prod(\Delta_i +1) = [x \wedge y, x \vee y] $	cells	title page
$\circ$	composition of transitions — Λ_9 is a category	cells	§12.11
seed	least G with $\mathcal{R}(G) = X$ — NP-hard	sets of cells	§14.5.7
S	envelope-step count; $2S$ = the tight-pair count	sets of cells	§14.5.9
ent	the entrant operator: argmax over frontier (n, ℓ) of $ D(n, \ell) $ in the ion's self-consistent field	subshells	§35; register 1701
π	the shape map $\mathbb{C}^3 \rightarrow \mathbb{R}^3$, $w = (1/2(Z_1 ^2 - Z_2 ^2), \text{Re } Z_1 \bar{Z}_2, \text{Im } Z_1 \bar{Z}_2)$	configurations	Montgomery 2014; Hopf 1931
N₈	the norm $\prod_{\varepsilon \in \{\pm 1\}^3} (U - \varepsilon \cdot u)$ — the minimal polynomial of a sum of three square roots	potentials	Lagrange 1770

III · THE LANGUAGES, AND WHY EACH IS NECESSARY

A language is a coordinate system. Translation is re-coordinatisation. E is the cost.

Six languages agree on Λ at 976, and **all ten pairs hold** — $C(5,2)$, a complete graph rather than a ladder. Binary is adjacent to every other, not only to logic. **The agreement is the result; the list is not.** What follows is one object stated seven ways, with the quantity each language alone supplies.

language	its statement of Λ	value	what only it can say
analysis	$F(1)$, the enumerator at $z = 1$	976	the cell count without enumerating
analysis	$F(-1)$, the alternating sum	2	the parity imbalance — invisible to a set
order	$ \mathcal{R}(X) $, the closure	976	what the envelopes admit
order	$E = \mathcal{R}(X) - X $	0	the defect — no other language states one
geometry	the void, per interval	0.464	what an interval admits and Λ refuses
binary	the box, one bit per cell	6,912	the space before any constraint
binary	density	0.1412	the fraction a constraint set keeps
information	bits to print the cells	12,449	the cost of stating it in full
information	bits to print the seed	89	139× compression
statistics	recovered from pairwise marginals	976	the object from its margins alone
statistics	its defect	0	— <i>and this is its limit</i>
algebra	constraints generating the ideal	8	derivability — what follows from what
documentary	<i>has anyone said this before</i>	—	no algorithm exists

Why none is redundant

Analysis gives $F(-1) = 2$. That is an alternating sum over the cells, and **a set has no alternating sum** — order, geometry and binary cannot express it. It is the parity imbalance, and it is why §12.1 can state its two-column law — ascending and descending quantities, exchanged under $v \mapsto v^{-3}$, a bipartite sign structure that admits no odd number of orientation reversals.

Order is the only language that states a DEFECT. E has no counterpart in the others: the generating function does not know what it fails to enumerate, and a polyhedron

does not know which of its lattice points are absent. **Every result in this work that begins $E =$ begins here.**

Geometry gives the void — 46.4% of a typical interval is admitted by the box and refused by Λ . The same quantity read in order is slack and in calculus is V , **and §12.11 shows the three are one number.** But only geometry states it per interval, which is where the exclusion principle shows.

Binary gives the box. **Nothing else in this work says what the space was before a constraint touched it**, and every density, every defect and every compression ratio is a fraction of it.

Information gives the compression: **139 cells printed for every one in the seed.** That number exists in no other language — order knows the seed is seven, and only description length says what seven buys.

Statistics recovers Λ exactly from its pairwise marginals, and **that is the whole of what it can do.** Tested on the periodic table it reports **$E = 0$ where $\mathcal{R}$ reports 36** — *marginals cannot see a hole.* **It agrees with the other languages only where there is nothing to disagree about**, which makes it necessary for one purpose and useless for another. **Knowing which is the point of having it.**

Algebra gives derivability. Whether a bound follows from the others is a question about an ideal, and **§14.5.7's anti-exchange failure and register 618's rule-15-backs-rule-5 are both statements in this language** — one constraint implied by another, which no count can express.

Documentary has no closure operator at all. *Has anyone said this before* cannot be bounded by computation, which is why §29's precedent question can be bounded and never settled, and why two searches returning nothing is evidence of a kind the other six never have to produce.

The test of necessity is not that each language is available. It is that each states a quantity the others cannot, and this table is that check run.

IV · THE OBJECTS, BY FAMILY

A. The operators — closure, envelope, orientation — 43 objects

A.E — closure defect

$$E(X) = |\mathcal{R}(X)| - |X|$$

none

grade **DEFINITIONAL**· source *M §6.1 / T A4; Moore 1910*· depends on A.R· 14 objects depend on it· depth 3

PRIOR ART: the gap between a set and its closure. That it is worth naming as a DEFECT is the books move; the quantity is the closure minus the set.

A.EW — the metric defect

$E_W(X) = |W(X)| - |X|$: the METRIC defect, what the steps reach and the index has not valued; distinct from $E(X) = |R(X)| - |X|$, the ORDINAL defect

an index whose cells carry values

grade **DEFINITIONAL**· source *R 1133; Dijkstra 1959*· depends on A.W, A.R· 1 object depends on it· depth 5

for Λ E_W is empty, because a cell of Λ is an arrangement and carries no number — which is why the book needed only E . For $\Lambda_spectra$ the two differ, and the gap between them is the cells that are placed and unvalued

PRIOR ART: the reachable set under a step relation, with accumulated cost and a least-cost route, is single-source shortest paths — Dijkstra, A note on two problems in connexion with graphs, Numer. Math. 1 (1959) 269-271.

A.R — the recovery operator

$R(X) = \{ x \in \prod_i \hat{A}_i(X) : x_i \leq \phi_{ij}(x_j) \forall i \neq j \}$

none beyond A.alph, A.env

grade **DEFINITIONAL**· source *M §6.1 / T A3; Moore 1910; Deville et al. 1999*· depends on A.alph, A.env· 20 objects depend on it· depth 2

PRIOR ART: R is a closure operator (Moore 1910) whose constraints are monotone staircases (Deville, Barette & Van Hentenryck, Artif. Intell. 109 (1999) 243-271). Neither the operator form nor the constraint class is new; what is new is reading E as a defect.

A.S — the step operator

the STEP operator: for each coordinate direction, the ratio between adjacent VALUED cells, its median and its scatter; a direction earns a step when the scatter is tight enough to carry a value

an index whose cells carry values

grade **MEASURED**· source *R 1132-1137; Edlen 1964*· depends on A.R· 1 object depends on it· depth 3

derived unaided from the spectra index it finds iso at s (1.233, scatter 1.13, 43 pairs), iso at p (1.185, 1.16, 25) and l at s (1.416, 1.43, 62). It must be derived from MEASURED cells ONLY: re-deriving from the walked result tightens iso at s to 1.08 on 116 pairs, because the walked values were generated by that step, and the step count goes 3 to 8

PRIOR ART: measuring a step as the ratio between adjacent valued cells is the isoelectronic and isonuclear method of Edlen, Atomic spectra, Handbuch der Physik XXVII (1964) 80-220.

A.W — the walk

the VALUATION closure: every cell reachable from a valued one by steps in S, accumulated error below a threshold, by the least-error route

an index, a step set S, and a tolerance

grade **PROVED**· source R 1132-1136; Dijkstra 1959· depends on A.S· 2 objects depend on it· depth 4

EXTENSIVE and IDEMPOTENT with exactly zero drift, since the least-error route from the seed leaves nothing tighter to find. MONOTONE except at one cell in 1,345: Mg II ni is measured at +0.0000, the Seaton step is multiplicative, and zero blocks a route a walked positive value would open — so adding a measurement can REMOVE a walked cell

PRIOR ART: the valuation closure is a shortest-path closure over a multiplicative cost. Extensivity, idempotence and monotone-except-at-zero follow from Dijkstra (1959).

A.alph — projection onto coordinate i

$\hat{A}_i(X) = \{ x_i : x \in X \}$

X a finite set of integer tuples

grade **DEFINITIONAL**· source M §6.1 / T A1; Codd 1970· 2 objects depend on it· depth 0

PRIOR ART: projection onto a coordinate is the relational projection operator — Codd, A relational model of data for large shared data banks, CACM 13 (1970) 377-387.

A.anchor — the anchor

REDUNDANCY ANCHORING: keep every route to a cell, not the best one; the spread between independent arrivals MEASURES the error rather than estimating it

a valuation closure with more than one route to a cell

grade **MEASURED**· source R 1136; Gauss 1809· depends on A.W· depth 5

956 cells of the spectra index are reached by two or more distinct routes. Observed spread 1.05 against a propagated estimate of 1.14 — ratio 0.92, so the walk's error bars are honest and slightly conservative. The check comes from the index disagreeing with itself, not from an outside formula

PRIOR ART: keeping every route to an estimate and reading their SPREAD as the error, rather than taking the best route, is the method of combining independent determinations — Gauss, Theoria motus corporum coelestium (1809), sec. 3.

A.blind — the three blindnesses

each language is blind in its own way: statistics cannot state a closure defect, analysis cannot state an ordering, a multiplicative algebra cannot state an exception

an index described in more than one language

grade **MEASURED**· source *R 1169; Shannon 1948; Freuder 1978*· depends on Q.bridge· depth 13

register 761 established the first. The second: least squares minimises squared error and has no rank in it — an equation fitting 311 values to $R^2 = 0.924$ got every ordering wrong. The third: a product of positive factors is monotone in each, so a multiplicative fit returns 205/205 where the data has eleven genuine exceptions

PRIOR ART: that a representation cannot state what its own coordinates cannot distinguish is the general fact behind all three blindnesses. Statistics sees marginals only (Shannon 1948 for what a marginal carries); analysis has no order in it; a multiplicative algebra is monotone in each factor by construction. Register 1170 measures them.

A.bound — the defect bounds

$0 \leq E(X) \leq |\text{box}| - |X|$

$\text{box} = \prod_i |\hat{A}_i(X)|$

grade **PROVED**· source *T audit 27; Moore 1910*· depends on A.E, A.ext· depth 4

PRIOR ART: the bounds are extensivity below and the ambient product above; both follow from the closure axioms.

A.bpc — binary path consistency / 2-decomposability (Kimura et al. 2024)

$X \subseteq \text{BPC}(X) \subseteq \mathcal{R}(X)$, $\text{BPC}(X) = \{ x : (x_i, x_j) \in \text{proj}_{ij}(X) \forall i < j \}$

none

grade **PROVED**· source *T A5; Cooper 1989; Janssen et al. 1989*· depends on A.R· 1 object depends on it· depth 3

PRIOR ART: binary path consistency and its relation to global consistency — Cooper, An optimal k-consistency algorithm, Artif. Intell. 41 (1989) 89-95; Janssen, Jegou, Nougier & Vilarem, A filtering process for general constraint-satisfaction problems, IEEE (1989), which gives a polynomial algorithm for pairwise consistency.

A.cert — the certificate condition

an open index does not close itself except where a CERTIFICATE exhibits an operation on its coordinate system reaching a fixed point of R

the admitted operations: relabel, re-coordinatise, refine a fibration, drop a coordinate

grade **PROVED**· source *M §18.4.1, promoted; Tarski 1955*· depends on A.clos, A.modeA, A.modeB· 1 object depends on it· depth 8

PRIOR ART: a closure operator does not close an open set without an added operation; exhibiting one is the fixed-point construction of Tarski, A lattice-theoretical fixpoint theorem, Pacific J. Math. 5 (1955) 285-309.

A.clos — closure operator (Moore family)

$\mathcal{R}$ is extensive, monotone, idempotent; the closed sets form a Moore family

none

grade **PROVED**· source *M §14.2 / T 1.3 (App. G); Moore 1910; Ward 1942*· depends on A.ext· 2 objects depend on it· depth 4

PRIOR ART: an extensive, monotone, idempotent operator is a CLOSURE OPERATOR and its fixed sets form a Moore family — E. H. Moore, Introduction to a Form of General Analysis (1910); M. Ward, The closure operators of a lattice, Ann. Math. 43 (1942) 191-196. $\mathcal{R}$ is one; nothing about $\mathcal{R}$ is needed to know its fixed sets are closed under intersection.

A.dechter — Dechter 1992

strong (w^*+1) -consistency on induced width w^* gives decomposability

*induced width w^**

grade **CITED**· source *Dechter 1992*· depends on A.gc· depth 5

A.dechter

A.define — the defining letter

a rung-1 coordinate contributes no envelope: same cells, box, E, envelope-step count and seed with or without it — it is invisible to $\mathcal{R}$ and still counts as a letter

it is the index's own name carried as a coordinate, taking one value because the object is what does not vary in it

grade **PROVED**· source *M §21.6.1; Birkhoff 1937*· depends on A.env· depth 2

PRIOR ART: a coordinate with a single value contributes no join-irreducible, so it changes neither the lattice nor its envelopes.

A.dens — density and informativeness

$E = 0$ is informative at any density; $E > 0$ at low density measures sparsity

density = $|X| / |\text{box}|$

grade **COMPUTED**· source *T 1.9 (App. G); Erdos & Renyi 1960*· depends on A.E· 1 object depends on it· depth 4

PRIOR ART: that a structural property can hold at any density, while its failure measures sparsity, is the random-graph threshold picture of Erdos & Renyi, On the evolution of random graphs, Publ. Math. Inst. Hung. Acad. Sci. 5 (1960) 17-61.

A.derived — adjunction never repairs (M Thm 10.1)

a derived coordinate cannot repair closure: the box grows by its value count and $|X|$ is fixed

the coordinate is a function of the others

grade **PROVED**· source *T 1.7 (App. G) / M §17.2 Thm 10.1; Birkhoff 1940*· depends on A.E· 1 object depends on it· depth 4

PRIOR ART: a derived coordinate is a function of the others, so it adds no join-irreducibles and cannot enlarge the closed family; it only enlarges the ambient box.

A.ebits — description length

$E_bits(X) = \log_2 C(|\mathcal{R}(X)|, E(X))$

counting measure

grade **COMPUTED**· source *M §25.3; Shannon 1948; Rissanen 1978*· depends on A.E· 1 object depends on it· depth 4

PRIOR ART: description length as a measure of structure — Shannon, A mathematical theory of communication, Bell Syst. Tech. J. 27 (1948); Rissanen, Modeling by shortest data description, Automatica 14 (1978) 465-471. $\log_2 C(|R|, E)$ is the cost of naming which admitted cells are absent.

A.env — monotone upper envelope / staircase bound

$\phi_{ij}(v) = \max\{ x_i : x \in X, x_j \leq v \}; \max \emptyset = -\infty$

$i \neq j$

grade **DEFINITIONAL**· source *M §6.1 / T A2; Deville et al. 1999*· depends on A.alph· 7 objects depend on it· depth 1

PRIOR ART: the monotone upper envelope of a relation is its staircase bound — the connected row-convex class of Deville, Barette & Van Hentenryck (1999) and the row-convex networks of van Beek & Dechter, J. ACM 42 (1995) 543-561.

A.erel — E is coordinate-relative

$E = 0$ is relative to coordinates: any X with $|X| = a \cdot b$ relabels onto an $a \times b$ rectangle, which is a full box and closed — so every index with a composite cell count has a coordinate system in which $E = 0$

the coordinates must be fixed by the subject, not chosen; every coordinate in this book is

grade **PROVED**· source *M §21.6; Birkhoff 1940; Birkhoff 1937; Dushnik & Miller 1941*· depends on A.R· depth 3

PRIOR ART: $E = 0$ is a statement about a COORDINATISATION, not a set. Any set of size ab relabels onto an $a \times b$ grid, which is closed. This is the standard observation that lattice properties are not invariants of the underlying set.

PRIOR ART: a lattice property belongs to a COORDINATISATION and not to the underlying set — the same reason order dimension is not an invariant of cardinality. Birkhoff (1937); Dushnik & Miller, Partially ordered sets, Amer. J. Math. 63 (1941) 600-610.

A.expr — the expression always exists

a closed index equals the set its own envelopes admit, so its expression always exists and is recoverable

$$X = R(X)$$

grade **PROVED**· source *M §15.2 corollary / §2.23; Moore 1910*· depends on A.R, S.bounds· depth 4

PRIOR ART: a closed set equals the fixed point of its own closure operator, so its defining expression always exists. Immediate from idempotence.

A.ext — extensivity

$$X \subseteq R(X), \text{ hence } E(X) \geq 0$$

none

grade **PROVED**· source *T 1.2 (App. G) / M A.2; Moore 1910*· depends on A.R· 2 objects depend on it· depth 3

PRIOR ART: extensivity is the first Moore axiom. That $E(X) \geq 0$ is its immediate corollary, not a separate result.

A.fix — Bergman double-projection; Baker-Pixley 1975 (majority term)

$$X \text{ is closed} \iff X = R(X)$$

each coordinate presented as a chain

grade **PROVED**· source *M §14.1 (Thm 9.1, A.2); Moore 1910; Baker & Pixley 1975*· depends on A.clos· 1 object depends on it· depth 5

PRIOR ART: X closed iff $X = R(X)$ is the definition of a fixed point of a closure operator. That the binary projections suffice is Baker & Pixley, Polynomial interpolation and the Chinese remainder theorem for algebraic systems, Math. Z. 143 (1975) 165-174 — the majority-term / (2,d) interpolation theorem.

A.freuder — Freuder 1982

a tree-structured constraint network is globally consistent after arc consistency

constraint graph acyclic

grade **CITED**· source *T 1.10 (App. G); Freuder 1982*· depends on A.gc· depth 5

PRIOR ART: Freuder, A sufficient condition for backtrack-free search, J. ACM 29 (1982) 24-32 — a tree-structured constraint network is globally consistent after arc consistency. This is the reason Λ closes, and it is his.

A.gc — global consistency (CSP)

$E(X) = 0 \iff$ the binary constraint network is globally consistent

the constraints are the monotone binary projections

grade **CITED**· source *T 1.4 (App. G) / A5c; Freuder 1978; Dechter 1992*· depends on A.bpc, A.E· 3 objects depend on it· depth 4

A.gc

PRIOR ART: global consistency and its k-consistency ladder — Freuder, Synthesizing constraint expressions, CACM 21 (1978) 958-966; Dechter, From local to global consistency, Artif. Intell. 55 (1992) 87-107.

A.intext — interior and exterior

E is INTERIOR and E_W is EXTERIOR: the order operator is bounded by its sample and the step operator is not

an index with valued cells and a step set

grade **MEASURED**· source *R 1196-1199; Beeri, Fagin, Maier & Yannakakis 1983; Dijkstra 1959*· depends on A.E, A.EW· 1 object depends on it· depth 6

on Λ spectra in-region, R places 1,925 from 277 held ($E = 1,648$) and W values 277 ($E_W = 0$) at the book's tolerance, because every step's scatter exceeds it — multiplicity 1.664, isoelectronic 1.674, l 2.872. Raising tau does not close the gap: every cell W gains, R mostly refuses, and ALL 218 fail one envelope, charge given Z. The measured set runs Z 2-83 and charge 1-9; the refused cells run Z 1-85 and charge 1-10. The overlap never exceeds 12% of E at any tolerance

PRIOR ART: the order operator is a closure bounded by its sample (BFMY 1983 for what local closure can reach); the step operator is a reachability closure unbounded by it (Dijkstra 1959). That E is interior and E_W exterior follows from the two being different kinds of operator.

A.logic — the logic level

LOGIC is not a language: it is the mechanism binary -> language -> binary by which any language answers a question about a cell

a language with a closure operator

grade **DEFINITIONAL**· source *R 1173; Boole 1854; Tarski 1936*· depends on A.R· 1
object depends on it· depth 3

three levels: BINARY is the type, a cell is admitted or not; a LANGUAGE is a coordinate system with a closure operator; LOGIC is the map. A language earns a row when logic can operate on it and return a binary, which is why documentary has none — it returns a citation. The book's $C(5,2) = 10$ combinations is then exact: five operator-bearing languages, plus statistics as a sixth and documentary as a seventh

PRIOR ART: that logic is the mechanism by which a language answers a binary question, rather than a language itself, is the object-language/metalanguage distinction — Boole, An Investigation of the Laws of Thought (1854); Tarski, Der Wahrheitsbegriff in den formalisierten Sprachen, Studia Philos. 1 (1936) 261-405.

A.modeA — failure mode A — ordering

sub-case A, ORDERING: the term is present and determined but non-monotone; repairable by re-ordering

the term is a function of an existing coordinate

grade **COMPUTED**· source *T 1.6 (App. G); Freuder 1978*· depends on A.rule· 1 object
depends on it· depth 7

PRIOR ART: a non-monotone constraint is not captured by an envelope, and higher consistency is needed — Freuder, Synthesizing constraint expressions, CACM 21 (1978) 958-966.

A.modeB — failure mode B — arity

sub-case B, ARITY: every term is monotone; the constraint names more coordinates than an envelope has arguments; not repairable by any operation on coordinates

the minimal failing support has size ≥ 3

grade **COMPUTED**· source *T 1.6 (App. G); Freuder 1978*· depends on A.rule· 2 objects
depend on it· depth 7

PRIOR ART: a constraint naming more coordinates than the envelope pairs is a higher-arity constraint; the k-consistency ladder is Freuder (1978).

A.montanari — Montanari 1974

for monotone constraints, path consistency implies global consistency

constraints monotone

grade **CITED**· source *T 1.10 (App. G); Montanari 1974*· depends on A.gc· depth 5

PRIOR ART: Montanari, Networks of constraints: fundamental properties and applications to picture processing, Inf. Sci. 7 (1974) 95-132 — for monotone constraints, path consistency implies global consistency. A constraints are monotone, so this applies directly.

A.morph — the cap as a morphism

an occupancy coordinate's cap is admissible iff it is a morphism for the operation preserved

meet-morphism preserves meets; join-morphism preserves joins

grade **COMPUTED**· source *M §18.4.1; Birkhoff 1940*· depends on A.rule· depth 7

PRIOR ART: a cap is admissible iff it is a lattice morphism for the operation — the standard homomorphism condition. Birkhoff, Lattice Theory (1940).

A.orient — orientation cost

E – E₄, the ORIENTATION COST: 0 on eight of ten indexed objects, 2 on the audits, 750 on the parity rule

R₄ defined

grade **COMPUTED**· source *M §14.5.5; Deville et al. 1999; Deville, Barette & Van Hentenryck 1999*· depends on A.r4· depth 4

PRIOR ART: the cost of choosing one orientation over the full class. The class is Devilles; the cost is measured here.

PRIOR ART: the four orientations of a staircase are the (alpha, beta)-monotone constraints of Deville, Barette & Van Hentenryck (1999). R uses one corner and R₄ the closure over all four; the ORIENTATION COST is the difference, measured at 0 on eight of ten indexed objects and 2 on the aufbau index.

A.prod — the product rule for closure

if no constraint links the factors, $\mathcal{R}(A \times B) = \mathcal{R}(A) \times \mathcal{R}(B)$

A, B on disjoint coordinate sets, no cross constraint

grade **PROVED**· source *T D9; Birkhoff 1940*· depends on A.R· 1 object depends on it· depth 3

PRIOR ART: a closure operator on a product with no cross-constraints factorises; the statement is the product form of a Moore family. Birkhoff, Lattice Theory (1940).

A.prodE — the product rule for the defect

$E(A \times B) = |A| \cdot E_B + |B| \cdot E_A + E_A \cdot E_B$

as A.prod

grade **PROVED**· source *T D10; classical; Euler 1748*· depends on A.prod, A.E· depth 4

PRIOR ART: the defect of a product expands as $|A|E_B + |B|E_A + E_A E_B$ — the inclusion-exclusion expansion of $(|A|+E_A)(|B|+E_B)$ minus $|A||B|$. Elementary.

PRIOR ART: expanding $(|A| + E_A)(|B| + E_B) - |A||B|$ gives the three terms. Elementary algebra; the point is that defects MULTIPLY as well as add across a product. **PRIOR ART:** expanding $(|A| + E_A)(|B| + E_B) - |A||B|$ gives the three terms. The point is that defects MULTIPLY as well as add across a product, which is the elementary product rule for counting — Euler, *Introductio in analysin infinitorum* (1748), ch. XVI.

A.r4 — the four-orientation operator

$\mathcal{R}_4$, the four-orientation closure over Deville's staircase class

idempotent, hence a closure operator

grade **COMPUTED**· source *M §14.5.5; Deville et al. 1999*· depends on A.stair, A.R· 2 objects depend on it· depth 3

PRIOR ART: the four orientations of a staircase constraint are the (alpha, beta)-monotone class of Deville, Barette & Van Hentenryck (1999); R uses one corner and $\mathcal{R}_4$ the closure over all four.

A.relax — $\mathcal{R}$ as a relaxation

$\mathcal{R}$ is not a k-wise closure for any k: k-wise defect is 0 at $k = d$ while $E(\mathcal{R})$ can be 750

$|\Delta\emptyset|=1$ at 750 against k-wise 0 for $k = 2,3,4$

grade **PROVED**· source *M §14.5.4; Beeri, Fagin, Maier & Yannakakis 1983*· depends on A.R· 1 object depends on it· depth 3

PRIOR ART: pairwise (k-wise) consistency implies global consistency exactly for ACYCLIC hypergraphs — BFMY, On the desirability of acyclic database schemes, J. ACM 30 (1983) 479-513. R is the global operator and the k-wise closures are the local ones; the gap between them is their theorem, and this object measures it.

A.rule — the tightening rule

a tightening preserves $E = 0$ iff it binds one coordinate by a monotone function of one other

the index is a sublattice of a product of chains

grade **COMPUTED**· source *T 1.8 (App. G) / M §14.4; Deville et al. 1999; Deville, Barette & Van Hentenryck 1999; van Beek & Dechter 1995*· depends on A.fix· 10 objects depend on it· depth 6

PRIOR ART: a tightening preserves closure iff it stays inside the monotone staircase class.

PRIOR ART: a tightening preserves closure exactly when it stays inside the monotone staircase class — the connected row-convex constraints of Deville, Barette & Van Hentenryck, Constraint satisfaction over connected row-convex constraints, *Artif. Intell.* 109 (1999) 243-271, and the row-convex networks of van Beek & Dechter, On the minimality and global consistency of row-convex constraint networks, *J. ACM* 42 (1995) 543-561. Binding one coordinate by a monotone function of one other is precisely a member of that class.

A.seaton — the polarisation formula

$\text{delta}(n) = (\alpha/K(l))(3 - l(l+1)/n^2)$ for a non-penetrating series, so $\text{delta}_0 = 3 \alpha/K$ and $\text{delta}_2 = -l(l+1) \alpha/K$

a Rydberg electron that does not enter the core

grade **CITED**· source *Seaton 1958; Drake & Swainson 1991*· 1 object depends on it· depth 0

tested on 23 adjacent- l pairs in its domain: median observed/predicted 1.12, improved from 1.19 by restoring the n -dependent term, which confirms the term belongs. It fails on 20 near-hydrogenic pairs at the noise floor and on Ba II ($\text{delta}_f = 0.756$) and Hg II (1.062), where the 4f orbital has collapsed into the core and the series penetrates

A.slack — slack

$\text{SLACK}(S) = \text{measure}(\text{ambient})/\text{measure}(S)$; $\log \text{SLACK} = \log|\mathcal{R}(X)| - \log|X|$

a measure on the ambient

grade **COMPUTED**· source *M §25.2; Shannon 1948*· depends on A.E, A.ebits· 1 object depends on it· depth 5

PRIOR ART: log of the ratio of ambient to actual is a bit count — the same quantity as A.ebits in another form. Shannon (1948).

A.stair — staircase / connected row-convex

ϕ -hat recovers a STAIRCASE constraint: (α, β) -monotone with α, β in $\{\leq, \geq\}$

binary

grade **CITED**· source *Deville, Barták, Van Hentenryck 1999*· depends on A.env· 2 objects depend on it· depth 2

A.stair

A.staircls — the one-corner characterisation

$E(\mathcal{R}) = 0$ iff X is an intersection of $(\leq, \leq)$ staircases — one corner of Deville's class

the anti-diagonal is a staircase with $E = 750$

grade **PROVED**· source *M §14.5.4; Deville, Barette & Van Hentenryck 1999; van Beek & Dechter 1995*· depends on A.relax, A.stair· depth 4

PRIOR ART: connected row-convex and monotone staircase constraint classes — Deville, Barette & Van Hentenryck, Constraint satisfaction over connected row-convex constraints, Artif. Intell. 109 (1999) 243-271; van Beek & Dechter, On the minimality and global consistency of row-convex constraint networks, J. ACM 42 (1995) 543-561.

A.stat2 — the statistics operator

the statistics operator is max-entropy on the PAIRWISE marginals, not the first-order ones

an index of dimension $d \geq 3$

grade **MEASURED**· source *R 1174*· depends on A.logic· 1 object depends on it· depth 4

on Λ , first-order marginals admit all 6,912 ambient cells; pairwise marginals admit exactly 976 and reproduce R cell for cell. IPF converges to the max-entropy distribution matching the SPECIFIED marginals (Deming and Stephan 1940; Ireland and Kullback 1968), so which marginals is the whole question

A.three — the three populations

the Method equation partitions an index into three populations: interior captures, the working overlap, and exterior predictions

an index, its order operator and its step set

grade **MEASURED**· source *R 1199-1200; Freuder 1978; Dijkstra 1959*· depends on A.intext· depth 7

at $\tau = 3.0$: INTERIOR 1,503 cells R places and no step values, median $l = 2$ and charge 4 — these are captures. BOTH 145 placed and valued, median $l = 3$, the polarisation regime. EXTERIOR 218 the steps reach past the sample edge. The exterior verifies where it reaches: Cs I np walked to 3.659 against a measured 3.5667, an error of 2.6% with a stated error factor of 1.67

PRIOR ART: partitioning by what a closure admits and what a propagation reaches — the k-consistency gap (Freuder 1978) crossed with shortest-path reachability (Dijkstra 1959). The three populations are the measurement.

A.two — the binary-projection criterion

a tightening preserves $E = 0$ iff each binary projection is the intersection of that projection's own two upper envelopes

phi-hat indexed over ORDERED pairs

grade **PROVED**· source *M §14.4 / §2.15.2; Baker & Pixley 1975*· depends on A.rule· depth 7

PRIOR ART: that binary projections decide membership is the (2,d) interpolation property — Baker & Pixley, Polynomial interpolation and the Chinese remainder theorem for algebraic systems, Math. Z. 143 (1975) 165-174.

S. The seed — generation, covering, structure — 18 objects

S.bits — the seed as binary

the five read as binary: corners 1 and 2 are exact complements (3 of 3), corner 3 and the unit cell are exact complements (3 of 3), corner 4 is fully specified at 11001100, and every coordinate receives both a 0 and a 1 with no column one-sided

measured at one cap setting only; whether the pairing survives other caps is untested

grade **COMPUTED**· source *M §14.5.14; Shannon 1938*· depends on S.unit· depth 9

PRIOR ART: reading a structure as binary words with complementation is Boolean algebra — Shannon, Trans. AIEE 57 (1938) 713-723.

S.bounds — the bounds are recoverable

S3: the bounds are recoverable from the cells

closure

grade **PROVED**· source *M §15.2; Moore 1910*· depends on A.R· 1 object depends on it· depth 3

PRIOR ART: if the envelopes are recoverable from the cells then the constraints are too, since R is determined by its envelopes.

S.box — the box seed law, $d + c - 2$

a full box c^d seeds at $d + c - 2$

exact by branch and bound at 3^3 and 4^3

grade **PROVED**· source *M §14.5.9; classical; Dilworth 1950; Birkhoff 1937*· depends on S.cover· depth 5

PRIOR ART: the corners of a d-dimensional box over c values generate it under coordinatewise max, and $d + c - 2$ is the count of extreme steps. Elementary.

PRIOR ART: as S.down, for a full box: the extreme steps in each coordinate generate it. PRIOR ART: the generators of a down-set closed under coordinatewise max are its maximal elements; for a box over c values in d coordinates that count is $d + c - 2$. Dilworth, A decomposition theorem for partially ordered sets, Ann. Math. 51 (1950) 161-166; Birkhoff (1937) for the representation.

S.car — the Carathéodory lower bound

seed $\geq$ the Carathéodory number = the breadth = d for a product of d chains

semilattice with subsemilattices as convex sets

grade **CITED**· source *Carathéodory / convexity spaces; Caratheodory 1911*· depends on S.cover· depth 5

PRIOR ART: Caratheodory, Über den Variabilitätsbereich der Fourierschen Konstanten, Rend. Circ. Mat. Palermo 32 (1911) 193-217 — every point of a convex hull in $\mathbb{R}^d$ is a combination of at most $d+1$ points. Its lattice analogue is the breadth, and it bounds the seed below.

S.channel — the channel conditions

six channel conditions hold in all 219 minimum covers of Λ_8 : an s-s, s-p, p-s and p-p transition, a null transition $q = 0$, and a full transfer $q = k$

the constraint is on the CHANNEL, not the cell — which is why every cover looks alike while nothing is forced

grade **COMPUTED**· source *M §14.5.12; Condon & Shortley 1935*· depends on S.core· 1 object depends on it· depth 7

PRIOR ART: the s-s, s-p and related conditions are statements about which subshell transitions the seed must contain; the subshell structure is Condon & Shortley (1935).

S.core — the seed's necessary core

four of the seed's seven cells are NECESSARY: each uniquely covers 12-23 envelope elements and has ZERO alternatives in Λ ; they cover 87 of 102

a stable attractor of greedy covering, not necessity; §14.5.7 stands unamended (register 600)

grade **COMPUTED**· source *M §14.5.10; Karp 1972; Chvatal 1979*· depends on S.lam· 2 objects depend on it· depth 6

PRIOR ART: an element covered by exactly one set forces that set into every cover — the standard reduction rule for set cover.

S.cover — the seed is a set cover

the minimum seed is a MINIMUM SET COVER: elements the envelope steps, sets the cells

S.seed

grade **PROVED**· source *M §14.5.9; Karp 1972; Johnson 1974*· depends on S.seed· 5 objects depend on it· depth 4

PRIOR ART: MINIMUM SET COVER is one of the 21 NP-complete problems of Karp, Reducibility among combinatorial problems (1972);

the greedy $\ln n$ approximation is Johnson, Approximation algorithms for combinatorial problems, JCSS 9 (1974) 256-278. The seed problem IS set cover, which is why five heuristics agree on 7 and the lower bound is 5.

S.down — the down-set seed law, $d + c - 1$

a down-set over c values in d coordinates seeds at $d + c - 1$

exact: $LB = UB$ at $d=4$ $c=4$

grade **PROVED**· source *M §14.5.9; Dilworth 1950*· depends on S.cover· depth 5

PRIOR ART: the minimum generating set of a down-set is its set of maximal elements; the $d + c - 1$ count follows. Dilworth, A decomposition theorem for partially ordered sets, Ann. Math. 51 (1950) 161-166.

S.erasure — the seed is an erasure structure

the remaining three seed positions admit 519 completing triples over 66 distinct cells, none in all 519, each appearing in 3 to 157 — a redundancy gradient of fifty to one

heavy-tailed: min 3, median 12, max 157, a ratio of thirteen to one

grade **COMPUTED**· source *M §14.5.11; Karp 1972*· depends on S.core· depth 7

PRIOR ART: the alternative completions of a partial cover; the count is the measurement.

S.ground — the ground-state derivation

the ground configuration determines every channel's existence and bounds its defect's integer part

an atom of atomic number Z

grade **MEASURED**· source *R 1139-1141; Madelung 1936; Janet 1929; Hund 1925*· depends on K.produce· 5 objects depend on it· depth 6

multiplicity from Hund on the core: 24 of 24 electron counts, exact containment. The cell set from aufbau: 8,488 cells admitting all 311 measured channels where the sampled index admits 296. And $\text{floor}(\delta) \leq \min(p, n_0-l-1)$ for 311 of 311, exact 57%, within two 97%

PRIOR ART: the ground configuration follows the $n+l$ ordering — Janet, La classification helicoidale des elements chimiques (1929); Madelung's rule as usually stated (1936). Hund's first rule (Z. Phys. 33, 1925) gives the term. Both are read, not derived.

S.lam — Λ 's seed — seven cells

seed(Λ_8) = 7 exactly — 976 cells from seven, 139 to 1; LB 5, five heuristics 12/7/7/7/40

branch and bound over 102 elements

grade **COMPUTED**· source *M §14.5.9; Karp 1972; Johnson 1974*· depends on S.cover·
1 object depends on it· depth 5

PRIOR ART: the seed problem is minimum set cover, NP-complete (Karp 1972), with a greedy $\ln n$ approximation (Johnson 1974). Five heuristics agreeing on 7 against a lower bound of 5 is the measurement.

S.open — the open-index form, seed + E

an open index is recovered as seed($\mathcal{R}(X)$) plus the E cells $\mathcal{R}(X)$ holds and X does not; cost seed + E

exact on four open indexes; compresses only where $E \ll |X|$

grade **PROVED**· source *M §21.5; Moore 1910*· depends on S.seed· 1 object depends on it· depth 4

PRIOR ART: an open index needs its closure seed plus the cells the closure adds and it does not hold — the decomposition follows from extensivity.

S.parent — the parent-term wall

an OPEN-SHELL core gives many parent terms and no separable Rydberg series, so its levels can be published and its defect cannot be extracted

an ion whose core has an open subshell

grade **MEASURED**· source *R 1180; Condon & Shortley 1935; Racah 1943*· depends on S.ground· 1 object depends on it· depth 7

Fe IV's 3d4 core carries sixteen LS terms. The capture shows 13 of them, 24 distinct (parent, l, term) series, and EVERY ONE has exactly one member — $n = 4$ only. Fe IV has ~1,000 analysed levels and no extractable defect. The compendium holds Ne-like and Na-like ions at charge 15 and 16 and no open-shell ion above charge 6, which is not a collection preference but the fact that an open-shell core does not produce the object a quantum-defect index holds

PRIOR ART: parent terms and the fractional-parentage decomposition are Condon & Shortley, The Theory of Atomic Spectra (1935), ch. VII, and Racah's coefficients of fractional parentage (1943). That an open-shell core gives one series per parent is their structure; the measurement is that Fe IV's 24 series each hold one member.

PRIOR ART: parent terms and coefficients of fractional parentage are Condon & Shortley, The Theory of Atomic Spectra (1935), ch. VII, and Racah, Theory of complex spectra III, Phys. Rev. 63 (1943) 367-382. That an open-shell core gives one series per parent is their structure; the measurement is that the 24 Fe IV series each hold one member.

S.regime — the regime coordinate

the SIGN of d2 is a coordinate the ground state cannot supply, and the MAGNITUDE of d2 predicts the channel's own scatter

a Rydberg channel with a fitted Ritz curve

grade **MEASURED**· source *R 1157-1162; NIST Atomic Spectroscopy compendium; Ritz 1903*· depends on S.ritz· 2 objects depend on it· depth 8

Seaton requires $d2 < 0$ for polarisation, so a positive d2 says the channel is not what the formula describes. By orbital: s 76% positive, p 74%, d 37%, f 6%, g 7%, h 9%. The two populations differ in median raw spread 0.0159 vs 0.0029 and median fit residual 0.0058 vs 0.0003, U-test $p < 1e-5$. The MAGNITUDE class predicts the spread at $R^2 = 0.761$ against l alone at 0.230 and the sign at 0.267, and the sign adds nothing to the magnitude. The three quantities do not separate: $d0$ against $|d2|$ gives $r^2 = 0.270$

PRIOR ART: the SIGN rule is NIST's own — its compendium states that the Ritz coefficient a is usually positive for core-penetration series and negative for core-polarization series. What is measured here is the distribution across 274 channels and that $|d2|$ predicts a channel's scatter at $R^2 = 0.761$.

PRIOR ART: the SIGN rule is NIST own — its compendium states that the Ritz coefficient a is usually positive for core-penetration series and negative for core-polarization series. What is measured here is the distribution across 274 channels and that $|d2|$ predicts a channel scatter at $R^2 = 0.761$.

S.ritz — the channel curve

a channel is a CURVE, not a number: $\delta(n) = d0 + d2/(n-d0)^2$, and the index reaches $d0$ while nothing in it reaches $d2$

a Rydberg channel with four or more members

grade **MEASURED**· source *R 1150-1156; Ritz 1903; Hartree 1928*· depends on S.ground· 1 object depends on it· depth 7

the $d2$ term removes 48% of what the compendium called scatter (0.0199 $\rightarrow$ 0.0104 across 274 channels). $d0$ against $\ln$ Ne, l and charge gives $R^2 = 0.549$; the curvature gives 0.018. The ISOELECTRONIC step moves both coordinates by the same factor at every l — 1.194/0.859 at s, 1.150/0.979 at p, 0.858/0.907 at d, 0.561/0.627 at f — while the l step moves them differently: 1.427/0.752, 3.696/1.373, 8.536/5.264. Along a sequence the channel translates; along l it deforms, and fitting l as a 2×2 map improves on the scalar by only 14% at $s \rightarrow p$ and 1% beyond

PRIOR ART: the extended Ritz formula $\delta = \delta_0 + a/(n - \delta_0)^2 + b/(n - \delta_0)^4$ is Ritz, Zur Theorie der Serienspektren, Ann. Phys. 12 (1903) 264-310, with Hargreaves and Hartree (Proc. Camb. Phil. Soc. 24, 1928) supplying the foundation. NIST's compendium states it in this form.

S.seed — a seed

$G \subseteq X$ is a seed of X under $\mathcal{R}$ iff $\hat{\phi}(G) = \hat{\phi}(X)$

X closed; $\mathcal{R}$ depends on G only through $\hat{\phi}$

grade **PROVED**· source *M §14.5.7; Birkhoff 1937; Moore 1910*· depends on A.env, A.R.
3 objects depend on it· depth 3

PRIOR ART: a generating set of a closure system is any G with $\text{cl}(G) = \text{cl}(X)$. That the envelope alone decides it is the specific form here; the notion is Moore-Birkhoff.

S.status — the existence partition

the index partitions by EXISTENCE STATUS, and nothing in it is impossible

the aufbau survey

grade **MEASURED**· source *R 1191-1192; Theodosiou, Inokuti & Manson 1986*·
depends on S.ground, S.parent· 1 object depends on it· depth 8

101,328 cells. *** THE STATUS AXIS WAS REPLACED AT REGISTER 1578:
VERIFIED/POSSIBLE/IMPROBABLE became WITNESSED/UNWITNESSED plus a
NAMED BOUND, because improbable is a judgement about the future rather than a
fact about the record — the same object-versus-observer fault register 1287 found in
standing. *** Now: 337 WITNESSED (0.333% of the index) and 100,991
UNWITNESSED, of which 929 are exact by symmetry and need no measurement.
Bounds on the rest: series above ng 28,527; no long-lived isotope 24,312; no analysis
located at this charge 17,428; open-shell cores 26,641; not naturally occurring 225;
and 2,929 with NO bound at all — separable series simply not yet measured. The
charge bound is contradicted by twelve measured cells above it in this index's own file
(R 1577). Every accuracy figure must be quoted against the witnessed count, not
101,328.

PRIOR ART: the distinction between what a survey admits and what is measurable is exactly the distinction Theodosiou, Manson & Inokuti make in their 1986 table across all ionisation stages.

S.unit — the unit cell

all 219 covers contain a cell matching (3,0,1,1,*,0,1,1) — 3s1 to e-s ending 1, doublet, e free at 1 or 3; its function is to carry the value ONE where the corners carry extremes

three of the fifteen elements the corners miss are the alphabet values $q=1$, $g=1$, $2S=1$; $e=3$ witnesses the target's ceiling and $e=1$ the source's

grade **COMPUTED**· source *M §14.5.13; Karp 1972; Chvatal 1979*· depends on
S.channel· 1 object depends on it· depth 8

PRIOR ART: a cell present in every minimum cover is forced; the standard set-cover argument.

PRIOR ART: a set present in every minimum cover is forced by an element it uniquely covers — the same reduction rule. Chvatal (1979). That all 219 covers contain this cell is the measurement.

F. The family of closed indexes — 4 objects

F.exact — the family's defect, exactly

$\mathcal{R}(\text{Cl}(U)) = 2^U$, so $E(\text{Cl}(U)) = 2^{|U|} - |\text{Cl}(U)|$ exactly

$\text{Cl}(U)$ holds $\emptyset$ and U , and every ordered pair of points is separated by some closed set, making every envelope constant at 1

grade **PROVED**· source *M §14.6.2; Moore 1910*· depends on F.open· depth 6

PRIOR ART: the closure of a set of generators over an unconstrained alphabet is the full power set, so the defect is the exact complement count. A corollary of extensivity and idempotence.

F.moore — the family of closed sets

the $\mathcal{R}$ -closed subsets of a closed index form a Moore family: $\cap$ -closed 100%, $\cup$ -closed 33-68%; 73, 146, 731 members

exhaustive at 8, 9, 16 cells

grade **PROVED**· source *M §14.5; Moore 1910; Ward 1942*· depends on S.seed· 1 object depends on it· depth 4

PRIOR ART: the closed sets of a closure operator form a Moore family — intersection-closed with a top. Moore, *Introduction to a Form of General Analysis* (1910); Ward, *The closure operators of a lattice*, *Ann. Math.* 43 (1942) 191-196. That the $\mathcal{R}$ -closed subsets of a closed index form one is that theorem applied at one level up.

F.open — the family theorem

THEOREM: the family of all closed indexes is itself an OPEN index — $E = 182, 365, 64,804$

every member has $E = 0$

grade **PROVED**· source *M §14.6; Tarski 1955; Ward 1942*· depends on F.moore· 2 objects depend on it· depth 5

PRIOR ART: the lattice of closed sets of a closure operator is complete (Tarski, *A lattice-theoretical fixpoint theorem*, *Pacific J. Math.* 5 (1955) 285-309), but completeness is not the same as being CLOSED under the same operator one level up. That the family of closed indexes is itself open is measured here; the setting is Tarski-Ward.

F.unstable — the unstable family

and it cannot be stated: a closed index has a seed, an open one costs seed + E, and E is 64,804 at 16 cells

corollary of F.open and S.open

grade **PROVED**· source *M §14.6.1; Kolmogorov 1965*· depends on F.open, S.open· depth 6

PRIOR ART: the cost of stating an object is its description length — Kolmogorov, Three approaches to the quantitative definition of information, Probl. Inf. Transm. 1 (1965) 1-7. A closed index has a seed; an open one costs seed plus defect, and the difference is the statement cost.

G. Graphs, constraints and languages — 12 objects

G.allcons — the index of all constraints

every genuine constraint in the book, indexed over five coordinates: 22 carried by $\mathcal{R}$, 2 by $\mathcal{R}_4$, NONE by neither; $E = 21$, $E_4 = 6$, orientation cost 15

eight rows of a first attempt were coordinate systems entered as constraints; the two that no operator carried were both of those

grade **COMPUTED**· source *M §21.5.5; Freuder 1978*· depends on A.r4, G.cons· depth 4

PRIOR ART: indexing constraints by the coordinates they name is the constraint-hypergraph view — Freuder, CACM 21 (1978) 958-966.

G.book — the book as an index

E(book) = 578 at chapter resolution, 0 at part resolution; a certificate exists claim-bearing paragraphs with a § citation

grade **COMPUTED**· source *M §30.1; Rota 1964*· depends on A.E, A.cert· depth 9

PRIOR ART: that a defect depends on the RESOLUTION at which an object is indexed is the coarsening question; Moebius inversion over a refinement lattice is Rota (1964).

G.cons — the constraint index

Λ 's seven constraints form a TREE — 8 nodes, 7 edges — with ZERO of 35 triples spanning three nodes

no 3-body among the constraints

grade **COMPUTED**· source *M §21.5.2; Freuder 1982; Dechter & Pearl 1989*· depends on L.c1· 5 objects depend on it· depth 1

PRIOR ART: that a tree-structured constraint graph is globally consistent after arc consistency is Freuder, A sufficient condition for backtrack-free search, J. ACM 29 (1982) 24-32; the width/induced-width machinery is Dechter & Pearl, Tree clustering for constraint networks, Artif. Intell. 38 (1989) 353-366.

G.graph — the constraint graph

Λ_{13} 's constraint graph: 12 nodes, 13 edges, girth 5, diameter 6, radius 3, treewidth 2, hub k at degree 4, cycle rank 2, no triangle at any stage

treewidth 2 at Λ_{13} keeps Freuder's bound and §21.5.1's one-level shortfall

grade **COMPUTED**· source *M §21.5.3 / Figure 21.1; Euler 1736; Diestel 2017*· depends on G.cons· depth 2

PRIOR ART: girth, diameter, radius and treewidth are standard graph invariants — see Diestel, *Graph Theory* (5th ed., 2017). Graph theory itself begins with Euler, *Solutio problematis ad geometriam situs pertinentis* (1736).

G.near — the seven near-misses

Λ is one edge from a 3-body in exactly seven places, one per existing edge
every one a bond the physics does not make

grade **COMPUTED**· source *M §21.5.2; Berge 1962*· depends on G.cons· depth 2

PRIOR ART: adding one edge to a tree creates exactly one cycle; the number of places is the number of existing edges. Berge, *Theorie des graphes* (1958/1962).

G.prot — the protocol index

the 24 protocols occupy 19 cells in four coordinates; §2.8 and §2.24 share one
§2.24 is §2.8 specialised to heuristics

grade **COMPUTED**· source *computed; Freuder 1978*· depends on G.reg· depth 5

PRIOR ART: as G.allcons, applied to the protocol index.

G.ref — the reference index

the reference index: 38 cells, box 144, $E = 0$, and NOT a tree — access closes a cycle

access, referent, checked, verdict

grade **COMPUTED**· source *M §16.6.1; Freuder 1982*· depends on A.E· depth 4

PRIOR ART: a non-tree constraint graph that nonetheless closes shows treeness is sufficient and not necessary — the converse direction of Freuder (1982).

G.reg — the register as an index

$E(\text{register}) = 6$ at 68.8% density, and R recovers $\text{repair} \leq \phi(\text{corroboration})$

coordinates assigned from the entry text

grade **COMPUTED**· source *M §28.9; Shannon 1948*· depends on A.E, A.R· 1 object depends on it· depth 4

PRIOR ART: recovery under a noisy channel is bounded by the channel capacity; that $\text{repair} \leq \phi(\text{corroboration})$ is that bound in this setting. Shannon (1948).

G.shape — shape decides the seed

the constraint graph's shape decides the seed: path 8, star 6, balanced trees 6, forest 5 on six nodes over one alphabet; neither extreme is cheapest

seed is monotone in S with no exception, all six exact by branch and bound

grade **COMPUTED**· source *M* §21.5.4; Freuder 1982; Dechter & Pearl 1989· depends on S .cover, G .cons· depth 5

PRIOR ART: that the constraint graph shape decides what a local method achieves is Freuder 1982 for trees and Dechter & Pearl 1989 for the general induced-width bound.

G.stat — statistics as a language

statistics is a sixth language: max-entropy closure on the pairwise marginals recovers Λ at 976, $E = 0$, restores a deleted cell, and GROWS on an added one, 976 to 1,048

its signature matches the information language, not order/geometry/analysis; the five split three-two between absorbing and growing

grade **COMPUTED**· source *M* §20; Deming & Stephan 1940; Csiszar 1975· depends on $A.R$ · depth 3

PRIOR ART: the max-entropy distribution matching given marginals is reached by iterative proportional fitting — Deming & Stephan, Ann. Math. Stat. 11 (1940) 427-444; its information-geometric characterisation is Csiszar, I-divergence geometry, Ann. Prob. 3 (1975) 146-158.

G.tower — the cycle rank of the tower

the tower's cycle rank rises by one at each two-parent axis — 0,0,1,1,2,2 — while triangles stay 0

the tree breaks at Λ_{10}

grade **COMPUTED**· source *M* §21.5.2; Berge 1962· depends on G .cons, T .tower· depth 7

PRIOR ART: the cycle rank $|E| - |V| + c$ is the first Betti number of a graph — Berge, *Theorie des graphes et ses applications* (1958/1962). That it rises by one at each two-parent axis is the measurement.

G.trip — independence is a triple property

independence is a property of TRIPLES: surviving tests go as $(1-f)^3$, not $(1-f)$

the bracket relates $T(n-1)$, $T(n)$, $T(n+1)$

grade **COMPUTED**· source *M* §24.9; Dawid 1979; Pearl 1988· depends on B .brk· depth 1

PRIOR ART: conditional independence is a relation on TRIPLES and obeys the graphoid axioms — Dawid, Conditional independence in statistical theory, J. R. Stat. Soc. B 41 (1979) 1-31; Pearl, Probabilistic Reasoning (1988), ch. 3.

L. Λ 's own constraints — 41 objects

L.E0 — the closure theorem

$E(\Lambda) = 0$

at the stated caps; verified at four settings and through the f shell

grade **COMPUTED**· source M §7.3, reg. 236; Moore 1910; Freuder 1982· depends on L.closed, A.E· depth 8

PRIOR ART: $E = 0$ says the index equals its own closure. Moore (1910) for the operator; Freuder (1982) for why a tree-structured constraint graph gives it.

L.F — the generating function

$$F = \sum_n z_1^n \sum_{\{\ell \leq n-1\}} z_2^\ell \sum_{\{k \leq 4\ell+2\}} z_3^k [\sum_{\{S \leq k\}} z_8^S] \sum_{\{q \leq k\}} z_4^q \sum_e z_5^e \sum_{\{f < e\}} z_6^f \sum_{\{g \leq \min(q, 4f+2)\}} z_7^g$$

at stated caps; the tree has no cycles so the sum factorises

grade **COMPUTED**· source M §11.3; Euler 1748; Stanley 1986· depends on L.tree, L.def· 5 objects depend on it· depth 3

PRIOR ART: a multivariate generating function over a constrained region, written as nested sums. Euler, Introductio (1748); Stanley, Enumerative Combinatorics Vol. 1 (1986), ch. 1.

L.F1 — the cardinality

$F(1, \dots, 1) = |\Lambda| = 976$

as L.F

grade **COMPUTED**· source M §11; Euler 1748; Stanley 1986· depends on L.F· depth 4

PRIOR ART: setting all variables to one recovers the cardinality — the elementary specialisation of a generating function.

PRIOR ART: $F(1, \dots, 1) = |X|$ is the elementary specialisation of a generating function. Euler, Introductio (1748), ch. XVI; Stanley, Enumerative Combinatorics Vol. 1 (1986), ch. 1.

L.Fm1 — the alternating specialisation

$F(-1) = 2$, because ℓ and f each have exactly two consecutive values but are coupled to n and e

at stated caps

grade **COMPUTED**· source *M §11.8.1; Euler 1748; Stanley 1986*· depends on L.rankpoly· depth 5

PRIOR ART: $F(-1)$ counts the difference between even- and odd-rank cells; the standard alternating specialisation.

PRIOR ART: $F(-1)$ is the difference between even- and odd-rank counts, the standard alternating specialisation. Euler (1748); Stanley (1986), ch. 3, where it is related to rank symmetry.

L.alpha — the join-irreducible count

$$|J(\Lambda)| = \sum_i (|A_i| - 1)$$

each generator is $\min\{x \in \Lambda : x_c \geq v\}$ for one coordinate and one value

grade **COMPUTED**· source *M reg. 268; Birkhoff 1937*· depends on L.birk· depth 10

PRIOR ART: the join-irreducibles of a product of chains are the coordinate steps, so $|J| = \sum (|A_i| - 1)$. Immediate from Birkhoff representation.

L.amp — the amplification

$$A(y) = N[\varphi(\Lambda \cup \{y\})] - N[\varphi(\Lambda)] - 1; \text{ median } 340, \text{ minimum } 59 \text{ over } 220 \text{ trials}$$

y a single inserted cell

grade **COMPUTED**· source *M §16.8.4; Rota 1964*· depends on A.R, L.def· depth 3

PRIOR ART: the change in a closure count on adding a generator; the amplification is measured, the closure is Moore-Rota.

L.arith — divisor lattice embedding

$$N(x) = \prod_i p_i^{x_i}; x \leq y \iff N(x) | N(y); v \mapsto \text{lcm}, \wedge \mapsto \text{gcd}, \text{rank} \mapsto \Omega(N)$$

p_i the *i*-th prime

grade **COMPUTED**· source *M §9; Birkhoff 1937*· depends on L.def· 2 objects depend on it· depth 2

PRIOR ART: the divisor lattice with lcm as join and gcd as meet is the classical arithmetic model of a product of chains. Birkhoff, *Lattice Theory* (1940), ch. II.

L.birk — Birkhoff representation 1937

$$\Lambda \cong \text{down-sets of } J(\Lambda); |J(\Lambda)| = 17, 20 \text{ covering relations}$$

Λ finite distributive

grade **COMPUTED**· source *M §8.3; Birkhoff 1937*· depends on L.dist· 5 objects depend on it· depth 9

PRIOR ART: Birkhoff's representation theorem, Duke Math. J. 3 (1937)
 — every finite distributive lattice is the down-sets of its join-irreducibles. $|J| = 17$ is a computation inside that theorem.

L.bits — the Boolean representation

Λ is the 976 words in $\{0,1\}^{17}$ that are down-sets of $J(\Lambda)$; join = OR, meet = AND; 17 bits carried, 9.93 needed, 7.07 surplus

Birkhoff correspondence

grade **COMPUTED**· source *M §11.1.1; Birkhoff 1937; Shannon 1938*· depends on L.birk· 1 object depends on it· depth 10

PRIOR ART: Birkhoff (1937) gives the down-set representation; encoding down-sets as Boolean words with OR as join and AND as meet is Shannon (1938).

L.box — the box factorises

$|\text{box} \cap \Lambda|$ factorises because the constraint graph is a tree; no Möbius sieve needed

constraint graph acyclic

grade **COMPUTED**· source *M §10.4; Freuder 1982; Rota 1964*· depends on L.tree· depth 3

PRIOR ART: the count factorises over a tree — no Moebius sieve is needed because there are no cycles to inclusion-exclude over. Freuder (1982) for the tree property; Rota (1964) for what the sieve would otherwise cost.

L.c1 — node counting

$$\ell \leq n - 1$$

hydrogenic radial solution

grade **CITED**· source *M §7.1; Bohr 1913; Schroedinger 1926*· 4 objects depend on it· depth 0

PRIOR ART: $l \leq n-1$ is the angular-momentum constraint of the hydrogen solution — Bohr, On the constitution of atoms and molecules, Phil. Mag. 26 (1913) 1-25; Schroedinger, Quantisierung als Eigenwertproblem, Ann. Phys. 79 (1926) 361-376.

L.c2 — Pauli

$$k \leq 2(2\ell+1) = 4\ell+2$$

Pauli exclusion

grade **CITED**· source *M §7.1; Pauli 1925; Stoner 1924*· 2 objects depend on it· depth 0

PRIOR ART: the subshell capacity $2(2l+1)$ is Stoner, The distribution of electrons among atomic levels, Phil. Mag. 48 (1924) 719-736, made exclusive by Pauli, Z. Phys. 31 (1925) 765-783.

L.c3 — the transfer bound

$$q \leq k$$

counting

grade **DEFINITIONAL**· source *M §7.1; Pauli 1925*· 2 objects depend on it· depth 0

PRIOR ART: $q \leq k$, a transferred count cannot exceed the occupancy. Pauli (1925).

L.c4 — node counting

$$f \leq e - 1$$

hydrogenic radial solution

grade **CITED**· source *M §7.1; Bohr 1913*· 2 objects depend on it· depth 0

PRIOR ART: as L.c1, applied to the second shell pair.

L.c5 — Pauli

$$g \leq 4f+2$$

Pauli exclusion

grade **CITED**· source *M §7.1; Stoner 1924; Pauli 1925*· 2 objects depend on it· depth 0

PRIOR ART: as L.c2, applied to the second shell pair.

L.c6 — the second transfer bound

$$g \leq q$$

counting

grade **DEFINITIONAL**· source *M §7.1; Pauli 1925*· 2 objects depend on it· depth 0

PRIOR ART: $g \leq q$, as L.c3 on the second shell pair.

L.c7 — vector coupling

$$2S \leq k$$

vector coupling on the source

grade **CITED**· source *M §7.1; Hund 1925; Pauli 1925*· 3 objects depend on it· depth 0

PRIOR ART: $2S \leq k$ because at most k electrons can align their spins — Pauli exclusion (1925) with Hund first rule, Z. Phys. 33 (1925) 345-371.

L.c8 — the occupancy floor

$k \geq 1$ (definitional restriction, not a bound)

a cell is a transition, not a state

grade **DEFINITIONAL**· source *M §10.2, reg. 301; Pauli 1925*· 2 objects depend on it· depth 0

PRIOR ART: $k \geq 1$ restricts to occupied subshells; definitional rather than a bound.

L.chains — the linear extensions

maximal chains of Λ = linear extensions of $J(\Lambda) = 1,113,045,672$

$\Lambda \cong J(P)$

grade **COMPUTED**· source *M §12.9; Stanley 1986*· depends on L.birk· depth 10

PRIOR ART: maximal chains of $J(P)$ correspond to linear extensions of P — Stanley, Enumerative Combinatorics Vol. 1, Prop. 3.5.2. The count is the computation; the bijection is his.

L.chi — the membership function

$\chi(x) = H(n-1-\ell)H(4\ell+2-k)H(k-q)H(k-2S)H(e-1-f)H(4f+2-g)H(q-g)$; the coefficient function of F is χ

H the Heaviside step

grade **COMPUTED**· source *M §11.1; Heaviside 1893*· depends on L.F, L.def· 1 object depends on it· depth 4

PRIOR ART: the membership function as a product of step functions — Heaviside, Electromagnetic Theory (1893), where the unit step is introduced.

L.circuit — the implication circuit

the 20 covering relations, as implications, cut $2^{17} = 131,072$ words to exactly 976, at depth 5

monotone: AND, OR, implication; no NOT, no feedback

grade **COMPUTED**· source *M §11.1.1; Shannon 1938*· depends on L.bits· depth 11

PRIOR ART: implications as a monotone Boolean circuit — Shannon, A symbolic analysis of relay and switching circuits, Trans. AIEE 57 (1938) 713-723.

L.closed — sublattice of a product

Λ is closed under coordinatewise $\vee$ and $\wedge$

every constraint of the form $x_i \leq \varphi(x_j)$ with φ non-decreasing

grade **PROVED**· source *M §7.3; Birkhoff 1940*· depends on L.def, A.rule· 2 objects depend on it· depth 7

PRIOR ART: closure under coordinatewise join and meet is the definition of a sublattice of a product (Birkhoff, Lattice Theory, 1940). What makes Λ one is that all eight constraints have the form $x_i \leq f(x_j)$ with f monotone.

L.def — the definition of Λ

$\Lambda = \{ (n, \ell, k, q, e, f, g, 2S) \in \mathbb{Z}^8 : \text{L.c1}.. \text{L.c8} \}, \text{caps } (n, e, \ell, k, f) = (3, 3, 1, 3, 1)$

caps stated; figures at other caps must say so (§7.4)

grade **DEFINITIONAL**· source *M §7; Bohr 1913; Pauli 1925*· depends on L.c1, L.c2, L.c3, L.c4, L.c5, L.c6, L.c7, L.c8· 10 objects depend on it· depth 1

PRIOR ART: the eight constraints are the shell-structure rules of Bohr (1913) and Pauli (1925), written as inequalities on integer coordinates.

L.dim — order dimension (Dushnik-Miller)

order dimension of Λ_8 is 8, rising by one per adjoined axis

at the stated caps

grade **PROVED**· source *M §8.6; Dushnik & Miller 1941*· depends on L.def· 1 object depends on it· depth 2

PRIOR ART: Dushnik & Miller, Partially ordered sets, Amer. J. Math. 63 (1941) 600-610. That a product of n chains has dimension n follows from the definition; what is measured here is that Λ 's dimension equals its coordinate count at every stage of the tower.

L.dist — Birkhoff / product of chains

Λ is distributive (a sublattice of a product of chains)

none

grade **PROVED**· source *M §8.1; Birkhoff 1937*· depends on L.closed· 5 objects depend on it· depth 8

PRIOR ART: a lattice is distributive iff it embeds in a product of chains (Birkhoff, Rings of sets, Duke Math. J. 3, 1937). Λ is such a sublattice by construction, so distributivity is INHERITED, not proved here.

L.metric — the lattice metric

$\log d$ is an ℓ^1 metric; $d(x, z) \leq d(x, y) + d(y, z)$

as L.occ

grade **COMPUTED**· source *M §9.2; Monjardet 1981*· depends on L.occ· depth 4

PRIOR ART: Monjardet, Metrics on partially ordered sets - a survey, Discrete Math. 35 (1981) 173-184.

L.mobius — Möbius function of a distributive lattice

$\mu(x,y) = (-1)^{|y \setminus x|}$ if $y \setminus x$ is an antichain in $J(\Lambda)$, else 0

$\Lambda \cong J(P)$

grade **COMPUTED**· source *M §9.3; Rota 1964*· depends on L.birk· depth 10

PRIOR ART: Rota, On the foundations of combinatorial theory I: theory of Moebius functions, Z. Wahrscheinlichkeitstheorie 2 (1964) 340-368. The antichain form of mu on a distributive lattice is his.

L.modular — modularity (equality, not submodularity)

$\text{rank}(a \vee b) + \text{rank}(a \wedge b) = \text{rank}(a) + \text{rank}(b)$, $\text{rank} = \sum x_i$

graded lattice

grade **COMPUTED**· source *M §8.2; Dedekind 1900*· depends on L.dist· depth 9

PRIOR ART: the modular rank identity holds in any modular lattice and every distributive lattice is modular. Dedekind, Math. Ann. 53 (1900).

L.occ — the interval measure

$d(x,y) = \tau(\text{lcm}(N(x), N(y)) / \text{gcd}(N(x), N(y))) = \prod_i (|x_i - y_i| + 1) = |[x \wedge y, x \vee y]|$

τ the divisor count; $d(x,x)=1$

grade **COMPUTED**· source *M §9.2; Monjardet 1981*· depends on L.arith· 3 objects depend on it· depth 3

PRIOR ART: the interval size as a product of coordinate spans, and its arithmetic form via lcm and gcd. Monjardet, Metrics on partially ordered sets, Discrete Math. 35 (1981) 173-184.

L.omega — distinct-prime-counting function

$\omega(N(x)) \leq \dim(\Lambda)$

one prime per coordinate

grade **PROVED**· source *M §9.1; Birkhoff 1937*· depends on L.arith, L.dim· depth 3

PRIOR ART: in the divisor representation, the number of distinct primes dividing $N(x)$ is the number of coordinates in which x is non-minimal, bounded by the dimension.

L.pal — palindromic rank polynomial $\Leftrightarrow$ self-dual

F palindromic $\Leftrightarrow$ the poset is self-dual; Λ 's F is not palindromic

graded poset

grade **PROVED**· source *M §11.8; Stanley 1986*· depends on L.rankpoly, L.skew· depth 5

PRIOR ART: Stanley, Enumerative Combinatorics Vol. 1, ch. 3 — a graded poset's rank polynomial is palindromic iff it is rank-symmetric.

L.pushback — join-prime and meet-prime

no cell of Λ_8 is both join-prime and meet-prime, so pushback $\geq 16 > 0$ for every cell

18 join-irreducibles and 18 meet-irreducibles, intersection empty

grade **PROVED**· source *M §16.8.5; Birkhoff 1937; Gratzner 1978*· depends on L.birk· 1 object depends on it· depth 10

PRIOR ART: join-prime and meet-prime are dual notions in a distributive lattice, and a cell cannot generally be both. Gratzner, General Lattice Theory (1978).

L.rankpoly — the rank polynomial

$F(z) = z^3(z^{17} + 4z^{16} + 10z^{15} + \dots + 122z^8 + 121z^7 + \dots + 5z + 1)$; $F'(1)/F(1) = 11.0666$

single-variable specialisation

grade **COMPUTED**· source *M §11.1; Stanley 1986*· depends on L.F· 2 objects depend on it· depth 4

PRIOR ART: Stanley, Enumerative Combinatorics Vol. 1 (1986), ch. 3. $F'(1)/F(1)$ = mean rank is the standard first-moment identity.

L.real — the index against the real subshells

Λ 's eight constraints on 247 real subshells across ALL 118 elements incl. 19 anomalies and 15 predicted superheavies: 18,288 tests, 0 failures

IUPAC ground states

grade **COMPUTED**· source *computed; Madelung 1936; NIST*· depends on L.c1· depth 1

PRIOR ART: the 247 real subshells across 118 elements are the observed ground configurations, tabulated by NIST; the ordering is Janet-Madelung. Testing Λ constraints against them checks the index, not the table.

L.skew — the rank skew

centre of mass 11.0666 vs midpoint 11.5, skew -0.43 ; only 8 of 976 cells fixed by $x \mapsto \max - x$

at the stated caps

grade **COMPUTED**· source *M §8.4; Gauss 1809*· depends on L.def· 1 object depends on it· depth 2

PRIOR ART: the third standardised moment of a rank distribution. Gauss (1809).

L.sperner — Sperner property / Dilworth

the largest antichain equals the largest rank level; max 122 at rank 11

rank sequence log-concave hence unimodal

grade **COMPUTED**· source *M §8.4; Sperner 1928; Stanley 1980*· depends on L.dist· depth 9

PRIOR ART: Sperner 1928 for the Boolean lattice; Stanley, Weyl groups, the hard Lefschetz theorem, and the Sperner property, SIAM J. Alg. Disc. Meth. 1 (1980) 168-184, proves the order-ideal lattice of a product of chains is Peck — rank-symmetric, rank-unimodal, strongly Sperner. Λ is of that form, so the property is INHERITED.

L.step — the interval-removal step

$X \setminus [a, b]$ is a sublattice iff a is join-prime and b is meet-prime; $\text{step}(\Lambda_s) = 4$

distributive

grade **PROVED**· source *M §16.8.5; Birkhoff 1937; Gratzner 1978*· depends on L.pushback· depth 11

PRIOR ART: removing an interval leaves a sublattice iff the endpoints are join-prime and meet-prime — the standard interval-removal criterion. Gratzner, General Lattice Theory (1978), ch. II.

L.total — the membership function is total

$\chi_\Lambda : \prod A_i \rightarrow \{0, 1\}$ is total: membership decided for every ambient point

Λ a finite intersection of decidable comparisons

grade **PROVED**· source *M §16.5; Birkhoff 1940*· depends on L.chi· depth 5

PRIOR ART: a membership function defined by inequalities on coordinates is total on the ambient product by construction.

L.tree — the constraint tree

constraint graph $n-\ell-k-q-g-f-e$ with 2S pendant at k : 8 nodes, 7 edges, a caterpillar; treewidth 1

every constraint binds exactly two coordinates

grade **COMPUTED**· source *M §8.5, §11.4; Freuder 1982*· depends on L.def· 5 objects depend on it· depth 2

PRIOR ART: that a constraint graph is a tree, and what follows from it, is Freuder, A sufficient condition for backtrack-free search, J. ACM 29 (1982) 24-32.

L.void — the void

void(x,y) = $||_i(|\Delta_i|+1) - |[x\wedge y, x\vee y] \cap \Lambda|$

none

grade **DEFINITIONAL**· source *M §10; Rota 1964*· depends on L.occ· 1 object depends on it· depth 4

PRIOR ART: the difference between a box count and the cells it actually contains is what Moebius inversion would compute. Rota (1964).

L.voidfrac — the void-free fraction

void-free fraction 27.7-30.1% across 776M pairs; joint 30.13% vs product 20.19%, factor 1.49

at stated caps

grade **COMPUTED**· source *M §10.2; Rota 1964*· depends on L.void· depth 5

PRIOR ART: as L.void; the fraction is the measurement.

T. The tower and its couplings — 16 objects

T.a10 — Racah 1943 seniority

$2S' \leq v \leq g, v \equiv g \pmod{2}$

Racah seniority for ℓ^N

grade **CITED**· source *M §12.11.1 / T A14; Racah 1943*· depends on T.a9· 2 objects depend on it· depth 2

PRIOR ART: seniority v , with $2S \leq v \leq g$ and v congruent to $g \pmod{2}$, is Racah, Phys. Rev. 63 (1943) 367-382 — the classification of states of ℓ^n by the number of unpaired electrons.

T.a11 — the parent bound

$2J_c \leq \hat{\phi}(k), \hat{\phi} = \max 2J$ over terms of $\ell^k = \{1:3, 2:4, 3:5\}$

$\hat{\phi}$ read off the realised extent, not from a law

grade **COMPUTED**· source *M §12.11.1 / T A15; Condon & Shortley 1935; Racah 1942*· depends on T.a10· 2 objects depend on it· depth 3

PRIOR ART: the term structure of a subshell ℓ^k , and the maximum J it carries, is Condon & Shortley, The Theory of Atomic Spectra (1935), ch. VII; Racah, Theory of complex spectra II, Phys. Rev. 62 (1942) 438-

462, gives the general classification. The values {1:3, 2:4, 3:5} are read from that table, not derived here.

T.a12 — the recoupling bound

loose: $2K \leq 2J_c + 2f_{\max}$ (one parent). **exact:** $|2J_c - 2f| \leq 2K \leq 2J_c + 2f$ step 2 (two parents)

jK pair coupling

grade **CITED**· source *M §12.11.1 / T A16; Wigner 1931; Racah 1942*· depends on T.a11· 3 objects depend on it· depth 4

PRIOR ART: the triangle condition $|j_1 - j_2| \leq J \leq j_1 + j_2$ in steps of one is the Clebsch-Gordan series — Wigner, *Gruppentheorie* (1931); its application across parents is Racah, *Phys. Rev.* 62 (1942) 438-462.

T.a13 — the spin-half bound

$|2J - 2K| \leq 1$

outer electron carries spin 1/2

grade **CITED**· source *M §12.11.1 / T A17; Condon & Shortley 1935*· depends on T.a12· 1 object depends on it· depth 5

PRIOR ART: $|2J - 2K| \leq 1$ is the coupling of a spin-half to K in the jK scheme; Condon & Shortley (1935), ch. X.

T.a9 — the seniority floor

$2S' \leq g$

vector coupling on the target

grade **CITED**· source *M §12.10.1; Racah 1943*· depends on L.c7· 5 objects depend on it· depth 1

PRIOR ART: $2S \leq g$ bounds the total spin by the occupancy — Pauli exclusion in Racah seniority form, *Theory of complex spectra III*, *Phys. Rev.* 63 (1943) 367-382.

T.a9p — the admissible Pauli cut

$2S' \leq 2f+1$ (the admissible Pauli cut); $\Lambda_g' = 1,561$ cells, 93 removed; closes the cycle f-g-2S'

$\min(g, 4f+2-g) \leq 2f+1$ by averaging

grade **COMPUTED**· source *M §12.11.1 / T 2.2 (App. G); Pauli 1925; Racah 1943*· depends on T.a9· depth 2

PRIOR ART: the Pauli cut on a subshell's allowed terms is Pauli's exclusion principle (*Z. Phys.* 31, 1925) as applied to equivalent electrons; the seniority classification that makes it computable is Racah, *Theory of complex spectra III*, *Phys. Rev.* 63 (1943) 367-382.

T.dens — the axis density

density(axis) = $\Sigma_{\text{parents}} |\text{exact fibre}| / \Sigma_{\text{parents}} |\text{admissible fibre}| = 63.7, 67.5, 44.7, 17.0, 31.4, 64.4\%$

three non-obvious exact sets required

grade **COMPUTED**· source *M §12.11.1 / T A18; Racah 1942; Wigner 1931*· depends on T.tower· 2 objects depend on it· depth 7

PRIOR ART: the ratio of exact to admissible fibre sizes measures how much a triangle condition tightens a Pauli bound. Both bounds are standard; the ratio is measured.

PRIOR ART: the ratio of exact fibre to admissible fibre measures how much the triangle condition (Wigner 1931, Clebsch-Gordan) tightens the Pauli cap (Racah 1942). Both bounds are theirs; the density is measured.

T.dich — the counting-coupling dichotomy

counting coordinates close exactly; coupling coordinates close as envelopes; no third kind

the exact coupling bound is made of reflection, congruence and triangle

grade **COMPUTED**· source *M §12.11.3; Racah 1942*· depends on T.dens· 1 object depends on it· depth 8

PRIOR ART: the split between coordinates that COUNT (occupancy, capacity) and coordinates that COUPLE (J, K, S) is the organising distinction of Racah's algebra. Counting coordinates carry Pauli caps; coupling coordinates carry triangle conditions, and the two close differently.

T.excl — the three excluded forms

three excluded forms: REFLECTION (particle-hole conjugation), CONGRUENCE (fermion parity), TRIANGLE (one sum and one difference)

each is non-monotone or multi-parent

grade **COMPUTED**· source *M §12.11.2; Racah 1943; Wigner 1931*· depends on T.dich, A.rule· 2 objects depend on it· depth 9

PRIOR ART: particle-hole conjugation, fermion parity and the triangle rule are all standard: conjugation from the complementary-shell theorem (Racah 1943), parity from the antisymmetry of the wavefunction, and the triangle inequality $|j_1 - j_2| \leq J \leq j_1 + j_2$ from the Clebsch-Gordan series (Wigner 1931).

T.invariant — basis independence of the level set

the exact J multiset is identical in jK, LS, LK and jj; the four cell counts price four ROUTES

coupling schemes are basis changes

grade **COMPUTED**· source *M §12.11.4; Wigner 1931; Racah 1942*· depends on T.scheme, T.dens· depth 8

PRIOR ART: the J multiset of a configuration is independent of coupling scheme because the schemes are unitary recouplings of one space — Wigner, *Gruppentheorie* (1931); Racah's 6-j and 9-j coefficients are the transformation matrices. What is measured here is the CELL COUNT each scheme costs, not the invariance.

T.para — parastatistics

capacity(ℓ) = $m(4\ell+2)$ for parastatistics of order m ; $E = 0$ through Λ_{13} for $m = 1, 2, 3$

$m(4\ell+2)$ monotone in ℓ for every m , so A.rule is preserved

grade **COMPUTED**· source *T 2.1 (App. G) / D3; Green 1953*· depends on T.tower, A.rule· depth 7

PRIOR ART: parastatistics of order m , in which a state holds up to m particles, is H. S. Green, *A generalized method of field quantization*, *Phys. Rev.* 90 (1953) 270-273. The capacity $m(4\ell+2)$ is its shell-model form.

T.real — the bounds against real terms

the four coupling bounds on the exact term structure of every subshell: 85,829 tests, 0 failures

$\ell = 0..3, k = 1..4\ell+2$

grade **COMPUTED**· source *microstate enumeration; Condon & Shortley 1935*· depends on T.a9· depth 2

PRIOR ART: the exact term structure of every subshell is tabulated in Condon & Shortley (1935), ch. VII, and in the NIST compendium. The 85,829 tests check the four coupling bounds AGAINST that table; the table is not this work.

T.scheme — the four coupling schemes

closure holds in LS, LK, jK, jj: $\Lambda_{13} = 431,050 / 341,150 / 199,130 / 206,520$, $E = 0$ throughout

a uniform one-parameter looseness convention

grade **COMPUTED**· source *T 2.1b (App. G); Condon & Shortley 1935; Racah 1942*· depends on T.tower· 1 object depends on it· depth 7

NOT CHECKABLE and the book says why: the looseness convention is named and not printed, and two committed readings BRACKET rather than determine the figures — LS in [383,065, 597,325] contains 431,050, jj in [160,380, 244,060] contains 206,520 (R 1010)

PRIOR ART: LS, LK, jK and jj are the four standard angular-momentum coupling schemes — Condon & Shortley (1935), ch. X; the jK and LK intermediate schemes are Racah's. That the same physical states are counted in each is the content of the recoupling theory, and the four cell counts price the SCHEME rather than the physics.

T.tight — constraint tightness as a count

the tight-pair count of a base is exactly 2S, S the envelope-step count

van Beek & Dechter's measure

grade **COMPUTED**· source *M §14.5.9; Deville et al. 1999; Deville, Barette & Van Hentenryck 1999*· depends on A.env· depth 2

PRIOR ART: the tight pairs of a staircase constraint are its envelope steps — Deville, Barette & Van Hentenryck, Artif. Intell. 109 (1999) 243-271.

T.tower — the tower

$\Lambda_8 \dots \Lambda_{13} = 976, 1654, 2535, 13585, 22275, 64290$; E = 0 at every stage, swept against every ambient cell to 47,775,744

at stated caps

grade **COMPUTED**· source *M §12.11.0.10, reg. 249; Racah 1942; Condon & Shortley 1935; Racah 1942, 1943*· depends on T.a9, T.a10, T.a11, T.a12, T.a13· 5 objects depend on it· depth 6

CORRECTED 2026-08-11, re-applying register 1399 whose source fix was LOST in the 1.6.1 rollback — the register crossed as prose, the code did not. This object STATED 70905, 199130 at Λ_{12} and Λ_{13} while the exact triangle GIVES 22275, 64290. The first four agreed; only the last two differed, and register 541 named the cause without connecting it: the exact triangle against the loose one-parent bound differ by 3.18 at Λ_{12} . A sequence is a single claim and its terms must come from one convention, so the computed (exact-triangle) values now stand. Note also that 199,130 was simultaneously T.scheme's jK figure — one number doing two jobs in two objects with neither saying so.

PRIOR ART: the tower adjoins the coupling quantum numbers in the standard order — seniority, then J of the core, then K, then J. Each is Racah or Condon & Shortley; the cell counts and the closure at each stage are the measurement.

T.trad — the tree or the tightness

the tree or the tightness: imposing the exact triangle at axis 12 gives 22,275 cells and E = 35,570

the exact bound has two parents

grade **COMPUTED**· source *M §12.11.5 / T 2.4 (App. G); Freuder 1982; Racah 1942*· depends on T.a12, A.rule· depth 7

PRIOR ART: the trade between tree structure and constraint tightness is Freuder (1982) for the tree side; the exact triangle is Racah/Wigner. The pricing is the measurement.

E. Empirical indexes — table, nuclide, layout — 4 objects

E.ioniz — the refusal map

first ionisation energies given to the bracket as bare numbers refuse at Be→B, N→O, Mg→Al and P→S — four steps of twelve, 67% admissible

the same two positions in both periods: the s2-p1 subshell opening and the p3-p4 first pairing

grade **COMPUTED**· source *M §6.2; Mendeleev 1869; NIST*· depends on B.brk· depth 1

PRIOR ART: first ionisation energies as a periodic property date to Mendeleev; the values are NISTs. That bare numbers refuse the bracket is a statement about what a bracket needs, not about the data.

E.layout — the price of a layout

hydrogen's placement is free and helium's costs 16, and they are not additive: -16 and 0 apart, -1 together

E set by the largest group used in period 1

grade **COMPUTED**· source *M §6.1.1; Janet 1929*· depends on E.table· depth 3

PRIOR ART: hydrogen and helium are the classic placement anomalies of the periodic table, and Janets left-step form resolves them differently from the classroom table. The cost measured here is of the CHOICE.

E.nuclide — the nuclide chart's defect

the measured nuclide chart indexed by (Z, N) closes at E = 9, stable across four proton-number cutoffs; the nine cells are the mass formula's pairing and clustering terms

cells named at one cutoff persist at every larger one

grade **COMPUTED**· source *M §6.2; Segre 1945*· depends on A.env· depth 2

*** COUNT NOT REPRODUCED 2026-08-11 (R 1550). *** AME2020 Table I was captured from the published paper (Chin. Phys. C 45, 030003, Table I, 3558 rows, Z = 0-118, A = 1-295) and E recomputed with the same operator: E = 2, NOT 9, and stable at 2 across cutoffs $Z \leq 20, 50, 82, 92$ and 118. Six variants of the cell set were tried - axes swapped, neutron excluded, measured-only, Z and N both positive - and none gives 9; measured-only gives 95. WHAT DOES REPRODUCE: the STABILITY across cutoffs, the cells being NAMEABLE PHYSICS, and the reading of them as the mass formula pairing term - the two defect cells are the empty cell (0,0) and Z=2 N=0, THE DIPROTON, which is unbound and is the textbook pairing failure. Three of four claims hold and the number does not. Cause UNDETERMINED: a different source edition, a different inclusion rule, or an arithmetic error in the original.

PRIOR ART: the chart of nuclides indexed by (Z, N) is Segres chart, in use since the 1940s.

E.table — the thirty-six decomposed

the periodic table's 36 decompose 25 + 11, not 26 + 10; E is placement-sensitive, 36 at group 18 and 20 at group 2

all 118 elements

grade **COMPUTED**· source *M §6.1.1; Mendeleev 1869; Janet 1929*· depends on A.env· 1 object depends on it· depth 2

PRIOR ART: the periodic tables arrangement is Mendeleev, *Über die Beziehungen der Eigenschaften zu den Atomgewichten der Elemente*, *Z. Chem.* 12 (1869) 405-406; the left-step form on n+1 is Janet (1929). That E is placement-sensitive is a statement about which arrangement, not about the elements.

B. The bracket — 21 objects

B.V — the price V — the bracket's width against its error

$V = w/e$ where $w = |T(n+1) - T(n-1)|$, $e = |T(n) - \frac{1}{2}(T(n-1) + T(n+1))|$

three consecutive members

grade **DEFINITIONAL**· source *M §21.1; Milne-Thomson 1933*· depends on B.brk· 3 objects depend on it· depth 1

PRIOR ART: the ratio of a first difference to a second is the standard curvature-to-slope measure of a finite-difference scheme. Milne-Thomson, *The Calculus of Finite Differences* (1933), ch. I-II.

B.V43 — the four-thirds law

$V = 4\nu/3$ for a Rydberg series; in general $V(x,p) = 4x/(h|p-1|)$ for $y = x^p$

$T = Z^2R/\nu^2$

grade **COMPUTED**· source *M §21.1; Rydberg 1890*· depends on B.V· 3 objects depend on it· depth 2

PRIOR ART: for a Rydberg series $T \sim \nu^{-2}$, so the ratio of first to second differences is $4\nu/3$ by direct expansion. Rydberg, *Recherches sur la constitution des spectres d emission*, *K. Sven. Vetensk. Akad. Handl.* 23 (1890).

B.Vexact — the exact price

$V = 4\nu^3/(h(3\nu^2 - h^2))$; with $r = \nu/h$, $V = 4r^3/(3r^2 - 1)$. Z and R cancel identically

step h free

grade **PROVED**· source *M §21.4; Rydberg 1890*· depends on B.V43· 2 objects depend on it· depth 3

PRIOR ART: the exact form follows from the Rydberg term $T = Z^2 R / \nu^2$ by finite differencing; Z and R cancel because the ratio is scale-free.

B.adm — the five-sigma admissibility threshold — where the bracket may be applied at all

$$r = 2Z^2R/(\nu^3\sigma) \geq 5$$

levels separated by more than 5σ

grade **DEFINITIONAL**· source *M §22.5; Gauss 1809*· 1 object depends on it· depth 0

PRIOR ART: a five-sigma admissibility threshold is a statement about signal against measurement error, in the Gaussian error model of Gauss, *Theoria motus corporum coelestium* (1809).

B.aitken — Aitken Δ^2 / Seki Kōwa

Aitken's Δ^2 on a Rydberg series lands at $I - T/3$, because the correction is $(\Delta T)^2/\Delta^2 T \rightarrow (2/3)T$

algebraic not geometric convergence

grade **PROVED**· source *M §24.6; Aitken 1926*· depends on B.newton· depth 3

PRIOR ART: Aitken, On Bernoulli's numerical solution of algebraic equations, *Proc. R. Soc. Edinb.* 46 (1926) 289-305 — the delta-squared process. That it lands at $I - T/3$ on a Rydberg series is the measurement; the process is his.

B.brk — monotone interpolation bracket

$T(n)$ lies between $T(n-1)$ and $T(n+1)$

T monotone in n within a channel

grade **PROVED**· source *M §20.1; Leibniz 1682; Milne-Thomson 1933*· 6 objects depend on it· depth 0

measured at 789 of 789 interior cells — and it is the TRIVIAL bracket, which cannot fail (R 797, 830)

PRIOR ART: that a term of an alternating or monotone sequence lies between its neighbours is the classical bracketing of finite differences — Leibniz's alternating-series test in its difference form; Milne-Thomson, *The Calculus of Finite Differences* (1933), ch. I. The book's contribution is applying it to Rydberg terms cell by cell, not the bracket.

B.coll — the assembly rule — a composite observable's scaling exponent is $2a + 3b$

an observable $\langle r \rangle^{a/(\Delta E)} b$ scales as $\nu^{\{2a+3b\}}$

$$\langle r \rangle \propto \nu^2, \Delta E \propto \nu^{-3}$$

grade **PROVED**· source *M §24.1; Bohr 1913; Bethe & Salpeter 1957*· 1 object depends on it· depth 0

PRIOR ART: the scaling of hydrogenic expectation values, $\langle r^a \rangle \sim n^{2a}$ and $\Delta E \sim n^{-3}$, is Bohr's correspondence scaling (1913) in its quantum form; tabulated in Bethe & Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms* (1957), sec. 3.

B.fail — when the bracket fails

the bracket fails $\Leftrightarrow |\Delta T| > 2Z^2R/n^3$

a perturber reorders only if the shift exceeds half the local spacing

grade **PROVED**· source *M §23.3; Rydberg 1890*· depends on B.brk· 1 object depends on it· depth 1

the yardstick verified: $2Z^{2R/n^3}$ against measured adjacent spacing over 742 pairs gives median 1.201 (R 832)

PRIOR ART: the local spacing of a Rydberg series is $2Z^2R/n^3$, the derivative of the term formula. A shift exceeding half of it reorders the levels.

B.floor2 — the floor at two

$V > 2$ for any monotone sequence; $V \rightarrow 2$ only as one step vanishes

$d_0, d_1 > 0$; $V = 2(d_0 + d_1)/|d_0 - d_1|$

grade **PROVED**· source *M §21.2 Prop 14.1; Jensen 1906*· depends on B.V· 1 object depends on it· depth 2

PRIOR ART: $V > 2$ for a monotone sequence is a form of Jensen's inequality — the chord lies above the curve for a convex function, so the second difference cannot exceed half the first. Jensen, *Sur les fonctions convexes*, *Acta Math.* 30 (1906) 175-193.

B.floor32 — the Rydberg floor

$V \geq 32/11 = 2.909$ for a Rydberg series specifically

hydrogenic form

grade **COMPUTED**· source *M §21.2; Rydberg 1890; Jensen 1906*· depends on B.floor2, B.V43· depth 3

PRIOR ART: the floor $V > 2$ is Jensen convexity (1906); the sharper $32/11$ for a Rydberg series follows from its specific n^{-2} form.

B.frac — the fractional widths

$w/T = 4(h/\nu) + 8(h/\nu)^3$ and $e/T = 3(h/\nu)^2$; their ratio is V

Rydberg

grade **COMPUTED**· source *M §21.6; Taylor 1715*· depends on B.Vexact· depth 4

PRIOR ART: expanding the first and second differences of ν^{-2} in powers of h/ν is Taylor series with a step. Taylor, Methodus incrementorum directa et inversa (1715).

B.hstar — the optimal step

$h^* = \sqrt{(2\beta/(\alpha y''))}$ minimises $\alpha w + \beta V$, valid for $h^* \ll \nu$

leading order

grade **COMPUTED**· source M §21.5.3; Lagrange 1797; Curtis & Reid 1974· depends on B.pareto· depth 5

NOT CHECKABLE here: its source M §21.5.3 does not appear in the book's text and the objective it minimises is not stated, so the stationary point cannot be verified (R 1010)

PRIOR ART: minimising a weighted sum of two competing costs by setting the derivative to zero is elementary optimisation; the $h^* \sim \sqrt{(2\beta/(\alpha y''))}$ form is the standard step-size optimum of finite-difference practice.

PRIOR ART: the optimal finite-difference step balances truncation error against roundoff, giving h^* proportional to $\sqrt{(\epsilon/y)}$ — Curtis & Reid, The choice of step lengths when using differences to approximate Jacobian matrices, J. Inst. Math. Appl. 13 (1974) 121-126. Here the two costs are width and price rather than truncation and roundoff, and the same balance applies.

B.newton — Newton decrement, Nesterov-Nemirovskii 1994

$8y''/y' = 8\lambda^2$ with $\lambda^2 = \nabla f^T [\nabla^2 f]^{-1} \nabla f$; for a Rydberg series $\lambda^2 = (2/3)T$

one dimension: $\lambda^2 = f^2/f''$

grade **PROVED**· source M §23.8.1; Newton 1669; Nesterov & Nemirovskii 1994· depends on B.V· 2 objects depend on it· depth 2

PRIOR ART: the Newton decrement $\lambda^2 = \text{grad } f^T [\text{Hess } f]^{-1} \text{grad } f$ is the standard measure of proximity to a minimum in interior-point theory (Nesterov & Nemirovskii 1994); Newton's method itself is De analysi (1669).

B.nuV — the value-one crossing — where the bracket's price V passes unity

$\nu_V = (3Z^2R/5q)^{1/4}$

curvature resolvable at quotation granularity q

grade **PROVED**· source M §23.15; Rydberg 1890· depth 0

the crossing where V passes unity, computed from the quotation granularity q

PRIOR ART: solving $V = 1$ for ν against a quotation granularity q gives the quartic root; the term formula is Rydberg (1890).

B.ordbr — the ordered bracket

sign($f(n) - p(n)$) = $(-1)^{\{k+1\}(-1)^m}$ with m nodes above n ; a two-sided deductive bracket at every order

$$\text{sign}(f^{\wedge\{j\}}) = (-1)^j$$

grade **PROVED**· source *M §23.10.1; Newton 1687; Milne-Thomson 1933*· depends on B.brk· depth 1

PRIOR ART: the sign of the error of a Newton interpolating polynomial alternates with the number of nodes above the evaluation point — the standard remainder theorem for finite differences, Newton's divided-difference form (*Principia*, Book III, Lemma V); Milne-Thomson (1933), ch. VIII.

B.ordk — the admissible order

order k admissible while $|\Delta^{\{k+1\}}T| > 5 \cdot 2^{\{k+1\}} \cdot \sigma$

the $(k+1)$ -th difference is a signed sum of $2^{\{k+1\}}$ levels

grade **PROVED**· source *M §23.10.4; Milne-Thomson 1933*· depends on B.adm· depth 1

PRIOR ART: the admissible order of a difference scheme is set by where the next difference falls below the noise, and the $2^{\{k+1\}}$ growth of noise under k -fold differencing is standard. Milne-Thomson (1933), ch. II.

B.pareto — the width-price front

$dw/dh > 0$ and $dV/dh < 0$ for every monotone convex y and every h ; the family $\{(w,V)\}$ is a Pareto frontier

$$w \sim h, e \sim h^2$$

grade **PROVED**· source *M §21.5.2; Pareto 1896*· depends on B.Vexact· 1 object depends on it· depth 4

PRIOR ART: a family in which no member improves both objectives is a **PARETO FRONT** — Pareto, *Cours d'economie politique* (1896). That $dw/dh > 0$ and $dV/dh < 0$ for every monotone convex y makes $\{(w,V)\}$ one is the proof here; the notion is his.

B.pole — the linear pole

V has a pole at $p = 1$: a linear observable has no curvature to price

$$y = x^p$$

grade **PROVED**· source *M §18.5; classical; Taylor 1715; Milne-Thomson 1933*· depends on B.V43· depth 3

PRIOR ART: a linear function has vanishing second difference, so any ratio measuring curvature against slope diverges there. The pole at $p = 1$ is that statement.

B.rank1 — the rank-one factorisation

rank(log q) = 1 across fifteen Rydberg observables; second singular value 1.8×10^{-14}

every observable is ν to a power

grade **COMPUTED**· source *M §24.2; Eckart & Young 1936*· depends on B.coll· depth 1

PRIOR ART: that a matrix has rank one is read from its singular values — Eckart & Young, The approximation of one matrix by another of lower rank, *Psychometrika* 1 (1936) 211-218. A second singular value of $1.8e-14$ is numerical zero, so log q factorises exactly.

B.selfconc — self-concordance

T(ν) is self-concordant, $|f'''| \leq 2(f'')^{\{3/2\}}$, for $\nu \leq (\sqrt{6}/2) \cdot Z\sqrt{R}$

$T = Z^2 R / \nu^2$

grade **PROVED**· source *M §23.8.2; Nesterov & Nemirovskii 1994*· depends on B.newton· depth 3

PRIOR ART: self-concordance, $|f'''| \leq 2(f'')^{\{3/2\}}$, is Nesterov & Nemirovskii, Interior-Point Polynomial Algorithms in Convex Programming (1994), the condition under which Newton's method converges at a rate independent of the problem's conditioning.

B.silence — what silence implies

the bracket holding $\Rightarrow |\Delta T| < 2Z^2 R / \nu^3$ at that cell

contrapositive of B.fail

grade **PROVED**· source *M §23.5; Rydberg 1890*· depends on B.fail· depth 2

849 of 849 cells satisfy it; median $|dT|/\text{spacing}$ 0.0015, largest 0.25 (R 831)

PRIOR ART: the contrapositive of B.fail, on the same term formula.

K. Transit and information — 24 objects

K.arrow — the arrow index

$X_w = \{a : w(\text{tgt } a) \leq w(\text{src } a)\}$ is closed; one arrow per coordinate and no others; sums, max and min all fail

w a single coordinate

grade **COMPUTED**· source *M §12.11.0.2; Birkhoff 1937*· depends on K.clockfail, A.rule· depth 8

PRIOR ART: a set of the form $\{a : w(\text{target}) \leq w(\text{source})\}$ for monotone w is a down-set of the induced order, hence closed. Birkhoff, Rings of sets (1937).

K.axis — the axis test

an axis earns a COORDINATE when it is independent of the others AND its distinct values are few relative to what it adds; both conditions are necessary

a closed index and a candidate axis

grade **MEASURED**· source *R 1127-1131; Dushnik & Miller 1941; Shannon 1948*· depends on K.redun· 1 object depends on it· depth 4

on the same elements, $Z \leq 16$: (Z,charge,l) gives 0% redundancy, +multiplicity gives 82%, +2J gives 0% again. 2J is NOT derived — no combination determines it — but $|L-S| \leq J \leq L+S$ constrains it to $2\min(l,S)+1$ values of seventeen. A coordinate must be free, not merely undetermined. The entropy account is PARTIAL: retained entropy is 26% for multiplicity and 27% for 2J, which does not separate them

PRIOR ART: an axis earns a coordinate when it is order-independent of the others (Dushnik & Miller 1941) and carries information (Shannon 1948). The test combines both.

K.cat — the tower is a category

Λ_9 is a category: 41,682 composable pairs, zero failures, associative

as K.comp

grade **COMPUTED**· source *M §12.11.0; Mac Lane 1971*· depends on K.comp· 2 objects depend on it· depth 3

PRIOR ART: associativity of composition and the existence of identities are the category axioms — Mac Lane, Categories for the Working Mathematician (1971). Verifying them on 41,682 composable pairs checks that Λ_9 IS a category; the axioms are not this work.

K.clock — the composition clock

occupancy never rises along composition: 0 of 739 steps; the tick is $k - g = (k-q) + (q-g)$

destinations begin empty — an assumption, not a constraint

grade **COMPUTED**· source *M §12.11.0.1; classical; Floyd 1967*· depends on K.comp· 1 object depends on it· depth 3

PRIOR ART: a quantity that never increases along composition is a MONOVARIANT, the standard tool for proving termination. That occupancy is one here is the measurement; the technique is old.

K.clockfail — the clock that does not tick

index (g,G) with $g \leq G \leq 4f+2$ and $g \leq q$: 13,775 cells, closed, and 31.6% of composable pairs raise occupancy

prior occupancy carried as a coordinate

grade **COMPUTED**· source *M §12.11.0.2; Floyd 1967*· depends on K.clock, A.rule· 1 object depends on it· depth 7

PRIOR ART: a candidate variant that is closed but does not decrease is a failed termination measure. Floyd (1967).

K.comp — composition

b•a defined when $\text{tgt}(a) = \text{src}(b)$; the composite transfers $\min(q_a, q_b)$

Λ_9 's target $(e, f, g, 2S')$ has the source's constraint forms

grade **COMPUTED**· source *M §12.11.0; Mac Lane 1971*· depends on T.a9· 6 objects depend on it· depth 2

PRIOR ART: composition defined when target meets source is the category axiom; what is measured is that the composite transfers $\min(q_a, q_b)$, which is a property of this index and not of categories.

K.corner — corners and faces

the whole-object defect decomposes into FACES and CORNERS, and the corners are an artefact of loose coordinatisation

an index of dimension d and its $d-1$ axis slices

grade **MEASURED**· source *R 1177; Freuder 1978*· depends on K.langclose· depth 7

of the 6,195 cells R admits and Λ_{spectra} does not hold, 3,993 are admitted by some three-axis slice and 2,202 by none. Statistics admits ONE of the 2,202 and geometry 28%, against 20% and 70% of the faces. And every available fifth coordinate makes the corners worse — by 70% to 652% — because all four candidates are DERIVED (register 1122)

PRIOR ART: cells admitted by the full closure and by no lower-arity projection are exactly the k -consistency gap — Freuder, Synthesizing constraint expressions, CACM 21 (1978) 958-966.

K.coupling — the refuted conjecture

coordinate COUPLING does not predict redundancy

a closed index

grade **MEASURED**· source *R 1120; Dechter & Pearl 1989*· depends on K.redun· depth 4

across six indices — Λ , Λ_{spectra} , the periodic table, Janet, the calendar, a box ordering — redundancy against coupling gives $r^2 = 0.002$, $p = 0.94$. Janet and the box ordering both have 50% coupling and 0% redundancy; Λ_{spectra} has 17% coupling and 20% redundancy

PRIOR ART: that coupling alone does not predict what a local method achieves — the induced width, not the edge count, is the parameter. Dechter & Pearl, Tree clustering for constraint networks, Artif. Intell. 38 (1989) 353-366.

K.deadend — the dead ends, defined

the non-composable cells are defined exhaustively: at Λ_{10} all 485 have $g = 0$ and every $g = 0$ cell is non-composable; at Λ_{13} , 11,188 at $g = 0$, 5,116 with $2J$ in $\{6,7,8\}$ which no $2J_c$ can start, and 8,214 failing only in combination

L.c8 is $k \geq 1$, so a target ending empty can never be a source; and $2J$ inherits $2K$'s range while $2J_c$ is bounded by $\varphi(k)$ alone

grade **COMPUTED**· source *M §12.11.1.5; Mac Lane 1971*· depends on K.peak· depth 6

PRIOR ART: cells that compose with nothing are those whose target is no other cell source — the non-composable part of a quiver.

K.decay — data-processing inequality + Pinsker

for a tree-structured index, $I(u;v)$ is non-increasing in tree distance d , and $|P(A_u \cap A_v) - P(A_u)P(A_v)| \leq \sqrt{(I(u;v) \cdot \ln 2 / 2)}$

Markov random field on a tree; data processing; Pinsker

grade **PROVED**· source *M §12.11.0.7; Lauritzen 1996; Pearl 1988*· depends on K.markov· depth 5

PRIOR ART: mutual information is non-increasing in tree distance because every path is separated — the data-processing inequality on a Markov tree. Cover & Thomas, Elements of Information Theory (1991), Thm 2.8.1; Lauritzen, Graphical Models (1996), ch. 3.

K.girth — the girth is four

the girth of the unit-step graph on Λ_9 is exactly 4

unit-step adjacency = covering relation, 6,658 edges

grade **PROVED**· source *T 4.3 (App. G); classical graph theory; Berge 1962*· depends on L.dist· depth 9

PRIOR ART: girth is the length of the shortest cycle, standard since Euler. That a unit-step graph on a product of chains has girth 4 follows from the commuting square of any two coordinate moves.

K.helly — Helly number

molecular transit is an intersection question; the Helly number is ≥ 5 and ≤ 144

a molecule moves as one fibre; a circuit is a square; ground set $C(9,2) \cdot 4 = 144$

grade **COMPUTED**· source *T 4.4 (App. G); Helly 1923*· depends on K.comp· depth 3

PRIOR ART: the Helly number of a family is the least h such that every h -wise intersecting subfamily intersects — Helly, Über Mengen konvexer Körper mit gemeinschaftlichen Punkten, Jahresber. DMV 32 (1923) 175-176.

K.jump — the categorial jump

the axis that makes the tower a category is worth twenty points: Λ_8 is 50.3% composable and Λ_9 , adding $2S' \leq g$, is 70.7%

the companion's index sits between them at 59.5% — three transition indexes, three fractions, so this is not a norm

grade **COMPUTED**· source *M §12.11.1.3; Mac Lane 1971*· depends on K.twocol· depth 5

PRIOR ART: the axis that closes composition is the one that supplies identities and associativity — the category axioms. What is priced is the cell cost of adding it.

K.langclose — closure as agreement

$E(X) = 0$ if and only if the languages agree

an index of dimension ≥ 3 and two or more operators

grade **MEASURED**· source *R 1176; Beeri, Fagin, Maier & Yannakakis 1983*· depends on K.three· 1 object depends on it· depth 6

six indexes, three operators — order, statistics, geometry. Λ and a box ordering: $E = 0$ and every pair agrees, 0 cells differing. The periodic table, Janet, the calendar and Λ _spectra: $E > 0$ and every pair differs. No exception. Closure is therefore a language-theoretic property as well as an order-theoretic one, and the language test is cheaper: R enumerates the ambient product, pairwise consistency needs only the marginals

PRIOR ART: global and local closures coincide exactly on acyclic structures — BFMY, J. ACM 30 (1983) 479-513. That $E = 0$ iff the languages agree is that theorem read as an equivalence.

K.markov — the Markov property

the future is conditionally independent of the past given the transfer: every past gives the same future set at each q

the constraint graph is a tree, q the cut vertex

grade **COMPUTED**· source *M §12.11.0.8; Lauritzen 1996; Pearl 1988*· depends on C.cut, L.tree· 1 object depends on it· depth 4

PRIOR ART: the future being conditionally independent of the past given the present is the MARKOV PROPERTY. Its graphical form — separation in the graph implies conditional independence — is Lauritzen, Graphical Models (1996), ch. 3, and Pearl, Probabilistic Reasoning in Intelligent Systems (1988).

K.peak — the composability peak

composability peaks at Λ_{10} and falls after: 0.0000, 0.7068, 0.8087, 0.6956, 0.6394, 0.6186 across Λ_8 to Λ_{13}

counting axes raise the fraction and coupling axes lower it, with no exception — §12.11.3's dichotomy predicts the sign as well as the price

grade **COMPUTED**· source *M §12.11.1.4; Berge 1962*· depends on K.twocol· 1 object depends on it· depth 5

PRIOR ART: the composability fraction is the arc density of the composition quiver; its rise and fall with the tower is the measurement.

K.produce — the production rule

an axis PRODUCED by another gives a dimension exactly when its production is NOT MONOTONE

a closed index and an axis derived from its coordinates

grade **MEASURED**· source *R 1143; Birkhoff 1937*· depends on K.axis· 2 objects depend on it· depth 5

multiplicity is produced by Ne and cycles $2, \{1,3\}, 2, \{1,3\}, \{2,4\}$ — NOT monotone — and adding it takes redundancy from 20% to 82%. Ne is produced by (Z,charge) and monotone: 0%. n0 is produced by (Z,charge,l) and monotone: 0%. R's envelopes are cumulative maxima, so a monotonically-produced axis is already inside them; a non-monotone one is invisible to them. This resolves the apparent contradiction between R 1122 and R 1123

PRIOR ART: a monotone function of existing coordinates adds no join-irreducibles and so no dimension; a non-monotone one does. Birkhoff (1937).

K.quiver — the composition quiver

composable cells are the arcs of a quiver Q on 33 objects; the composition graph is the line digraph L(Q), $|E| = \sum_v \text{in}(v) \cdot \text{out}(v) = 27,027$

a loop at every vertex

grade **COMPUTED**· source *T 4.2 (App. G); Gabriel 1972*· depends on K.cat· depth 4

PRIOR ART: a quiver is a directed graph whose paths generate an algebra — Gabriel, Unzerlegbare Darstellungen I, Manuscripta Math. 6 (1972) 71-103. The composable cells form the arcs of one.

K.redun — the projection ladder

redundancy under R rises with the number of INDEPENDENT coordinates; a DERIVED coordinate lowers it

a closed index X of dimension d

grade **MEASURED**· source *R 1119-1122; Shannon 1948*· depends on A.R· 2 objects depend on it· depth 3

Λ PROJECTED onto its own first d coordinates, same constraints: d=8 gives 61% removable with exact recovery, d=7,6,5 give 30%, d=4 gives 5%, d=3 gives 0%. And

adding the electron count $N_e = Z - c + 1$ to the spectra index — a FUNCTION of two coordinates it already has — drops redundancy from 20% to 0% while doubling the envelopes and raising coupling 17% to 33%

PRIOR ART: redundancy is the excess of a representation over its entropy. That it rises with the number of INDEPENDENT coordinates is the measurement; the notion is Shannon (1948).

K.three — the three-coordinate rule

an index needs THREE coordinates before its languages can disagree

an index and two languages

grade **PROVED**· source *R 1175; Beeri, Fagin, Maier & Yannakakis 1983*· depends on A.stat2· 1 object depends on it· depth 5

at $d = 2$ there is one coordinate pair, so pairwise consistency and the cell coincide and every language returns the same answer. The periodic table, Janet and the calendar were all tested at two and all reported agreement; rebuilt at three — with block, l and weekday, each non-monotone in the second coordinate — the periodic table gives order $E = 100$ against statistics 0, reproducing the book's recorded caveat. All three match the book at 2-D first: 36, 0 and 7

PRIOR ART: pairwise consistency is a statement about PAIRS, so an object with one pair cannot make it. BFMY (1983) for the general local-to-global framework.

K.transit — the transit profile

the transit profile $MI/H = 0.51, 0.93, 0.08, 0.13, 0.77$ at Λ_9 - Λ_{13} : reduced at 10, a label through 11 and 12, re-attaching at 13

Λ_{12} built with the loose bound

grade **COMPUTED**· source *M §12.11; Shannon 1948*· depends on T.tower· depth 7

PRIOR ART: MI/H is the normalised mutual information, standard since Shannon (1948).

K.twocol — space is what an index holds, time is what it composes

an index has a time column exactly when its cells are moves: Λ_8 976/0, Λ_9 1,654/1,169, the companion's index 2,370/1,410, while the periodic table and the calendar have no second column and cannot have one

a transition cell has two ends so composability is askable; a state cell has one position and the question does not arise

grade **COMPUTED**· source *M §12.11.1.3; Mac Lane 1971*· depends on K.cat· 3 objects depend on it· depth 4

PRIOR ART: an index has a composition structure exactly when its cells are morphisms rather than objects — the distinction is the category axioms. Mac Lane (1971). CORRECTED 2026-08-11: this statement printed Λ_8 as 976/491. It is 976/0. The composition table

gives 0 with fraction 0.0000, Λ_8 composes not at all — four source coordinates against three target — and K.window states the reason as a dependency: ' Λ_8 target has 3 coordinates against 4', so $\text{tgt}(a) = \text{src}(b)$ is not even askable at Λ_8 and Λ_9 is the FIRST composable level. The 491 was a hand-authored figure in an object graded COMPUTED, contradicting the computation it depends on.

K.window — the composable window

Λ_9 is the first composable level and the last tree level

Λ_8 target has 3 coordinates against 4; Λ_{10} has 10 nodes and 10 edges

grade **COMPUTED**· source *T 4.1 (App. G); Beerli, Fagin, Maier & Yannakakis 1983*· depends on K.comp, L.tree· depth 3

PRIOR ART: that treeness and composability are different properties, and that an index can be the last of one and the first of the other, is the acyclicity/local-consistency distinction of BFMY, J. ACM 30 (1983) 479-513.

K.zcross — the periodic table as a coordinate

composability is 22.71% within one element and 85.23% across the 118, while E is 28,503 without Z and 972,862 with it

composability wants the whole table and closure wants one atom; Λ closes because it has no Z

grade **COMPUTED**· source *M §12.11.1.6; Edlen 1964*· depends on K.twocol· 1 object depends on it· depth 5

PRIOR ART: that composability is far higher across elements than within one is the isoelectronic structure — Edlen, Atomic spectra, Handbuch der Physik XXVII (1964).

M. The modular chain — 9 objects

M.C1 — the presymplectic potential

the presymplectic potential $\theta = \delta\varphi \, \mathbb{F}_\ell \varphi \, \eta$ carries no transverse derivative, so Ω is block diagonal in y and $\{\varphi(u,y), \varphi(u',y')\} = (1/4\sqrt{q(y)})\text{sgn}(u-u')\delta^{d-2}(y-y')$

u -independent transverse metric, i.e. $\Theta = 0$; non-derivative interactions only

grade **COMPUTED**· source *T 10.4c (App. G), F1, F2; Wald & Zoupas 2000*· depends on M.hsml· 2 objects depend on it· depth 3

PRIOR ART: the presymplectic potential and its ambiguities are Wald & Zoupas, General definition of conserved quantities in general relativity, Phys. Rev. D 61 (2000) 084027.

M.C2 — the open object — half-sided modular inclusion on a non-expanding horizon · GRADED OPEN

$\Delta_{\{M(u_1)\}^{\{it\}} M(u_2) \Delta_{\{M(u_1)\}^{\{-it\}}} \subseteq M(u_2)$ for $t \leq 0$, $u_1 < u_2$, WITH Ω cyclic and separating for BOTH $M(u_1)$ and $M(u_2)$

N a non-expanding horizon with no Killing field, ω Hadamard

grade **OPEN**· source *T 10.4d* (App. G), *F6*· depends on M.C1, M.hsmi· 1 object depends on it· depth 4

REGRADED CONDITIONAL 2026-08-11 (R 1507). This object is NOT open. It is PROVED on one existence hypothesis, in nine steps NONE of which uses a Killing field: $P_\lambda \geq 0$ from the ANEC on ACHRONAL generators (R 1471); $U(s)$ unitary by Stone; $U(s) B U(s)^\dagger$ inside B because the cut translates into itself, which is CAUSAL structure (R 1506); $U(s)\Omega = \Omega$ from the hypothesis; $C_\lambda = U(1) B U(1)^\dagger$, the source own equation; Ω cyclic for C_λ since $C_\lambda \Omega = U(B \Omega)$ with U unitary and $B \Omega$ dense (R 1501); separating since C_λ is inside B ; hence the four Borchers conditions, hence HSMI by Borchers 1992 and Wiesbrock 1993. THE HYPOTHESIS: there exists a state annihilated by every smeared ANE operator and cyclic and separating for the exterior algebra. A Killing horizon supplies it (Hartle-Hawking); a non-expanding horizon is not known to. So the isometry supplies a WITNESS, not a step. The open question is a CONTAINMENT between two classes of spacetimes, not a parameter. STATEMENT COMPLETED 2026-08-11 (R 1481): the standardness clause was missing. A half-sided modular inclusion is N inside M with a vector Ω CYCLIC AND SEPARATING FOR BOTH, and $\sigma_t(N)$ inside N for $t \leq 0$ — Lechner-Scotford Def 2.1, equivalent to the Borchers triple by Borchers 1992 and Wiesbrock 1993. This object stated only the inclusion clause. The omitted clause is exactly the one Faulkner and Speranza ASSUME rather than derive — arXiv 2405.00847 sec 3.1 says, of the smaller algebra, assuming Ω is cyclic for it — and Reeh-Schlieder does not supply it, because their Ω is a vacuum for the average null energy operators and, in their own words, may not coincide with a global minimal energy state. Araki and Zsido, Rev. Math. Phys. 17 (2005) 491-543, extend Wiesbrock to weights and fill a gap in the 1993 proof. OPEN on the GEOMETRIC route only. M.sorce closes it by construction — a geometric modular flow needs a conformal Killing vector and a non-expanding horizon has none; Chandrasekaran & Flanagan (arXiv:2601.07915) have the Killing case. The ALGEBRAIC route is not blocked: M.hsmi characterises HSMI by a one-parameter unitary group with positive generator and mentions no Killing field, and M.C1's commutator is already an algebraic object. That route was ATTEMPTED (R 1018-1022): its free half goes through — M.C1's commutator gives a positive generator per null generator, negative/positive spectral weight $5e-5$, so a standard pair exists per generator and block diagonality keeps them unmixed. The obstruction is in the CORNER EDGE MODES, where Chandrasekaran & Flanagan show the null translation generator is necessarily two-sided. A proposal that $\Theta = 0$ supplies the missing relative boost was tested and FAILS: $\theta_{ab} \rightarrow a(y) \theta_{ab}$ under $l \rightarrow a(y)l$, so $\theta = 0$ is invariant under the rescaling and cannot fix its parameter. The paper's OWN route (Sec 7.3) supplies the boost from SMOOTHNESS instead — local Rindler frames give $\chi = \kappa(u \, d_u - v \, d_v)$ with $\text{grad}(\mu \chi_\nu) = O(u, v)$, and footnote 44 says a local boost Killing field is all that is needed. It establishes the modular flow's geometric action TO FIRST ORDER at any cut and states explicitly that these CANNOT be patched together. M.C2 needs finite $u_1 < u_2$. The reduction bottoms out at POSITIVITY of the null translation generator, and that is NOT implied by Hadamard: Hadamard is

microlocal (a wavefront-set condition) while positivity is a global spectral one, and a smooth deformation of the vacuum raises negative/positive spectral weight from $4.8e-05$ to $2.6e-03$ while changing the UV tail by 0.0000%. Thermal states are Hadamard and carry both signs. Positivity is a SELECTION CRITERION assumed by everyone who needs it — Chandrasekaran-Flanagan Sec 8 and Dappiaggi-Moretti-Pinamonti both — and Kay-Wald 1991 get uniqueness only for states INVARIANT UNDER THE KILLING FLOW, the hypothesis M.C2 drops. CORRECTION (R 1038-1042): the hypothesis is NOT purely geometric — the standard NEH definition's third condition is 'Einstein field equations hold on Δ and $-T^a{}_b l^a l^b$ is future causal', an energy condition on the state, which via Raychaudhuri forces $T_{ab} l^a l^b = 0$ and $\sigma_{ab} = 0$, hence $L_l q_{ab} = 0$ — which is M.C1's hypothesis, so M.C1 assumes a consequence of the definition. The shortfall is ORDER: the condition fixes the background, while half-sidedness is spectral on the perturbations

M.hsmi — Borchers 1992 / Wiesbrock 1993

half-sided modular inclusion is CHARACTERISED by a one-parameter unitary group with positive generator

common cyclic separating vector; extended to weights by Araki-Zsidó 2004

grade **CITED**· source *T E2, E3; Borchers 1992; Wiesbrock 1993*· depends on M.tt· 2 objects depend on it· depth 2

PRIOR ART: half-sided modular inclusion and its characterisation by a one-parameter group with positive generator — Borchers, The CPT theorem in two-dimensional theories of local observables, Commun. Math. Phys. 143 (1992) 315-332; Wiesbrock, Half-sided modular inclusions of von Neumann algebras, Lett. Math. Phys. 28 (1993) 107-114.

M.ledger — the modular ledger

$d - 1 = 1 + (d - 2)$ ON THE FREE ALGEBRA: one HSMI per generator supplies the affine line; the transverse direct integral is C1 and no HSMI supplies it. INCOMPLETE ON THE DRESSED ALGEBRA, which carries doubled corner modes besides

a null hypersurface in d dimensions — THIS IS A MANIFOLD COUNT, NOT AN ALGEBRA COUNT (R 1484)

grade **COMPUTED**· source *T 10.4e (App. G), F7; Borchers 1992; Wiesbrock 1993*· depends on M.C1, M.C2· depth 5

COMPLETED 2026-08-11 (R 1484): the sum is TWO accountings. LEVEL ONE, the manifold, is what this object counts and it is correct. LEVEL TWO, the algebra, carries the corner content, and register 1020 writes it exactly: the residual freedom is $l \rightarrow a(y)l$, acting on the affine parameter as $u \rightarrow u/a(y) + b(y)$. Those are two arbitrary FUNCTIONS ON THE CUT, not DIMENSIONS OF A MANIFOLD, so no arithmetic sum can hold them and none should be attempted. R 1022 named what would close M.C2 — a canonical scaling of the affine parameter — which is $a(y)$. QUALIFIED 2026-08-11 (R 1483): the sum is the FREE count and was stated without the qualifier. Register 1019, which M.C2 own check already cites, records that half-sidedness fails on the DRESSED algebra and not the free one, and that Chandrasekaran and Flanagan recover it by

EXTENDING THE PHASE SPACE WITH DOUBLED CORNER MODES — relative boosts AND null translations of the respective corners. Those modes belong to the affine family and the transverse family at once, so they are not a summand but an OVERLAP, and $1 + (d-2)$ does not count them. This object mentioned neither corner nor edge mode nor two-sided nor either author. dimensional ledger — and the ledger IS the kinematic/stateful decomposition (R 1035): the STATEFUL object (HSML, hence M.C2) supplies exactly ONE dimension, the affine line; the KINEMATIC one (M.C1, Theta = 0) supplies the other $d-2$. Read as a counting argument until then

PRIOR ART: one half-sided modular inclusion per generator supplies the affine line — the Borchers-Wiesbrock theorem, Borchers, Commun. Math. Phys. 143 (1992) 315-332; Wiesbrock, Lett. Math. Phys. 28 (1993) 107-114.

M.rs — Reeh-Schlieder

the vacuum is cyclic and separating for local algebras

Hadamard state; region with nonempty causal complement

grade CITED· source *T E0; Reeh & Schlieder 1961*· 1 object depends on it· depth 0

PRIOR ART: the vacuum is cyclic and separating for local algebras — Reeh & Schlieder, Bemerkungen zur Unitätsäquivalenz von Lorentzinvarianten Feldern, Nuovo Cim. 22 (1961) 1051-1068.

M.semi — semifinite carries a trace

a semifinite factor carries a trace, hence an entropy; type III carries none

von Neumann classification

grade CITED· source *T E5; Murray & von Neumann 1936*· depends on M.takesaki· depth 3

PRIOR ART: the type classification of factors, and that only semifinite ones carry a trace, is Murray & von Neumann, On rings of operators, Ann. Math. 37 (1936) 116-229, and its sequels.

M.sorce — Sorce 2024

any geometric modular flow must be generated by a conformal Killing field

general

grade CITED· source *T 10.4 (App. G); Sorce 2024*· depends on M.tt· depth 2

PRIOR ART: that a geometric modular flow must be generated by a conformal Killing field — Sorce, Analyticity and unitarity for cosmological correlators (2024) and related work on geometric modular flow.

M.takesaki — Takesaki duality 1973

$N = M \rtimes_{\{\sigma^\varphi\}} \mathbb{R}$ is type II_∞ with trace τ , $\tau \circ \theta_s = e^{-s} \tau$, and $M = N \rtimes_0 \mathbb{R}$ uniquely

M type III, φ a faithful semifinite normal weight; STATED FOR $\mathbb{R}$

grade **CITED**· source *T E4; Takesaki 1973*· depends on M.tt· 1 object depends on it· depth 2

PRIOR ART: the structure theorem for type III factors as crossed products — Takesaki, Duality for crossed products and the structure of von Neumann algebras of type III, Acta Math. 131 (1973) 249-310.

M.tt — Tomita-Takesaki

for (M, Ω) there exist Δ, J with $\Delta^{\{it\}} M \Delta^{\{-it\}} = M$

M a von Neumann algebra, Ω cyclic and separating

grade **CITED**· source *T E1; Tomita 1967; Takesaki 1970*· depends on M.rs· 3 objects depend on it· depth 1

PRIOR ART: Tomita-Takesaki modular theory — Tomita, Quasi-standard von Neumann algebras (1967, unpublished); Takesaki, Tomita Theory of Modular Hilbert Algebras and its Applications, Springer LNM 128 (1970).

W. The violation index — 9 objects

W.core15 — the core at fifteen letters

at fifteen letters: 18,072 cells, $E = 816 = 1 \times 816$, core ($X_{\text{exp}}=0$, $U_{\text{ghost}}=0$, $NEC_{\text{pt}}=3$, $EOM=2\text{nd}$)

the split alphabet of T §6.5 (App. G)

grade **COMPUTED**· source *T 7.1 (App. G); Chinneck & Dravnieks 1991*· depends on W.core9· 1 object depends on it· depth 5

PRIOR ART: as W.core9 at a finer alphabet. That the core is unchanged while the multiplicity grows is the measurement; the notion of an irreducible infeasible subsystem is theirs.

W.core9 — the core at nine letters

at nine letters: 2,370 cells, $E = 30 = 1 \times 30$, core ($X=0$, $U=0$, $NEC=3$)

the coordinate set of T §5.1 (App. G)

grade **COMPUTED**· source *T 5.3 (App. G); Chinneck & Dravnieks 1991*· depends on A.E· 3 objects depend on it· depth 4

cell count RECOMPUTED: the violation index's 17 rules applied to the $4 \cdot 3 \cdot 3 \cdot 3 \cdot 5 \cdot 2 \cdot 2 \cdot 3 \cdot 3$ box give exactly 2,370. $E=30$ is the arity-4 CONSTRAINT-LANGUAGE defect ($NEC \geq 3 - > IC \vee U \vee X$), not the envelope defect of the cell set, which is 3,040 (R 1009)

PRIOR ART: a minimal set of conditions whose joint failure is irreducible is a MINIMAL UNSATISFIABLE SUBSET — Chinneck & Dravnieks, Locating minimal infeasible constraint sets in linear programs, ORSA J. Comput. 3 (1991) 157-168; in SAT the same object

is the MUS. The conditions themselves are physics: the null energy condition (Penrose 1965), ghost states (Pais & Uhlenbeck 1950) and the equations of motion.

W.face — the null surface is a face

the companion's local null surface is a FACE of the 6-cube, not a parity class: 32 of 64 with every rung 2, and a parity class gives $E = 32$ where the printed E is 0

deducible from three printed numbers with no access to any cell; THETA = 0 at 16 and STAT = 0 at 8 nest Killing horizons inside non-expanding ones

grade **COMPUTED**· source *T §10.1 (App. G), §10.4; Coxeter 1948*· depends on A.R· depth 3

PRIOR ART: a face of the n -cube is the set fixing some coordinates and freeing the rest — Coxeter, *Regular Polytopes* (1948), ch. VII. A parity class is not a face, which is what distinguishes them here.

W.frontier — the frontier metric — distance and status on the fifteen-letter alphabet

$d(c) = X_{\text{exp}} + U_{\text{ghost}} + |\text{NEC}_{\text{pt}} - 3| + \text{EOM} + d_P$; $s(c) = 1\{\cdot\} + 1\{\cdot\} + 1\{\cdot\} + 1\{\cdot\} + s_P$

the fifteen-letter alphabet; d_P the L1 distance on the eleven free letters

grade **DEFINITIONAL**· source *T D7, D8; classical; Hamming 1950*· depends on W.core15· depth 6

PRIOR ART: a defect built as a sum of independent violation counts, with a support counted separately, is the standard form of a penalty function.

PRIOR ART: a defect built as an L1 distance on free letters plus a count of violated conditions is a Hamming-type distance with weights — Hamming, Error detecting and error correcting codes, *Bell Syst. Tech. J.* 29 (1950) 147-160.

W.jur — jurisdiction does not change the defect

a jurisdicted forcing and an unjurisdicted disjunction give identical defects; arity ≥ 3 makes a defect possible, jurisdiction narrowness makes it small

the scope condition is a coordinate

grade **COMPUTED**· source *T 8.2 (App. G); Freuder 1978*· depends on W supp· depth 9

PRIOR ART: that a constraint of arity 3 or more is not captured by binary projections is the k -consistency ladder — Freuder, Synthesizing constraint expressions, *CACM* 21 (1978) 958-966. Whether a scope condition is jurisdicted or disjunctive does not change the arity, hence not the defect.

W.onehot — the one-hot partition

status is one bit per vocabulary and the partition forces one-hot: 35 of 40 terms weight 1, 2 weight 0, 3 weight 2

weight 2 = relation or unsplit conflation; weight 0 = ungrounded or constructible

grade **COMPUTED**· source *T 10.5 (App. G); classical; Shannon 1938*· depends on W.rel· depth 4

partition RECOMPUTED: $35 + 2 + 3 = 40$ terms, total weight 41 (R 1009)

PRIOR ART: a partition of a set forces exactly one indicator to be set — the one-hot encoding. Elementary, and the measurement is that 35 of 41 statuses satisfy it.

PRIOR ART: a partition forces exactly one indicator — the one-hot encoding of Boolean algebra, Shannon, Trans. AIEE 57 (1938) 713-723. Weight 2 signals a relation or a conflation; weight 0 an ungrounded term.

W.rel — Brunetti-Fredenhagen-Verch 2003

a term that is a relation between two vocabularies cannot be a coordinate of either

the BFV partition $A : Loc \rightarrow Alg$ has four parts

grade **COMPUTED**· source *T 8.4 (App. G); Brunetti, Fredenhagen & Verch 2003*· depends on A.R· 1 object depends on it· depth 3

PRIOR ART: a term that is a relation between two vocabularies is a morphism, not an object — the locally covariant framework of Brunetti, Fredenhagen & Verch, The generally covariant locality principle, Commun. Math. Phys. 237 (2003) 31-68, makes the distinction precise.

W.scale — the defect does not scale

the defect does not scale: core = 1 at 4, 5, 7 and 9 coordinates; only the multiplicity moves (3,5,10,30)

projections of one index

grade **COMPUTED**· source *T 5.7 (App. G); Chinneck & Dravnieks 1991*· depends on W.core9· depth 5

PRIOR ART: that the core of an infeasible system is invariant under refining the encoding is the well-definedness of the MUS; only the multiplicity of witnesses changes.

W.supp — the core is minimal

of 414 coordinate subsets excluding {X,U,NEC}, 0 fail: the triple is the unique minimal failing subset, arity exactly 3. THE COUNT RECONCILES — subsets of

size 2..6 at d = 9 number exactly 414 — but the convention is unprinted and the failure count needs the cells. Arity 3

as *W.core9*

grade **COMPUTED**· source *T 5.4 (App. G); Chinneck & Dravnieks 1991*· depends on *W.core9, A.modeB*· 1 object depends on it· depth 8

combinatorics RECOMPUTED: subsets of size 2..6 from 9 coordinates number 456; those containing all of {X,U,NEC} number 42; $456-42 = 414$ exactly as stated (R 1009)

PRIOR ART: testing all 414 subsets and finding none fails without the triple establishes MINIMALITY in their sense — no proper subset is infeasible.

EM. The electromagnetic quotient — 7 objects

EM.cross — the selection-rule crossing

EM-allowed cells compose at 11.6% within one element against the forbidden 40.7%, and the order reverses across the table at 89.7% against 77.9%

the rule that forbids composition inside an atom is the rule that enables it between atoms; parity repeats it at 38.4% against 20.1%

grade **COMPUTED**· source *M §12.11.1.7; Laporte 1924*· depends on *K.zcross*· depth 6

PRIOR ART: the composability contrast between allowed and forbidden cells is a consequence of the parity rule; the measurement is the fraction.

EM.image — the complete rectangle

Λ_9 's image on (multipole, ΔS) is the COMPLETE rectangle at every cap; E = 0 vacuously

as *EM.map*

grade **COMPUTED**· source *M §12.11.8; Wigner 1927*· depends on *EM.map, A.dens*· depth 5

PRIOR ART: that the selection rules cut a complete rectangle in (multipole, ΔS) is the product structure of the space-spin decomposition. Wigner, Z. Phys. 43 (1927) 624.

EM.map — electric/magnetic multipole selection rules

the multipole is determined by $|\Delta l|$ and parity alone: 0->M1, 1->E1, 2->E2, 3->E3

one-electron jump; no source J exists in Λ

grade **COMPUTED**· source *M §12.11.8; Laporte 1924; Condon & Shortley 1935*· depends on *L.def*· 5 objects depend on it· depth 2

PRIOR ART: the multipole order of a transition is fixed by $|\Delta l|$ and parity — Laporte, Z. Phys. 23 (1924) 135; the full multipole classification is Condon & Shortley (1935), ch. IV.

EM.notcomp — selection and composition are unrelated

composability and the EM condition share 0.0004 bits of a possible 0.633

on Λ 's cells

grade **COMPUTED**· source M §12.11.8; Shannon 1948· depends on EM.map, K.comp· depth 3

PRIOR ART: shared information between two binary properties, in bits. Shannon (1948).

EM.parity — the parity hole

$|\Delta l| = 1$ is $\delta^{-1}(\{-1, +1\})$, a hole at zero, not convex; E = 750

as EM.spin

grade **COMPUTED**· source M §12.11.8; Laporte 1924; Wigner 1927· depends on EM.map, I.convex, T.excl· depth 11

PRIOR ART: the parity selection rule $|\Delta l| = 1$ for electric dipole radiation is Laporte, Z. Phys. 23 (1924) 135, and its group-theoretic ground is Wigner, Z. Phys. 43 (1927) 624.

EM.quotient — the electromagnetic quotient

the EM index is a QUOTIENT of Λ , not an extension: adjoining its coordinates gives E = 3,900

every EM coordinate is a function of Λ 's own

grade **COMPUTED**· source M §12.11.8; Noether 1918· depends on EM.map, A.derived· depth 5

PRIOR ART: a selection rule is a quotient by a symmetry, not an extension of the state space. Noether, Invariante Variationsprobleme, Nachr. Ges. Wiss. Goettingen (1918) 235-257, is the general statement of the correspondence between symmetry and conserved structure.

EM.spin — the spin diagonal

$\Delta S = 0$ is a diagonal, hence two monotone one-parent bounds; imposing it preserves E = 0 at four cap settings

LS coupling

grade **COMPUTED**· source M §12.11.8; Russell & Saunders 1925; Wigner 1931· depends on EM.map, I.convex· depth 11

PRIOR ART: $\Delta S = 0$ for electric dipole transitions in LS coupling is the spin selection rule — Russell & Saunders, Astrophys. J. 61

(1925) 38; Wigner, Gruppentheorie (1931), for the representation-theoretic statement.

C. Protocol and audit objects — 9 objects

C.Aq — the first factor

$$A_q(z) = \sum_{n=1}^{\infty} z^n \sum_{\ell=0}^{\min(n-1,1)} z^\ell \sum_{k=\max(q,1)}^{\min(4\ell+2,3)} z^k (1-z^{\min(k,3)+1})/(1-z)$$

caps as stated; q held fixed

grade **COMPUTED**· source *M §12.7 / T C2eq; Euler 1748; Stanley 1986*· depends on L.F· 1 object depends on it· depth 4

PRIOR ART: a nested sum over a constrained region, written as a polynomial in z, is the standard rank-generating construction. Euler (1748) for the method; Stanley, Enumerative Combinatorics Vol. 1 (1986), ch. 1, for the modern treatment.

C.Bq — the second factor

$$B_q(z) = \sum_{e=1}^{\infty} z^e \sum_{f=0}^{\min(e-1,1)} z^f (1-z^{\min(q,4f+2)+1})/(1-z)$$

as C.Aq

grade **COMPUTED**· source *M §12.7 / T C3eq; Euler 1748; Stanley 1986*· depends on L.F· 1 object depends on it· depth 4

PRIOR ART: as C.Aq — the second factor of the fibred count, built the same way.

C.box — the box generating function

$$\text{Box}(a,b)(z) = \prod_i z^{a_i} (1-z^{b_i-a_i+1})/(1-z)$$

every fibre bottoms out in a product of chains

grade **DEFINITIONAL**· source *M §12.7.2 / T C4eq; Euler 1748*· depends on C.Aq, C.Bq· depth 5

PRIOR ART: the generating function of a box is a product of finite geometric series — Euler, *Introductio in analysin infinitorum* (1748), ch. XVI, where partition generating functions are introduced.

C.compare — the comparison audit

the COMPARISON AUDIT: index, math and language cypher checked against one another, not each against itself

any two indexes and the cypher

grade **MEASURED**· source *R 1169-1172; Tarski 1936; Beeri, Fagin, Maier & Yannakakis 1983*· depends on A.R, K.produce· depth 6

three checks. COVERAGE: every index coordinate has a math object and every family an index part — 0 and 0. LANGUAGE: five index/language pairs never run — Λ _spectra in geometry, algebra, information and statistics, Λ _alpha in analysis. CONTRADICTION: three standing, all on Λ _spectra between ORDER and ANALYSIS — delta falls with l (194/205 vs 102/205), triplet exceeds singlet (55/66 vs 33/66), delta falls along a sequence (89/143 vs 135/143). The audit names the pair and does NOT adjudicate

PRIOR ART: comparing two formal readings of one object and requiring agreement is the metalanguage move (Tarski 1936); that global and local readings agree exactly on acyclic structures is BFMY (1983).

C.cut — every cut closes

every two-sided cut of the tree gives defect zero, not only q

the tree is doing the work, not the transfer

grade **COMPUTED**· source *M §12.11.0.8; Freuder 1982*· depends on C.fib, L.tree· 1 object depends on it· depth 3

PRIOR ART: every two-sided cut of a tree separates it, so the defect vanishes at each — Freuder, J. ACM 29 (1982) 24-32.

C.fib — the fibred count

$$|\Lambda| = \sum_q |A(q)| \cdot |B(q)| = 33 \cdot 5 + 33 \cdot 10 + 23 \cdot 15 + 8 \cdot 17 = 976$$

conditioned on the transfer q

grade **COMPUTED**· source *M §12.6.1 / T C1eq; Fubini 1907*· depends on L.def· 4 objects depend on it· depth 2

PRIOR ART: summing a product over a fibration of the index set is the discrete Fubini theorem. That the fibres here are independent is what makes $|\Lambda|$ factor as a sum of products.

C.local — local closure is not implied

$E(\Lambda) = 0$ and $E(A_q) = E(B_q) = 0$ for every q; the second does not follow from the first

cross-sections with induced coordinates

grade **COMPUTED**· source *M §12.8.5 / T C5eq; Rota 1964*· depends on C.fib, A.E· depth 4

PRIOR ART: that a global count factorises over a decomposition does not imply each part is separately closed — the failure of naive Moebius inversion over non-independent parts. Rota (1964).

C.pareto — the fibre trade

$|A(q)|$ falls 33,33,23,8 while $|B(q)|$ rises 5,10,15,17; no q improves both

at stated caps

grade **COMPUTED**· source *M §12.8.1; Pareto 1896*· depends on C.fib· depth 3

PRIOR ART: no q improves both factors — a Pareto front in the two counts. Pareto, Cours d économie politique (1896).

C.qmean — the transfer distribution

$\langle q \rangle = 1.4631$, sd 0.930; peak at $q = 2$ with 345 cells (35.3%)

at stated caps

grade **COMPUTED**· source *M §12.8.2; classical; Gauss 1809*· depends on C.fib· depth 3

PRIOR ART: the first and second moments of a distribution over a coordinate; elementary.

PRIOR ART: first and second moments of a distribution over a coordinate. Gauss, Theoria motus (1809).

I. The intake — interval maps and convexity — 3 objects

I.convex — the convexity criterion

$\delta^{-1}(T)$ is a sublattice iff T intersect $\text{range}(\delta)$ is convex in $\text{range}(\delta)$

as I.interval

grade **PROVED**· source *M §17.3; Birkhoff 1937*· depends on I.interval, T.excl· 2 objects depend on it· depth 10

PRIOR ART: the preimage of a set under a lattice homomorphism is a sublattice iff the set is convex in the image — the standard sublattice criterion. Birkhoff, Lattice Theory (1940), ch. II.

I.interval — the interval map

$\delta = f - l$ is an INTERVAL MAP: $\min(\delta a, \delta b) \leq \delta(a \vee b)$, $\delta(a \wedge b) \leq \max(\delta a, \delta b)$

a sublattice of a product of chains

grade **PROVED**· source *M §17.3; Birkhoff 1940*· depends on L.dist· 1 object depends on it· depth 9

PRIOR ART: a difference of two coordinates is an INTERVAL MAP on a distributive lattice — it need not be a homomorphism but it is bounded above and below by the coordinatewise extremes. Standard; Birkhoff, Lattice Theory (1940).

I.shape — the shape of an interval

between any two points there is an interval and the method returns its measure; three objects, one shape

cells, states under composition, measurements

grade **COMPUTED**· source *M §9.2; Birkhoff 1940; Monjardet 1981*· depends on L.occ, K.comp, B.brk, A.slack· depth 6

PRIOR ART: between any two points of a lattice lies the interval [a and b, a or b]; the method returns its measure. Elementary lattice geometry.

PRIOR ART: between any two lattice points lies the interval [a and b, a or b]; its measure on a product of chains is the product of coordinate spans. Birkhoff, Lattice Theory (1940), ch. II; Monjardet, Discrete Math. 35 (1981) 173-184.

P. The spectral mechanisms — what moves a defect and by how much — 17 objects

P.buildlimit — constructed limit

a limit can be built from two spectra and tested by convergence

a series converging on a state above its own ionisation threshold

grade **MEASURED**· source *R 711, 712; Edlen 1964*· depends on P.converge, P.collapse· depth 2

Ne I 2s.2p6.np on 173,929.75 + 217,047.598 gives +0.8408 +/- 0.0191 over ten

PRIOR ART: constructing a limit by summing the ionisation energies of successive stages is standard spectroscopic practice. Edlen, Handbuch der Physik XXVII (1964).

P.charge — the same-element ladder

at s and p the quantum defect falls with charge at FIXED ELEMENT

a fixed element with two or more charge states

grade **MEASURED**· source *R 963; Edlen 1964*· depends on P.qdt· depth 1

37 of 37 monotone at $l \leq 1$, no exceptions; all nine failures are at $l \geq 2$ where P.dcollapse governs

PRIOR ART: the fall of the defect with ionisation stage along an isonuclear sequence is standard spectroscopy — Edlen, Atomic spectra, Handbuch der Physik XXVII (1964) 80-220.

P.converge — self-determined limit

a Rydberg series measures its own ionisation limit

enough members that the fit is determined

grade **MEASURED**· source *R 684, 685, 686, 698, 699; Rydberg 1890; Ritz 1903*· depends on P.qdt· 1 object depends on it· depth 1

38 of 58 put the published limit within 3x the fit own error; only 15 of 58 within the PUBLISHED error (R 812-813)

PRIOR ART: extrapolating a Rydberg series to its limit is the classical method of determining an ionisation energy — Rydberg (1890), Ritz (1903).

P.coreblind — parent-term independence

delta depends on l and the core's charge, not on the core's STATE

two parent terms of one species

grade **MEASURED**· source R 709, 710, 743; Seaton 1958· depends on P.qdt· depth 1

TWO instances; and CONTRADICTED IN PRINCIPLE by MQDT, which defines the defect as $\mu(l, \lambda, \alpha +)$ depending on the core state (R 947)

PRIOR ART: single-channel quantum defect theory treats the core as a fixed phase shift, so delta depends on l and the core charge only. Its failure for multi-channel cases is Seaton MQDT (Rep. Prog. Phys. 46, 1983).

P.dcollapse — orbital collapse

an orbital collapses at the onset of its shell and leaves the Rydberg series

3d across the transition row, 4f across the lanthanides

grade **MEASURED**· source R 719, 734, 735, 736; Goepfert-Mayer 1941; Griffin, Andrew & Cowan 1969· depends on P.iso· depth 2

all ten P.iso exceptions are d and small; Ba III 4f gives 1.0 against neon's 0.006

PRIOR ART: orbital collapse at the onset of a shell — Goepfert-Mayer, Phys. Rev. 60 (1941) 184-187; Griffin, Andrew & Cowan, Phys. Rev. 177 (1969) 62-71.

P.iso — the isoelectronic ladder

for one element and l, delta falls as the core charge rises

defect large enough to exceed its own scatter

grade **MEASURED**· source R 708, 716, 717, 718· depends on P.qdt· 1 object depends on it· depth 1

8 of 9 s/p ladders exact; and a form $\delta = a + b \cdot \ln(c+1)/c$ fits the Mg-like ns ladders to 2% of range and predicts out-of-sample, but FAILS on nd, nf and all He-like ladders (47-65%); the method is Edlen 1964 (R 942-945)

P.jj — the coupling-scheme marker

J-inconsistency within one series marks where LS coupling has failed

same series, different J

grade **MEASURED**· source R 703; Condon & Shortley 1935· depends on P.qdt· depth 1

114 of 129 consistent on raw levels, interval 82-93%; heavy elements fail at 23% against light at 8% (R 818-819)

PRIOR ART: the transition from LS to jj coupling as the spin-orbit interaction grows with Z is Condon & Shortley, The Theory of Atomic Spectra (1935), ch. X.

P.jsplit — the j-splitting

delta splits by the outer electron's own j when there is something to couple to an OPEN-SHELL core, or Z large enough for the electron's spin-orbit

grade **MEASURED**· source R 754, 756, 757, 758, 759; Sommerfeld 1916· depends on P.qdt· depth 1

median 0.0454 over 19 open/heavy pairs against 0.0002 over 10 closed/light — 227x

PRIOR ART: fine-structure splitting of a term by J is Sommerfeld, Zur Quantentheorie der Spektrallinien, Ann. Phys. 51 (1916) 1-94, in its relativistic form; Dirac (1928) supplies the exact theory.

P.lcollapse — the l-collapse

delta falls monotonically with l and reaches zero by f

unperturbed series

grade **MEASURED**· source R 693, 702, 713, 724; Hartree 1928; Seaton 1958· depends on P.qdt· 2 objects depend on it· depth 1

229 of 230 adjacent-l pairs correct, judged against EACH PAIR'S OWN 2 sigma rather than a fixed tolerance. The test carried an unstated slack of 0.02 until R 1072; at a fixed 0.02 it reads 251/251, strictly 242/251. The one failure is He I p->d, whose p defect is NEGATIVE — the Appendix B omission of R 871, which could not be recomputed when the reduced-mass Rydberg was corrected (R 1072-1074)

PRIOR ART: the defect falls with l because the centrifugal barrier keeps the electron out of the core — the standard penetration argument, Hartree, Proc. Camb. Phil. Soc. 24 (1928) 89, and Seaton (1958).

P.lens — the precision lens

near the limit delta is measured through a lens worsening as n^3

delta depends on (limit - E), which shrinks as n^2 while level error does not

grade **MEASURED**· source R 749, 750, 751, 752; Rydberg 1890· depends on P.qdt· depth 1

WEAK at scale: only 44 of 79 channels show a wider high half (56%), and the n^3 prediction of 4.2x is observed at 1.3x (R 827)

PRIOR ART: near the limit $dT/d\nu \sim \nu^{-3}$, so a fixed energy uncertainty maps to a defect uncertainty growing as ν^3 . Direct from the term formula.

P.mono — the monotonicity of the defect

the defect approaches δ_0 monotonically: PENETRATION series fall, POLARISATION series rise

steps resolvable against their own uncertainty

grade **MEASURED**· source R 806, 821-824; NIST Atomic Spectroscopy compendium; Ritz 1903· depends on P.qdt· depth 1

penetration ($|d| \geq 0.1$) falls 146 of 163 = 92%; polarisation ($|d| < 0.01$) RISES 52 of 52 = 100%; combined 94% against 66% as a single claim (R 985-987)

PRIOR ART: the extended Ritz formula has a positive second coefficient for penetration and a negative one for polarisation, so the approach to δ_0 is monotone from above or below. Stated in NIST compendium.

P.perturb — series perturbation

a large spread measures a perturber, not bad data

a state of the same symmetry crossing the series

grade **MEASURED**· source R 696, 704, 741, 742; Fano 1961; Lu & Fano 1970· depends on P.qdt· depth 1

Si I nd spreads 0.15-0.21 where ASD's own leading-percentage column names 3s3p3 at 14%

PRIOR ART: a perturbed Rydberg series is a series crossed by a level of another channel — Fano, Effects of configuration interaction on intensities and phase shifts, Phys. Rev. 124 (1961) 1866-1878; Lu & Fano, Graphic analysis of perturbed Rydberg series, Phys. Rev. A 2 (1970) 81-86.

P.polar — core polarisation

beyond $l=3$ the defect follows core POLARISABILITY in direction, not in magnitude

$l \geq 4$, where the orbital does not reach the core

grade **MEASURED**· source R 731-733, 788, 792-795; Born & Heisenberg 1924; Seaton 1958· depends on P.lcollapse· depth 2

grows with Z at +0.0005/unit over seven elements; a quantitative α_d fit was attempted and did NOT work, cause undiagnosed (R 794)

PRIOR ART: core polarisation as the source of the high- l defect is Born & Heisenberg, Über den Einfluss der Deformierbarkeit der Ionen

auf optische und chemische Konstanten, Z. Phys. 23 (1924) 388-410,
and Mayer & Mayer, Phys. Rev. 43 (1933) 605.

P.qdt — core penetration

the quantum defect measures how far a Rydberg orbital reaches into the ionic core

a Rydberg series with three or more members, or two with a published limit

grade **MEASURED**· source R 693, 707, 713; Seaton 1958, 1983· 12 objects depend on it· depth 0

four independent confirmations, none encoded

PRIOR ART: quantum defect theory — Seaton, The quantum defect method, MNRAS 118 (1958) 504-518, and Quantum defect theory, Rep. Prog. Phys. 46 (1983) 167-257.

P.selfsame — series self-consistency

one series in disjoint n windows gives one defect

windows where P.lens's amplification is comparable

grade **MEASURED**· source R 747, 748, 751; Ritz 1903· depends on P.qdt· depth 1
67 of 72 channels agree between disjoint halves to better than 0.05 (R 811)

PRIOR ART: one unperturbed series has one defect; disjoint n-windows must agree. The consistency test of the Ritz form (1903).

P.termsplit — core angular structure

at l=3 delta splits by the core's TERM while its J-pairs stay together

j-K coupled series on one core

grade **MEASURED**· source R 739; Condon & Shortley 1935· depends on P.qdt· depth 1

12 of 15 groups have terms separating by more than their J-pairs — median within 0.0011 vs between 0.0462 (R 826)

PRIOR ART: the splitting of a Rydberg series by the core term, with J-pairs staying together, is the parent-term structure of Condon & Shortley (1935), ch. VII.

P.trunc — truncation loss

truncation removes most channels and degrades those it leaves

a series cut at low n

grade **MEASURED**· source R 675, 676, 677, 705, 706; Inglis & Teller 1939· depth 0
Ne I: 33 Handbook levels give 0 channels, the full table gives 4

PRIOR ART: a Rydberg series is truncated in practice by field ionisation and by plasma microfields — Inglis & Teller, Ionic depression of series limits in one-electron spectra, Astrophys. J. 90 (1939) 439-448.

Q. The channel equation — the closed form and its terms — 11 objects

Q.alpha — the polarisability index

Λ_{α} : the polarisability index over cores, coordinates Z . Ne(core) . n_{out} . l_{out}

the set of cores appearing in the spectra index

grade **MEASURED**· source *R 1165; Born & Heisenberg 1924; Mayer & Mayer 1933*· depends on S.regime· 1 object depends on it· depth 9

18 cores held, $E = 10$. Sorted by $(n_{\text{out}}, l_{\text{out}})$ the isoelectronic ordering is a physical requirement — alpha must FALL with Z along a sequence — and the Ne-like and Mg-like rows obey it while the Ar-like row does not: K I 5.490 published, Ca II 6.665 extracted, Sc III 4.705. The index makes the violation visible where a table would not

PRIOR ART: core dipole polarisability as the origin of the high- l defect is Born & Heisenberg, Z. Phys. 23 (1924) 388-410; the systematic values are Mayer & Mayer, The polarizabilities of ions from spectra, Phys. Rev. 43 (1933) 605-611.

Q.anchor — the two-ended anchor

anchoring the equation at BOTH ends of Z fixes the extrapolation at almost no in-region cost

the channel equation and a far-field literature value

grade **MEASURED**· source *R 1201-1205; Theodosiou, Inokuti & Manson 1986*· depends on Q.region, Q.collapse· 1 object depends on it· depth 13

seven far anchors — Cs I np measured at 3.5667, and Th/Ac ns, np, nd, nf from actinide theory at 5.2, 4.75, 3.8, 2.0 — extend the sample from $Z \leq 83$ to $Z = 90$. Far-anchor median error falls 0.754 to 0.064, a factor of twelve; in-region rises 0.0637 to 0.0665. The saturating exponent returns for real: $e(\text{Ne}) = 0.8297 - 0.0900 \ln \text{Ne}$ against -0.0266 from the in-region sample alone. CAVEAT: four of the seven anchors are THEORETICAL, so the high- Z arm is calibrated against another calculation and moves if that is revised

PRIOR ART: asymptotic quantum defects for all ionisation stages of all ions with $Z \leq 50$ are tabulated in Theodosiou, Inokuti & Manson, At. Data Nucl. Data Tables 35 (1986) 473-486, Hartree-Slater. The far anchors used here are Cs I np (measured) and actinide values from arXiv:2508.06733.

Q.bound — the Pauli bound

$B = \min(p, n_0 - l - 1)$: the integer part's exceptionless upper bound, from aufbau alone

a channel and its core's ground configuration

grade **PROVED**· source R 1141; Pauli 1925; Janet 1929· depends on S.ground· 5 objects depend on it· depth 7

$\text{floor}(\delta) \leq B$ for 311 of 311 measured channels. As an equality it is right 57%; every error is negative; within two of the bound, 303 of 311. p is the core's orbital count at that l and n_0 the first Pauli-allowed n , both from aufbau with no spectrum

PRIOR ART: the bound counts core orbitals of the same l and the first Pauli-allowed principal number, both read from the ground configuration. Paulis exclusion principle, Z. Phys. 31 (1925) 765-783, fixes the occupancies; Janets $n+l$ ordering (1929) fixes which subshells are filled.

Q.bridge — the bridge equation

the bridge between the metric and ordinal descriptions is MULTIPLICATIVE with a REGIME factor: every factor positive so each preserves rank by itself, and the regime factor differs between two channels only when they differ in regime

a channel equation required to respect both values and orderings

grade **MEASURED**· source R 1169-1171; Pareto 1896; Spearman 1904· depends on Q.delta, S.regime· 1 object depends on it· depth 12

Stated on the channel equation then current, Q.delta, which Q.final superseded at register 1205; the claim was re-tested there on the standing form and holds (one ℓ -ordering violation in 328 channels). The dependence line keeps Q.delta because Q.final descends from this object, and the graph records the order of establishment, not present support (register 1750).

an additive fit gets P.lcollapse 102/205 against a measured 194 — chance. A purely multiplicative fit gets 205/205, 143/143, 66/66 where the measurements give 194, 89, 55 — it cannot represent an exception because a product of positives is monotone in each factor. With a regime factor: 204/205, 129/143, 61/66, and $\ln-R^2$ improves 0.8026 to 0.8164

PRIOR ART: that a multiplicative form preserves rank and an additive one does not is elementary; that no single form does both well is a Pareto statement about the two objectives.

PRIOR ART: a product of positive factors is monotone in each and so preserves rank; a sum of signed terms need not. Rank preservation as an objective distinct from least squares is Spearman, The proof and measurement of association between two things, Amer. J. Psychol. 15 (1904) 72-101. That no single form does both well is a Pareto statement (1896).

Q.collapse — the Janet collapse

the ORBITAL COLLAPSE threshold is a Janet block boundary, exactly

a channel with $p = 0$ at $l \geq 2$

grade **MEASURED**· source *R 1187-1190; Goeppert-Mayer 1941; Griffin, Andrew & Cowan 1969*· depends on S.ground, Q.bound· 2 objects depend on it· depth 8

the $n+l = 5$ block opens at $Z = 21$ and 3d collapses at 21; $n+l = 7$ opens at 57 and 4f at 57; $n+l = 8$ opens at 89 and 5f at 89. Three exact matches. Across 116 $p = 0$ channels at $l = 2$ or 3: collapsed median 0.637, uncollapsed 0.036, U-test $p = 9.8e-4$. Adding the term takes Ti IV nd from -0.620 to -0.056 and the overall rms from 0.1825 to 0.1411. The largest outliers are atoms APPROACHING a boundary — Ca I nd = 0.908 at $Z = 20$ against a threshold of 21, Ba II nf = 0.756 at 56 against 57 — so the collapse is a rapid transition, not a step

PRIOR ART: orbital collapse — the sudden contraction of the 3d and 4f wavefunctions as Z crosses a threshold — is Goeppert-Mayer, Rare-earth and transuranic elements, Phys. Rev. 60 (1941) 184-187, and Griffin, Andrew & Cowan, Theoretical calculations of the d-, f- and g-electron transition series, Phys. Rev. 177 (1969) 62-71. What is measured here is that the threshold coincides with the Janet block boundary at $Z = 21, 57$ and 89.

Q.delta — the channel equation

delta = [B - q e[^](-a(Ne)l) e[^](k/Ne) c[^]g + P] (1 + s [triplet]): a closed form for any Rydberg channel from atomic-index quantities alone

a Rydberg channel (Z, charge c, l, multiplicity)

grade **MEASURED**· source *R 1145-1168; Seaton 1958; Fermi 1928; Pauli 1925*· depends on Q.bound, Q.pen, Q.pol, Q.exch· 2 objects depend on it· depth 11

311 measured channels: rms 0.2449, R^2 0.924, median |error| 0.090. By l: s 0.283, p 0.276, d 0.280, f 0.119, g 0.0096. Against TWELVE published values never fitted on: median error 0.00155 at $l \geq 3$ and 0.368 below. 50% within 0.090, 90% within 0.414, 99% within 0.826

PRIOR ART: the superseded form of the channel equation. Its terms are the Pauli bound (Pauli 1925), the Thomas-Fermi deficit (Fermi 1928) and Seaton polarisation (1958). Superseded by Q.final at register 1205.

Q.exch — the exchange factor · GRADED OPEN

the exchange factor (1 + s [triplet]), s = -0.0782

a channel of a two-valence-electron system

grade **ASSERTED**· source *R 987, withdrawn R 1168*· depends on Q.bound· 1 object depends on it· depth 8

WITHDRAWN (R 1168). The claim was that the fitted s recovers register 987's sign and size. It does not: register 987 measures the triplet defect EXCEEDING the singlet

in 22 of 22 ns cases, and the additive fit returned $s = -0.0782$, the right magnitude and the WRONG SIGN. Cross-checked on 66 singlet/triplet pairs the equation gets 33, exactly chance. The parameter absorbed something else

PRIOR ART: the exchange splitting between singlet and triplet is Heisenberg, Mehrkoerperproblem und Resonanz in der Quantenmechanik, Z. Phys. 38 (1926) 411-426; that the higher multiplicity lies lower is Hund first rule (1925). WITHDRAWN at register 1168: the fitted sign is opposite to the measured one.

Q.final — the channel equation

$\delta = a p^e(\text{Ne}) \text{Ne}^k \ln(c+1)/c$ where $p > 0$, and $h C(Z) ((\text{Ne}-1)/\text{Ne}) \text{Ne}^k \ln(c+1)/c$ where $p = 0$

any Rydberg channel (Z , charge, l , multiplicity)

grade **MEASURED**· source *R 1205-1206; Seaton 1958; Fermi 1928; Janet 1929*· depends on Q.anchor, Q.bound· depth 14

$a = 0.3772$, $e(\text{Ne}) = 0.8297 - 0.0900 \ln \text{Ne}$, $k = 0.4942$, $h = 0.5415$. 284 channels from $Z = 2$ to 90 and charge 1 to 10: rms 0.1610, R^2 0.9741, $l \geq 4$ rms 0.0150, hydrogenic output EXACTLY zero, Pauli bound 328/328, one l -ordering violation against a measured zero. Four fitted numbers and every input from the ground state or the periodic table. It values all 1,648 cells R places, where the walk valued none

PRIOR ART: every term of the form has a source. The $\ln(c+1)/c$ charge dependence is the isoelectronic behaviour of Edlen 1964; the Ne^k factor is Thomas-Fermi (Fermi 1928); p is the Pauli orbital count (Pauli 1925); $C(Z)$ is orbital collapse (Griffin, Andrew & Cowan 1969) at the Janet boundary (Janet 1929). The equation assembles them; it does not introduce any.

Q.pen — the deficit term

the penetration deficit, $-q e^{(-a(\text{Ne}) l)} e^{(k/\text{Ne})} c^g$, with $a(\text{Ne}) = a_0 + a_1 \text{Ne}^{(-1/3)}$

a channel below the polarisation regime

grade **COMPUTED**· source *R 1147; Hartree 1928; Fermi 1928*· depends on Q.bound· 1 object depends on it· depth 8

the decay a is set by the atom's SIZE: a against $\ln \text{Ne}$ gives $r^2 = 0.437$ across 56 species, against electrons-beyond-closure 0.143, against charge nothing ($p = 0.34$). K I's measured d/s ratio implies $a = 1.03$, Sr I's implies 0.157 — a factor of seven, and a universal a fits neither

PRIOR ART: the penetration deficit and its $\text{Ne}^{(1/3)}$ scaling are the Thomas-Fermi picture — Fermi, Eine statistische Methode zur Bestimmung einiger Eigenschaften des Atoms, Z. Phys. 48 (1928) 73-79; Hartrees self-consistent field, Proc. Camb. Phil. Soc. 24 (1928) 89-110, gives the orbital form. Fermi 1928 applied it to Rydberg corrections directly.

Q.pol — Seaton's term

the polarisation term, $3\alpha c^2 / K(l)$ for $l \geq 4$ with α from Λ_α , $K(l) = l(l+1)(2l-1)(2l+1)(2l+3)$

a non-penetrating channel and a core polarisability

grade **CITED**· source R 1165-1168; Seaton 1958; Born & Heisenberg 1924· depends on A.seaton, Q.alpha· 1 object depends on it· depth 10

Seaton's formula. With α supplied as a species LABEL from Λ_α rather than a universal constant, the $l \geq 4$ rms falls from 0.01504 to 0.00958 — 36%. Below $l = 4$ it must be gated OFF: applied at s and p it drives the fit to rms 39.3 because $K(l)$ is small there

PRIOR ART: the polarisation term $3\alpha c^2 / K(l)$ is Seaton (1958); the physical origin is core polarisability, Born & Heisenberg, Z. Phys. 23 (1924) 388-410. SUBSUMED at register 1204 by the Janet collapse coordinate.

Q.region — the validated domain

the equation is VALIDATED in the verified-plus-possible region and EXTRAPOLATED outside it

the channel equation and the existence partition

grade **MEASURED**· source R 1193-1194; Theodosiou, Inokuti & Manson 1986· depends on Q.delta, S.status, Q.collapse· 1 object depends on it· depth 12

Stated on the channel equation then current, Q.delta, which Q.final superseded at register 1205; the claim was re-tested there on the standing form and holds (in-region error 0.147, far-anchor 0.064). The dependence line keeps Q.delta because Q.final descends from this object, and the graph records the order of establishment, not present support (register 1750).

277 in-region channels give rms 0.1329 and R^2 0.9747 — better than the 311-channel fit on fewer points, because the 51 excluded were 24 three-parent cores, 15 sixteen-parent cores and 12 ions above charge 10. And the saturating exponent falls from 1.3257 - 0.2751 ln Ne to 0.5828 - 0.0266 ln Ne: nine-tenths of the Ne-dependence was open-shell contamination. Validated on 277, extrapolated to 98,078

PRIOR ART: the distinction between where a defect is measurable and where it is calculated is exactly the distinction their 1986 table makes — they compute Hartree-Slater values everywhere and note where experiment exists.

LS. The Löwdin solution — the order derived from the equation — 8 objects

Chapter 35 and its companion paper; register 1701-1712. These objects are verified by the solution's own sealed instruments and receipts, not yet by this volume's own checks; that check is outstanding, and until it is done this page states the verification

that exists rather than implying one that does not. The mathematics of the challenge (Chapter 34, registers 1249–1357) is stated separately below under THE MATHEMATICS OF THE LÖWDIN WORK.

LS.ent — the entrant operator

ent(Z) = argmax over frontier (n,ℓ) of |D(n,ℓ)|, D the converged one-channel depth in the self-consistent field of the ion (Z, cfg(Z–1)); scalar-relativistic Koelling-Harmon; c = 137.035999 the only entered number. Chained, cfg(Z) = cfg(Z–1) + ent(Z): 107 of 107 against the measured ground configurations, Z = 2–108

the operator Chapter 27 said a non-closing index requires; sign-exact by construction, and the margin |D(ent)| – |D(runner-up)| is its own error bar, recorded at every step

grade **COMPUTED**· source M §35; register 1701, 1702· depends on L.closed, Q.final· depth 15

LS.law — the ordering law, five clauses

Clause 1 (ordering): smaller n+ℓ opens first — 107/107. Clause 2 (tie-break): equal n+ℓ, smaller n first — exceptions exactly {La, Ac, Th}, derived by LS.coll. Clause 3 (correlation): at the five contested rows the second-order differential is positive — every competition widens. Domain clause: $Z \leq 112$; spin-orbit worst case 0.083 Ha under every margin. Relativistic clause: the $c \rightarrow \infty$ twin disagrees at eleven elements

statement and proof form in the companion paper

grade **PROVED**· source M §35; register 1701–1706· depends on LS.ent, LS.coll, LS.twin· depth 17

LS.coll — the collapse condition

The double-well criterion, stated from the field, for which f channel is collapsed at which Z: decides exactly the Clause-2 exception set and occupies exactly the domain where Chapter 34's corridor is silent ($L = -\infty$ at f openings)

adjacent to the transition the mean-field equations admit two stationary solutions of one configuration with distinct converged operators; a sign at SCF tolerance there is branch content, not noise

grade **COMPUTED**· source M §35; register 1703; Griffin–Andrew–Cowan 1969, 1971· depends on LS.ent· depth 16

PRIOR ART: the orbital-collapse double well is Griffin, Andrew & Cowan, Phys. Rev. 177 (1969) 62; the placement of it against the corridor's silence is this work's.

LS.pin — the pinned-channel theorem (no g block)

Every g channel offered by the walk sits at $-1/(2n^2)$ to storage precision: 5g over 65 elements, 6g over 70, 7g over 57, 8g over 28. $dn^*/dZ = 0$ across a hundred protons

the absence of collapse, not its slow approach; a relation the data cannot violate is defending something — here, the nonexistence of a g period below $Z = 121$

grade **MEASURED**· source *M §35; register 1704*· depends on LS.ent· depth 16

LS.asym — the state-dependent multiplier identity

For u, v eigenstates of different self-consistent operators, the two eigen-relation evaluations of $\langle u|T|v \rangle$ differ by exactly $\text{asym}(u,v) = (\epsilon_v - \epsilon_u)\langle u|v \rangle - \langle u|(V_v - V_u)|v \rangle + [\langle u|X_v \rangle - \langle v|X_u \rangle]$; verified to machine precision, four of four elements, worst $2.6 \cdot 10^{-15}$, independent of SCF tolerance

a residual of large cancelling terms, not a convergence artefact; it reclassified a registered instrument fault as derived content

grade **PROVED**· source *M §35; register 1708, 1709; Löwdin 1950*· depends on LS.ent· depth 16

PRIOR ART: an instance of the non-orthogonality problem, Löwdin, J. Chem. Phys. 18 (1950) 365.

LS.quart — the exact-quartic decomposition

The energy functional with one shell's orbital varying along a fixed direction is exactly quartic in the path parameter (one-body quadratic; two-body quartic); five evaluations at $t \in \{0, \pm 1/2, \pm 1\}$ determine every Taylor coefficient with zero truncation error

the floating-point floor at $|E| \sim 10^4 \text{ Ha}$ ($\sim 10^{-12}$) is the only limit, and it is stated, not implied; protocol

grade **PROVED**· source *M §35*· depends on LS.ent· depth 16

LS.chord — the defect closed: chord = rot + perp

The walk's one systematic internal discrepancy — the Hellmann-Feynman-in- q defect — is a Pulay term of the occupation parameter. In the one-shell-frozen gauge it splits exactly: rot = linear gradient law + $(q \cdot s/dq^2) \cdot (\text{asym})$ + endpoint-Hessian term; perp = the first-order perturbed-HF response on the orthogonal complement. Balance: zero unexplained residue

the signed trace verified at ratios 0.999992 and 1.000103; the endpoint-Hessian term ($+1.7 \cdot 10^{-7}$, $+6.1 \cdot 10^{-9}$) is twenty to five hundred times too small for the residual it was hypothesised to explain — a hypothesis falsified by the clause that scored HIT, and kept

grade **PROVED**· source *M §35; register 1707, 1709, 1711; Pulay 1969; Gerratt-Mills 1968*· depends on LS.asym, LS.quart· depth 17

PRIOR ART: the occupation-parameter term is Pulay, Mol. Phys. 17 (1969) 197; the orthogonal-complement response is Gerratt & Mills, J. Chem. Phys. 49 (1968) 1719.

LS.twin — the twin operator, $c \rightarrow \infty$

The identical entrant operator with the constant removed disagrees with the $c = 137.035999$ operator at eleven elements — Mn, Zn, Ag, Cd, Nd, Pm, Sm, Lu, Hg, Lr, Rf — every disagreement an error against nature, since the relativistic operator scores 107/107

the relativistic clause of LS.law, measured rather than asserted

grade **MEASURED**· source *M §35; register 1706*· depends on LS.ent· depth 16

3B. Three bodies — the index of families — 9 objects

Chapter 36 and its companion paper; register 1713–1724. One new root, 3B.shape — shape space, standard mathematics cited as found (Montgomery 2014; Hopf 1931; register 1739) — and otherwise the family derives from L.c1–L.c8, B.brk and the operators, and the seed did not move — the cycle’s falsifier passing on a new subject.

3B.shape — shape space and the shape sphere

$\mathbb{R}^3 = \mathbb{C}^3 / (\text{translations} \times \text{rotations})$; $S^2 = \text{that} / \text{scale}$. **Onto; identifies exactly the oriented-congruent triangles; only triple collision maps to 0; w_3 = signed area up to a mass constant; $\|w\| = I/2$**

the coordinates in which three bodies become one point

grade **PROVED**· source *Montgomery 2014, Thm 1; Hopf 1931*· 2 objects depend on it· depth 0

3B.metric — the shape metric

$ds^2 = |dw|^2 / (2\sqrt{\|w\|})$; **every plane through 0 totally geodesic; cone metric $dr^2 + \frac{1}{4}r^2 d\theta^2$**

as 3B.shape

grade **PROVED**· source *Montgomery 2014, Thm 3*· depends on 3B.shape· depth 1

3B.JM — the Jacobi-Maupertuis metric

$g_E = (E+U) \cdot ds^2$; **trajectories at energy E are its geodesics; $dt = ds_E / \sqrt{2(E+U)}$**

licence in the index: §12.11.1.3, §12.11.0.2

grade **PROVED**· source *Maupertuis 1744; Jacobi 1837*· depends on 3B.metric, 3B.pot· depth 2

3B.pot — the potential on shape space

$U(w) = \sum c_{ij}/d_{ij}$, $c_{ij} = (m_i m_j)^{3/2} / \sqrt{(m_i + m_j)}$, $d_{ij}^2 = \|w\| - w \cdot b_{ij}$; **identity $r_{ij}^2 = d_{ij}^2 / \mu_{ij}$. No hyper-radius factor**

650 triangles, 13 mass cases, error $< 10^{-10}$; the first audit’s uniform 13/13 failure was a hyper-radius divisor inherited from working material — register 1718, protocol 1722

grade **PROVED**· source *Montgomery 2014 §11*; register 1718; main §D.5.9· depends on 3B.shape· depth 1 · *checked numerically on 650 random triangles at 10^{-10} — corroboration, not the verification*

3B.norm — the algebraic variety

With $u_{ij} = c_{ij}/d_{ij}$, $p = \sum u^2$, $q = \sum u_i^2 u_j^2$, $r = \prod u^2$: $U^8 - 4pU^6 + (6p^2 - 8q)U^4 - 4(p^3 - 4pq + 16r)U^2 + (p^2 - 4q)^2 = 0$; constant term $\prod (u_1 \pm u_2 \pm u_3)^2$

proved symbolically (sympy) and measured at 13 mass cases; an inherited form with U^4 coefficient “ $6S_2^2 - 4S_4 + 8S_{11}$ ” = $2p^2 + 16q$ was wrong and is withdrawn — register 1719

grade **PROVED**· source *Lagrange 1770*; register 1719, 1720· depends on 3B.pot· depth 2

PRIOR ART: N_8 is the norm over $(\mathbb{Z}/2)^3$ — Lagrange’s resolvent method, 1770. Not new mathematics.

3B.five — the five fixed points

Three Euler roots (one positive root of the quintic per ordering) and two Lagrange points, for every mass triple — 13/13

seed of the family index in §14.5’s sense


grade **PROVED**· source *Euler 1767*; *Lagrange 1772*; register 1717; main §D.5.9· depends on 3B.pot· depth 2 · *the thirteen-order-type check is corroboration, not the verification*

3B.tri — the triangle form on K_3

**$\{|a-b| \leq c \leq a+b\}$: join-closed, meet-broken. Cap 8: 344 cells, 0 join failures, 8,385 meet failures; caps 3-12 meet failures
12·111·477·1,488·3,780·8,385·16,812·31,227·54,555·90,705; two-body chain 0 throughout**

confirms §12.11.2 on a grid the book never ran

Fig. 2 §12.11.2 on three bodies: join-closed, meet-broken; two bodies close.



meet and join failures against cap; the two-body chain at zero throughout

grade **MEASURED**· source M §36; register 1716· depends on L.c1–L.c8· depth 1

3B.def — the deficit

**K_3 has treewidth 2; strong 3-consistency is required; $\mathcal{R}$ reaches 2 (§32.3).
Shortfall exactly one level**

the separation hypothesis of §7.1 broken by the smallest graph that can break it

Fig. 5 The blind is exactly on the level (§21.5.1); the tree closes the triangle does not. $\mathcal{R}$ reaches level 2 = needed needs strong 3-consistency; $\mathcal{R}$ gives 2



K_2 against K_3 , and the consistency level each requires

grade **PROVED** · source M §21.5.1; Freuder 1982 · depends on 3B.tri, G.graph, L.tree · depth 3

3B.index — the index Λ_3

$\Lambda_3 = \{\text{KAM, per, chaos, erg, coll}\}$ on $\mathcal{M}_{\{E,L\}}$. $E(\Lambda_3) = 0$: exhaustive (Chazy classes), disjoint up to measure zero (Saari 1971/73; Painlevé, $n = 3$). By §25.6, predictions = 0 — Brudno's theorem on the chaotic stratum

cited theorems assembled; the assembly is this work's. The maximal forbidden case of Chapter 18 is the solved case, because what is forbidden is exactly what Poincaré excludes — register 1724

grade **PROVED** · source M §36; register 1713, 1714, 1724; Chazy 1922; Saari 1971; Brudno 1983 · depends on 3B.def, 3B.five · depth 4

V · THE CHAINS

The longest derivation paths in the register — what rests on what.

depth 14 · A.alph $\leftarrow$ A.env $\leftarrow$ A.R $\leftarrow$ K.redun $\leftarrow$ K.axis $\leftarrow$ K.produce $\leftarrow$ S.ground $\leftarrow$ S.ritz $\leftarrow$ S.regime $\leftarrow$ Q.alpha $\leftarrow$ Q.pol $\leftarrow$ Q.delta $\leftarrow$ Q.region $\leftarrow$ Q.anchor $\leftarrow$ Q.final

depth 13 · A.alph $\leftarrow$ A.env $\leftarrow$ A.R $\leftarrow$ K.redun $\leftarrow$ K.axis $\leftarrow$ K.produce $\leftarrow$ S.ground $\leftarrow$ S.ritz $\leftarrow$ S.regime $\leftarrow$ Q.alpha $\leftarrow$ Q.pol $\leftarrow$ Q.delta $\leftarrow$ Q.region $\leftarrow$ Q.anchor

depth 13 · A.alph $\leftarrow$ A.env $\leftarrow$ A.R $\leftarrow$ K.redun $\leftarrow$ K.axis $\leftarrow$ K.produce $\leftarrow$ S.ground $\leftarrow$ S.ritz $\leftarrow$ S.regime $\leftarrow$ Q.alpha $\leftarrow$ Q.pol $\leftarrow$ Q.delta $\leftarrow$ Q.bridge $\leftarrow$ A.blind

depth 12 · A.alph ← A.env ← A.R ← K.redun ← K.axis ← K.produce ← S.ground ← S.ritz ← S.regime ← Q.alpha ← Q.pol ← Q.delta ← Q.bridge

depth 12 · A.alph ← A.env ← A.R ← K.redun ← K.axis ← K.produce ← S.ground ← S.ritz ← S.regime ← Q.alpha ← Q.pol ← Q.delta ← Q.region

depth 11 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← L.birk ← L.bits ← L.circuit

depth 11 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← L.birk ← L.pushback ← L.step

depth 11 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← I.interval ← I.convex ← EM.spin

depth 11 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← I.interval ← I.convex ← EM.parity

depth 11 · A.alph ← A.env ← A.R ← K.redun ← K.axis ← K.produce ← S.ground ← S.ritz ← S.regime ← Q.alpha ← Q.pol ← Q.delta

depth 10 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← L.birk ← L.alpha

depth 10 · A.alph ← A.env ← A.R ← A.ext ← A.clos ← A.fix ← A.rule ← L.closed ← L.dist ← L.birk ← L.mobius

depth 17 · A.alph ← A.env ← A.R ← K.redun ← K.axis ← K.produce ← S.ground ← S.ritz ← S.regime ← Q.alpha ← Q.pol ← Q.delta ← Q.region ← Q.anchor ← Q.final ← LS.ent ← LS.asym ← LS.chord · *the Löwdin solution's longest path, through the channel equation; the walk's one defect closed at zero residue*

depth 4 · L.c1 ← G.cons ← G.graph ← 3B.def ← 3B.index · *the three-body index; the family's own links rest on L.c1-L.c8, G.graph and L.tree, roots already in the register — closes to §25.6 and the Brudno rate*

VI · WHAT IS UNFINISHED

Neither the Löwdin solution nor the three-body work adds an open object; the two below are untouched by both.

Two objects. *Neither is unfinished for want of a proof attempt. Each has a diagnosis, and the diagnosis names what is missing rather than restating that something is.*

M.C2 — half-sided modular inclusion on a non-expanding horizon

$$\Delta_{\{M(u_1)\}^{\wedge\{it\}} M(u_2) \Delta_{\{M(u_1)\}^{\wedge\{-it\}}} \subseteq M(u_2) \quad \text{for } t \leq 0, u_1 < u_2$$

What is settled. Half-sided modular inclusion is CHARACTERISED — Borchers, *The CPT theorem in two-dimensional theories of local observables*, Commun. Math. Phys. **143** (1992) 315–332, and Wiesbrock, *Half-sided modular inclusions of von Neumann*

algebras, Lett. Math. Phys. **28** (1993) 107–114: the inclusion holds exactly when a one-parameter unitary group with positive generator implements the translation. The vacuum is cyclic and separating for local algebras (Reeh & Schlieder 1961), and modular theory supplies Δ and J (Tomita 1967; Takesaki 1970). **None of that is in question.**

What is missing, precisely. The hypothesis — *a non-expanding horizon with no Killing field, ω Hadamard* — asks the expansion $\Theta = 0$ to select what only a STATE can select, and offers Hadamard, a microlocal condition, as the selector.

The horizon’s own definition supplies a state condition at BACKGROUND order — Einstein’s equation with future-causality forces vanishing shear, vanishing null flux and $\mathbf{f}_\ell \mathbf{q}_{ab} = 0$ — and leaves a gap at PERTURBATION order. *That gap is the object.*

Where the obstruction sits. The free half is confirmed numerically: the spectral weight ratio is of order 10^{-5} . The obstruction is in the corner edge modes, and the proposal $\Theta = 0$ was refuted by the transformation law. Sorce (2024) closes the geometric route by construction — a geometric modular flow must be generated by a conformal Killing field — so a horizon with no Killing field cannot have one, and the route that remains is algebraic.

What would settle it. The covariance of conditional expectations across the full family of cuts, in the manner of an over-determined joint fit. *Chandrasekaran & Flanagan (arXiv:2601.07915) is the nearest published treatment and was read into the diagnosis at registers 1037–1042.*

Q.exch — the exchange factor

$$\delta = [\dots] \cdot (1 + s \cdot [\text{triplet}]), \quad s = -0.0782$$

What is settled. Exchange splitting between singlet and triplet is Heisenberg, *Mehrkörperproblem und Resonanz in der Quantenmechanik*, Z. Phys. **38** (1926) 411–426; that the higher multiplicity lies lower is Hund’s first rule, Z. Phys. **33** (1925) 345–371. **The physics is not in doubt.**

What is missing, precisely. *The fitted sign is opposite to the measured one.* The factor was withdrawn at register 1168 for that reason and the grade left as ASSERTED rather than removed, because the term is real and the form is wrong.

A uniform s cannot work. *Exchange acts through the overlap of the Rydberg orbital with the core, and that overlap falls sharply with ℓ . A single multiplicative constant can only give 0 of 66 triplet-above-singlet pairs or 66 of 66; the measurement is neither.*

What would settle it. An ℓ -dependent exchange term, fitted against the singlet–triplet pairs the compendium holds. The Pauli bound (Q.bound) already carries the orbital count p that the overlap should follow.

VII • THE BIBLIOGRAPHY

Every object of this compendium names a work. What follows is those works, ordered by year, with the objects each carries. *The book claims nothing new where an earlier result will do; where a measurement is this work's, the object says so.*

162 works, 1669-2026. Ten rows the generator had read out of callout text — status words, a possessive, a method name — were removed on 2026-08-24 (register 1736); the count was 172.

year	work	objects
1669	Newton	B.newton
1682	Leibniz	B.brk
1687	Newton	B.ordbr
1715	Taylor	B.frac B.pole
1736	Euler	G.graph
1744	Maupertuis	3B.JM
1748	Euler	A.prodE C.Aq C.Bq C.box L.F L.F1 L.Fm1
1767	Euler	3B.five
1770	Lagrange	3B.norm
1772	Lagrange	3B.five
1797	Lagrange	B.hstar
1809	Gauss	A.anchor B.adm C.qmean L.skew
1837	Jacobi	3B.JM
1842	Jacobi	3B.JM
1854	Boole	A.logic
1869	Mendeleev	E.ioniz E.table
1878	Hill	3B.index
1890	Rydberg	B.V43 B.Vexact B.fail B.floor32 B.nuV B.silence P.converge P.lens
1890	Poincaré	3B.index
1893	Heaviside	L.chi
1896	Pareto	B.pareto C.pareto Q.bridge
1900	Dedekind	L.modular
1903	Ritz	P.converge P.mono P.selfsame S.regime S.ritz
1904	Spearman	Q.bridge
1906	Jensen	B.floor2 B.floor32

year	work	objects
1907	Fubini	C.fib
1910	Moore	A.E A.R A.bound A.clos A.expr A.ext A.fix F.exact F.moore L.E0 S.bounds S.open S.seed
1911	Caratheodory	S.car
1913	Bohr	B.coll L.c1 L.c4 L.def
1913	Pauli	L.def
1916	Sommerfeld	P.jsplit
1918	Noether	EM.quotient
1922	Chazy	3B.index
1923	Helly	K.helly
1924	Born & Heisenberg	P.polar Q.alpha Q.pol
1924	Laporte	EM.cross EM.map EM.parity
1924	Stoner	L.c2 L.c5
1925	Hund	L.c7 Q.exch S.ground
1925	Pauli	L.c2 L.c3 L.c5 L.c6 L.c7 L.c8 L.def Q.bound Q.delta Q.final T.a9p
1925	Russell & Saunders	EM.spin
1926	Aitken	B.aitken
1926	Schroedinger	L.c1
1927	Wigner	EM.image EM.parity
1928	Dirac	P.jsplit
1928	Fermi	Q.delta Q.final Q.pen
1928	Hartree	P.lcollapse Q.pen S.ritz
1928	Sperner	L.sperner
1929	Janet	E.layout E.table Q.bound Q.final S.ground
1931	Wigner	EM.spin T.a12 T.dens T.excl T.invariant
1931	Hopf	3B.shape
1933	Mayer & Mayer	Q.alpha
1933	Milne-Thomson	B.V B.brk B.ordbr B.ordk B.pole
1935	Condon & Shortley	EM.map P.jj P.termsplit S.channel S.parent T.a11 T.a13 T.real T.scheme T.tower
1936	Eckart & Young	B.rank1

year	work	objects
1936	Madelung	L.real S.ground
1936	Tarski	A.logic C.compare
1936	Murray & von Neumann	M.semi
1937	Birkhoff	A.define A.erel I.convex K.arrow K.produce L.alpha L.arith L.birk L.bits L.dist L.omega L.pushback L.step S.box S.seed
1937	Shannon	L.bits
1938	Shannon	L.circuit S.bits W.onehot
1939	Inglis & Teller	P.trunc
1940	Birkhoff	A.derived A.erel A.morph A.prod I.convex I.interval I.shape L.arith L.closed L.total
1940	Deming & Stephan	A.stat2 G.stat
1941	Dushnik & Miller	A.erel K.axis L.dim
1941	Goeppert-Mayer	P.dcollapse Q.collapse
1942	Racah	T.a11 T.a12 T.dens T.dich T.invariant T.scheme T.tower
1942	Ward	A.clos F.moore F.open
1943	Racah	S.parent T.a10 T.a9 T.a9p T.excl T.tower
1945	Segre	E.nuclide
1948	Coxeter	W.face
1948	Shannon	A.blind A.ebits A.slack EM.notcomp G.reg K.axis K.redun K.transit
1950	Dilworth	S.box S.down
1950	Hamming	W.frontier
1950	Pais & Uhlenbeck	W.core9
1950	Löwdin	LS.asym
1953	Green	T.para
1954	Kolmogorov	3B.index
1955	Tarski	A.cert F.open
1957	Bethe & Salpeter	B.coll
1958	Pauli	Q.delta
1958	Seaton	A.seaton P.coreblind P.lcollapse P.polar P.qdt

year	work	objects
		Q.delta Q.final Q.pol
1959	Dijkstra	A.EW A.W A.intext A.three
1960	Erdos & Renyi	A.dens
1961	Fano	P.perturb
1961	Reeh & Schlieder	M.rs
1962	Berge	G.near G.tower K.girth K.peak
1963	Arnold	3B.index
1964	Edlen	A.S K.zcross P.buildlimit P.charge P.iso Q.final
1964	Moebius	G.book
1964	Rota	C.local G.book L.amp L.box L.mobius L.void L.voidfrac
1965	Kolmogorov	F.unstatable
1965	Penrose	W.core9
1967	Floyd	K.clock K.clockfail
1967	Tomita	M.tt
1968	Ireland and Kullback	A.stat2
1968	Alekseev	3B.index
1968	Gerratt & Mills	LS.chord
1969	Andrew & Cowan	P.dcollapse Q.collapse Q.final
1969	Griffin, Andrew & Cowan	LS.coll
1969	Pulay	LS.chord
1970	Codd	A.alph
1970	Fano	P.perturb
1970	Takesaki	M.tt
1971	Lane	K.cat K.comp K.deadend K.jump K.twocol
1971	Griffin, Andrew & Cowan	LS.coll
1971	Saari	3B.index
1972	Gabriel	K.quiver
1972	Karp	S.core S.cover S.erasure S.lam S.unit
1973	Takesaki	M.takesaki
1973	Saari	3B.index
1974	Curtis & Reid	B.hstar
1974	Johnson	S.cover S.lam

year	work	objects
1974	Montanari	A.montanari
1974	McGehee	3B.index
1975	Baker & Pixley	A.fix A.two
1975	Csiszar	G.stat
1975	Marchal & Saari	3B.index
1976	Monaghan	3B.index
1978	Freuder	A.blind A.gc A.modeA A.modeB A.three G.allcons G.prot K.corner W.jur
1978	Gratzer	L.pushback L.step
1978	Rissanen	A.ebits
1979	Chvatal	S.core S.unit
1979	Dawid	G.trip
1980	Stanley	L.sperner
1981	Monjardet	I.shape L.metric L.occ
1982	Freuder	A.freuder C.cut G.cons G.ref G.shape L.E0 L.box L.tree T.trad 3B.def
1982	Racah	T.trad
1982	Marchal & Bozis	3B.index
1983	Beer, Fagin, Maier & Yannakakis	A.intext C.compare K.three
1983	Maier & Yannakakis	A.intext A.relax C.compare K.langclose K.three K.window
1986	Inokuti & Manson	Q.anchor Q.region S.status
1986	Stanley	C.Aq C.Bq L.F L.F1 L.Fm1 L.chains L.pal L.rankpoly
1988	Pearl	G.trip K.decay K.markov
1989	Cooper	A.bpc
1989	Dechter & Pearl	G.cons G.shape K.coupling
1989	Nouguier & Vilarem	A.bpc
1991	Chinneck & Dravnieks	W.core15 W.core9 W.scale W.sup
1991	Cover & Thomas	K.decay
1991	Drake & Swainson	A.seaton
1991	Kay-Wald	M.C2

year	work	objects
1992	Borchers	M.C2 M.hsmi M.ledger
1992	Dechter	A.dechter A.gc
1993	Wiesbrock	M.C2 M.hsmi M.ledger
1994	Nesterov & Nemirovskii	B.newton B.selfconc
1995	Beek & Dechter	A.rule A.staircls
1996	Lauritzen	K.decay K.markov
1998	Montgomery	3B.shape
1999	Deville	A.env A.orient A.r4 T.tight
1999	Hentenryck	A.orient A.rule A.stair A.staircls T.tight
2000	Wald & Zoupas	M.C1
2000	Chenciner & Montgomery	3B.five
2002	Montgomery	3B.metric
2003	Fredenhagen & Verch	W.rel
2006	Hsiang & Straume	3B.shape
2014	Montgomery	3B.shape 3B.metric 3B.pot
2017	Diestel	G.graph
2019	Fleischer & Knauf	3B.index
2021	Kol	3B.index
2023	Kol	3B.index
2024	Sorce	M.sorce
2026	Lach, <i>The Löwdin Solution</i>	LS.ent LS.law LS.coll LS.pin LS.asym LS.quart LS.chord LS.twin
2026	Lach, <i>The Three-Body Problem for Unknown Masses</i>	3B.tri 3B.def 3B.index

The works this compendium leans on most

work	objects
Birkhoff, 1937	15
Moore, 1910	13
Pauli, 1925	11
Condon & Shortley, 1935	10
Birkhoff, 1940	10
Freuder, 1978	9
Freuder, 1982	9
Seaton, 1958	8

work	objects
Rydberg, 1890	8
Shannon, 1948	8
Stanley, 1986	8
Euler, 1748	7

THE MATHEMATICS OF THE LÖWDIN WORK

Registers 1249–1357. Each object is stated with its provenance: what is standard and whose it is, what is arithmetic, and what this work introduces.

The twelve objects the closing list names are entered in the main volume's Appendix D at §D.5.10 (register 1734), where they close at $E = 0$ with the corridor's precedent read as this work's.

The corridor — a system of linear inequalities

The object. Requiring an observed subshell g to have least v against every admissible rival r gives one inequality per rival,

$$a(\sqrt{p_r} - \sqrt{p_g}) < n_r - n_g, \text{ with } p = n - \ell - 1$$

whose solution is an interval with endpoints

$$L, U = \Delta n(\sqrt{p_g} + \sqrt{p_r})/(p_g - p_r)$$

Standard. Linear programming in one variable; the feasible set of a finite system of linear inequalities in $\mathbb{R}^1$ is an interval. *Fourier 1826; Motzkin 1936.*

This work. *That the coefficients are node counts, so the endpoints lie in $\mathbb{Q}(\sqrt{\mathbb{N}})$ and take only nineteen distinct values across the periodic table. And that all 106 intervals are non-empty — the system is consistent.*

The closed form for every $ns/(n-1)d$ competition:

$$a_{\text{cross}} = (\sqrt{(n-1)} + \sqrt{(n-4)})/3$$

giving 0.5773503, 1.0000000, 1.2168450, 1.3938270 at $n = 4, 5, 6, 7$ — four exact hits, and the 3 in the denominator is $p_g - p_r$, invariant across the table.

The staircase algebra of a closed index

A.staircls states that $E(\mathcal{R}) = 0$ iff the held set is an intersection of monotone staircases. **A closed index is therefore a system of inequalities and its cells are the lattice points satisfying them.**

Read off Λ_{PCA} , whose cells satisfy $L(s) \leq d \leq U(s)$ with

$$L(s) = \lfloor s/2 \rfloor \text{ and } U(s) = s + \lfloor s/3 \rfloor$$

both exact on $s \in \{0,1,2,3\}$. **And off Λ_{amp}** , whose twenty cells satisfy the single inequality $0 \leq i \leq \ell$.

This work. *That the algebra of a closed index can be read back out of it as floor functions — the staircases are not merely asserted to exist but written.*

The Slater triangle, and a parity defect resolved

Standard. The multipole expansion of $1/r_{12}$ gives, for a pair of subshells, F^k for k even up to $2\min(\ell, \ell')$ and G^k for $k \equiv \ell + \ell' \pmod{2}$ from $|\ell - \ell'|$ to $\ell + \ell'$. Slater 1929; Condon & Shortley 1935; the angular factors are Racah's (1942–1949).

The angular factor of G^k is the $3j$ symbol squared, and for an s electron against an ℓ electron the sole exchange integral has angular factor **exactly $1/(2\ell + 1)$** — verified 1, 1/3, 1/5, 1/7 to machine precision.

This work. *That indexing on the multipole rank k gives $E = 1$ with the single defect F^1 , a term parity forbids — and that reindexing on position within sequence closes it at $E = 0$. The rank's parity is fixed by the kind and the ℓ pair, so rank is not a free coordinate.*

The falsification of a selection rule

The object. Given the corridor at every element, a walk needs a rule for where in each interval to place the carried value. **Eight rules were tested; seven give 106/106**, including uniformly random interior points on **200 of 200 seeds**. Only “move to the farther endpoint” fails, at 74/106, because it can overshoot.

This work. *That the ordering data constrains only membership of the interval, not position within it — so the eighteen resets landing on exact surds is a property of one chosen rule and not of the table. A different rule reproduces the periodic table with a disjoint trajectory and a going negative at five elements.*

Recorded because the surd trajectory was reported as a result before the falsification was run, and the falsification withdrew it.

The necessity of state

The object. For each admissible subshell, set a to that subshell's own crossing value and ask whether it is then least- v . **104 of 106 steps admit two to four self-consistent subshells** — the observed one is always among them, never uniquely determined.

This work. *That the periodic table is not computable from a single atom's configuration. It requires one number carried forward: the arithmetic supplies the values, the walk supplies the selection.*

The observability boundary

The object. Across eleven closed indexes, the quantity each closure cannot fix was identified. **Nine of eleven are expectation values $\langle \Psi | \hat{O} | \Psi \rangle$** . The exception is Λ_{cross} , whose value is a node count needing no measurement — **and it is the only index with no statistics language**.

An index closes when its cells are enumerable. What it cannot supply is exactly what requires an operator.

This work. *That the compendium's own language test detects this: the statistics language is silent precisely where no observation is needed. One confirming instance; the prediction is falsifiable and has been tested once.*

Λ _spectra's closure, and what the limit leaves

The object. The index runs to the LAST AVAILABLE SPECIES and stops. Λ is complete at 976 because its coordinates are bounded by the physics; Λ _spectra is infinite unless capped, and *all cappings are closed when complete.*

Indexing every subshell any atom holds, (n, ℓ, k) over the observed ground configurations: **98 cells, E = 58**. Of those 58, **38 require more electrons than any atom has** — $6f^1$ through $6f^{14}$ and beyond, needing Z past the table. Past the limit, not defects. Applying the limit:

Corrected at register 1426: the split was first reported as 47 / 11, using the ground-state table's own edge of $Z = 108$ as the limit while the synthesised table runs to 118. The nine cells between — $6d^7$ - $6d^{10}$ and $7p^2$ - $7p^6$ — are real elements, not absences. The eleven named exceptions below are unaffected; only the count of beyond-limit cells was wrong.

E = 11, and every cell is named.

absent	Madelung predicts at	observed
$3d^4 \cdot 3d^9$	Cr 24 · Cu 29	$3d^5 \cdot 3d^{10}$
$4d^3 \cdot 4d^6 \cdot 4d^9$	Nb 41 · Ru 44 · Ag 47	$4d^4 \cdot 4d^7 \cdot 4d^{10}$
$4f^2 \cdot 4f^8$	Ce 58 · Gd 64	$4f^1 \cdot 4f^7$
$5d^8$	Pt 78	$5d^9$
$5f^1 \cdot 5f^5 \cdot 5f^8$	Ac 89 · Np 93 · Cm 96	$5f^0 \cdot 5f^4 \cdot 5f^7$

The Madelung exceptions, recovered as closure defects rather than looked up — E.nuclide's shape, a defect whose every missing cell can be named.

This work — rival = donor iff the donor is not full. Twelve walk steps have a subshell empty as another fills. A subshell at capacity has nowhere to put an electron, so it is not Pauli-admissible and cannot be a rival in anyone's bracket, including its own. An s shell holds 2 and is full when it donates:

donor	occupancy before	binding rival
4s, 5s, 6s (Cr, Cu, Nb, Ru, Pt)	2, full	4p, 5p, 6p
5s (Pd)	1, partial	5s
5d, 6d, 7p (Pr, Tb, Pa, Pu, Bk, Rf)	partial	itself

Holds 12 of 12, with no fitted term. *Pd is its own control: the only s donor that is not full, and it sits with the d and f donors. Were the rule about angular momentum it would sit with Cr.* The anomalous steps differ from ordinary ones in WHICH RIVALS

EXIST, and that is fixed by occupancy — which ν already carries in q . Registers 1395–1398.

Attributions

The $\frac{2}{3}$ scaling. *Thomas–Fermi’s.* The quadratic in nuclear charge along an isoelectronic sequence is **Krug & von Lilienfeld, arXiv:2406.18416 (2024)**, fitted on $Z = 1\text{--}86$. **Carcassés & González, Phys. Rev. A 80 (2009) 024502**, give $E_{\text{ioniz}} = Z^2 N^{(-2/3)} g(N/Z)$ and flag that it fails for neutrals — the same domain this work’s residuals bow in.

Configuration crossings along isoelectronic sequences are a computed object: **Berengut et al., arXiv:1204.0603** locate the 6p–5f crossing in the thallium sequence by Dirac–Fock.

The node theorem, Pauli’s exclusion principle and the **centrifugal term $\ell(\ell+1)$** are standard and this work introduces none of them.

Seaton’s ratio $\delta_2/\delta_0 = -\ell(\ell+1)/3$ is **Seaton’s**; the domain restriction to $p = 0$ is this work’s.

Slater’s rules and integrals are Slater’s; the observation that the count of discarded exchange integrals equals the count of bracket failures is this work’s.

What this work introduces, in one list

object

the corridor	106 consistent linear inequalities with node-count coefficients
the nineteen surds	the complete set of endpoints across the table
a_cross = $(\sqrt{(n-1)} + \sqrt{(n-4)})/3$	closed form for the ns/(n–1)d family
the staircase algebra	$\lfloor s/2 \rfloor \leq d \leq s + \lfloor s/3 \rfloor$, read out of a closed index
the sequence-index fix	rank parity is not a free coordinate
the selection-rule falsification	the corridor is forced, the path is not
the necessity of state	104 of 106 steps ambiguous without memory
the observability boundary	closure enumerates; observation values
the singleton-output rule	an index is closed when its reading is unique
the domain prohibition	no parameter of this work is universal, so no pooled fit across regions is admissible
the limit	Λ_{spectra} closes at the last available species; $E = 11$, all named
rival = donor iff not full	the twelve anomalous steps, from Pauli alone

The Löwdin indexes — Λ_{law} and Λ_{const}

Owed since register 1296; written. Nothing here answers Löwdin's challenge — the amplitude law remains three fitted numbers.

Λ_{law} — seven laws on (law, carrier). $E = 0$.

An arbitrary ordering gave $E = 7$. Across all $7! \times 4! = 120,960$ orderings the defect runs 0 to 21, and **288 reach zero**.

The closing carrier order is $u < p < l - l_{\text{core}} < Z - T$, monotone in **how LOCAL the variable is**: u is the whole atom, p counts one l 's shells, $l - l_{\text{core}}$ compares two angular momenta, $Z - T$ is one distance to one threshold. Two laws sit on u , three on p , one on each of the others.

The held cells form a **staircase**, which by A_{staircls} is exactly the condition for $E(R) = 0$.

What only this index contributes: THE CARRIER. Fitting a law in the wrong carrier is what took the l -spread from a spurious r^2 of 0.868 to a real 0.356.

Λ_{const} — fourteen constants on (role, carrier). $E = 0$ at 2 of 576.

Role order **exponent < centre < width < scale**. Standing: three fitted, six measured, five derived or attributed.

Every CENTRE is a small integer or half-integer - 2 for the free shells, 2 for the gate, -1.5 for the switch, 2.5 for the l -validity, $u_0 = 4$. **No SCALE is**. The role axis orders by how much physics a number has absorbed.

beta = 2/3 is load-bearing: the only constant of arity three, in the amplitude, the exponent and the validity, **and the one that is attributed**.

The refusal, which is the more useful half

standing cannot be a coordinate. With attributed/derived/measured/fitted on an axis the index closes at NO ordering; without it, at once.

An index whose coordinates mix the OBJECT with the OBSERVER cannot close, because the observer's axis has no order the object respects.

The same fault is origin in Λ_{var} , kind in Λ_{phys} and state in Λ_{ladder} - four occurrences of one mistake, stated here as a rule.