General Relativity

Leonard Susskind

Core track, 2008 Fall Einsteins General Theory Of Relativity
Lecture notes organized by LazyingArt LLC with Video2Book

Original lectures by Leonard Susskind. Lecture notes organized by LazyingArt LLC with Video2Book.

Contents

1 Gravity, Free Fall, and Newton's Theorem **1**

1.1 Gravity as a Special Force and Why We Start With Newton . . . 1

1.2 Galileo's Flat-Earth Gravity and the First Equivalence-Principle Lesson . . . 2

1.2.1 Question & Answer . . . 3

1.3 Newton's Inverse-Square Force: From Magnitude to Direction . . . 4

1.3.1 Question & Answer . . . 5

1.4 From Two Bodies to Many Bodies . . . 6

1.5 Tidal Forces as the Repair to Uniform Free Fall . . . 7

1.5.1 Question & Answer . . . 7

1.6 The Gravitational Field . . . 8

1.7 Divergence, Gauss's Theorem, and Newton's Theorem . . . 8

1.7.1 Question & Answer . . . 11

1.8 Summary . . . 12

2 Newtonian Gravity, Gauss's Law, and the Equivalence Principle **13**

2.1 Dark Energy as a Prelude, Not the Main Road . . . 13

2.1.1 Question & Answer . . . 14

2.2 A Little Formalism: Del, Fields, and the Gravitational Acceleration . . . 14

2.3 Gauss's Law as Physics, Gauss's Theorem as Mathematics . . . 15

2.4 Putting Them Together: Exterior Fields and the Shell Theorem . . . 17

2.4.1 Question & Answer . . . 17

2.5 While We Are At It: Inside the Earth . . . 18

2.5.1 Question & Answer . . . 19

2.6 The Equivalence Principle as Elevator Intuition and Coordinate Transformation . . . 20

2.7 Einstein's Move: Light, Locality, and the Metric . . . 21

2.7.1 Question & Answer . . . 23

2.8 Summary 24

Chapter 1

Gravity, Free Fall, and Newton's Theorem

Gravity enters this course already carrying a double burden. It is one force among others, but it is also the force that seems to know something about geometry. Susskind therefore begins with a deliberate narrowing of scope. We are not yet doing the full general theory of relativity. We are doing Newtonian gravity, but we are doing it in a way that will later make the passage to general relativity feel motivated rather than abrupt: first the Newtonian language of forces and vectors, then the cancellation of mass in gravitational motion, then the full inverse-square law, and finally the vector calculus that explains why spherical gravitating bodies behave as if all their mass were concentrated at a point.

Source note. These companion notes follow Leonard Susskind's first general relativity lecture and preserve its order of argument. This draft was prepared with support from LazyingArt LLC and Video2Book.

1.1 Gravity as a Special Force and Why We Start With Newton

Susskind opens by saying, in effect, that gravity is unusual. Electric and magnetic forces do not behave like it, and somehow gravity is tied to the geometry of space and time. But precisely because of that, he does not begin with curved spacetime. He begins with Newton.

Definition 1.1. An inertial frame is a coordinate system, equipped with clocks, in which a force-free test object moves with uniform velocity and therefore has zero acceleration.

This is the lecture's first piece of discipline. Before we speak about gravity, we must know what we mean by force, acceleration, coordinates, and vectors. The preserved frame below catches the board at the moment when Susskind has written only the bare scalar-looking equation $F = ma$. The transcript makes clear, however, that he is already speaking of a three-dimensional vector equation.

Thus we write

$$\vec{F} = m\vec{a}, \tag{1.1}$$

$$F_i = ma_i = m\ddot{x}_i, \qquad i = 1, 2, 3. \tag{1.2}$$



Figure 1.1: The lecture begins from the stripped-down Newtonian law. The board still shows the scalar handwriting $F = ma$, but the spoken lecture already treats it as the vector equation $\vec{F} = m\vec{a}$.

The mass m is a scalar. In Newtonian physics it is conserved: it may be redistributed from one place to another, but it is not itself a directional object.

A particle has coordinates x_1, x_2, x_3, which together form the position vector

$$\vec{r} = (x_1, x_2, x_3). \tag{1.3}$$

Its first time derivative is velocity and its second time derivative is acceleration:

$$v_i = \dot{x}_i, \qquad a_i = \ddot{x}_i. \tag{1.4}$$

This is the quiet mathematical framework within which the special feature of gravity will soon appear. If the gravitational force carries the same mass that already sits in Newton's law, then that mass may cancel, and when it cancels the motion stops depending on the nature of the falling object.

1.2 Galileo's Flat-Earth Gravity and the First Equivalence-Principle Lesson

Now the lecture deliberately steps backward from Newton to Galileo. This is not antiquarianism. Galileo isolates the simplest approximation in which gravity reveals what is peculiar about it.

In the flat-Earth approximation two simplifications hold at once. First, the direction of gravity is everywhere the same: downward. Second, its magnitude does not depend on height. If we choose the x_2-axis to be vertical and take upward to be positive, then the gravitational force has only one nonzero component:

$$F_2 = -mg. \tag{1.5}$$

The minus sign merely says that the force points downward. Here g is the near-surface gravitational acceleration. On Earth it is about $10\,\mathrm{m/s^2}$; on the Moon it is smaller, on Jupiter larger. What

matters in the present argument is that for a given planet it does not depend on the mass of the falling body.

Combining Galileo's force law with Newton's equation gives the first decisive cancellation:

$$F_2 = m\ddot{x}_2, \tag{1.6}$$

$$m\ddot{x}_2 = -mg, \tag{1.7}$$

$$\ddot{x}_2 = -g. \tag{1.8}$$

This is the first serious mathematical signal of the equivalence principle in the lecture. The acceleration of free fall is independent of the mass of the object. Heavy bodies and light bodies obey the same equation of motion. However Galileo performed the experiment, whether from a tower or on an inclined plane, this is the content of the result: absent air resistance, different masses fall alike.

The crucial point is that this is not true for generic forces. Electric forces couple to charge, not mass. Gravity is special because its force is proportional to the same mass that Newton's law then multiplies by the acceleration. The mass cancels, and with it goes any dependence of the motion on the identity of the falling body.

1.2.1 Question & Answer

Why can a freely falling cloud fail to detect gravity even though gravity is present?

Take a diffuse cloud of particles and let them all go in the same uniform gravitational field. Some particles may be heavy, some light, some large, some small. None of that matters. Every particle obeys $\ddot{x}_2 = -g$. Therefore every particle falls in exactly the same way.

The consequence is stronger than the slogan "all bodies fall alike." The cloud does not deform. Neighboring particles keep the same relative separation. If the cloud had been held together by struts, those struts would feel no stress from the gravitational field itself, because all parts of the structure share the same motion. In the flat-Earth approximation, gravity produces no internal stretching, no compression, and no shear.

That is why Susskind says that free fall is locally indistinguishable from being far away from all gravitating bodies. The point is not that one cannot tell the difference between standing on the Earth and falling. Of course one can. The point is subtler: within a self-contained falling system, and without using an external reference such as the floor, one cannot detect the gravitational field by internal mechanical measurements.

The student questions in the lecture push this further. Is the effect optically detectable? Suppose the particles in the cloud carry lasers. The answer is still no, because the relative distances do not change. Is the peculiar feeling of stepping off a building evidence of gravitational acceleration? Again no. That feeling comes from the disappearance of the support forces that normally act on the bottoms of our feet. In Susskind's phrase, what we ordinarily feel is not falling but *not* falling.

This is the lecture's first version of the equivalence principle: in a perfectly uniform gravitational field, free fall erases gravity locally.



Figure 1.2: The lecture first writes Newton's law as a statement about force magnitude. The point-source sketch labeled M prepares the immediate transition to vector form.

1.3 Newton's Inverse-Square Force: From Magnitude to Direction

The flat-Earth approximation has now done its work. The lecture turns to Newton's full force law. Every object in the universe exerts a gravitational force on every other object. Start with two masses, a lighter mass m and a heavier mass M. The forces are attractive, equal, and opposite.

The board first presents the law as a statement about magnitude, not yet about direction. In the preserved frame the denominator is written as R^2; we keep that equation visible because it is legible on the board, but in the polished formulas below we normalize the notation to $r = |\vec{r}|$.

On the board one sees

$$F = \frac{mMG}{R^2}. \tag{1.9}$$

In cleaned notation we write

$$F = \frac{GmM}{r^2}, \tag{1.10}$$

where G is Newton's constant,

$$G \approx 6.7 \times 10^{-11}\,\mathrm{N\,m^2/kg^2}. \tag{1.11}$$

The lecture pauses over the smallness of this number. Gravity is weak. That is not merely rhetoric. For two 1 kg masses separated by 1 m,

$$F = G \approx 6.7 \times 10^{-11}\,\mathrm{N}. \tag{1.12}$$

A bit of electrostatic charge or a modest magnet easily beats that. Gravity dominates ordinary life only because the Earth is so heavy.

There is another special feature here as well. Electric forces may attract or repel. Gravitational forces, in Newton's theory, are always attractive.



Figure 1.3: Clean reconstruction of the geometry behind the vector law. The radial vector points from the source mass M to the test mass m, while the force points inward, opposite to $\vec{r}$.

Now Susskind sharpens the scalar law into a vector law. Put the heavy mass M at the origin and let $\vec{r}$ point from M to the lighter particle m. Then the force on m must point along the radial direction, but opposite to $\vec{r}$. Introducing the unit vector

$$\hat{r} = \frac{\vec{r}}{r}, \tag{1.13}$$

we obtain

$$\vec{F} = -\frac{GmM}{r^2}\hat{r} = -\frac{GmM}{r^3}\vec{r}. \tag{1.14}$$

The extra factor of r downstairs is exactly what keeps the magnitude at $1/r^2$ while inserting the direction.

1.3.1 Question & Answer

Equal forces, unequal accelerations. If the force on the Earth due to a falling body is equal and opposite to the force on the body due to the Earth, why does the Earth hardly accelerate at all?

Because equality of forces is not equality of accelerations. Newton's equation says

$$\vec{a} = \frac{\vec{F}}{m}.$$

For the same force, the larger mass has the smaller acceleration. The Earth and the falling body exert equal forces on one another, but the Earth's enormous mass suppresses its acceleration.

Why inverse square? The lecture's historical motivation runs through Kepler. For circular motion of radius r and angular frequency ω, the inward acceleration is

$$a_{\text{circ}} = \omega^2 r. \tag{1.15}$$

If the force per unit mass behaves as $f(r) \propto 1/r^2$, then

$$\omega^2 r \propto \frac{1}{r^2}, \qquad \text{so} \qquad \omega^2 \propto \frac{1}{r^3}. \tag{1.16}$$

Since $T = 2\pi/\omega$, we obtain

$$T^2 \propto r^3. \tag{1.17}$$

This is Kepler's law relating period and orbital radius. In the lecture this is not meant as a full historical reconstruction, but as the right Newtonian clue: Kepler's law points strongly toward the inverse-square force, and the inverse-square force is then the law that extends correctly beyond perfect circles to the ellipse.



Figure 1.4: Documentary witness for the many-body discussion. The board is already being erased and the sign convention is being corrected live, so the final notes immediately replace the unstable blackboard notation by a clean standard form.

What if the bodies get too close? The lecture also inserts a caution. When bodies touch, or even come sufficiently close, gravity is no longer the whole story. Electrostatic, atomic, and nuclear forces enter. Newton's inverse-square law is the right long-range gravitational law, not a complete short-distance theory of matter.

1.4 From Two Bodies to Many Bodies

Now we pass from one pair of gravitating bodies to a whole assembly of particles. Newtonian gravity works pairwise, but the total force on a given particle is the vector sum of all the pairwise forces exerted by the others. Susskind pauses to remark that this superposition rule is not logically forced by the idea of force itself; it is a fact about Newtonian gravity.

The available board witness for this part of the lecture is weak but useful. It catches the many-body formula in the middle of a live correction, just as the sign convention is being discussed and the board is already being erased.

Let $\vec{r}_i$ be the position of the i-th particle. If we define the vector from particle i to particle j by $\vec{r}_j - \vec{r}_i$, then the clean Newtonian force law is

$$\vec{F}_i = \sum_{j \neq i} G\, m_i m_j \, \frac{\vec{r}_j - \vec{r}_i}{|\vec{r}_j - \vec{r}_i|^3}. \tag{1.18}$$

This is the simplest way to encode attraction without importing the lecture's temporary sign ambiguity. The student objection in the lecture is exactly that one may put the minus sign in

front or absorb it into the choice of whether the vector points from i to j or from j to i. Once the convention is fixed, the formula is fixed.

Now combine this with Newton's equation for the same particle:

$$\vec{F}_i = m_i \vec{a}_i, \tag{1.19}$$

$$m_i \vec{a}_i = \sum_{j \neq i} G\, m_i m_j \, \frac{\vec{r}_j - \vec{r}_i}{|\vec{r}_j - \vec{r}_i|^3}. \tag{1.20}$$

The mass m_i cancels:

$$\vec{a}_i = \sum_{j \neq i} G\, m_j \, \frac{\vec{r}_j - \vec{r}_i}{|\vec{r}_j - \vec{r}_i|^3}. \tag{1.21}$$

Once again the motion of the particle we are tracking does not depend on its own mass. It depends on the masses of all the other particles, but not on its own. This is the same structural fact we already saw in the flat-Earth problem, now stated for the general Newtonian field.

The lecture also inserts an important warning here. This formula tells us the instantaneous force if we know the positions of all the particles. Once the entire system is allowed to evolve, every particle changes the motion of every other particle, and the full dynamics quickly becomes a complicated many-body problem. That complication is real, but it is not the point of the present lecture. The present point is the persistence of mass-independence in the motion of a test object.

1.5 Tidal Forces as the Repair to Uniform Free Fall

At this point the lecture returns to the earlier statement about free fall and tightens it. In the flat-Earth approximation we said that a freely falling object does not detect gravity internally. That statement was deliberately a little too strong. It is true only for a perfectly uniform field.

Once the force depends on distance from the source, not all parts of an extended body accelerate in exactly the same way. If a very tall body falls feet-first toward the Earth, the feet are closer to the Earth than the head, so the lower part experiences a stronger acceleration. The body is stretched vertically. In differential language, nearby parts separated by $\delta\vec{x}$ experience different accelerations because

$$\delta\vec{a} \approx \vec{g}(\vec{x} + \delta\vec{x}) - \vec{g}(\vec{x}) \tag{1.22}$$

is no longer negligible.

There is a second effect. If the same extended body lies horizontally, then its left and right ends are pulled toward the Earth's center along slightly different directions. The field is radial, so those forces are not parallel. The body is therefore squeezed horizontally even if the magnitudes of the forces are nearly the same. In the lecture this becomes the vivid picture of the absurdly tall man who is stretched in one orientation and compressed in another.

These are tidal forces. They are real. They are not an artifact of coordinates. They arise because the Newtonian gravitational field varies in both magnitude and direction from point to point.

1.5.1 Question & Answer

If free fall hides gravity, why do sufficiently large falling bodies get stretched and squeezed?

Because free fall hides only the common part of the acceleration. If the field were perfectly uniform, every part of the body would share the same motion and nothing internal would reveal gravity. But a real gravitational field is not uniform. Different parts of the body sample different values and different directions of the field, and those differences are exactly what produce stress and strain.

So the lecture keeps two statements at once. A sufficiently small freely falling object does not notice gravity, because the gradient of the field across its size is negligible. A sufficiently large freely falling object does notice gravity, because that gradient is not negligible. Then one part is pulled a little harder than another, or one part is pulled in a slightly different direction than another, and the body is stretched or squeezed.

This is also why the forces are called *tidal*. The Moon exerts such forces on the Earth. The solid Earth is comparatively rigid, but the oceans are not. The water layer is therefore slightly stretched along the Earth–Moon line and squeezed in the transverse directions, producing the familiar tidal bulges.

1.6 The Gravitational Field

Only after this correction does Susskind define the gravitational field. The definition is operational. We imagine adding one more particle, a very light particle, light enough that it does not appreciably influence the motion of the others. This extra particle is the test particle.

Definition 1.2. The gravitational field at a point $\vec{x}$ is the acceleration of a sufficiently light test particle placed at $\vec{x}$.

This definition is tailored to the special property of gravity. Since the force is proportional to the mass of the test particle, the ratio of force to mass is independent of which test particle we use. We may therefore think directly in terms of acceleration rather than force. The field is the vector acceleration assigned to each point in space.

If the source masses are at positions $\vec{r}_j$, a clean Newtonian expression is

$$\vec{g}(\vec{x}) = \sum_j G\, m_j \, \frac{\vec{r}_j - \vec{x}}{|\vec{r}_j - \vec{x}|^3}. \tag{1.23}$$

The lecture temporarily denotes this acceleration by $\vec{A}$. We keep $\vec{g}$ for the gravitational field itself and reserve $\vec{A}$ for the generic vector field used in the vector-calculus discussion after the break.

The field is a vector field. It has magnitude and direction, it depends on position, and if the source masses move it depends on time as well. That is the right place for the lecture to pause. Once the field has been laid out over space, we need the mathematics that describes how such a field varies from point to point.

1.7 Divergence, Gauss's Theorem, and Newton's Theorem

After the break Susskind says openly that we now need a little fancy mathematics. The first object is a generic vector field $\vec{A}(\vec{x})$, not yet necessarily the gravitational field. We want a scalar quantity that measures whether the field is locally spreading out from a point or converging into it.



Figure 1.5: The board formula for divergence. The key distinction is between the vector field $\vec{A}$ and the scalar quantity $\nabla \cdot \vec{A}$.



Figure 1.6: Schematic field used in the lecture to explain local divergence. The point is to compare the amount of flow entering and leaving a small control region.



Figure 1.7: Reconstruction of the control-region picture described in the lecture. More field leaves the small circle on the right than enters from the left, so the region has positive divergence.

The board below preserves the lecture's definition.

In cleaned notation we write

$$\nabla \cdot \vec{A} = \partial_x A_x + \partial_y A_y + \partial_z A_z = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}. \tag{1.24}$$

The lecture is careful about the point that this is a scalar, not a vector. The arrows drawn on the other board are the field $\vec{A}$ itself. The divergence is the scalar quantity extracted from the spatial derivatives of its components.

To make this intuitive, the lecture turns to water flow. We imagine a shallow lake of fixed depth, or more generally an incompressible fluid whose density is fixed. Water pumped in from below creates a local source, so the velocity field spreads outward and has positive divergence. Water removed from below creates a sink, so the field converges inward and has negative divergence. The lecture later notes that one could complicate the analogy by allowing compression or thinning, but for pedagogy it keeps the picture simple: fixed depth, fixed density, visible source and sink language.

The accompanying arrow-field sketch is preserved below.

The second tool is Gauss's theorem, which ties local source strength to total flux through a boundary.

Theorem 1.3 (Gauss's theorem)**.** *For a vector field $\vec{A}$,*

$$\int_V (\nabla \cdot \vec{A})\, dV = \oint_{\partial V} A_\perp \, d\sigma = \oint_{\partial V} \vec{A} \cdot \hat{n} \, d\sigma. \tag{1.25}$$

The meaning is direct. The integral on the left adds up all sources and sinks inside the region V. The integral on the right adds up the net outward flow through the boundary ∂V. Only the normal component matters; motion tangent to the boundary slides along it and contributes no outward flux.

This is the theorem the lecture has been building toward. We now apply it to a spherically symmetric source of divergence concentrated inside some finite sphere. Let the total enclosed divergence be

$$Q = \int_V (\nabla \cdot \vec{A})\, dV. \tag{1.26}$$

If we surround that source by a larger sphere of radius r, then Q is independent of the radius of the larger sphere, provided only that the larger sphere encloses the whole source region.

By spherical symmetry, the field outside must be radial and its magnitude must be constant over the Gaussian sphere. Therefore the flux integral collapses to

$$Q = \oint_{\partial V} A_\perp \, d\sigma = A(r) \oint_{\partial V} d\sigma \tag{1.27}$$

$$= 4\pi r^2 \, A(r). \tag{1.28}$$

Hence

$$A(r) = \frac{Q}{4\pi r^2}, \qquad \vec{A}(r) = \frac{Q}{4\pi r^2}\hat{r}. \tag{1.29}$$

This is the lecture's clean derivation of a $1/r^2$ field from spherical symmetry and Gauss's theorem. It is here that Susskind briefly says "four-thirds πr^2" and immediately corrects it. Only the corrected surface area $4\pi r^2$ belongs in the notes.

Now gravity reenters. A positive source in the water analogy gives an outward field. Real Newtonian gravity points inward. So the gravitational analog is a negative source, or a convergence. For a point mass M,

$$\vec{g}(r) = -\frac{GM}{r^2}\hat{r}. \tag{1.30}$$

The lecture emphasizes that this is only a mathematical analogy. Space is not literally a fluid. But the analogy is useful because it explains why a point source naturally produces a $1/r^2$ field.

This brings us to Newton's theorem.

Proposition 1.4 (Newton's theorem)**.** *Outside a spherically symmetric mass distribution, the gravitational field is exactly the same as that of a point mass with the same total mass concentrated at the center.*

This is not a small convenience. It is what makes planetary gravity tractable. The exterior field of a spherical planet does not care whether the mass is concentrated at the center, spread out smoothly, denser in the middle, or denser near the surface. Outside the body, only the total mass matters.

1.7.1 Question & Answer

Why is the field outside a spherical body point-mass-like, while the field inside a spherical shell vanishes?

Outside the body, the answer is Gauss plus symmetry. A Gaussian sphere enclosing the entire spherically symmetric source sees only the total enclosed mass. The flux is therefore fixed. Since the field has the same magnitude everywhere on that sphere and points radially, the only possible exterior field is the $1/r^2$ field of a point mass with that same total mass.

Inside a hollow spherical shell, the answer is sharper still. A Gaussian surface entirely inside the shell encloses no source at all. Therefore the enclosed divergence is zero, and the net flux through



Figure 1.8: A spherical shell. Outside the shell, the field is the same as that of a point mass at the center. Inside the shell, the field vanishes.

the Gaussian surface is zero. But by symmetry there is no distinguished interior direction. The only field compatible with zero enclosed source and full spherical symmetry is

$$\vec{g}_{\text{inside shell}} = 0.$$

The lecture also insists on the corresponding limitation. This is an exterior theorem. Once we go inside a gravitating body, we are asking a different question. In the mine shaft thought experiment the field near the center of the Earth is certainly not what it would be if the entire Earth's mass were concentrated at the center. The exterior theorem does not survive unchanged at interior points.

Susskind is right to linger here. This theorem is not just elegant; it is the remarkable piece of luck that made Newtonian celestial mechanics workable.

1.8 Summary

The lecture begins with the most elementary Newtonian equation and ends with the shell theorem, but the movement is carefully staged. First we learn why gravity is special: its force is proportional to the same mass that appears in Newton's law, so in a uniform field the mass cancels and free fall becomes universal. Then we leave the flat-Earth approximation, write Newton's inverse-square law, sharpen it into vector form, and generalize it to many bodies. The motion of a test particle still ignores its own mass, which leads naturally to the gravitational field.

Only then do we stop and repair the earlier slogan. A real gravitational field varies from place to place, so large freely falling bodies are stretched and squeezed by tidal forces. After the break the lecture introduces divergence and Gauss's theorem, and these become the machinery that explains why spherical symmetry produces a $1/r^2$ law and why spherical shells are so remarkable.

That is the real payoff of Lecture 1. Already within Newtonian gravity, the mathematics begins to organize itself in a way that points beyond force law and toward geometry.

Chapter 2

Newtonian Gravity, Gauss's Law, and the Equivalence Principle

These notes follow Leonard Susskind's second lecture on general relativity in the order in which the argument is actually built. The lecture opens with a short detour about dark energy, then deliberately returns to Newtonian gravity, sharpens that old physics into field language, and only then turns toward Einstein. Where the blackboard itself carries part of the mathematical story, we keep the original lecture frames visible as documentary evidence and place the cleaned equations or diagrams nearby. Credit for the lecture belongs to Leonard Susskind; the present curation also owes explicit credit to LazyingArt LLC and Video2Book.

2.1 Dark Energy as a Prelude, Not the Main Road

The lecture begins by answering a worry that had been left hanging over from earlier exchanges: if dark energy acts like a repulsive component of gravity, why does it not tear ordinary bound systems apart? Susskind does not begin with cosmological field equations. He begins with a spring, because for the limited point he wants to make, that is the cleanest picture.

For an ordinary restoring force we may write, schematically,

$$F(r) \sim -kr. \tag{2.1}$$

A positive cosmological constant is then modeled, at this opening level, as a tiny repulsive correction of the same distance-dependent type,

$$F(r) \sim -kr + \lambda r. \tag{2.2}$$

The point is not that an atom literally is a spring. The point is that if a bound system already has a strong restoring force, then a very small repulsive term proportional to distance merely shifts the equilibrium position slightly. It weakens the binding a little; it does not destroy it.

That is the practical content of the opening discussion. The coefficient λ is so small that its effect is negligible on laboratory scales, negligible on the scale of the solar system, and still tiny compared with ordinary binding effects inside galaxies. Only on cosmological scales does such a term compete with anything else.

Susskind then sharpens the contrast by mentioning the "Big Rip" scenario. The ordinary cosmological constant does not do this. The rip would require something more violent: effectively, that the repulsive coefficient grow with time. He treats that not as sober mainstream physics, but as a bad speculative idea.

2.1.1 Question & Answer

Question. Why does dark energy not tear atoms, solar systems, or galaxies apart?

Answer. Because in the lecture's opening model the repulsive contribution is fantastically small. In a bound system it produces, to leading order, only a slight shift of the equilibrium configuration. The atom becomes a tiny bit larger; the orbit becomes a tiny bit more spread out. The "Big Rip" picture would require the repulsive effect to grow with time and eventually overwhelm binding forces. Susskind explicitly treats that as a different and much less credible story.

With the conceptual debris cleared away, the lecture returns to where it meant to go all along: we still need a little more Newton before we can earn Einstein.

2.2 A Little Formalism: Del, Fields, and the Gravitational Acceleration

Susskind introduces this next step with exactly the right tone: first a little mathematics, though really it is only formalism, a notation for organizing equations. The operator

$$\nabla = (\partial_x, \partial_y, \partial_z) \tag{2.3}$$

behaves symbolically like a vector, even though it is really a differential operator.

If $\phi(x, y, z)$ is a scalar field, then applying ∇ produces a vector field,

$$\nabla\phi = (\partial_x\phi,\, \partial_y\phi,\, \partial_z\phi)\,. \tag{2.4}$$

If instead $\vec{v}(x, y, z)$ is a vector field, then the divergence is the scalar

$$\nabla \cdot \vec{v} = \partial_x v_x + \partial_y v_y + \partial_z v_z. \tag{2.5}$$

We should keep the bookkeeping completely straight here. The lecture contains a fleeting verbal slip, but the mathematics is unambiguous: the gradient of a scalar is a vector, while the divergence of a vector is a scalar.

A scalar field is just a scalar quantity depending on position. A vector field is a vector quantity depending on position. The Newtonian gravitational field is such a vector field.

Now Susskind defines it operationally. Take a distribution of source masses m_i at positions $\vec{x}_i$. Introduce a tiny test mass, release it at the field point $\vec{x}$, and observe its acceleration. That acceleration is the gravitational field:

$$\vec{A}(\vec{x}) = \text{acceleration of a test mass released at } \vec{x}. \tag{2.6}$$

There is a brief and telling hesitation over the sign convention. That hesitation should not be erased, because it shows exactly what must be fixed. We want the field to point *toward* the source mass. Therefore we choose

$$\vec{R}_i := \vec{x}_i - \vec{x}, \tag{2.7}$$

so that $\vec{R}_i$ points from the test mass toward the ith source. With that choice the Newtonian field is

$$\vec{A}(\vec{x}) = G\sum_i m_i \frac{\vec{R}_i}{R_i^3} = G\sum_i m_i \frac{\hat{R}_i}{R_i^2}, \qquad \hat{R}_i = \frac{\vec{R}_i}{R_i}. \tag{2.8}$$

That is simply the inverse-square law rewritten as a field at a point. The magnitude scales like $1/R_i^2$; the direction is carried by $\vec{R}_i$.



Figure 2.1: Field vector and source-point displacement. The board frame is kept because it is the clearest documentary evidence for the source-point/field-point geometry, while the redraw fixes the final sign convention chosen in the lecture.

Up to this point the field has been assembled one source at a time. The next move is the one Susskind explicitly advertises: there is a neat way to summarize all this.

2.3 Gauss's Law as Physics, Gauss's Theorem as Mathematics

To pass from discrete particles to a field equation, we introduce the mass density. If a small volume ΔV contains a small mass Δm, then

$$\rho = \frac{\Delta m}{\Delta V}. \tag{2.9}$$

In the continuum picture $\rho(\vec{x})$ is itself a scalar field.

Now comes the summary of Newtonian gravity:

$$\nabla \cdot \vec{A} = -4\pi G\rho. \tag{2.10}$$

This is Gauss's law for gravity. The minus sign records the fact that the field points inward toward positive mass. The field converges. A convergence is a negative divergence.

Susskind is especially careful here to separate two statements that are often blurred together. Gauss's law is physics. It is a law of nature, equivalent to Newtonian gravity. Gauss's theorem is mathematics. It says that for any sufficiently nice region V with boundary ∂V,

$$\int_V (\nabla \cdot \vec{A})\, dV = \oint_{\partial V} A_\perp \, d\sigma. \tag{2.11}$$

On the left we integrate the divergence throughout the volume. On the right we integrate the component of the field normal to the boundary. The theorem translates volume divergence into boundary flux.

The blackboard frame for this moment is only partially legible, but the intent is clear enough to justify writing the standard form.





Figure 2.2: Surface-flux term in Gauss's theorem. The lecture frame shows the half-written theorem and the boundary patch labeled $A_\perp$; the redraw makes explicit that the theorem uses the normal component through the surface element $d\sigma$.

This is still abstract. The lecture does not linger there. Susskind immediately says, in effect, let us put them together and see what they do.

2.4 Putting Them Together: Exterior Fields and the Shell Theorem

Suppose the mass distribution is spherically symmetric, or isotropic, about a center. Then we surround it by a concentric Gaussian sphere of radius r. Now the theorem and the law meet each other.

From Gauss's law,

$$\nabla \cdot \vec{A} = -4\pi G\rho, \tag{2.12}$$

so inside the volume integral we have

$$\int_V (\nabla \cdot \vec{A})\, dV = -4\pi G \int_V \rho\, dV. \tag{2.13}$$

But the integral of ρ over the volume is simply the enclosed mass,

$$\int_V \rho\, dV = M_{\text{enc}}. \tag{2.14}$$

Meanwhile, on the surface side, symmetry tells us something equally important. The field must be radial, and its radial component has the same value at every point of the Gaussian sphere. Therefore

$$\oint_{\partial V} A_\perp\, d\sigma = 4\pi r^2 A_r. \tag{2.15}$$

Putting these together gives

$$4\pi r^2 A_r = -4\pi G M_{\text{enc}}, \tag{2.16}$$

and therefore

$$A_r = -\frac{GM_{\text{enc}}}{r^2}. \tag{2.17}$$

If the sphere encloses the whole body, then $M_{\text{enc}} = M$, and we recover the familiar inverse-square field

$$A_r = -\frac{GM}{r^2}. \tag{2.18}$$

This is also the point in the lecture where Susskind remarks on the appearance of the factor 4π. In this normalization it is chosen so that the surface area of the sphere cancels cleanly. One could normalize the equations differently, but this choice makes spherical problems fall open at once.

What has happened is more than a recovery of Newton's formula. We have also learned something new in this language: the exterior field of any spherically symmetric mass distribution is the same as the field of a point mass at the center.

2.4.1 Question & Answer

Question. Why does only the mass inside the Gaussian sphere matter?



Gaussian sphere



Figure 2.3: Gauss law with enclosed mass. The lecture frame shows the shift from the abstract flux theorem to the gravitational mass integral; the redraw isolates the concentric spherical geometry used in the derivation.

Answer. Because the flux through the Gaussian sphere is fixed by the integral of the density *inside* that sphere. Once spherical symmetry is assumed, the flux determines the radial field on the boundary. Matter outside contributes zero net field there. This is the shell theorem in flux form.

That answer has a remarkable corollary. If the entire mass were placed on a thin spherical shell, then the field inside the shell would vanish identically. Outside, the field would be exactly the same as that of a point mass at the center. Susskind emphasizes that Newton knew this and proved it geometrically, without calculus. In these notes we see it fall directly out of Gauss's law.

2.5 While We Are At It: Inside the Earth

At this point Susskind turns, in the same spirit, to a question from the previous meeting: what does the gravitational field look like *inside* the Earth? The same formalism answers it almost for free.

We again choose a Gaussian sphere, but now the sphere lies inside the Earth. The flux side is

unchanged:

$$\oint_{\partial V} A_\perp \, d\sigma = 4\pi r^2 A_r, \tag{2.19}$$

where r is now the radius of the *inner* Gaussian sphere. What changes is the mass enclosed.

To make the calculation explicit, Susskind adopts a simplified Earth: uniform density ρ throughout the interior, abruptly dropping to zero at the surface. Then

$$M_{\text{enc}}(r) = \frac{4\pi}{3} r^3 \rho. \tag{2.20}$$

Substituting this into Gauss's law gives

$$4\pi r^2 A_r = -4\pi G M_{\text{enc}}(r), \tag{2.21}$$

$$4\pi r^2 A_r = -4\pi G \left(\frac{4\pi}{3} r^3 \rho \right). \tag{2.22}$$

The board algebra in the lecture becomes momentarily unstable here, and the audience catches a missing factor of 4π. The corrected result, which is the one we should preserve, is

$$A_r = -\frac{4\pi G \rho}{3} \, r. \tag{2.23}$$

This also matches the exterior formula at the surface $r = R$ once we use

$$M = \frac{4\pi}{3} R^3 \rho. \tag{2.24}$$

Now the lecture cashes out the result physically. Since $\vec{A}$ is an acceleration field, multiplying by the mass m of the test particle gives

$$m\ddot{r} = -\frac{4\pi G \rho}{3} \, m r. \tag{2.25}$$

This is the equation of a harmonic oscillator. Inside the idealized uniform Earth, gravity becomes a linear restoring force.

That is a lovely Newtonian payoff. A particle displaced from the center is pulled back exactly as though attached to a spring. In the tunnel-through-the-Earth thought experiment, the particle would oscillate through the planet.

2.5.1 Question & Answer

Question. How can the field be linear in r if the enclosed mass grows like r^3?

Answer. Because Gauss's law fixes the *flux*, not the field directly. The flux contains an area factor $4\pi r^2$. Thus the scaling is

$$A_r \, r^2 \sim r^3, \tag{2.26}$$

so only one power of r survives:

$$A_r \sim r. \tag{2.27}$$

This is exactly the local conceptual snag that surfaces in the classroom, and it is useful because it makes visible the role of the Gaussian sphere itself.

The lecture briefly announces gravitational potential here, but then defers it. That deferral is important. The Newtonian recap has reached the point it needed to reach, and Susskind is ready to say so: this has been old physics. Now we move to the new idea.





Figure 2.4: Concentric spheres and enclosed-mass factor. The preserved lecture frame supplies the documentary evidence for the inner Gaussian sphere and the visible $\frac{4\pi}{3}r^3\rho$ factor, while the redraw gives the cleaned geometry for the corrected derivation.

2.6 The Equivalence Principle as Elevator Intuition and Coordinate Transformation

The equivalence principle first appears in the lecture in the form everybody remembers: the elevator. Suppose we have a cabin accelerating horizontally with acceleration g. A passenger releases an object inside. In the inertial frame outside the elevator, the object simply continues its motion. But to the observer riding inside, the object accelerates backward toward the floor of the elevator with magnitude g. It looks like gravity, sounds like gravity, smells like gravity.

Susskind insists that this alone would not make Einstein Einstein. The real step is not merely to say the words. The real step is to use them. But before he uses them, he rebuilds the elevator mathematically.

He chooses the horizontal coordinate x precisely because he wants to draw an x–t diagram. First consider uniform velocity. If the floor of the elevator moves with constant speed v, then the passenger's coordinate relative to the floor is

$$x' = x - vt, \qquad t' = t. \tag{2.28}$$



Figure 2.5: Horizontal elevator version of the equivalence principle. In the passenger's frame, the released object accelerates backward toward the floor exactly as if a uniform gravitational field were present.

This is still Newtonian physics: all observers use the same time. If an object is at rest in the inertial frame, then $x =$ constant and hence

$$\dot{x}' = -v, \qquad \ddot{x}' = 0. \tag{2.29}$$

So a uniformly moving frame creates no fictitious gravitational field.

Now let the elevator accelerate uniformly. Its floor then follows

$$x_{\text{floor}}(t) = \frac{1}{2}gt^2, \tag{2.30}$$

and the corresponding transformed coordinate is

$$x' = x - \frac{1}{2}gt^2. \tag{2.31}$$

For an object at rest in the inertial frame, we differentiate once and then again:

$$\dot{x}' = -gt, \qquad \ddot{x}' = -g. \tag{2.32}$$

This is the mathematical statement of the elevator argument. In the accelerated frame, freely moving bodies behave as though there were a uniform gravitational field.

This is not yet full relativity. Susskind says so directly. But it already gives the conceptual bridge he wants: acceleration corresponds to a change to curvilinear coordinates in space-time. That is why the discussion is heading toward geometry.

2.7 Einstein's Move: Light, Locality, and the Metric

Now comes the genuinely Einsteinian use of the principle. If gravity and acceleration are equivalent only for falling stones, then the principle is not deep. Einstein's claim is stronger: the laws of physics in a gravitational field are the same as the laws of physics in an accelerated frame. Therefore the principle must apply to light as well.

Imagine an elevator standing still. A light beam sent horizontally across it travels in a straight line and hits the opposite wall directly across. Now imagine the elevator accelerating upward. The light beam still begins horizontally, but by the time it reaches the far wall the elevator has moved upward. So in the elevator frame the light lands lower than before. Its path is curved. In fact, in this approximation, it is parabolic.



Figure 2.6: x–t sketches for the moving and accelerating elevator. The accelerating floor follows a parabola, and the transformed coordinate produces the effective field $\ddot{x}' = -g$.

That simple argument leads to Einstein's early estimate for light bending by the Sun. The lecture keeps the estimate rough and historical, and the notes should do the same. Consider a light ray that just skims the solar surface. Near closest approach, take the downward acceleration to be roughly the surface gravity,

$$g_\odot \approx \frac{GM}{R^2}. \tag{2.33}$$

The time spent in the region where this matters is estimated by the light-crossing time of about two solar radii:

$$\Delta t \approx \frac{2R}{c}. \tag{2.34}$$

Hence the downward component of the velocity picked up by the ray is

$$\Delta v_\perp \approx g_\odot \Delta t \approx \frac{2GM}{Rc}. \tag{2.35}$$

For a small deflection angle,

$$\theta \approx \frac{\Delta v_\perp}{c} \approx \frac{2GM}{Rc^2}. \tag{2.36}$$

This is Einstein before the full apparatus of general relativity. Susskind keeps the historical caution in view: the final relativistic answer differs by a factor of two. The point of the lecture here is not precision but the conceptual use of the equivalence principle in a place where nobody had used it before.



Figure 2.7: Light in an upward-accelerating elevator. This is the elevator version of the statement that light falls in a gravitational field.



Figure 2.8: Crude sun-skimming-ray estimate. The lecture treats the deflection qualitatively and then estimates it by combining surface gravity with a light-crossing time.

2.7.1 Question & Answer

Question. Can free fall completely eliminate gravity, or only locally?

Answer. Only locally. A sufficiently small elevator, observed for a sufficiently short time, can replace the Earth's field by a single accelerating frame. But the real Earth's field is not uniform. Its direction varies from place to place, and its magnitude varies with distance. A large enough freely falling body notices this immediately: its feet are pulled more strongly than its head, and neighboring free-fall lines converge sideways. Those are tidal forces. They are the obstruction to eliminating gravity globally.

That is the limiting correction Susskind insists on before the lecture ends. Gravity can be transformed away only in a local neighborhood. The real, invariant content of a gravitational field lies in what survives that attempt, namely tidal effects.

He then steps back from mechanics and turns to the blackboard itself. How does one describe the geometry of a flat two-dimensional surface? In rectangular Cartesian coordinates, for neighboring points,

$$ds^2 = dx^2 + dy^2. \tag{2.37}$$

But in curvilinear coordinates on the same flat board, the distance between neighboring points is more generally

$$ds^2 = g_{xx}\,dx^2 + 2g_{xy}\,dx\,dy + g_{yy}\,dy^2. \tag{2.38}$$

The cross term tells us that the coordinate lines need not be perpendicular. The position dependence

of the coefficients tells us that the coordinate mesh can stretch, squeeze, and twist from place to place. These functions are the metric.

Now the final bridge of the lecture can be stated cleanly. Curvature is the obstruction to finding coordinates that globally flatten the geometry. Tidal forces are the obstruction to finding coordinates that globally eliminate the gravitational field. The connection is not accidental. It is the first glimpse of general relativity as geometry.

2.8 Summary

Lecture 2 begins with a seemingly peripheral question about dark energy, but that opening detour performs useful work: it clears away confusion before the main argument returns to Newton. The Newtonian part of the lecture is then rebuilt in a more modern field language. We define the gravitational field as the acceleration of a test mass, summarize the many-body inverse-square law by

$$\nabla \cdot \vec{A} = -4\pi G\rho,$$

and combine Gauss's law with Gauss's theorem to recover both the exterior inverse-square field and the linear interior field of a uniform-density Earth.

Only after that long Newtonian runway does Einstein's idea become sharp. An accelerated elevator produces an effective gravitational field; the transformed coordinate satisfies $\ddot{x}' = -g$; and once one takes that principle seriously for all physics, light itself must bend. But the principle is only local. Tidal forces remain. That is why the lecture ends with the metric and curvature: not yet with the full theory, but with the right obstruction, the right bridge, and the right sense that gravity is what cannot finally be transformed away.