Advanced Quantum Mechanics: Advanced Quantum Mechanics Lecture 2

Leonard Susskind

Transcript-derived notes curated by LazyingArt LLC

Original lecture by Leonard Susskind. Transcript-derived notes curated by LazyingArt LLC with Video2Book.

Chapter 1

Symmetry, Degeneracy, and Angular Momentum for Quantum Rotational Systems

This chapter develops a symmetry-first view of rotational quantum systems. We define symmetries as operators commuting with the Hamiltonian, distinguish generic level coincidences from symmetry-protected multiplets, and build angular-momentum structure from the commutator algebra of generators. The particle on a circle provides an explicit worked model where quantized m-values, reflection pairing, and ladder operators emerge from operator algebra rather than ad hoc assumptions.

1.1 Symmetry as an invariance principle

A symmetry is an operation that leaves the quantum description unchanged. For a unitary transformation S, invariance of dynamics is the condition

$$SHS^\dagger = H$$

equivalently

$$[H, S] = 0.$$

Because S commutes with H, Schrödinger evolution is covariant under S: if $|\psi(t)\rangle$ evolves by H, then $S|\psi(t)\rangle$ evolves in the same way. In particular, E-eigenvalues are mapped to the same E-eigenspaces.

Symmetries can be exact (invariance of the full Hamiltonian), approximate (effective in a regime), or accidental (valid only for very specific parameter choices). Our focus is on exact symmetries because they control spectral structure.

1.2 Symmetry examples and the organizing role of operators

The same formal rule $[H, S] = 0$ unifies many familiar examples:

- spatial translations and rotations,

- interchange of identical particles,
- relabeling symmetries of a periodic lattice (discrete translation in crystal index),
- discrete mirror or inversion operations.

In practice, this is less about geometry alone and more about algebraic leverage: a set of commuting operators gives you a good quantum-number basis, while noncommuting families often generate nontrivial multiplets. Symmetry language gives a compact way to classify allowed degeneracies before solving differential equations.



Figure 1.1: Discrete translation and relabeling symmetry on an idealized crystal lattice, one of the lecture's opening examples of an operation that leaves the energy spectrum unchanged.

1.3 Degeneracy, symmetry, and when degeneracy is protected

For a bounded discrete spectrum, exact equality of two nearby energies is rare unless there is structure behind it. We call this *accidental* coincidence when no symmetry explanation exists. In contrast, symmetry-protected degeneracies are stable under small perturbations that preserve the same symmetry.

Proposition 1.1. *Let A and B be Hermitian symmetry generators commuting with H: $[A, H] = [B, H] = 0$. Define*

$$C := \frac{1}{i\hbar}[A, B].$$

Then C is Hermitian and $[C, H] = 0$.

Proof. Since A and B are Hermitian, $[A, B]^\dagger = -[A, B]$, so C is Hermitian. Then

$$[C, H] = \frac{1}{i\hbar}([AB, H] - [BA, H])$$

and expanding each commutator gives zero because both A and B commute with H. Hence $[C, H] = 0$. $\square$

Thus one can grow a closed set of symmetry generators by commutators. If that set closes to a Lie algebra with nontrivial structure constants, the representation theory is what drives degeneracy and spectral multiplets.

1.4 Particle on a circle: quantized L_z from single-valued wave functions

Let $\theta \in [0, 2\pi)$ label the position of a particle constrained to a circle. A small rotation by ϵ acts as

$$(\mathcal{R}(\epsilon)\psi)(\theta) = \psi(\theta - \epsilon).$$

Its infinitesimal form defines the generator L_z:

$$\mathcal{R}(\epsilon)\psi(\theta) \approx \left(1 - \frac{i\epsilon}{\hbar} L_z\right)\psi(\theta), \qquad L_z = -i\hbar\frac{d}{d\theta}.$$

Hence

$$L_z \left|\psi_m\right\rangle = m\hbar \left|\psi_m\right\rangle \quad \Longleftrightarrow \quad -i\hbar\frac{d\psi_m}{d\theta} = m\hbar\psi_m$$

and therefore

$$\psi_m(\theta) = \frac{1}{\sqrt{2\pi}} e^{im\theta}.$$

Single-valuedness on $0 \sim 2\pi$ gives

$$\psi_m(\theta + 2\pi) = \psi_m(\theta) \iff e^{i2\pi m} = 1 \iff m \in \mathbb{Z}.$$

So the orbital angular-momentum quantum numbers are $m = 0, \pm 1, \pm 2, \ldots$.



Identification of θ and $\theta + 2\pi$: $\psi(\theta + 2\pi) = \psi(\theta)$

Figure 1.2: Circle coordinate, opposite angles, and periodicity condition.

1.5 Mirror symmetry and the $m \leftrightarrow -m$ pairing

Introduce a mirror (reflection) operator $\mathcal{M}$ with

$$(\mathcal{M}\psi)(\theta) = \psi(-\theta).$$

Its action on a momentum eigenmode is

$$\mathcal{M}\psi_m(\theta) = \psi_{-m}(\theta).$$

On a circle eigenstate:

$$\mathcal{M}L_z\psi_m = m\hbar\,\mathcal{M}\psi_m, \qquad L_z\mathcal{M}\psi_m = -m\hbar\,\mathcal{M}\psi_m,$$

so

$$[\mathcal{M}, L_z]\psi_m = 2m\hbar\,\mathcal{M}\psi_m \neq 0 \quad (m \neq 0).$$

Thus $\mathcal{M}$ does not commute with L_z, even though both may commute with H in a symmetric setup. When both symmetries are present (no mirror-breaking field), states m and $-m$ are paired at equal energy.

Important warning from the formalism: equal probabilities $|\psi_m(\theta)|^2 = |\psi_{-m}(\theta)|^2$ does not imply identical wave functions. Degeneracy concerns states and operators, not only densities.

A uniform magnetic field is a standard counterexample: it can split $\pm m$ levels while preserving some rotational structure of orbital motion in other settings. Symmetry arguments apply to what the Hamiltonian does, not to geometric intuition alone.

1.6 Generators, commutator algebra, and closure

For continuous symmetries a small transformation is written as

$$U(\epsilon) = 1 - \frac{i\epsilon}{\hbar}G + \mathcal{O}(\epsilon^2),$$

with Hermitian generator G. The commutator of two generators is another generator of another direction in the same symmetry structure if closure holds.

A useful worked pattern is:

$$\begin{aligned} &[\,\mathcal{M}, L_z\,] \neq 0, \\ &[H, L_z] = 0, \qquad [H, \mathcal{M}] = 0 \quad \text{(if mirror symmetric)}. \end{aligned}$$

Hence $\mathcal{M}$ and L_z generate two operations whose noncommutativity forces additional structure.

1.7 Three-dimensional angular momentum and ladder structure

For a particle in 3D,

$$\begin{aligned} L_x &= yp_z - zp_y, \\ L_y &= zp_x - xp_z, \\ L_z &= xp_y - yp_x. \end{aligned}$$

with canonical $[x_i, p_j] = i\hbar\delta_{ij}$. A compact expansion gives

$$[L_x, L_y] = i\hbar L_z,$$

and cyclic permutations. Equivalently,

$$[L_i, L_j] = i\hbar \sum_k \epsilon_{ijk} L_k.$$

For a rotationally invariant Hamiltonian, $[H, L_i] = 0$, so L_x, L_y, L_z are conserved quantities.

Define raising and lowering operators

$$L_\pm := L_x \pm iL_y.$$

Using the algebra above,

$$[L_z, L_\pm] = \pm\hbar L_\pm, \qquad [L_+, L_-] = 2\hbar L_z.$$

If $|\ell, m\rangle$ is an eigenvector of L_z,

$$L_z|\ell, m\rangle = m\hbar|\ell, m\rangle,$$

then

$$L_z(L_\pm|\ell, m\rangle) = (m \pm 1)\hbar(L_\pm|\ell, m\rangle),$$

so $L_\pm$ create a state with adjacent m-value when nonzero.

In a finite ladder representation, the chain terminates:

$$L_+|\ell, \ell\rangle = 0, \qquad L_-|\ell, -\ell\rangle = 0,$$

giving $m = -\ell, -\ell+1, \ldots, \ell$, hence $2\ell+1$ states in one rotational multiplet. For orbital wave functions on configuration space, $\ell, m \in \mathbb{Z}$ and $m \in \{-\ell, \ldots, \ell\}$. For more general angular-momentum types (including spin), half-integer m spectra are also possible; in this lecture the orbital circle argument selects the integer branch.



Figure 1.3: Illustrative ladder picture: $L_\pm$ connect adjacent m-levels inside a symmetry multiplet.

1.8 Summary

Symmetry enters as a commutation criterion with the Hamiltonian. Single symmetries can explain invariant spectral structure, but genuine multiplets require a noncommuting symmetry set whose commutator algebra closes. The circle provides the cleanest demonstration: rotational invariance gives $L_z = -i\hbar\partial_\theta$, single-valuedness gives integer m, and mirror symmetry links m to $-m$. In three dimensions, the algebra $[L_i, L_j] = i\hbar\epsilon_{ijk}L_k$ yields ladder operators $L_\pm$, a finite m-tower for each ℓ, and equal-energy multiplets in rotationally invariant systems.