String Theory

Leonard Susskind

Supplementary track, 2010 Fall String Theory And M Theory
Transcript-derived notes curated by LazyingArt LLC

Original lectures by Leonard Susskind. Transcript-derived course notes curated by LazyingArt LLC.

Contents

1 String Theory and M-Theory **1**

1.1 Hadrons, Towers, and Regge Trajectories . . . 1

1.2 Meson Scattering and the Duality Tension . . . 2

1.3 From Quark Pictures to World-Sheets . . . 3

1.3.1 Question & Answer . . . 3

1.4 Flux Tubes, Stretchability, and the Historical Turn . . . 4

1.5 Relativity Reorganized in the Infinite-Momentum Frame . . . 4

1.6 Open Strings as Masses and Springs . . . 6

1.7 Internal Energy as M^2 . . . 8

1.7.1 Question & Answer . . . 9

1.7.2 After-Break Clarifications . . . 9

1.8 Summary . . . 10

2 Lecture 2: String Theory and M-Theory **12**

2.1 Mathematical Preliminaries on the Interval $[0, \pi]$ . . . 12

2.2 What Makes a String a Particle? . . . 14

2.2.1 Question & Answer . . . 14

2.3 Review of the Infinite-Momentum / Light-Front Trick . . . 15

2.4 A String as Mass Points and Springs . . . 16

2.5 Continuum Energy, Lagrangian, and Wave Interpretation . . . 18

2.6 Endpoint Dynamics and the Boundary-Condition Puzzle . . . 20

2.6.1 Question & Answer . . . 20

2.7 Fourier Modes, Harmonic Oscillators, and the First Quantum Spectrum . . . 21

2.8 Summary . . . 24

Chapter 1

String Theory and M-Theory

These notes follow Leonard Susskind's opening lecture on string theory, with curation by LazyingArt LLC. The lecture begins where string theory actually began: not with quantum gravity, but with hadron physics. We first follow an experimental pattern, then a scattering puzzle, then a geometric re-description, and only after that do we turn to the mathematics of strings. The order is essential. The lecture is not presenting a finished formalism and then sprinkling in some history; it is letting the need for the formalism emerge.

1.1 Hadrons, Towers, and Regge Trajectories

String theory entered physics through hadrons. In the late 1960s one did not yet speak in the settled modern language of quarks and gluons, but one did have spectroscopy. Protons, neutrons, mesons, and their excitations did not appear as isolated accidents. They came in families, and the families showed a remarkably simple pattern.

The organizing device was the Chew–Frautschi plot: angular momentum vertically, mass squared horizontally. Susskind moves rather freely between angular momentum, spin, and the symbol L; for the written notes it is cleaner to denote the vertical axis by J. The experimental fact was that hadronic families lay approximately on straight lines,

$$J \propto M^2, \tag{1.1}$$

and, more strikingly, different families had approximately the same slope. In this sense there was something like a universal Regge slope. One may summarize the observed quantization schematically as

$$\Delta J = 1 \qquad \Longrightarrow \qquad \Delta M^2 \approx \text{constant}. \tag{1.2}$$

For mesons the angular momenta are integers; for baryons they are half-integers. But in both cases the same broad phenomenon appears: higher and higher spin states, marching upward in nearly parallel lines on the J versus M^2 plane.

This is already a strong clue. One can spin up a basketball or an atom, but the resulting trajectory is not a straight line in M^2, and in any case the thing eventually breaks apart or ionizes. The hadronic pattern is too simple, too linear, and too persistent. It tells us that hadrons are not point particles. They are composite objects that can be excited rotationally, and they must have a rather special kind of stretchability.



Figure 1.1: Regge trajectory and dual scattering sketches.

This is the first beat of the lecture. It is historical, but not decorative. The data are already suggesting the right sort of object: something extended, excitable, and apparently not eager to give out as the spin increases.

1.2 Meson Scattering and the Duality Tension

That was one observation. Then there was another. Susskind now pivots from spectroscopy to scattering, and he does it concretely: take pion–pion scattering. In conventional field-theoretic language, two pions may come together, form a ρ-like resonance, and then emerge again as two pions. But if that diagram exists, then the crossed diagram exists as well: the same ingredients also describe the exchange of a ρ-like state between the pions.

So far there is no mystery. The mystery arrives when we remember that the ρ is not a single isolated particle. It sits on a whole Regge trajectory. Therefore in the annihilation channel one should not insert only the ρ, but also the ρ', the ρ'', and so on. Likewise in the exchange channel. The lecture never turns this into a detailed amplitude analysis, but the conceptual statement can be written schematically as

$$\mathcal{A}_{\pi\pi}(s,t) \sim \sum_n \mathcal{A}^{(s)}_{\rho_n}(s) \qquad \text{and} \qquad \mathcal{A}_{\pi\pi}(s,t) \sim \sum_n \mathcal{A}^{(t)}_{\rho_n}(t). \tag{1.3}$$

The point is not that these are exact formulas in the lecture. The point is that when people



Figure 1.2: A clean redraw of the board cluster: a Regge plot at left and the two channel pictures at right.

numerically summed the known resonances in one channel, they seemed to be getting, approximately and already, what the other channel would have contributed.

That is the duality tension. Ordinary quantum field theory says: if this diagram exists and that diagram exists, then we must add them. The hadronic data seemed instead to be saying that the sum over a full tower in one channel was already reproducing the other. Naively adding both looked like overcounting.

The board itself makes this transition visible: the Regge plot and the two channel sketches are drawn as one conceptual cluster, as if the spectral regularity and the dual-channel puzzle were already different faces of the same thing.

1.3 From Quark Pictures to World-Sheets

At this point the lecture becomes exploratory in exactly the right way. Susskind does not say: here is the formal object and here is its theorem. He says, in effect, something is going on here, and the Feynman-diagram bookkeeping is no longer the right way to see it.

The next move is to draw quark diagrams with a topological character. Imagine quark and antiquark lines running upward in time. If we view the whole history as a two-dimensional sheet, then one slice through the sheet looks like annihilation and recreation, while another slice looks like exchange. The two channel pictures are no longer two separate dynamical mechanisms; they are two ways of cutting one underlying history.

This is where the jargon arrives:

Definition 1.1. A world sheet is the two-dimensional spacetime history traced by a string, just as a world line is the one-dimensional spacetime history traced by a point particle.

The lecture is careful here. It does not claim that a full string theory has been derived. It claims something more primitive and, historically, more accurate: a two-dimensional object can naturally account for the duality between the channels.

1.3.1 Question & Answer

Why think of a string-like bridge between the quark and antiquark rather than an ordinary exchanged particle?

Because the object in question is not localized at a single intermediate point. If we take an instant of time, we do not see two endpoints and one exchanged quantum. We see a quark, an antiquark, and a whole bunch of stuff in between. Susskind explicitly leaves open whether that intervening stuff is truly continuous or only an approximation to continuity; quantum mechanics makes the distinction subtle. But whatever it is, it fills the interval between the endpoints. Moreover, the Regge trajectories showed no sign of ending. A shoelace with two golf balls at its ends eventually breaks when spun hard enough. The hadronic object, as far as the data then showed, did not. That is what made the one-dimensional extended picture feel more basic than ordinary particle exchange.

Once one says this, the first hadronic clue returns in a new form. A hadron that is a string with two quarks attached to it is exactly the kind of object one can spin up, deform, and analyze energetically.

1.4 Flux Tubes, Stretchability, and the Historical Turn

The lecture now pauses the crude historical picture and reinterprets it in modern terms. Today we would say that the string between a quark and an antiquark is a flux tube of the gluon field. The most useful contrast is with electrodynamics. Between opposite electric charges the field lines spread out, so the field weakens as the charges are pulled apart. That is not a string.

In QCD the nonlinearities of the gauge field change the story. The effective field lines do not simply spread; they are drawn into a tube. As the endpoints separate, the tube gets longer rather than thinner. One may think of the tube as made of gluonic structure. Then the key property is that the energy of stretching does not merely increase the distance between fixed constituents; it goes into creating more constituents in between, so that the density along the tube stays roughly fixed.

This is why Susskind leans on the analogy with a rubber band whose number of molecules is *not* fixed. A normal rubber band stretches by separating the same molecules further and further until it breaks. The hadronic string, in the lecture's modernized picture, stretches by producing more stuff in the gaps. That is why it can remain string-like over a long range of excitation.

The same physical lesson appears in the superconductor analogy. A confined line of magnetic flux in a superconductor has approximately uniform field strength along its length. Uniform field strength means uniform energy density, and uniform energy density means energy proportional to length. We do not need the full superconducting construction here; the moral is enough. Field theory really does produce string-like objects with tension.

Only after making this point does Susskind step back historically. The mathematics that grew out of these hadronic ideas did not, in its simplest form, become the final theory of hadrons. It kept producing, embarrassingly, a massless spin-two particle. What had looked like a nuisance in hadron physics would later be recognized as the graviton. So the mathematical theory born from hadron phenomenology turns out to look more like quantum gravity than like hadronic spectroscopy. But before one gets to that reinterpretation, one still needs the basic mechanics of the string itself, and for that the lecture turns sharply to kinematics.

1.5 Relativity Reorganized in the Infinite-Momentum Frame

Before talking about string mathematics, the lecture talks about relativity versus nonrelativistic kinematics. This is not a digression. It is the license for everything that follows.

For a slow particle the energy is

$$E_{\mathrm{nr}} = \frac{p^2}{2m} + B, \tag{1.4}$$

where B is an additive internal or binding term that does not depend on the state of motion. In relativity the exact energy is

$$E = \sqrt{p^2c^2 + m^2c^4}. \tag{1.5}$$

If $p \ll mc$, then we can factor out the large term and expand:

$$E = mc^2\sqrt{1 + \frac{p^2}{m^2c^2}} \tag{1.6}$$

$$\approx mc^2 + \frac{p^2}{2m}. \tag{1.7}$$

This is the familiar nonrelativistic limit. But Susskind insists that it is only an approximation, and it is a good approximation only when all relevant particles are moving slowly, not merely the center of mass of the system.

Now comes the other expansion, the one that matters for strings. Boost the entire system so that every constituent carries a very large positive momentum along the z-axis. From here on we set $c = 1$. Then

$$E_i = \sqrt{P_{zi}^2 + p_{\perp i}^2 + m_i^2}, \qquad p_{\perp i}^2 = p_{xi}^2 + p_{yi}^2. \tag{1.8}$$

The board notation shifts between P_3 and P_z; we standardize to P_z. Since P_{zi} is large while $p_{\perp i}$ and m_i remain fixed, we expand in powers of $1/P_{zi}$:

$$E_i = P_{zi}\sqrt{1 + \frac{p_{\perp i}^2 + m_i^2}{P_{zi}^2}} \tag{1.9}$$

$$\approx P_{zi} + \frac{p_{\perp i}^2 + m_i^2}{2P_{zi}}. \tag{1.10}$$

Summing over all constituents gives

$$E = \sum_i P_{zi} + \sum_i \frac{p_{\perp i}^2}{2P_{zi}} + \sum_i \frac{m_i^2}{2P_{zi}}. \tag{1.11}$$

The first term is just the total momentum down the z-axis. It is huge, but it is also conserved. As an additive constant in the energy it may be dropped without affecting the dynamics. The remaining expression can be read as a Hamiltonian:

$$H = \sum_i \left(\frac{p_{\perp i}^2}{2P_{zi}} + \frac{m_i^2}{2P_{zi}} \right). \tag{1.12}$$

The board then reinterprets the same formula as the generator of time evolution,

$$i\,\partial_t = H, \tag{1.13}$$

with $\hbar = 1$.

Now the lecture makes the crucial interpretation. In this transverse description the quantity in the denominator plays the role that mass usually plays in nonrelativistic mechanics. So P_z acts



Figure 1.3: Large-P_z expansion of the relativistic energy.

as transverse inertia. Meanwhile the term involving m_i^2 is independent of the transverse state of motion, and so it plays the role of internal or binding energy. This is the exact structural reason that a relativistic system, viewed in the infinite-momentum frame, behaves nonrelativistically in the transverse plane.

Susskind then adds the physical explanation. Because the Hamiltonian scales like $1/P_z$, the evolution becomes slow when P_z is large. That is just time dilation. One is not denying relativistic physics; one is reorganizing it into a form that is calculable.

1.6 Open Strings as Masses and Springs

By what chutzpah do we now use nonrelativistic physics to describe a relativistic string? By the chutzpah just earned. In the infinite-momentum frame the transverse dynamics really are nonrelativistic, and that is enough to write down the simplest model.

Take an open string. Introduce a coordinate σ along it, with

$$0 \leq \sigma \leq \pi.$$

The choice of π is conventional; Susskind chooses it because later closed strings will naturally run from 0 to 2π.

Start discretely. Think of the string as a chain of point masses joined by springs. If $X_i(t)$ denotes



Figure 1.4: Hamiltonian form of the light-cone energy.

the transverse position of the i-th mass point, then the energy is

$$E = \sum_i \left(\frac{\mu}{2} \dot{X}_i^2 + \frac{k}{2} (X_i - X_{i+1})^2 \right). \tag{1.14}$$

Here $\dot{X}_i^2$ abbreviates the sum of the two transverse components, and the second term is simply Hooke's law.

The lecture is careful about the continuum limit. As the beads become denser, the mass of each bead must go to zero so that the total mass remains finite. At the same time the spring constant must scale upward: stretching a tiny segment by a fixed amount is harder than stretching a long segment by the same amount. When the spacing goes to zero, the sum becomes an integral and the nearest-neighbor difference becomes a derivative:

$$\sum_i \longrightarrow \int_0^\pi d\sigma, \qquad X_i - X_{i+1} \longrightarrow \partial_\sigma X. \tag{1.15}$$

Thus the continuum energy takes the form

$$E \sim \int_0^\pi d\sigma \left[\frac{1}{2} \dot{X}^2(\sigma, t) + (\partial_\sigma X(\sigma, t))^2 \right]. \tag{1.16}$$

The exact coefficients are not fully fixed by the board, and Susskind explicitly says he has absorbed constants by convention. What matters is the structure: kinetic energy from $\partial_t X$, potential energy from the stretching measured by $\partial_\sigma X$.



Figure 1.5: Discrete string energy and continuum σ-derivative limit.

The lecture pauses here for a useful distinction. If we were writing the Lagrangian, the potential term would come with a minus sign. But Susskind wants the energy, the Hamiltonian, so the terms appear with plus signs. That is the object we will need when we ask what the internal motion of the string contributes to the observable particle.

1.7 Internal Energy as M^2

We now return to the key interpretive question. The string may vibrate and stretch, but as a whole it is still one object. We may call it a particle if we like. Its center of mass describes the overall motion of the particle, while the relative stretching and internal vibration describe its internal energy.

What, then, does the Hamiltonian of the previous section correspond to? In ordinary rest-frame reasoning one would identify internal energy with mass. But the whole point of the infinite-momentum analysis was that in this frame internal energy enters as mass squared. Thus the string Hamiltonian should be read as contributing to M^2, not to M:

$$H_{\text{int}} \propto M^2. \tag{1.17}$$

Susskind is quite emphatic about this. It is not just an analogy; it is the precise way relativity reorganizes itself in the light-cone frame.

This immediately explains why string excitations naturally come in evenly spaced units of M^2. The string is a collection of coupled oscillators. We do not quantize it fully in this lecture, but

the principle is already clear: oscillator energies are quantized in equal steps, so string excitations contribute equal steps to the internal Hamiltonian, hence to mass squared. This is exactly the kind of spacing the Regge trajectories seemed to demand.

1.7.1 Question & Answer

Why does the same string look Hooke-like in one frame but tension-like in its rest frame?

Because the Hooke-like energy is the effective internal energy of the boosted description, and that quantity maps to M^2, not directly to the rest energy M. The rest-frame string is recovered only after we translate the light-cone result back into ordinary relativistic language.

Let us go through the argument the way the lecture does. Consider a uniformly stretched string of physical length L. Along the parameter interval $0 \leq \sigma \leq \pi$ we then have, schematically,

$$\partial_\sigma X \sim \frac{L}{\pi}. \tag{1.18}$$

Insert this into the stretching term:

$$H_{\text{stretch}} \sim \int_0^\pi d\sigma \, (\partial_\sigma X)^2 \tag{1.19}$$

$$\sim \int_0^\pi d\sigma \left(\frac{L}{\pi}\right)^2 \tag{1.20}$$

$$= \pi \left(\frac{L}{\pi}\right)^2 \propto L^2. \tag{1.21}$$

So in the effective nonrelativistic description the stretching energy behaves like Hooke's law:

$$H_{\text{stretch}} \propto L^2. \tag{1.22}$$

But this energy is to be identified with M^2. Therefore

$$M^2 \propto L^2, \qquad \text{so} \qquad M \propto L. \tag{1.23}$$

In the rest frame the energy of the string is its mass, and so

$$E_{\text{rest}} = M = T\,L. \tag{1.24}$$

Thus the rest-frame string is not an ordinary Hooke-law spring at all. It is a constant-tension object, precisely of the flux-tube type discussed earlier.

Susskind is careful here: this is an indirect deduction. We are not claiming to know how an arbitrary ordinary rubber band would behave if it were driven relativistically. What we have done is use the frame in which the transverse dynamics become simple, derive the L^2 behavior there, and then translate back. That is why the result is trustworthy even though the route is indirect.

1.7.2 After-Break Clarifications

After the break the lecture revisits the same point more slowly, because this is where the sleight of hand can be missed. The same relativistic formula

$$E = \sqrt{p^2 + m^2} \tag{1.25}$$

admits two different expansions.

If p is small and m is large, then

$$E \approx m + \frac{p^2}{2m}. \tag{1.26}$$

This is the ordinary rest-frame logic: internal energy is read as mass. If, on the other hand, p is very large, then

$$E \approx p + \frac{m^2}{2p}. \tag{1.27}$$

After the conserved momentum term is removed, the remaining energy is proportional to m^2. That is the whole conceptual switch. In one frame the relevant internal quantity is M; in the boosted frame it is M^2.

The lecture also addresses a natural question about residual motion along the boost direction. In light-cone string theory one does not introduce independent longitudinal degrees of freedom in the same way one introduces the transverse ones. The description is transverse, and yet it remains Lorentz invariant. In the lecture this is presented as a remarkable fact rather than something to prove immediately, and that is the right level to preserve here.

Finally, Susskind returns to the electron question. Why do hadrons show low-lying higher-spin excitations while electrons do not? The answer is the rotational estimate

$$E_{\text{rot}} \sim \frac{L^2}{2I}, \qquad I \sim MR^2. \tag{1.28}$$

Since angular momentum comes in quanta, the first excited rotational state costs roughly

$$\Delta E \sim \frac{\hbar^2}{2I}. \tag{1.29}$$

If the object is tiny, then R is tiny, hence I is tiny, and the excitation energy is enormous. Electrons are experimentally exceedingly small, so any such excitations would lie far beyond accessible collision energies. Hadrons are large on the scale of their own internal dynamics, so their moments of inertia are large enough that spinning them up by one unit costs only hadronic energies.

This closes the loop back to the opening observation. The Regge plots were not merely classification devices. They were telling us that hadrons behave as extended objects with accessible rotational and vibrational excitations.

1.8 Summary

The lecture begins with hadron phenomenology and lets string theory emerge from it. Straight, parallel Regge trajectories suggest that hadrons are composite, stretchable objects rather than points. The duality puzzle in pion scattering then suggests that what field theory wants to count as separate channels may really be two slicings of one underlying two-dimensional history, a world sheet.

From there the lecture pauses to reinterpret the old picture in modern QCD terms: a flux tube has constant energy per unit length, and so it behaves like a string with tension. Only then does the mathematical turn arrive. In the infinite-momentum frame a relativistic system becomes nonrelativistic in the transverse plane, with P_z playing the role of inertia and the internal term

behaving like mass squared. That allows us to model an open string as masses and springs, pass to the continuum Hamiltonian built from $\dot{X}^2$ and $(\partial_\sigma X)^2$, and then read that Hamiltonian as contributing to M^2. The payoff is immediate: Hooke-law stretching in the light-cone description becomes linear-tension behavior in the rest frame, and string excitations naturally organize themselves in evenly spaced quanta of mass squared.

Chapter 2

Lecture 2: String Theory and M-Theory

These notes follow Leonard Susskind's second lecture on string theory, curated in this archive by LazyingArt LLC. The lecture proceeds in a very deliberate order. First it stocks the blackboard with the interval calculus and Fourier facts that would otherwise interrupt the story later. Then it pauses over a conceptual question that matters more than it first appears to: in what sense can a string, made of many pieces, still count as a particle? Only after that does it return to the infinite-momentum or light-front point of view, build the open string as a chain of masses and springs, take the continuum limit, solve the endpoint puzzle, and quantize the resulting modes.

2.1 Mathematical Preliminaries on the Interval $[0, \pi]$

Susskind begins by saying explicitly why he is doing this: if these preliminaries were not written down now, they would have to be inserted later in the middle of the string theory, and that would be a nuisance. So we begin with the reservoir of tools.

Take a function $x(\sigma)$ on the interval $0 \leq \sigma \leq \pi$. The choice of letters is not accidental. He uses x and σ because these are the notations string theory will keep returning to. The interval $[0, \pi]$ is likewise chosen in advance because it is the natural interval for the open string in this lecture.

We approximate the continuous function by values on a lattice of N points. If

$$\Delta\sigma = \frac{\pi}{N}, \tag{2.1}$$

then the standard replacements are

$$x(\sigma) \longrightarrow x_i, \tag{2.2}$$

$$\Delta x_i \equiv x_i - x_{i-1} \approx \frac{\partial x}{\partial \sigma}\,\Delta\sigma, \tag{2.3}$$

$$\sum_i x_i\,\Delta\sigma \longrightarrow \int_0^\pi x(\sigma)\,d\sigma. \tag{2.4}$$

This is just calculus, but it is precisely the calculus we will soon use for the string.

The next distinction is between endpoint behaviors. Here the lecture is still mathematical rather than physical, but the physical pictures are already visible.

Dirichlet boundary conditions. These fix the value of the function at the endpoints:

$$x(0) = x(\pi) = 0. \tag{2.5}$$

A violin string with its ends pinned down is the elementary model.

Neumann boundary conditions. These instead fix the slope at the endpoints:

$$\left.\frac{\partial x}{\partial \sigma}\right|_{\sigma=0} = \left.\frac{\partial x}{\partial \sigma}\right|_{\sigma=\pi} = 0. \tag{2.6}$$

The function need not vanish there; it is only required to be flat.

The lecture briefly wanders through organ pipes and other examples, but the moral is simple. Dirichlet means fixed endpoints. Neumann means free endpoints. Susskind even gives a mechanical picture of a string whose ends pass through very light rings sliding on poles: not held down, but free to move. That is the kind of system for which Neumann conditions are natural.

There is also a small caution here. Piecewise continuous functions are allowed in Fourier analysis, but for the purposes of string theory we do not want jumps in the string itself: a jump would mean the string had broken.

Once the endpoint behavior is fixed, the Fourier basis is fixed with it. On $[0, \pi]$ one writes

$$x_D(\sigma) = \sum_{n=1}^{\infty} x_n \sin n\sigma, \tag{2.7}$$

$$x_N(\sigma) = \sum_{n=0}^{\infty} x_n \cos n\sigma. \tag{2.8}$$

Sines vanish at the endpoints, which is why they go with Dirichlet. Cosines have zero derivative at the endpoints, which is why they go with Neumann. The lecture also emphasizes a fact we will use later without apology: differentiation swaps the two classes. The derivative of a Dirichlet function is Neumann; the derivative of a Neumann function is Dirichlet.

To make these expansions useful, we need the orthogonality integrals. In the form relevant later,

$$\int_0^{\pi} \cos n\sigma \cos m\sigma \, d\sigma = 0 \qquad (n \neq m), \tag{2.9}$$

$$\int_0^{\pi} \cos^2 n\sigma \, d\sigma = \frac{\pi}{2} \qquad (n \geq 1), \tag{2.10}$$

$$\int_0^{\pi} d\sigma = \pi, \tag{2.11}$$

$$\int_0^{\pi} \sin n\sigma \sin m\sigma \, d\sigma = \frac{\pi}{2}\delta_{nm} \qquad (n, m \geq 1). \tag{2.12}$$

Remark 2.1. The transcript becomes garbled around the exceptional $n = 0$ cosine mode, but the intended mathematics is clear. The nonzero cosine modes have norm $\pi/2$, while the constant mode $\cos 0\sigma = 1$ integrates to π.

The last item placed on the board is the harmonic oscillator. After rescaling the coordinate so that the mass is one, the energy reads

$$E_{\text{osc}} = \frac{1}{2}\dot{x}^{\,2} + \frac{1}{2}\omega^2 x^2, \tag{2.13}$$

and the corresponding Lagrangian is

$$L_{\text{osc}} = \frac{1}{2}\dot{x}^{\,2} - \frac{1}{2}\omega^2 x^2. \tag{2.14}$$

The lecture contains a verbal slip here, but the mathematics is standard: the Lagrangian is $T - V$, not "kinetic energy squared minus potential energy squared." The quantum fact we keep in reserve is that oscillator levels come in steps of ω, up to the overall ground-state offset.

2.2 What Makes a String a Particle?

Now the lecture deliberately interrupts its own mathematical flow. Before building the string, it asks a question that could otherwise become a source of confusion for the rest of the course: what do we mean by a particle at all?

A string, after all, is about to be modeled as a large number of pieces connected by springs. If that is so, why should it behave like a particle rather than like an ordinary extended object?

The first part of the answer is negative. Pointlikeness is not the criterion. Electrons are surrounded by virtual clouds. Protons and neutrons are obviously composite. The lecture insists that what we call a particle need not be literally structureless.

Nor is it enough merely to have a mass and a center-of-mass position. A box of gas has those. A cup of coffee has those. Yet no physicist would naturally call a cup of coffee a particle.

2.2.1 Question & Answer

Question. When does a composite object count as a particle rather than "mush"?

Answer. A composite object counts as a particle when the excitation spectrum above its ground state is discrete and well separated. If there are large enough gaps to the first few excited states, the object behaves as an isolated quantum species. If instead there are innumerable nearby states at arbitrarily small separations, the object behaves like continuum matter.

This is the lesson of the lecture's coffee example. A cup of coffee can be excited by tiny disturbances: one can stir it slightly, excite a sound mode, disturb a molecule, or create a phonon with an absurdly small amount of energy. The nearby states are so dense that they practically form a continuum. That is mush.

A proton is different. It certainly has excitations, but it takes a substantial amount of energy to get to them. The nearby spectrum is not continuous. An electron is more isolated still. One does not have a ladder of electron states lying immediately above the electron.

So the criterion becomes spectral rather than ontological. The question for string theory is not whether the string is made of parts. Of course it is. The question is whether the internal excitations are separated by large enough gaps to make the lowest state behave as a particle. In the version of string theory under discussion, the gap is taken to be enormous, on the scale of the Planck mass, and that is why the string may still present itself as a particle in ordinary experiments.

This detour is not philosophical ornament. It prepares the central claim of the lecture's last third: once the string is quantized, its spectrum comes out discrete enough to justify calling the resulting states particles.

2.3 Review of the Infinite-Momentum / Light-Front Trick

Only after that conceptual pause does Susskind return to the framework from the previous lecture. The question is this: how can an extended relativistic system ever be described by something as simple as nonrelativistic mechanics?

He first emphasizes what does *not* work. A rapidly spinning baton is not well described by nonrelativistic physics just because we happen to sit in the frame where its center of mass is simple. The issue is the relative motion of its parts. If the ends are moving relativistically relative to each other, then the ordinary nonrelativistic approximation fails.

The trick is different. We boost the whole system to a very large momentum along some chosen axis, say the z-axis, and then examine only the motion in the transverse plane. In that description the transverse dynamics become exactly, not approximately, nonrelativistic in form.

Start from the relativistic energy

$$E = \sqrt{P_z^2 + p_x^2 + p_y^2 + M^2}, \tag{2.15}$$

where M is the rest mass of the full system and P_z is the large longitudinal momentum. When P_z is the dominant scale, we expand:

$$E \approx P_z + \frac{p_x^2 + p_y^2 + M^2}{2P_z}. \tag{2.16}$$

The first term is the trivial boost energy. It depends on the conserved longitudinal momentum but not on the internal state, so for the questions at hand we subtract it and write

$$E - P_z = \frac{p_x^2 + p_y^2 + M^2}{2P_z}. \tag{2.17}$$

The lecture then makes the crucial interpretive move. Energy is Hamiltonian. Hamiltonian governs time evolution. But if P_z becomes very large, the right-hand side tends to zero, which means that in the ordinary time variable the internal motion appears to freeze. This is just time dilation.

We do not want to throw away the internal motion. We want to follow it. So we rescale time. Equivalently, up to a conventional overall normalization, we multiply by P_z. Then the expression has a finite limit:

$$P_z(E - P_z) = \frac{1}{2}\left(p_x^2 + p_y^2 + M^2\right). \tag{2.18}$$

This now looks like a two-dimensional nonrelativistic Hamiltonian: transverse momentum squared divided by two, plus an internal contribution. The only conceptual twist is that the internal contribution is proportional to M^2, not M.

On the board, the line beginning with $H =$ records exactly this reinterpretation. The board itself does not finish writing $H = i\,d/dt$, but the lecture makes that identification in words: Hamiltonian is what governs time evolution.

Remark 2.2. Susskind briefly complains about the terminology "light cone" and says that "light front" would be better. We keep only the mathematical point. After the large boost and the time rescaling, the transverse dynamics are exactly nonrelativistic in form.

There is one more feature he flags, without dwelling on it. For strings, the longitudinal internal motion is constrained by the transverse motion. That is part of what makes the method so powerful. In the present lecture, however, this remains background motivation; the active calculation now moves to the transverse coordinates alone.



Figure 2.1: Hamiltonian form of the boosted energy. The board line beginning with $H =$ marks the shift from the boosted-energy expansion to its Hamiltonian interpretation.

2.4 A String as Mass Points and Springs

Now we come to strings themselves. In the boosted description the string moves only in two transverse directions, which we call X and Y. The lecture begins with the discrete model: N mass points joined by $N-1$ springs.

The coordinate σ labels position along the string and runs from 0 to π. This is not forced; it is chosen. Susskind even remarks that one could equally well choose 0 to 1 or 0 to 2. The reason for choosing $[0, \pi]$ is pedagogical. He wants to reserve $[0, 2\pi]$ for the closed string later on.

The discrete energy of the open string is

$$E = \sum_i \left[\frac{\mu}{2}(\dot{X}_i^2 + \dot{Y}_i^2) + \frac{k}{2}((\Delta X_i)^2 + (\Delta Y_i)^2) \right]. \tag{2.19}$$

The board writes the bead mass as m, but the lecture itself later corrects this and introduces μ as the nonrelativistic analog mass of each bead, so we use μ from the start in the cleaned notes. Likewise, the spring constant is called both k and κ in speech; we write k.

When Susskind writes the formula first for X, he immediately says: wherever I write X, you must also add the same thing with Y. That small spoken reminder matters. The string lives in two transverse directions, not one.

The next task is to choose the large-N scaling correctly. If the number of beads grows while each bead keeps a fixed mass, then the total mass diverges. That is not the continuum limit we want. We therefore choose

$$\mu = \frac{1}{N}, \tag{2.20}$$



Figure 2.2: Discrete string as mass points and springs. The lecture moves directly from the bead-chain sketch to the discrete energy formula.



Figure 2.3: A clean redraw of the bead-chain model used to take the continuum limit.

so that the total analog mass remains fixed.

The spring constant must scale in the opposite direction. Springs in series make a long chain easier to stretch, so if the overall stiffness of the string is to stay finite, the microscopic springs must become stiffer as the chain is refined. Susskind chooses

$$k = \frac{N}{\pi^2}. \tag{2.21}$$

The factor of π^2 is a convention inserted only to simplify later formulas. The essential point is the $k \sim N$ scaling.

This is the real payoff of the opening calculus detour. We now have a discrete mechanical chain and a precise prescription for taking its continuum limit.

2.5 Continuum Energy, Lagrangian, and Wave Interpretation

At this point the lecture uses, almost verbatim, the formulas it put on the board at the beginning. We have $\Delta\sigma = \pi/N$, we have sums, and we have nearest-neighbor differences. Now we turn them into integrals and derivatives.

A worked derivation of the continuum limit. For the X-sector kinetic term,

$$\sum_i \frac{\mu}{2}\dot{X}_i^2 = \sum_i \frac{1}{2N}\dot{X}_i^2 \tag{2.22}$$

$$= \frac{1}{2\pi}\sum_i \dot{X}_i^2\,\Delta\sigma \tag{2.23}$$

$$\longrightarrow \frac{1}{2\pi}\int_0^\pi d\sigma \left(\frac{\partial X}{\partial \tau}\right)^2. \tag{2.24}$$

For the spring term we use the continuum replacement

$$\Delta X_i \approx \frac{\partial X}{\partial \sigma}\,\Delta\sigma. \tag{2.25}$$

Then

$$\sum_i \frac{k}{2}(\Delta X_i)^2 \approx \sum_i \frac{N}{2\pi^2}\left(\frac{\partial X}{\partial \sigma}\right)^2 (\Delta\sigma)^2 \tag{2.26}$$

$$= \sum_i \frac{1}{2N}\left(\frac{\partial X}{\partial \sigma}\right)^2 \tag{2.27}$$

$$= \frac{1}{2\pi}\sum_i \left(\frac{\partial X}{\partial \sigma}\right)^2 \Delta\sigma \tag{2.28}$$

$$\longrightarrow \frac{1}{2\pi}\int_0^\pi d\sigma \left(\frac{\partial X}{\partial \sigma}\right)^2. \tag{2.29}$$

Exactly the same steps apply to Y.

So the continuum energy becomes

$$E = \frac{1}{2\pi}\int_0^\pi d\sigma \left[\left(\frac{\partial X}{\partial \tau}\right)^2 + \left(\frac{\partial Y}{\partial \tau}\right)^2 + \left(\frac{\partial X}{\partial \sigma}\right)^2 + \left(\frac{\partial Y}{\partial \sigma}\right)^2\right], \tag{2.30}$$

and the continuum Lagrangian is

$$L = \frac{1}{2\pi}\int_0^\pi d\sigma \left[\left(\frac{\partial X}{\partial \tau}\right)^2 + \left(\frac{\partial Y}{\partial \tau}\right)^2 - \left(\frac{\partial X}{\partial \sigma}\right)^2 - \left(\frac{\partial Y}{\partial \sigma}\right)^2\right]. \tag{2.31}$$



Figure 2.4: Continuum string Lagrangian. The board records the moment at which the minus sign of the potential term is being written.

The lecture pauses here to make an illuminating comparison. If we rename σ as a spatial coordinate on a line and rename X as a field ϕ, then this is just the energy of an ordinary wave system. Waves move left and right along the interval. When they hit the boundary, they reflect.

The boundary condition determines the character of the reflection. Under Dirichlet conditions the reflected wave flips sign. Under Neumann conditions it reflects without such inversion. Susskind says one could now write down the wave equation by varying the Lagrangian, but refuses to do so because the lecture will not need it. That refusal is part of the rhythm of the presentation, and we keep it. The wave viewpoint is useful; the full partial differential equation is not yet needed.

One more spoken clarification belongs here. The time variable τ is the rescaled time appropriate to the boosted description. Susskind loosely calls it proper time: the time carried along by the boosted clock so that we can continue to watch the internal motion instead of letting it freeze under time dilation.

Finally, the lecture reminds us why this energy matters. In the light-front description, the internal energy of the string will be identified, up to the same overall normalization convention as before, with the square of the relativistic mass of the string state.

2.6 Endpoint Dynamics and the Boundary-Condition Puzzle

Now the lecture returns to a question it deliberately postponed earlier. We have seen both Dirichlet and Neumann conditions. We have heard that free string endpoints should obey Neumann conditions. Why?

Here the lecture is at its cleanest. It raises the obstacle plainly and resolves it with a short Newton-law argument.



Figure 2.5: Discrete difference becomes derivative times spacing. This is the local step that converts the endpoint force into a boundary condition.



Figure 2.6: Endpoint geometry for the Newton-law argument. The last mass point feels a force from only one spring.

2.6.1 Question & Answer

Question. Why must the open-string endpoint satisfy Neumann rather than Dirichlet boundary conditions?

Answer. Because the endpoint is free rather than pinned, and for a free endpoint Newton's law forces the local stretch to vanish in the continuum limit.

Let the last bead be labeled n, and the previous one $n-1$. An interior bead is pulled from both sides. The endpoint is special: it feels only the force from the spring connecting it to the rest of the string. By Hooke's law,

$$F_{\text{end}} \sim k\,\Delta X. \tag{2.32}$$

Now we use the continuum replacement that the board is capturing:

$$\Delta X = \frac{\partial X}{\partial \sigma}\,\Delta\sigma. \tag{2.33}$$

Since $k \sim N$ and $\Delta\sigma \sim 1/N$, the N-dependence cancels, so

$$F_{\text{end}} \propto \frac{\partial X}{\partial \sigma}. \tag{2.34}$$

That is already the key step: the endpoint force is proportional to the slope of the string at the endpoint.

But Newton's law also says

$$F_{\text{end}} = \mu\,\frac{\partial^2 X}{\partial \tau^2}, \qquad \mu \sim \frac{1}{N}. \tag{2.35}$$

So if $\partial_\sigma X$ were nonzero at the endpoint, then the acceleration $\partial_\tau^2 X$ would grow like N as the continuum limit is taken. The endpoint would acquire an infinite acceleration. That is exactly the pathology the lecture wants to exclude.

The cure is immediate:

$$\left.\frac{\partial X}{\partial \sigma}\right|_{\sigma=0,\pi} = 0. \tag{2.36}$$

And the same argument applies to Y. Thus the free open-string endpoint satisfies Neumann boundary conditions.

This is a good example of the lecture's style. A boundary condition that could have been postulated as a formal rule is instead derived from a local mechanical argument. The mathematics is light, but the motivation is precise.

2.7 Fourier Modes, Harmonic Oscillators, and the First Quantum Spectrum

Now the system is finally ready to quantize. The first step is not yet quantum mechanics; it is a re-description of the classical string in terms of its harmonics.

Because the open string obeys Neumann conditions, we expand in cosines:

$$X(\sigma,\tau) = \sum_{n=0}^{\infty} X_n(\tau)\cos n\sigma, \qquad Y(\sigma,\tau) = \sum_{n=0}^{\infty} Y_n(\tau)\cos n\sigma. \tag{2.37}$$

The dependence on σ is carried by the cosines. The dependence on time τ sits in the coefficients $X_n(\tau)$ and $Y_n(\tau)$. That is where the string wiggles.

Differentiating with respect to time gives

$$\dot{X}(\sigma,\tau) = \sum_{n=0}^{\infty} \dot{X}_n(\tau)\cos n\sigma, \tag{2.38}$$

while differentiating with respect to σ gives

$$\frac{\partial X}{\partial \sigma} = -\sum_{n=1}^{\infty} n\, X_n(\tau) \sin n\sigma. \tag{2.39}$$

The board image suggests a sum beginning at $n = 0$, but the cleaned expression must begin at $n = 1$, since the differentiated zero mode vanishes.



Figure 2.7: Mode expansion of the σ-derivative. The visible board evidence is the minus sign, the factor of n, and the passage from cosines to sines.

Now we insert the mode expansions into the energy. This is where the orthogonality formulas from the beginning of the lecture are harvested. All cross-terms with $n \neq m$ vanish after integration over σ. The double sums collapse to single sums. The $n = 0$ mode must be handled separately because its cosine norm is π rather than $\pi/2$.

For the kinetic piece of the X-sector we find

$$\frac{1}{2\pi}\int_0^{\pi} d\sigma\, \dot{X}^2 = \frac{1}{2}\dot{X}_0^2 + \frac{1}{4}\sum_{n=1}^{\infty} \dot{X}_n^2, \tag{2.40}$$

and for the stretching piece,

$$\frac{1}{2\pi}\int_0^{\pi} d\sigma \left(\frac{\partial X}{\partial \sigma}\right)^2 = \frac{1}{4}\sum_{n=1}^{\infty} n^2 X_n^2. \tag{2.41}$$

So the full X-sector energy is

$$E_X = \frac{1}{2}\dot{X}_0^2 + \frac{1}{4}\sum_{n=1}^{\infty}\left(\dot{X}_n^2 + n^2 X_n^2\right), \tag{2.42}$$

and the corresponding X-sector Lagrangian is

$$L_X = \frac{1}{2}\dot{X}_0^2 + \frac{1}{4}\sum_{n=1}^{\infty}\left(\dot{X}_n^2 - n^2 X_n^2\right). \tag{2.43}$$

The Y-sector is identical:

$$L_Y = \frac{1}{2}\dot{Y}_0^2 + \frac{1}{4}\sum_{n=1}^{\infty}\left(\dot{Y}_n^2 - n^2 Y_n^2\right). \tag{2.44}$$

Several things are now visible at once.

First, X_0 and Y_0 are the average positions of the string, hence the center-of-mass coordinates. They have kinetic energy but no restoring force. The zero mode is a free particle mode, not an oscillator.

Second, each mode with $n \geq 1$ is an ordinary harmonic oscillator. Comparing with the earlier oscillator formula, we read off

$$\omega_n = n, \qquad n \geq 1. \tag{2.45}$$

So the open string is a countable infinity of independent oscillators in the X-direction, together with another countable infinity in the Y-direction. These are nothing but the harmonics of the string.

Third, the internal vibrational energy is

$$E_{\text{int}} = \frac{1}{4}\sum_{n=1}^{\infty}\left(\dot{X}_n^2 + n^2 X_n^2 + \dot{Y}_n^2 + n^2 Y_n^2\right). \tag{2.46}$$

Up to the same light-front normalization convention introduced earlier, this is the quantity that becomes the relativistic mass squared of the string state. That is the bridge from oscillator mathematics back to particle physics.

Quantization is now easy. The lecture does not yet introduce creation and annihilation operators systematically; it merely uses the oscillator spectrum. For the n-th oscillator, if Q_n quanta are present and $\hbar = 1$, then the excitation energy above the unspecified ground-state offset is

$$\Delta E_n = Q_n\,\omega_n = Q_n\,n. \tag{2.47}$$

Susskind is careful to rename the occupation number Q so that it is not confused with the mode label n.

The lowest levels can then be counted directly.

Ground state. All oscillators are in their lowest state. This is not empty space. It is one open string with no internal excitation.

First excited level. Excite the $n = 1$ oscillator once. One may do this in the X-tower or in the Y-tower, so the first excited level has multiplicity 2.

Second excited level. Now the total excitation energy is 2. There are five possibilities:

- two quanta in the $n = 1$ X-oscillator,
- two quanta in the $n = 1$ Y-oscillator,

- one quantum in each of the $n = 1$ X- and Y-oscillators,
- one quantum in the $n = 2$ X-oscillator,
- one quantum in the $n = 2$ Y-oscillator.

So the second excited level has multiplicity 5.

This is exactly the point the lecture wanted to set up from the beginning. The open string has a discrete spectrum with appreciable gaps. It therefore behaves like a particle in the precise spectral sense explained earlier.

The final remarks of the lecture are a preview, not a full development. Open strings have endpoints, and once interactions are allowed, endpoints can join and split. A fluctuating open string may bring one of its endpoints into contact with another endpoint and thereby produce a closed string. So one may consistently have a theory of closed strings without open strings, but not a theory of open strings without closed strings. As a rough guide only, the lecture says that open strings often behave like photons, while closed strings behave like gravitons.

2.8 Summary

The lecture unfolds by preparation, obstacle, and payoff. It first stocks the blackboard with the mathematics of functions on $[0, \pi]$: discretization, boundary conditions, Fourier expansions, orthogonality, and the harmonic oscillator. It then sharpens the notion of a particle by replacing pointlikeness with spectral isolation. With that conceptual and mathematical groundwork in place, the infinite-momentum or light-front viewpoint recasts the relativistic string as an exact two-dimensional mechanical system. The open string is built as a chain of masses and springs, passed to a continuum description, fixed by a Neumann endpoint condition derived from Newton's law, and finally rewritten as an infinite collection of uncoupled harmonic oscillators. The resulting spectrum is discrete, gapped, and therefore particle-like. That is the central gain of the lecture, and it is the point from which the next step, the closed string, naturally begins.