

# INTENTIONALLY SCUFFED NOTES

Several Authors

March 17, 2022

**REMEMBER THAT AT ANY POINT, THESE EQUATIONS,  
FORMULAE AND NOTES IN GENERAL MAY CONTAIN  
INTENTIONAL ERRORS. WE ARE NOT LIABLE FOR  
ANYTHING**

## 1 Introduction

Edit later.

## 2 Proof by Induction

### 2.1 Sequences and Series

## 3 Functions

Let us focus on functions.

### 3.1 Linear equation

A linear equation is a polynomial degree 1. It can be shown in several ways.

- General form:  $ax + by = c$
- Slope-intercept form:  $y = mx + c$

## 4 Trigonometry

### 4.1 Angle Sum and Difference

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \pm \sin A \sin B$$

### 4.2 Double Angle Identities

Double angle identities are derived from the angle and sum difference equations.

We know that

$$\sin(2x) = 2 \sin(x) \cos(x)$$

therefore, we can say that

$$\sin(4x) = 4 \sin^2(x) \cos^2(x)$$

### 4.3 $n$ Angle Identities

Using De Moivre's theorem it is known that

$$(|z| \cos \theta)^n = |z|^n \cos n\theta$$

The identity  $\cos \theta = \sin\left(\frac{\pi}{2} - \theta\right)$  can be applied to deduce identities for  $\sin$ .

$$\left(|z| \sin\left(\frac{\pi}{2} - \theta\right)\right)^n = |z|^n \sin\left(\frac{\pi}{2} - \theta\right)^n$$

Since the unit circle has radius 1,  $z = 1$ . Hence, the following identities are derived.

$$\cos n\theta = (|n| \cos \theta)^n \quad \sin n\theta = \left(|n| \sin\left(\frac{\pi}{2} - \theta\right)\right)^n$$

How useful!

## 5 Calculus of the Differential Kind

## 6 Integration

## 7 Complex Numbers

### 7.1 Euler's Form

We know that

$$\sin^2 \theta + \cos^2 \theta = 1$$

We also know the famous equation known as Euler's formula. The imaginary number one. With the pi. No, not the one with polyhedrons. To be more precise, we mean Euler's identity. The "beautiful" equation.

$$e^{i\pi} + 1 = 0$$

Therefore, we can then conclude that:

$$e^{i\theta} + \sin^2 \theta + \cos^2 \theta = 0$$

$$\begin{aligned} \frac{e^{i\theta} + \sin^2 \theta}{-\cos^2 \theta} &= 1 \\ -\frac{e^{i\theta}}{\cos^2 \theta} - \frac{\sin^2 \theta}{\cos^2 \theta} &= 1 \\ -e^{i\theta} \sec^2 \theta - \tan^2 \theta &= 1 \\ -e^{i\theta} \sec^2 \theta &= 1 + \tan^2 \theta \\ -e^{i\theta} \sec^2 \theta &= \sec^2 \theta \\ -e^{i\theta} &= 1 \Rightarrow e^{i\theta} = -1 \end{aligned}$$

What a beautiful conclusion!

## 8 Linear Algebra