

Chapter 3

RESULTS

[CHAPTER 3 DESCRIPTION]

1. What is the demographic profile of the Persons with Physical Disability in Montalban Rizal, along:

1.1 Age

Table 01

Frequency Distribution of Respondents in terms of Age

Age	Frequency	Percentage
Gen Z	43	15.5
Millennials	30.7	30.7
Gen X	149	53.85
Total	277	100

Table 01 shows a total of 277 individuals were surveyed and classified into three generational cohorts: Gen Z, Millennials, and Gen X. Of these respondents, 43 (15.5%) belong to Gen Z, approximately 85 (30.7%) are Millennials, and the largest segment—149 respondents—fall into Gen X, accounting for 53.85% of the sample. This distribution reveals that more than half of the participants are from Gen X, nearly one-third are Millennials, and the smallest proportion is Gen Z, providing a clear picture of the age makeup of our respondents.

1.2 Sex

Table 02

Frequency Distribution of Respondents in terms of Sex

Age	Frequency	Percentage
Male	106	38.3
Female	171	61.7
Total	277	100

Table 02 shows that the same 277 individuals surveyed were classified by sex: 106 respondents (38.3%) are male and 171 respondents (61.7%) are female. This distribution indicates that females comprise nearly two-thirds of the sample, while males make up just over one-third, offering a clear view of the gender composition of our study participants.

1.3 Monthly Income

Table 03

Frequency Distribution of Respondents in terms of Monthly Income

Age	Frequency	Percentage
Poor	212	76.5
Low Income	51	18.4

Low Middle	11	4.0
Middle Middle	3	1.1
Total	277	100

Table 03 shows that the same 277 individuals surveyed were classified into four monthly income categories: 212 respondents (76.5%) fall into the Poor bracket, 51 respondents (18.4%) are in the Low Income group, 11 respondents (4.0%) belong to the Low Middle category, and just 3 respondents (1.1%) are in the Middle Middle bracket. This distribution reveals that the vast majority of participants report a Poor income level, with progressively smaller shares in higher income brackets, thus clearly illustrating the economic profile of our sample.

1.4 Type of Physical Disability

Table 04

Frequency Distribution of Respondents in terms of Monthly Income

Age	Frequency	Percentage
Speech Impairment	10	3.6
Hearing Impairment	21	7.6
Orthopedic Impairment	137	49.5
Visually Impairment	109	39.4
Total	277	100

Table 04 shows that the 277 respondents were classified by type of impairment: 10 individuals (3.6%) have a speech impairment, 21 respondents (7.6%) have a hearing impairment, the largest group—137 respondents—suffers from an orthopedic impairment (49.5%), and 109 participants (39.4%) report a visual impairment. This distribution highlights that nearly half of the sample experience orthopedic impairments, while visual impairments account for just under two-fifths of respondents, and speech and hearing impairments together comprise just over one-tenth, offering a clear overview of the disability profile within our study population.

2. What levels are the self-care practices of Persons with Physical Disability

in Montalban, Rizal, along:

2.1 Physical Well Being

Table 05

Computed Weighted Mean for the level of the self-care practices of Persons with Physical Disability in Montalban, Rizal in terms of Physical Well Being

Items	Weighted Mean	Verbal Interpretation
1. Eat a whole foods-based diet rich in colorful fruits and vegetables?	4.27	Regularly
2. Drink enough water?	4.53	Always
3. Exercise for more than 20 minutes?	2.61	Often
4. Wake feeling refreshed from sleep?	3.67	Regularly
5. Sleep at least 7 hours per night?	3.61	Regularly
6. Make time to relax or nap?	3.64	Regularly

7. Take time to breathe deeply throughout the day?	3.62	Regularly
8. Engage in stress-reducing activities (excluding TV or screen time)?	3.23	Often
9. Spend time in nature?	2.99	Often
10. Feel nourished, healthy, and strong?	3.95	Regularly
Total	3.61	Regularly (High)

Table 05 shows that practice of drinking enough water emerged as the most consistently performed behavior, with a weighted mean of 4.53 (“Always”), while structured exercise—for more than 20 minutes—was the least frequent self-care activity, at 2.61 (“Often”). Across all ten items, the overall weighted mean is 3.61, placing the group in the “Regularly” category and indicating a generally high level of engagement in physical-well-being practices.

Six behaviors were rated “Regularly,” reflecting strong routines around nutrition, rest, and restorative habits. Respondents reported eating a whole foods-based diet rich in colorful fruits and vegetables (4.27), waking feeling refreshed from sleep (3.67), and sleeping at least seven hours per night (3.61). They also made time to relax or nap (3.64), took moments for deep breathing throughout the day (3.62), and felt nourished, healthy, and strong (3.95)—all practices that support sustained energy and stress management.

Three activities fell into the “Often” category, signaling opportunities for growth. While respondents do engage in stress-reducing activities apart from screen time (3.23) and occasionally spend time in nature (2.99), these habits are not yet fully entrenched.

2.2 Emotional and Spiritual Well Being

Table 06

**Computed Weighted Mean for the level of the self-care practices of Persons
with Physical Disability in Montalban, Rizal in terms of Emotional and
Spiritual Well Being**

Items	Weighted Mean	Verbal Interpretation
1. Make time to participate in things you enjoy?	3.60	Regularly
2. Give and receive affection regularly?	3.74	Regularly
3. Feel understood and valued by those who are close to you?	3.87	Regularly
4. Feel gratitude on a daily basis?	3.82	Regularly
5. Find meaning in life even during difficult times?	3.99	Regularly
6. Take an interest in or find joy in the world around you?	3.71	Regularly
7. Have hope that things will get better?	3.96	Regularly
8. Express yourself creatively?	3.12	Often
9. Treat yourself with kindness?	4.01	Regularly
10. Remember to make your dreams and goals a priority?	3.91	Regularly
Total	3.77	Regularly (High)

Table 06 shows that “Treat yourself with kindness” is the most consistently practiced behavior, with a weighted mean of 4.01 (“Regularly”), while “Express yourself creatively” is the least frequent, at 3.12 (“Often”). Overall, the total weighted mean across all ten items is 3.77, placing the group firmly in the “Regularly” category and reflecting a high level of engagement in emotional and spiritual self-care.

Nine of the ten practices are rated “Regularly,” highlighting robust routines of emotional and spiritual well-being. Respondents make time to participate in things they enjoy (3.60), give and receive affection regularly (3.74), and feel understood and valued by those close to them (3.87). They also report feeling gratitude daily (3.82), finding meaning in life even during difficult times (3.99), taking interest in or finding joy in the world around them (3.71), having hope that things will get better (3.96), treating themselves with kindness (4.01), and remembering to make their dreams and goals a priority (3.91).

Only one practice falls into the “Often” category: “Express yourself creatively” (3.12). While respondents do engage in creative expression sometimes, it is not yet a regular habit.

2.3 Professional file, Work and, Career

Table 07

Computed Weighted Mean for the level of the self-care practices of Persons with Physical Disability in Montalban, Rizal in terms of Professional file, Work and, Career

Items	Weighted Mean	Verbal Interpretation
1. Hold a work position in an area of your interest?	2.74	Often

2. Work in a position that matches your professional goals?	2.69	Often
3. Find a sense of meaning and enjoyment in your work?	2.88	Often
4. Empathize and connect with customers, clients, and work colleagues?	2.88	Often
5. Have confidence in your ability to address challenges in your professional life?	2.87	Often
6. Feel supported at work or in your professional life?	3.07	Often
7. Have someone you can rely on if you need help or guidance?	3.34	Often
8. Set limits at work, whether it be with clients or tasks?	2.81	Often
9. Disengage and leave pressures behind at the end of the day?	2.70	Often
10. Take vacation or holiday breaks to allow for some down time?	2.88	Often
Total	2.89	Often (Average)

Table 07 shows that “Have someone you can rely on if you need help or guidance” is the most consistently practiced professional self-care behavior, with a weighted mean of 3.34 (“Often”), while “Work in a position that matches your professional goals” is the least frequent at 2.69 (“Often”). Overall, the group’s total weighted mean is 2.89, placing their engagement squarely in the “Often” category and indicating a moderate level of attention to work- and career-related well-being.

All ten practices fall into the “Often” category, reflecting steady but not yet ingrained routines. Alongside relying on a trusted colleague or mentor (3.34), respondents report feeling supported at work (3.07) and finding meaning and enjoyment in their roles (2.88). They also empathize and connect with clients and colleagues (2.88), hold positions in areas of interest (2.74), and express confidence in addressing professional challenges (2.87). At the lower end, they are

least likely to work in roles aligned with their goals (2.69), disengage from work pressures at the end of the day (2.70), set clear limits on tasks or clients (2.81), and take vacation or holiday breaks (2.88).

This uniform “Often” pattern indicates a foundational level of professional self-care but underscores considerable potential for enhancement.

2.4 Social Life, Family and Relationships

Table 08

Computed Weighted Mean for the level of the self-care practices of Persons with Physical Disability in Montalban, Rizal in terms of Social Life, Family and Relationships

Items	Weighted Mean	Verbal Interpretation
1. Have a dependable person who listens to you?	3.83	Regularly
2. Have supportive family and friends close by?	4.19	Regularly
3. Get enough social time with people who make you happy?	3.88	Regularly
4. Participate in group activities with people who share a common interest?	3.48	Often
5. Spend time with people who make you laugh?	3.77	Regularly
6. Feel like your close relationships are loving and supportive?	3.99	Regularly
7. Have the ability to comfortably say no?	3.52	Regularly
8. Do something fun with family or friends at least once a week?	3.64	Regularly
9. Feel like your personal life brings balance to your professional life?	3.24	Often
10. Feel comfortable asking for help when you need it?	3.71	Regularly
Total	3.73	Regularly

Table 08 shows that “Have supportive family and friends close by” is the most consistently practiced behavior, with a weighted mean of 4.19 (“Regularly”), while “Feel like your personal life brings balance to your professional life” is the least frequent at 3.24 (“Often”). Overall, the group’s total weighted mean is 3.73, placing their engagement firmly in the “Regularly” category and indicating a high level of attention to social, familial, and relational well-being.

Eight of the ten practices are rated “Regularly,” reflecting robust social support and meaningful connections. Respondents report having a dependable person who listens to them (3.83), getting enough social time with people who make them happy (3.88), spending time with those who make them laugh (3.77), feeling that their close relationships are loving and supportive (3.99), comfortably saying no when needed (3.52), doing something fun with family or friends at least once a week (3.64), and feeling comfortable asking for help (3.71). Two activities fall into the “Often” category—participating in group activities with shared interests (3.48) and achieving a balance between personal and professional life (3.24)—indicating areas where consistency could be strengthened.

This overall pattern underscores a solid foundation of social and relational self-care but also highlights specific opportunities for enhancement.

3. What is the correlation of profile to self-care practices of Persons with Physical Disability in Montalban Rizal?

3.1 Age

Table 09

Test of Correlation of profile to self-care practices of Persons with Physical Disability in Montalban Rizal in terms of Age

Self-Care Practice	Computed Value	P-Value	Remarks
Physical Well Being	0.078	0.197	Not Significant
Emotional and Spiritual Well Being	-0.092	0.125	Not Significant
Professional life, work, and career	-0.062	0.286	Not Significant
Social life, family, and relationships	0.043	0.474	Not Significant

Table 09 illustrates that age bears no statistically significant relationship with any of the four self-care domains among persons with physical disabilities in Montalban, Rizal. Correlation coefficients between age and physical well-being ($r = 0.078$, $p = 0.197$), emotional and spiritual well-being ($r = -0.092$, $p = 0.125$), professional life, work, and career ($r = -0.062$, $p = 0.286$), and social life, family, and relationships ($r = 0.043$, $p =$

0.474) are all close to zero and fail to reach the conventional threshold for significance ($p < 0.05$). These findings indicate that, within this sample, age does not meaningfully predict variation in self-care practice levels across any of the measured domains.

3.2 Sex

Table 10

Test of Correlation of profile to self-care practices of Persons with Physical Disability in Montalban Rizal in terms of Sex

Self-Care Practice	Computed Value	P-Value	Remarks
Physical Well Being	-0.038	0.525	Not Significant
Emotional and Spiritual Well Being	-0.105	0.082	Not Significant
Professional life, work, and career	-0.031	0.605	Not Significant
Social life, family, and relationships	-0.057	0.343	Not Significant

Table 10 reveals that sex does not significantly correlate with any domain of self-care practices among persons with physical disabilities in Montalban, Rizal. The

association between sex and physical well-being is negligible ($r = -0.038$, $p = 0.525$), as is its relationship with professional life, work, and career ($r = -0.031$, $p = 0.605$) and social life, family, and relationships ($r = -0.057$, $p = 0.343$). Although the correlation with emotional and spiritual well-being is marginally larger ($r = -0.105$), it remains non-significant ($p = 0.082$). Since all p-values exceed the 0.05 threshold, these results indicate that male and female respondents engage in self-care practices at statistically comparable levels across all measured domains.

3.3 Monthly Income

Table 11

Test of Correlation of profile to self-care practices of Persons with Physical Disability in Montalban Rizal in terms of Monthly Income

Self-Care Practice	Computed Value	P-Value	Remarks
Physical Well Being	-0.044	0.461	Not Significant
Emotional and Spiritual Well Being	-0.014	0.812	Not Significant
Professional file, work, and career	0.276	<0.001	Significant
Social life, family, and relationships	-0.055	0.365	Not Significant

Table 11 demonstrates that monthly income is significantly correlated with one domain of self-care practices among persons with physical disabilities in Montalban, Rizal. A positive moderate association exists between income and professional life, work, and career ($r = 0.276$, $p < 0.001$), indicating that respondents with higher monthly incomes tend to engage more in career-related self-care activities. In contrast, income shows no significant relationship with physical well-being ($r = -0.044$, $p = 0.461$), emotional and spiritual well-being ($r = -0.014$, $p = 0.812$), or social life, family, and relationships ($r = -0.055$, $p = 0.365$). These non-significant correlations suggest that variations in monthly income do not meaningfully influence self-care practices in those three domains.

Regression of Monthly Income to Professional life, work, and career

Table 12

Overall Model Test

R	R²	P-value	Strength	Remarks
0.240	0.057	<0.001	Weak	Significant

Table 12 presents the overall model test for the regression predicting the professional life, work, and career self-care score from monthly income. The model is statistically significant ($p < 0.001$), indicating that income does reliably predict engagement in career-related self-care activities. However, the strength of the relationship is weak: the multiple correlation coefficient is $R = 0.240$, and the model

explains only 5.7 % of the variance in professional self-care ($R^2 = 0.057$). In practical terms, while higher income is a significant predictor of greater investment in professional self-care, most of the variability in that domain remains unexplained by income alone.

Table 13

Monthly Income Regression Model

Monthly Income	Coefficient	P-Value	Remarks
Lower Middle - Poor	0.595	0.103	Not Significant
Low Income - Poor	0.677	<0.001	Significant
Middle Middle - Poor	0.986	0.150	Not Significant

Table 13 examines the effect of income categories (with “Poor” as the reference group) on professional life, work, and career self-care scores. Compared to the Poor group, only the Low Income category shows a statistically significant increase in professional self-care ($\beta = 0.677$, $p < 0.001$), indicating that those in the Low Income bracket engage more in career-related self-care than those classified as Poor. In contrast, the Lower Middle versus Poor contrast ($\beta = 0.595$, $p = 0.103$) and the Middle Middle versus Poor contrast ($\beta = 0.986$, $p = 0.150$) both fail to reach significance, suggesting no reliable difference in professional self-care between those income levels and the Poor reference group. These results imply that modest rises into the Low Income category are

associated with enhanced professional self-care, but further income increases beyond that threshold do not confer additional significant benefits in this domain.

3.4 Type of Disability

Table 14

Test of Correlation of profile to self-care practices of Persons with Physical Disability in Montalban Rizal in terms of Type of Disability

Self-Care Practice	Computed Value	P-Value	Remarks
Physical Well Being	-0.004	0.953	Not Significant
Emotional and Spiritual Well Being	0.103	0.086	Not Significant
Professional life, work, and career	0.039	0.517	Not Significant
Social life, family, and relationships	0.040	0.508	Not Significant

Table 14 reveals that the type of disability does not significantly correlate with any domain of self-care practices among persons with physical disabilities in Montalban, Rizal. The relationship between type of disability and physical well-being is virtually nonexistent ($r = -0.004$, $p = 0.953$), indicating no meaningful association. Similarly,

correlations with professional life, work, and career ($r = 0.039$, $p = 0.517$), as well as social life, family, and relationships ($r = 0.040$, $p = 0.508$), are weak and statistically insignificant. Although the correlation with emotional and spiritual well-being ($r = 0.103$) is slightly higher, it still fails to reach significance ($p = 0.086$). Since all p -values exceed the 0.05 threshold, the results suggest that variations in the type of disability do not significantly influence the self-care practices of respondents across the measured domains.