

NCMHCE Quick Reference

92 Diagnoses · ICD-10 Codes · First-Line Treatments

PassReady Prep · GA Integrated Therapeutic Perspectives LLC

Depressive

Major Depressive Disorder, Moderate (F33.1)

First-line: CBT or IPT + consider SSRI

Major Depressive Disorder, Severe (F33.2)

First-line: SSRI/SNRI + psychotherapy; assess suicide risk

Persistent Depressive Disorder (F34.1)

First-line: CBT + SSRI (combination superior)

Premenstrual Dysphoric Disorder (F32.81)

First-line: SSRI (continuous or luteal-phase); CBT

Disruptive Mood Dysregulation Disorder (F34.81)

First-line: Parent training + CBT; stimulant if comorbid ADHD

Major Depressive Disorder, with Peripartum Onset (F53.0)

First-line: IPT or CBT preferred; SSRI if moderate-severe

Anxiety

Generalized Anxiety Disorder (F41.1)

First-line: CBT with relaxation training; SSRI/SNRI

Panic Disorder (F41.0)

First-line: CBT with interoceptive exposure

Social Anxiety Disorder (F40.10)

First-line: CBT with in vivo exposure; SSRI

Specific Phobia (F40.218)

First-line: In vivo graduated exposure therapy

Agoraphobia (F40.00)

First-line: CBT with graduated situational exposure

Separation Anxiety Disorder (F93.0)

First-line: CBT with graduated exposure + parent training

Selective Mutism (F94.0)

First-line: Behavioral intervention with graduated exposure

Trauma

Posttraumatic Stress Disorder (F43.10)

First-line: CPT, PE, or EMDR (trauma-focused CBT)

Acute Stress Disorder (F43.0)

First-line: Brief trauma-focused CBT

Adjustment Disorder, with Mixed Anxiety and Depressed Mood (F43.23)

First-line: Supportive counseling + CBT; time-limited

Adjustment Disorder, with Depressed Mood (F43.21)

First-line: Brief supportive therapy; CBT; address stressor

Reactive Attachment Disorder (F94.1)

First-line: Attachment-focused therapy with stable caregiver

Prolonged Grief Disorder (F43.81)

First-line: Complicated grief treatment (CGT); targeted CBT

Substance

Alcohol Use Disorder, Moderate (F10.20)

First-line: MI + CBT/12-step; naltrexone or acamprosate

Alcohol Use Disorder, Severe (F10.20)

First-line: Medically supervised detox + MAT + counseling

Alcohol Withdrawal (F10.230)

First-line: Medical management (benzodiazepines); CIWA monitoring

Opioid Use Disorder, Severe (F11.20)

First-line: MAT (buprenorphine/methadone) + behavioral counseling

Opioid Withdrawal (F11.23)

First-line: MAT initiation; symptom management

Stimulant Use Disorder (Cocaine) (F14.20)

First-line: CBT + CM (contingency management); no approved medication

Stimulant Intoxication (F14.129)

First-line: Consult clinical practice guidelines

Cannabis Use Disorder, Moderate (F12.20)

First-line: MI + CBT; no FDA-approved medication

Sedative Use Disorder (F13.20)

First-line: Medically supervised taper; CBT

Tobacco Use Disorder, Moderate (F17.200)

First-line: Nicotine replacement + behavioral counseling

Gambling Disorder (F63.0)

First-line: CBT + GA support; address cognitive distortions

Personality

Borderline Personality Disorder (F60.3)

First-line: DBT (Dialectical Behavior Therapy)

Narcissistic Personality Disorder (F60.81)

First-line: Schema therapy or transference-focused therapy

Antisocial Personality Disorder (F60.2)

First-line: CBT for criminogenic thinking; limited evidence

Avoidant Personality Disorder (F60.6)

First-line: CBT with graduated exposure + schema therapy

Dependent Personality Disorder (F60.7)

First-line: CBT building autonomy + assertiveness training

Obsessive-Compulsive Personality Disorder (F60.5)

First-line: CBT targeting rigidity; psychodynamic

Schizotypal Personality Disorder (F21)

First-line: CBT for social skills; low-dose antipsychotic if severe

Paranoid Personality Disorder (F60.0)

First-line: CBT building trust; avoid confrontation

Schizoid Personality Disorder (F60.1)

First-line: Supportive therapy; gradual social engagement

Histrionic Personality Disorder (F60.4)

First-line: Psychodynamic or CBT; address attention-seeking patterns

Bipolar

Bipolar I Disorder, Current Episode Manic (F31.13)

First-line: Mood stabilizer + atypical antipsychotic

Bipolar I Disorder, Current Episode Depressed (F31.31)

First-line: Quetiapine or lurasidone; careful with antidepressants

Bipolar II Disorder (F31.81)

First-line: Mood stabilizer + IPSRT or CBT; lamotrigine

Cyclothymic Disorder (F34.0)

First-line: Mood stabilizer (lower dose); psychoeducation + CBT

Neurodevelopmental

Attention-Deficit/Hyperactivity Disorder, Combined (F90.2)

First-line: Stimulant + behavioral strategies + accommodations

Attention-Deficit/Hyperactivity Disorder, Inattentive (F90.0)

First-line: Stimulant or atomoxetine + organizational skills

Autism Spectrum Disorder (F84.0)

First-line: ABA + social skills training + family support

Specific Learning Disorder (F81.0)

First-line: Specialized educational intervention; IEP/504

Intellectual Developmental Disorder, Mild (F70)

First-line: Adaptive skills training; supported living/employment

Tourette's Disorder (F95.2)

First-line: CBIT (behavioral); medication if severe (antipsychotics)

Neurocognitive

Delirium (F05)

First-line: Medical emergency — treat underlying cause

Major Neurocognitive Disorder due to Alzheimer's Disease (F02.80)

First-line: Cholinesterase inhibitors; caregiver support; safety planning

Major Neurocognitive Disorder due to Vascular Disease (F01.50)

First-line: Manage cardiovascular risk factors; cognitive rehab

Mild Neurocognitive Disorder (G31.84)

First-line: Cognitive rehab; lifestyle modifications; monitoring

OCD-Related

Obsessive-Compulsive Disorder (F42.2)

First-line: ERP + SSRI

Body Dysmorphic Disorder (F45.22)

First-line: CBT-BDD + SSRI

Hoarding Disorder (F42.3)

First-line: Specialized CBT for hoarding

Trichotillomania (F63.3)

First-line: Habit Reversal Training + stimulus control

Excoriation Disorder (L98.1)

First-line: Habit Reversal Training; SSRI

Psychotic

Schizophrenia (F20.9)

First-line: Antipsychotic + CBTp + psychosocial rehab

Schizoaffective Disorder, Bipolar Type (F25.0)

First-line: Antipsychotic + mood stabilizer + CBTp

Schizophreniform Disorder (F20.81)

First-line: Antipsychotic; monitor for progression to schizophrenia

Brief Psychotic Disorder (F23)

First-line: Short-term antipsychotic; resolves <1 month

Delusional Disorder (F22)

First-line: Antipsychotic (limited response); CBT for alliance

Eating

Anorexia Nervosa, Restricting Type (F50.01)

First-line: FBT (adolescents) or CBT-E; medical monitoring

Bulimia Nervosa (F50.2)

First-line: CBT-E; SSRI (fluoxetine)

Binge-Eating Disorder (F50.81)

First-line: CBT-E or IPT; lisdexamfetamine

Avoidant/Restrictive Food Intake Disorder (F50.82)

First-line: Gradual food exposure; family-based; nutritional support

Crisis

Suicidal Behavior / Acute Risk (R45.851)

First-line: C-SSRS + safety plan + means restriction + level of care

Nonsuicidal Self-Injury (R45.88)

First-line: DBT skills + functional assessment

Acute Crisis / Decompensation (F43.0)

First-line: Stabilize + assess safety + level of care determination

Homicidal Ideation / Threat Assessment (R45.850)

First-line: Duty to warn/protect + safety assessment + documentation

Disruptive

Oppositional Defiant Disorder (F91.3)

First-line: Parent Management Training + child CBT

Conduct Disorder, Adolescent-Onset Type (F91.2)

First-line: MST or Functional Family Therapy

Intermittent Explosive Disorder (F63.81)

First-line: CBT anger management; SSRI

Somatic

Somatic Symptom Disorder (F45.1)

First-line: CBT for health anxiety; scheduled visits

Illness Anxiety Disorder (F45.21)

First-line: CBT targeting health preoccupation

Functional Neurological Symptom Disorder (F44.4)

First-line: CBT + physical therapy; validate symptoms

Factitious Disorder (F68.10)

First-line: Gentle confrontation; address underlying needs

Dissociative

Dissociative Identity Disorder (F44.81)

First-line: Phase-oriented: stabilize, process, integrate

Dissociative Amnesia (F44.0)

First-line: Psychotherapy for underlying trauma

Depersonalization/Derealization Disorder (F48.1)

First-line: CBT + grounding; treat comorbid anxiety/depression

Sleep

Insomnia Disorder, Chronic (F51.01)

First-line: CBT-I (sleep restriction + stimulus control)

Nightmare Disorder (F51.5)

First-line: Image Rehearsal Therapy; prazosin if trauma-related

Hypersomnolence Disorder (F51.11)

First-line: Behavioral strategies; modafinil; treat underlying cause

Sexual-Gender

Gender Dysphoria in Adolescents/Adults (F64.0)

First-line: Affirmative therapy + medical referral

Female Sexual Interest/Arousal Disorder (F52.22)

First-line: CBT + sensate focus; address relationship factors

Erectile Disorder (F52.21)

First-line: CBT + behavioral techniques; medical eval

Ethics

Confidentiality / Duty-to-Warn Dilemma (Z65.3)

First-line: Ethical decision-making model; consult; document

Boundary / Dual-Relationship Dilemma (Z65.8)

First-line: Evaluate risk; consult; document; refer if needed

Mandated Reporting Decision (Z65.8)

First-line: Report per state law; document; inform client if safe

Informed Consent / Competence Dilemma (Z65.8)

First-line: Review capacity; document decision process