Text mining for exploration of COVID-19 severity factors

Khang Duy LAI - Mariia KLIMINA

University Paris Cité

UFR des Sciences Fondamentales et Biomédicales

May 12, 2022

Text mining for exploration of COVID-19 severity factors

Text mining for exploration of COVID-19 severity factors

Khang Duy LAI - Mariia KLIMINA
University Paris Clai
UFR des Sciences Fondamentales et Biomédicales
May 12, 2022

- 1 Introduction
- 2 State of the art
- 3 Data exploration
- 4 Data preprocessing
- **5** Data processing
- 6 Result

Text mining for exploration of COVID-19 severity factors $\,$

● Introduction

● State of the art

● Data exploration

● Data preprocessing

● Data processing

● Result

2022-05-12

Introduction

COVID-19 is the disease caused by the Sar-COV-2 virus that originated in China at the end of the year 2019. Over the time, studies have shown that there is some form of background diseases and risk factors that can hugely affect the severity cases rate of COVID-19. This project will apply NLP and text mining methods in order to explore the CORD-19 dataset and extract background diseases and risk factors.

Text mining for exploration of COVID-19 severity factors 2022-05 -Introduction

-Introduction

Introduction

COVID-19 is the disease caused by the Sar-COV-2 virus that originated in China at the end of the year 2019. Over the time, studies have shown that there is some form of background diseases and risk factors that can hugely affect the severity cases rate of COVID-19. This project will apply NLP and text mining methods in order to evolute the CORD-19 dataset and extract

duction State of the art Data exploration Data preprocessing Data processing

State of the art

In this project we used multiple state of the art NLP and Data Science libraries.

- Numpy, Pandas: Formatting the data and the calculations.
- Matplotlib: Library for drawing the charts and figures.
- Scikit-learn: LDA and T-SNE models.
- Spacy, Gensim, and NLTK: Important NLP libraries.
- Scispacy: NER, Spacy models for science papers.
- Bokeh: A library for visualising interacted charts.

Text mining for exploration of COVID-19 severity factors

State of the art

State of the art


In this project we used multiple state of the art NLP and Dat Science libraries.

- Numpy, Pandas: Formatting the data and the calculation
 Matplotlife: Library for drawing the charts and figures.
 Scikit-learn: LDA and T-SNE models.
- Spacy, Gensim, and NLTK: Important NLP librarie
 Scispacy: NER, Spacy models for science papers.
- -State of the art

Data exploration

2022-05

Data exploration



Text mining for exploration of COVID-19 severity factors Data exploration

-Data exploration



Data exploration

2022-05

General Information

The metadata consist of more than one millions articles.




Figure 2: Overlook of the meta data of the dataset

Text mining for exploration of COVID-19 severity factors Data exploration General Information

General Information Figure 2: Overlook of the meta data of the datase Data exploration

General Information

Columns in the metadata

```
['cord uid', 'sha', 'source x', 'title', 'doi',
 'pmcid', 'pubmed id', 'license', 'abstract',
 'publish time', 'authors', 'journal',
 'mag id', 'who covidence id', 'arxiv id',
 'pdf_json_files', 'pmc_json_files', 'url', 's2_id']
```

Text mining for exploration of COVID-19 severity factors -Data exploration

General Information

General Information

Introduction State of the art Data exploration Data prep

preprocessing Data pro

ta processing


ult

2022-05

Language status

As can be observed on a graph most of the papers are written in english. However, there were some exceptions.

During this part, we deleted all non-english articles by using languetect library.



Text mining for exploration of COVID-19 severity factors

-Data exploration

Language status

Language status

As can be observed on a graph monto of the papers are written in english. However, three were some exceptions.

During this part, we deleted all non-english articles by using langietest library.

Data preprocessing

- Converting JSON format into DataFrame format.
- Removing all non-english paper.
- Removing special characters
- Removing numbers
- Tokenizing.
- Removing stopwords.
- Stemming.
- Lemmatisation.

Text mining for exploration of COVID-19 severity
factors
Data preprocessing
Data preprocessing

Data preprocessing

- Converting JSON format into DataFrame format
 Removing all non-english paper.
- Removing special characters
- Removing numbers
 Tokenizing.
- Removing stopwords.
 Stemming.
 Lemmatisation.

troduction State of the art \Box

ta exploration

Data preprocessing Da

ta processing

sult

Text mining for exploration of COVID-19 severity

factors

Data preprocessing

LData preprocessing

Data preprocessing

Converting JSON format into DataFrame format.

- Merge body text into the same dataframe with metadata
- Add column to define language of paper
- Remove unecessary columns.

Remove

- Filter out only paper with English
- Using regex to remove special characters, numbers.
- Remove stopwords

State of the art Data exploration Data preprocessing Data processing Res

Data preprocessing

Stemming and Lemmatisation

- Lowers inflection in words to their root forms
- (connections, connected, connects, is connect)
- Using NLTK library

Text mining for exploration of COVID-19 severity factors

Data preprocessing

Data preprocessing



Data preprocessing

2-02-

202

Data preprocessing

[35]: 'quarantine isolation main containment strategy intended help protect public preventing spread contagious disease strategy primarily refer r estriction movement limitation personal contact quarantine definition person exposed disease isolation contagious person require separation person infected finding previous research pointed increased risk negative psychological outcome depression anxiety isolation quarantined per son equally heightened risk adverse mental health outcome rapid review brook reported increased negative psychological outcome including pos ttraumatic stress symptom confusion anger person quarantine author concluded important stressor longer quarantine electronic supplementary m aterial online version article doiorgs x contains supplementary material available authorized user duration infection fear frustration bored om inadequate supply inadequate information financial loss stigma finding suggest containment strategy quarantine isolation negative impact psychological outcome related broad spectrum psychosocial stressor need investigation mental health problem associated containment strategy highlighted rising implementation quarantine isolation worldwide currently ongoing covid pandemic unprecedented number people worldwide affe cted quarantine isolation identification individual elevated risk adverse mental health effect mandatory suggested vulnerable population ris k negative psychological outcome implementation containment strategy eg person mental illness low income lack social network particular grea ter risk quarantine isolation world health organization included covid list disease pathogen prioritized research development rd public heal th emergency context pose greatest public health risk epidemic potential insufficient countermeasure established containment strategy main of ountermeasure context systematic investigation evidence concerning psychological effect urgently need single study review suggest increased risk negative psychological outcome person quarantine isolation presented partially contradicting result furthermore prevalence estimate poi nt elevated level adverse outcome quarantined isolated population validity finding limited underlying uncontrolled study design conducted sy stematic literature review metaanalysis mental health effect quarantine isolation based controlled primary study data best knowledge metaana lysis including quarantine isolation exists date systematic literature review metaanalysis protocol project published prospero prospero regi strationno crd method followed quideline cochrane collaboration conduction systematic review searched pubmed psycinfo embase database study restriction beginning searched time period april assessing rate psychological effect guarantinedisolated person compared nonggrantinednonis olated person search entry described online supplement supplement database search entry broad specific search term combined increase likelih ood detecting eligible study research aim specific search term included list disease pathogen prioritized research development rd public hea Ith emergency context world health organization covid additional record identified manual search reference included study included language restriction translation native speaker acquired test eligibility criterion article language english study author contacted case missing data search carried endnote x clarivate analytics philadelphia usa trial considered appropriate test hypothesis included met following criterion observation person quarantine isolation described second quantitative assessment psychological outcome parameter performed comparators perso n quarantine isolation fourth data calculation effect size corresponding measure dispersion provided study observing psychological outcome p arameter qualitative assessment excluded study excluded focused specific subpopulation primary infection controlassociation isolated person prison study assessing correlation mental health outcome varying duration quarantine isolation excluded quantitative synthesis reported qual itative synthesis determinant entire literature search study screening carried independently reviewer f jvb consensus unclear case reached d iscussion additional member reviewing team lb jh testing eligibility criterion study selection classification coding data predefined excel s preadsheet microsoft excel mac version microsoft corporation usa followed recommendation cochrane collaboration handbook performed independe ntly reviewer lb jh reviewer jh lb independently extracted data characteristic study study sample quantitative data severity mean score freq uency incidence prevalence mental health outcome group comparison group eg relative risk odds ratio result determinant testing reported reac h statistical significance original study multiple measure outcome reported extracted data following hierarchy continuous measure mean score categorical measure highest cutoff defined author original study ie severe manifestation disorder risk bias study classified independently eviewer lb jh according newcastleottawa scale no recommended cochrane handbook table summary assessment study classified holding low unknown high risk bias taking account bias main domain selection comparability exposureoutcome disagreement resolved consensus additional review aut hor calculated standardized mean difference smd confidence interval ci outcome measure primary study respective measure dispersion available calculated ci p value recommended cochrane handbook stratified predefined mental health outcome effect size comparison quarantinedisolated n onguarantinedisolated group summarized forest plot table quantitative synthesis result possible heterogeneity included study methodology pop

Khang Duy LAI - Mariia KLIMINA

Université Paris Cité

ulation outcome restricted quantitative synthesis predefined outcome primary study provided data categorical outcome based validated diagnos

Text mining for exploration of COVID-19 severity factors Data preprocessing

-Data preprocessing

Data preprocessing

State of the art Data exploration Data preprocessing Data processing Result 0000 0000 0000

Data processing

- Data selection
 - Selecting articles with risk factors and severity key-words.
 - Clustering using Latent Dirichlet Allocation.
- NER (Named-entity recognition).

Text mining for exploration of COVID-19 severity
factors
Data processing
Data processing

Data processing

• Data selection
• Data selection
• Data selection
• Data selection with risk factors and neverly lay-words
• Cherters using Latent Dirichle Allocation.
• NER (Named-earlity recognition).

te of the art Data exploration Data preprocessing Data

Data processing

sult

Risk factors and severity paper filtering

- Create a dictionary of key words related to risk factors and severity.
- Filter out only the paper that contain words in the dictionary.

Text mining for exploration of COVID-19 severity factors

Data processing


☐Risk factors and severity paper filtering

Risk factors and severity paper filtering

Create a dictionary of key words related to risk factors an severity.
 Filter out only the paper that contain words in the

Data processing

Risk factors and severity paper filtering



Text mining for exploration of COVID-19 severity factors -Data processing

Risk factors and severity paper filtering



Data processing 00000000

LDA

In natural language processing, the latent Dirichlet allocation (LDA) is a generative statistical model that allows sets of observations to be explained by unobserved groups that explain why some parts of the data are similar.

The LDA algorithm structure:

- Providing to an algorithm a certain number of topics.
- The algorithm is assigning every word to a temporary topic.
- The algorithm is checking and updating topic assignments.

Text mining for exploration of COVID-19 severity factors 2022-05 Data processing

LDA

In natural language processing, the latent Dirichlet allocation (LDA) is a generative statistical model that allows sets of observations to be explained by unobserved groups that explain why some parts of the data are similar.


The LDA algorithm structure

LDA

- . Providing to an algorithm a certain number of tonics The algorithm is assigning every word to a temporary topic.
- The algorithm is checking and updating topic assignments.

ntroduction State of the art Data exploration Data preprocessing Data processing I

LDA



The coherence score measures how similar these words are to each other. The higher the coherence score is, the more suitable the topic number should be.

Université Paris Cité

Khang Duy LAI - Mariia KLIMINA

Text mining for exploration of COVID-19 severity factors

17 of 24

Text mining for exploration of COVID-19 severity factors

Data processing


 \sqsubseteq_{LDA}

2022-05



roduction State of the art Data exploration Data preprocessing Occoo Oc

LDA



Text mining for exploration of COVID-19 severity factors $\begin{tabular}{c} \begin{tabular}{c} \begin{tabu$



 $\sqsubseteq_{\mathrm{LDA}}$

2022-05-12

Data processing

NER

Named entity recognition (NER) is probably the extraction-method that seeks to locate and classify named entities in text into pre-defined categories such as the names of persons, organizations, locations, expressions of times, quantities, monetary values, percentages. In our case it's biomedical entities: diseases, chemicals etc.

We used Scispacy library with different SpaCy models for biomedical text processing.

Text mining for exploration of COVID-19 severity factors 2022-05 Data processing

└─NER

biomedical text processing

NER

Named entity recognition (NER) is probably the extraction-method that seeks to locate and classify names monetary values, percentages. In our case it's biomedical entities We used Scispacy library with different SpaCy models for

NER

Scispacy

- Science pretrain model
- en_ner_bc5cdr_md for biology, mark diseases and chemicals.

Text mining for exploration of COVID-19 severity factors

Data processing

NER



Result •000

Result

Example of table of result

Text mining for exploration of COVID-19 severity factors

Ü	chronic obstructive pulmonary disease copd	DISEASI
$\frac{-}{1}$	death	DISEASI
3	copd	DISEASI
9	dyspnea	DISEASI
10	cough	DISEASI
11	copd pulmonary function	DISEASI
13	respiratory tract infection	DISEASI
14	chronic unstable disease system malignancy	DISEASI
19	obstructive pulmonary disease	DISEASI
21	copd airflow	DISEASI
25	hypertension	DISEASI
26	atherosclerotic heart disease	DISEASI
Khang Duv	LAI - Marija KLIMINA	Université Paris Cité

Text mining for exploration of COVID-19 severity factors -Result └─Result

2022-05-12

21 of 24

Result

Example of table of result

1 death 3 copd 9 dyspnea DISEASI DISEASI 10 cough DISEASI 11 copd pulmonary function DISEASI 13 respiratory tract infection DISEASI 14 chronic unstable disease system malignancy

19 obstructive pulmonary disease 21 copd airflow 26 atherosclerotic heart disease

25 hypertension 27 bronchiectasis

DISEASI DISEASI DISEASI DISEASI DISEASI

Conclusion

- Basic diseases are filtered out
- The method have not sorted if the disease is a covid symtoms or related to the progress of severity cases.

Text mining for exploration of COVID-19 severity factors $\begin{array}{c} \begin{array}{c} \\ \\ \\ \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \\ \end{array} \\ \begin{array}{c} \\ \\ \\ \\ \end{array} \\ \\ \begin{array}{c} \\ \\ \\ \\ \\ \end{array} \\ \begin{array}{c} \\\\ \\ \\ \end{array} \\ \\ \\ \end{array} \\ \begin{array}{c} \\\\ \\\\ \\ \end{array} \\ \begin{array}{c} \\\\ \\\\ \\\\ \end{array} \\ \begin{array}{c} \\\\\\\\\\\\\\\\\\\\\\ \end{array} \\ \\ \end{array} \\$

Future improvement

- Create a knowledge graph
- Calculate the severity rate

Text mining for exploration of COVID-19 severity
factors
Result
Future improvement

Future improvement

Thank you

Thank you for your listening.

Text mining for exploration of COVID-19 severity
factors
Result
Thank you

Khang Duy LAI - Mariia KLIMINA