CNS Vital Signs

CNS Vital Signs Report	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
CNSVS Duration: 48:37 (min:secs)	CNSVS Online Version 2.0.6

Patient Profile	**Percentile Range**				**> 74**	**25 - 74**	**9 - 24**	**2 - 8**	**< 2**
	Standard Score Range				**> 109**	**90 - 109**	**80 - 89**	**70 - 79**	**< 70**
Domain Scores	**Patient Score**	**Standard Score**	**Percentile**	**VI****	**Above**	**Average**	**Low Average**	**Low**	**Very Low**
Neurocognition Index (NCI)	**NA**	**69**	**2**	**Yes**					**X**
Composite Memory	**88**	**81**	**10**	**Yes**			**X**		
Verbal Memory	**42**	**68**	**2**	**Yes**					**X**
Visual Memory	**46**	**103**	**58**	**Yes**		**X**			
Psychomotor Speed	**132**	**74**	**4**	**Yes**				**X**	
Reaction Time*	**950**	**56**	**1**	**Yes**					**X**
Complex Attention*	**15**	**73**	**4**	**Yes**				**X**	
Cognitive Flexibility	**17**	**63**	**1**	**Yes**					**X**
Processing Speed	**36**	**76**	**5**	**Yes**				**X**	
Executive Function	**18**	**63**	**1**	**Yes**					**X**
Social Acuity	**9**	**104**	**61**	**Yes**		**X**			
Reasoning	**13**	**126**	**96**	**Yes**	**X**				
Working Memory	**10**	**104**	**61**	**Yes**		**X**			
Sustained Attention	**31**	**108**	**70**	**Yes**		**X**			
Simple Attention	**37**	**61**	**1**	**Yes**					**X**
Motor Speed	**96**	**82**	**12**	**Yes**			**X**		

Domain Dashboard: Above average domain scores indicate a standard score (SS) greater than 109 or a Percentile Rank (PR) greater than 74, indicating a high functioning test subject. Average is a SS 90-109 or PR 25-74, indicating normal function. Low Average is a SS 80-89 or PR 9-24 indicating a slight deficit or impairment. Below Average is a SS 70-79 or PR 2-8, indicating a moderate level of deficit or impairment. Very Low is a SS less than 70 or a PR less than 2, indicating a deficit and impairment. Reaction times are in milliseconds. An * denotes that "lower is better", otherwise higher scores are better. Subject Scores are raw scores calculations generated from data values of the individual subtests.

VI** - Validity Indicator: Denotes a guideline for representing the possibility of an invalid test or domain score. "No" means a clinician should evaluate whether or not the test subject understood the test, put forth their best effort, or has a clinical condition requiring further evaluation.

Verbal Memory Test (VBM)	**Score**	**Standard**	**Percentile**	
Correct Hits - Immediate	7	65	1	Verbal Memory test: Subjects have to remember 15 words and recognize them in a field of 15 distractors. The test is repeated at the end of the battery. The VBM test measures how well a subject can recognize, remember, and retrieve words e.g. exploit or attend literal representations or attribute. "Correct Hits" refers to the number of target words recognized. Low scores indicate verbal memory impairment.
Correct Passes - Immediate	15	110	75	
Correct Hits - Delay	8	82	12	
Correct Passes - Delay	12	69	2	
Visual Memory Test (VSM)	**Score**	**Standard**	**Percentile**	
Correct Hits - Immediate	11	93	32	Visual Memory test: Subjects have to remember 15 geometric figures, and recognize them in a field of 15 distractors. The test is repeated at the end of the battery. The VSM test measures how well a subject can recognize, remember, and retrieve geometric figures e.g. exploit or attend symbolic or spatial representations. "Correct Hits" refers to the number of target figures recognized. Low scores indicate visual memory impairment.
Correct Passes - Immediate	11	98	45	
Correct Hits - Delay	12	106	66	
Correct Passes - Delay	12	107	68	
Finger Tapping Test (FTT)	**Score**	**Standard**	**Percentile**	
Right Taps Average	50	86	18	The FTT is a test of motor speed and fine motor control ability. There are three rounds of tapping with each hand. The FTT test measures the speed and the number of finger-taps with each hand. Low scores indicate motor slowing. Speed of manual motor activity varies with handedness. Most people are faster with their preferred hand but not always.
Left Taps Average	46	81	10	

CNS Vital Signs

CNS Vital Signs Report	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
CNSVS Duration: 48:37 (min:secs)	CNSVS Online Version 2.0.6

Symbol Digit Coding (SDC)	Score	Standard	Percentile	
Correct Responses	36	76	5	The SDC test measures speed of processing and draw upon several cognitive processes simultaneously, such as visual scanning, visual perception, visual memory, and motor functions. Errors may be due to impulsive responding, misperception, or confusion.
Errors*	0	110	75	
Stroop Test (ST)	**Score**	**Standard**	**Percentile**	
Simple Reaction Time*	590	54	1	The ST measures simple and complex reaction time, inhibition / disinhibition, mental flexibility or directed attention. The ST helps assess how well a subject is able to adapt to rapidly changing and increasingly complex set of directions. Prolonged reaction times indicate cognitive slowing / impairment. Errors may be due to impulsive responding, misperception, or confusion.
Complex Reaction Time Correct*	818	66	1	
Stroop Reaction Time Correct*	1081	57	1	
Stroop Commission Errors*	1	98	45	
Shifting Attention Test (SAT)	**Score**	**Standard**	**Percentile**	
Correct Responses	29	59	1	The SAT measures executive function or how well a subject recognizes set shifting (mental flexibility) and abstraction (rules, categories) and manages multiple tasks simultaneously. Subjects have to adjust their responses to randomly changing rules. The best scores are high correct responses, few errors and a short reaction time. Normal subjects may be slow but accurate, or fast but not so accurate. Attention deficit may be apparent.
Errors*	11	80	9	
Correct Reaction Time*	1536	60	1	
Continuous Performance Test (CPT)	**Score**	**Standard**	**Percentile**	
Correct Responses	38	24	1	The CPT measures sustained attention or vigilance and choice reaction time. Most normal subjects obtain near-perfect scores on this test. A long response time may suggest cognitive slowing and/or impairment. More than 2 errors (total) may be clinically significant. More than 4 errors (total) indicate attentional dysfunction.
Omission Errors*	2	24	1	
Commission Errors*	1	88	21	
Choice Reaction Time Correct*	698	31	1	
Perception Of Emotions Test (POET)	**Score**	**Standard**	**Percentile**	
Correct Responses	11	95	37	The POET measures how well a subject can perceive and identify specific emotions. The POET is an identification test of the positive emotions "happy" and "calm" and the negative emotions "angry" and "sad". "Social cognition" or "emotional acuity" has been defined as "the way in which people make sense of other people and themselves". It is the ability to perceive and understand social information. The reaction times in POET are much longer than in the other tests, indicating the complexity of central processes governing emotional acuity.
Average Correct Reaction Time*	1467	60	1	
Omission Errors*	1	95	37	
Commission Errors*	2	107	68	
Positive Emotions				
Correct Hits	6	106	66	
Reaction Time*	1422	54	1	
Negative Emotions				
Correct Hits	5	88	21	
Reaction Time*	1520	72	3	
Reasoning Test (RT)	**Score**	**Standard**	**Percentile**	
Correct Responses	13	123	94	The NVRT measures how well a subject can perceive and understand the meaning of visual or abstract information and recognizing relationships between visual-abstract concepts. The NVRT is comprised of 15 matrices, or visual analogies. The matrices are progressively more difficult. Each is presented for 14.5 seconds. Non-verbal or visual-abstract reasoning is the process of perceiving issues and reaching conclusions through the use of symbols or generalizations rather than concrete factual information.
Average Correct Reaction Time*	6600	78	7	
Commission Errors*	0	126	96	
Omission Errors*	2	90	25	

CNS Vital Signs

CNS Vital Signs Report	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
CNSVS Duration: 48:37 (min:secs)	CNSVS Online Version 2.0.6

Four Part Continuous Performance Test	Score	Standard	Percentile	
Part 1				The FPCPT test is a four part test that measures a subject's working memory and sustained attention. The FPCPT is a four part test: PART ONE - is a simple reaction time test, the subject must press the space bar when any stimulus is presented; PART TWO - is a variant of the continuous performance test, the subject is asked to respond to one stimulus, but not to any others. Discrimination is required, so the reaction times that are generated are "choice reaction times". PART THREE - is a "one back" CPT. The subject has to respond to a figure only if the figure immediately preceding was the same. PART FOUR - is a "two-back" CPT. It is a difficult task and is used to measure working memory. Parts two, three, and four of the tests are used to calculate sustained attention domain.
Average Correct Reaction Time*	708	50	1	
Part 2				
Correct Responses	6	103	58	
Average Correct Reaction Time*	802	-1	1	
Incorrect Responses*	0	104	61	
Average Incorrect Reaction Time*	0			
Omission Errors*	0	103	58	
Part 3				
Correct Responses	15	108	70	
Average Correct Reaction Time*	1020	43	1	
Incorrect Responses*	0	104	61	
Average Incorrect Reaction Time*	0			
Omission Errors*	1	108	70	
Part 4				
Correct Responses	10	99	47	
Average Correct Reaction Time*	956	76	5	
Incorrect Responses*	0	110	75	
Average Incorrect Reaction Time*	0			
Omission Errors*	6	99	47	

Generalized Anxiety Disorder 7-item (GAD-7) Scale	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 0:55 (min:secs)	CNSVS Online Version 2.0.6

GAD-7 Anxiety Severity:

This is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of "Not at all", "Several days", "More than half the days", and "Nearly every day" respectively. GAD-7 total score for the seven items ranges from 0 to 21.

Interpretation of Total Score:

0-4 None-Minimal anxiety

5-9 Mild anxiety

10-14 Moderate anxiety

15-21 Severe anxiety

1	Feeling nervous, anxious or on edge	3 - Nearly every day
2	Not being able to stop or control worrying	3 - Nearly every day
3	Worrying too much about different things	3 - Nearly every day
4	Trouble relaxing	3 - Nearly every day
5	Being so restless that it is hard to sit still	3 - Nearly every day
6	Becoming easily annoyed or irritable	3 - Nearly every day
7	Feeling afraid as if something awful might happen	3 - Nearly every day
	GAD-7 Anxiety Severity	**21**

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

CNS Vital Signs

NeuroPsych Questionnaire (NPQ) SF-45	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 4:03 (min:secs)	CNSVS Online Version 2.0.6

Domain	Score	Severity	Description
Attention	300	Severe	The Neuropsych Questionnaire Short Form asks patients (or an appropriate observer) a series of questions about their clinical state. The questions are about the symptoms of various neuropsychiatric disorders. The terminology is similar to that used in the diagnostic manuals, and in many familiar clinical questionnaires and rating scales; but it has been simplified, and all symptoms are scored on the same metric. Scores are reported on a scale of 0 (not a problem) to 300 (severe). As a rule, scores above 225 indicate a severe problem; scores from 150-224 indicate a moderate problem; and scores from 75-149, a mild problem. A high score on the Neuropsych Questionnaire Short Form means that the patient is reporting more symptoms of greater intensity. It doesn't necessarily mean that the patient has a particular condition; just that he or she (or their spouse, parent or caregiver) is saying that they have a lot of intense symptoms. Conversely, a low score simply means that the patient (or caregiver) is not reporting symptoms associated with a particular condition, at least during the period of time specified. It does not mean that the patient does not have the condition. Just as some people over-state their problems, others tend to under-state their problems. The Neuropsych Questionnaire Short Form is not a diagnostic instrument. The results it generates are only meant to be interpreted by an experienced clinician in the course of a clinical examination.
Impulsive	180	Moderate	
Memory	300	Severe	
Anxiety	283	Severe	
Panic	133	Mild	
Depression	300	Severe	
Mood Stability	300	Severe	
Aggression	175	Moderate	
Fatigue	300	Severe	
Sleep	300	Severe	
Suicide	0	Not a problem	
Pain	275	Severe	

Attention Questions		
1	Difficulty concentrating	3 - A severe problem
2	Easily distracted	3 - A severe problem
3	Feeling scattered, disorganized	3 - A severe problem
4	Forgetful, I need constant reminding	3 - A severe problem
5	Short attention span	3 - A severe problem
Impulsive Questions		
1	Feeling restless	3 - A severe problem
2	Fidgety, I can't sit still	2 - A moderate problem
3	Impatient	3 - A severe problem
4	Impulsive, act without thinking	1 - A mild problem
5	Overly active	0 - Not a problem
Memory Questions		
1	Forgetful, I need constant reminding	3 - A severe problem
2	My mind goes blank	3 - A severe problem
3	Problems with memory	3 - A severe problem
4	Putting something down and then forgetting where you put it.	3 - A severe problem
Anxiety Questions		
1	Feeling anxious	3 - A severe problem
2	Feeling nervous	3 - A severe problem
3	Feeling restless	3 - A severe problem
4	Feeling tense	3 - A severe problem
5	Fidgety, I can't sit still	2 - A moderate problem
6	Worrying too much	3 - A severe problem
Panic Questions		
1	Attacks of intense anxiety	2 - A moderate problem
2	Feeling so nervous it's hard to breathe	1 - A mild problem
3	Panic attacks	1 - A mild problem
Depression Questions		
1	Feeling depressed	3 - A severe problem
2	Feeling discouraged about the future	3 - A severe problem
3	Feeling irritable	3 - A severe problem
4	Feeling little or no interest in things	3 - A severe problem
5	Not enjoying things as much as before	3 - A severe problem
Mood Stability Questions		
1	Anger	3 - A severe problem
2	Easily frustrated	3 - A severe problem
3	Feeling irritable	3 - A severe problem
4	My moods change quickly	3 - A severe problem

CNS Vital Signs

NeuroPsych Questionnaire (NPQ) SF-45	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 4:03 (min:secs)	CNSVS Online Version 2.0.6

Aggression Questions		
1	Defiant or argumentative	3 - A severe problem
2	Destructive to property or things	1 - A mild problem
3	Hostile	3 - A severe problem
4	Physical aggression toward others	0 - Not a problem
Fatigue Questions		
1	Fatigue	3 - A severe problem
2	Feeling weak	3 - A severe problem
3	Low energy	3 - A severe problem
Sleep Questions		
1	Hard to fall asleep	3 - A severe problem
2	Restless or disturbed sleep	3 - A severe problem
Suicide Questions		
1	I feel like I would be better off dead	0 - Not a problem
2	I feel that my family would be better off if I were gone	0 - Not a problem
3	Thoughts about ending your life	0 - Not a problem
Pain Questions		
1	A lot of aches and pains	3 - A severe problem
2	Back pain	3 - A severe problem
3	Headache	2 - A moderate problem
4	Muscle soreness	3 - A severe problem

CNS Vital Signs

Patient Health Questionnaire (PHQ-9)	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 1:06 (min:secs)	CNSVS Online Version 2.0.6

Major Depressive Syndrome:
If 5 more responses are shaded ("More than half the days" or "Nearly every day") and at least one the shaded questions is #1 or #2.
Other Depressive Syndrome:
If 2 to 4 responses are shaded ("More than half the days" or "Nearly every day") and at least one the shaded questions is #1 or #2.

Note: the diagnoses of Major Depressive Disorder and Other Depressive Disorder requires ruling out normal bereavement (mild symptoms, duration less than 2 months), a history of a manic episode (Bipolar Disorder) and a physical disorder, medication or other drug as the biological cause of the depressive symptoms.

PHQ-9 Depression Severity:
This is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of "Not at all", "Several days", "More than half the days", and "Nearly every day" respectively. PHQ-9 total score for the nine items ranges from 0 to 27.

Interpretation of Total Score:
1-4 Minimal depression
5-9 Mild depression
10-14 Moderate depression
15-19 Moderately severe depression
20-27 Severe depression

1	Little interest or pleasure in doing things	3 - Nearly every day
2	Feeling down, depressed, or hopeless	3 - Nearly every day
3	Trouble falling or staying asleep, or sleeping too much	3 - Nearly every day
4	Feeling tired or having little energy	3 - Nearly every day
5	Poor appetite or overeating	3 - Nearly every day
6	Feeling bad about yourself - or that you are a failure or have let yourself or your family down	3 - Nearly every day
7	Trouble concentrating on things, such as reading the newspaper or watching television	3 - Nearly every day
8	Moving or speaking so slowly that other people could have noticed. Or the opposite - being so fidgety or restless that you have been moving around a lot more than usual	2 - More than half the days
9	Thoughts that you would be better off dead, or of hurting yourself in some way	0 - Not at all
	PHQ-9 Score	**23**

10. How difficult have the problems you reported made it for you to do your work, take care of things at home, or get along with other people? 4 - Extremely difficult

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

CNS Vital Signs

Pittsburgh Sleep Quality Index (PSQI) SF-10	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 7:39 (min:secs)	CNSVS Online Version 2.0.6

The self-rated PSQI assesses sleep quality and disturbances over the past month. Each Domain score has a range of 0 (better) to 3 (worse) with the exception of PSQI Total Score which has a range of 0 (better) to 21 (worse)

Domain	Score	Domain	Score
Duration Of Sleep	**1**	**Sleep Disturbance**	**2**
Sleep Latency	**3**	**Day Dysfunction Due To Sleepiness**	**2**
Sleep Efficiency	**3**	**Overall Sleep Quality**	**3**
Need Meds To Sleep	**3**	**PSQI Total Score**	**17**

1	During the past month, what time have you usually gone to bed at night?	10:00 pm
2	During the past month, how long (in minutes) has it usually taken you to fall asleep each night?	120 minutes
3	During the past month, what time have you usually gotten up in the morning?	7:30 am
4	During the past month, how many hours of actual sleep did you get at night? (This may be different than the number of hours you spent in bed.)	6 hours
5	During the past month, how often have you had trouble sleeping because you ...	
	a. Cannot get to sleep within 30 minutes.	3 - Three or more times a week
	b. Wake up in the middle of the night or early morning.	3 - Three or more times a week
	c. Have to get up to use the bathroom.	2 - Once or twice a week
	d. Cannot breathe comfortably.	1 - Less than once a week
	e. Cough or snore loudly.	0 - Not during the past month
	f. Feel too cold.	0 - Not during the past month
	g. Feel too hot.	1 - Less than once a week
	h. Had bad dreams.	3 - Three or more times a week
	i. Have pain.	2 - Once or twice a week
	j. If you have another reason(s) for having trouble sleeping, please describe: PTSD nightmares How often during the past month have you had trouble sleeping because of this?	 3 - Three or more times a week
6	During the past month, how would you rate your sleep quality overall?	3 - Very bad
7	During the past month, how often have you taken medicine to help you sleep? (prescribed or \"over the counter\")	3 - Three or more times a week
8	During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity?	0 - Not during the past month
9	During the past month, how much of a problem has it been for you to keep up enough enthusiasm to get things done?	3 - A very big problem
10	Do you have a bed partner or roommate?	Partner in same bed
	Is your roommate or bed partner available to answer a few question at this time?	Yes
	Ask your roommate or bed partner how often in the past month you have had ...	
	a. loud snoring?	2 - Once or twice a week
	b. long pauses between breaths while asleep?	0 - Not during the past month
	c. legs twitching or jerking while you sleep?	3 - Three or more times a week
	d. episodes of disorientation or confusion during sleep?	2 - Once or twice a week
	e. Ask your roommate or bed partner if you have any other restlessness while you sleep; please describe: bad dreams, jerk awake, gasp from nightmare, wakes in poor mood due to bad sleep Ask your roommate or bed partner how often in the past month you have had this restlessness while you sleep?	 3 - Three or more times a week

Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ: The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. Psychiatry Research 28:193-213, 1989.

CNS Vital Signs

The PTSD Checklist (PCL-5) SF-20	
Patient ID: NB19741101	Test Date: May 19, 2025 16:40:59
Age: 50	Administrator: David Burns
Total Test Time: 65:55 (min:secs)	Language: English (United States)
Duration: 3:19 (min:secs)	CNSVS Online Version 2.0.6

The PTSD Checklist (PCL-5) Is a list of problems and complaints that people sometimes have in response to stressful life experiences. The PCL-5 consists of 20 items which correspond to the DSM-5 symptom clusters for PTSD. Examinees are instructed to indicate how much they have been bothered by each symptom in the past month using a 5-point (1-5) scale. The anchors for the severity ratings are identical to those used on the SCL-90-R (Derogatis, 1983), and range from "Not at all" to "Extremely." Source: PCL-5 (9/15/2010) Weathers, Litz, Keane, Palmieri, Marx, & Schnurr -- National Center for PTSD. Used with Permission.

The event experienced: auto accident

The event occurred on: 6/16/24

Intrusion (Items 1 - 5)	**25**
Persistent Avoidance (Items 6 - 7)	**10**
Negative Alterations in Cognitions and Mood (Items 8 - 14)	**30**
Alterations in Arousal and Reactivity (Items 15 - 20)	**24**
Total Score (Items 1 - 20)	**89**

1	Repeated, disturbing, and unwanted memories of the stressful experience?	5 - Extremely
2	Repeated, disturbing dreams of the stressful experience?	5 - Extremely
3	Suddenly feeling or acting as if the stressful experience were actually happening again (as if you were actually back there reliving it)?	5 - Extremely
4	Feeling very upset when something reminded you of the stressful experience?	5 - Extremely
5	Having strong physical reactions when something reminded you of the stressful experience (for example, heart pounding, trouble breathing, sweating)?	5 - Extremely
6	Avoiding internal reminders of the stressful experience (for example, thoughts, feelings, or physical sensations)?	5 - Extremely
7	Avoiding external reminders of the stressful experience (for example, people, places, conversations, objects, activities, or situations)?	5 - Extremely
8	Trouble remembering important parts of the stressful experience?	2 - A little bit
9	Having strong negative beliefs about yourself, other people, or the world (for example, having thoughts such as: I am bad, there is something seriously wrong with me, no one can be trusted, the world is completely dangerous)?	3 - Moderately
10	Blaming yourself or someone else strongly for the stressful experience or what happened after it?	5 - Extremely
11	Having strong negative feelings such as fear, horror, anger, guilt, or shame?	5 - Extremely
12	Loss of interest in activities that you used to enjoy?	5 - Extremely
13	Feeling distant or cut off from other people?	5 - Extremely
14	Having trouble experiencing positive feelings (for example, being unable to have loving feelings for those close to you, or feeling emotionally numb)?	5 - Extremely
15	Feeling irritable or angry or acting aggressively?	5 - Extremely
16	Taking too many risks or doing things that cause you harm?	1 - Not at all
17	Being "superalert" or watchful or on guard?	4 - Quite a bit
18	Feeling jumpy or easily startled?	5 - Extremely
19	Having difficulty concentrating?	4 - Quite a bit
20	Trouble falling or staying asleep?	5 - Extremely