CNS Vital Signs

CNS Vital Signs Report	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
CNSVS Duration: 38:32 (min:secs)	CNSVS Online Version 2.0.6

Patient Profile	**Percentile Range**				**> 74**	**25 - 74**	**9 - 24**	**2 - 8**	**< 2**
	Standard Score Range				**> 109**	**90 - 109**	**80 - 89**	**70 - 79**	**< 70**
Domain Scores	**Patient Score**	**Standard Score**	**Percentile**	**VI****	**Above**	**Average**	**Low Average**	**Low**	**Very Low**
Neurocognition Index (NCI)	**NA**	**99**	**47**	**Yes**		**X**			
Composite Memory	**93**	**91**	**27**	**Yes**		**X**			
Verbal Memory	**46**	**80**	**9**	**Yes**			**X**		
Visual Memory	**47**	**106**	**66**	**Yes**		**X**			
Psychomotor Speed	**167**	**103**	**58**	**Yes**		**X**			
Reaction Time*	**715**	**93**	**32**	**Yes**		**X**			
Complex Attention*	**5**	**105**	**63**	**Yes**		**X**			
Cognitive Flexibility	**47**	**104**	**61**	**Yes**		**X**			
Processing Speed	**45**	**92**	**30**	**Yes**		**X**			
Executive Function	**49**	**106**	**66**	**Yes**		**X**			
Social Acuity	**8**	**97**	**42**	**Yes**		**X**			
Reasoning	**10**	**115**	**84**	**Yes**	**X**				
Working Memory	**13**	**114**	**82**	**Yes**	**X**				
Sustained Attention	**35**	**115**	**84**	**Yes**	**X**				
Simple Attention	**39**	**92**	**30**	**Yes**		**X**			
Motor Speed	**120**	**109**	**73**	**Yes**		**X**			

Domain Dashboard: Above average domain scores indicate a standard score (SS) greater than 109 or a Percentile Rank (PR) greater than 74, indicating a high functioning test subject. Average is a SS 90-109 or PR 25-74, indicating normal function. Low Average is a SS 80-89 or PR 9-24 indicating a slight deficit or impairment. Below Average is a SS 70-79 or PR 2-8, indicating a moderate level of deficit or impairment. Very Low is a SS less than 70 or a PR less than 2, indicating a deficit and impairment. Reaction times are in milliseconds. An * denotes that "lower is better", otherwise higher scores are better. Subject Scores are raw scores calculations generated from data values of the individual subtests.

VI** - Validity Indicator: Denotes a guideline for representing the possibility of an invalid test or domain score. "No" means a clinician should evaluate whether or not the test subject understood the test, put forth their best effort, or has a clinical condition requiring further evaluation.

Verbal Memory Test (VBM)	**Score**	**Standard**	**Percentile**	
Correct Hits - Immediate	12	98	45	Verbal Memory test: Subjects have to remember 15 words and recognize them in a field of 15 distractors. The test is repeated at the end of the battery. The VBM test measures how well a subject can recognize, remember, and retrieve words e.g. exploit or attend literal representations or attribute. "Correct Hits" refers to the number of target words recognized. Low scores indicate verbal memory impairment.
Correct Passes - Immediate	12	68	2	
Correct Hits - Delay	8	82	12	
Correct Passes - Delay	14	97	42	
Visual Memory Test (VSM)	**Score**	**Standard**	**Percentile**	
Correct Hits - Immediate	13	109	73	Visual Memory test: Subjects have to remember 15 geometric figures, and recognize them in a field of 15 distractors. The test is repeated at the end of the battery. The VSM test measures how well a subject can recognize, remember, and retrieve geometric figures e.g. exploit or attend symbolic or spatial representations. "Correct Hits" refers to the number of target figures recognized. Low scores indicate visual memory impairment.
Correct Passes - Immediate	11	98	45	
Correct Hits - Delay	11	100	50	
Correct Passes - Delay	12	107	68	
Finger Tapping Test (FTT)	**Score**	**Standard**	**Percentile**	
Right Taps Average	57	100	50	The FTT is a test of motor speed and fine motor control ability. There are three rounds of tapping with each hand. The FTT test measures the speed and the number of finger-taps with each hand. Low scores indicate motor slowing. Speed of manual motor activity varies with handedness. Most people are faster with their preferred hand but not always.
Left Taps Average	63	116	86	

CNS Vital Signs

CNS Vital Signs Report	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
CNSVS Duration: 38:32 (min:secs)	CNSVS Online Version 2.0.6

Symbol Digit Coding (SDC)	Score	Standard	Percentile	
Correct Responses	47	94	34	The SDC test measures speed of processing and draw upon several cognitive processes simultaneously, such as visual scanning, visual perception, visual memory, and motor functions. Errors may be due to impulsive responding, misperception, or confusion.
Errors*	2	87	19	
Stroop Test (ST)	**Score**	**Standard**	**Percentile**	
Simple Reaction Time*	300	101	53	The ST measures simple and complex reaction time, inhibition / disinhibition, mental flexibility or directed attention. The ST helps assess how well a subject is able to adapt to rapidly changing and increasingly complex set of directions. Prolonged reaction times indicate cognitive slowing / impairment. Errors may be due to impulsive responding, misperception, or confusion.
Complex Reaction Time Correct*	601	102	55	
Stroop Reaction Time Correct*	828	88	21	
Stroop Commission Errors*	2	85	16	
Shifting Attention Test (SAT)	**Score**	**Standard**	**Percentile**	
Correct Responses	51	102	55	The SAT measures executive function or how well a subject recognizes set shifting (mental flexibility) and abstraction (rules, categories) and manages multiple tasks simultaneously. Subjects have to adjust their responses to randomly changing rules. The best scores are high correct responses, few errors and a short reaction time. Normal subjects may be slow but accurate, or fast but not so accurate. Attention deficit may be apparent.
Errors*	2	112	79	
Correct Reaction Time*	1083	104	61	
Continuous Performance Test (CPT)	**Score**	**Standard**	**Percentile**	
Correct Responses	40	103	58	The CPT measures sustained attention or vigilance and choice reaction time. Most normal subjects obtain near-perfect scores on this test. A long response time may suggest cognitive slowing and/or impairment. More than 2 errors (total) may be clinically significant. More than 4 errors (total) indicate attentional dysfunction.
Omission Errors*	0	103	58	
Commission Errors*	1	88	21	
Choice Reaction Time Correct*	461	90	25	
Perception Of Emotions Test (POET)	**Score**	**Standard**	**Percentile**	
Correct Responses	10	79	8	The POET measures how well a subject can perceive and identify specific emotions. The POET is an identification test of the positive emotions "happy" and "calm" and the negative emotions "angry" and "sad". "Social cognition" or "emotional acuity" has been defined as "the way in which people make sense of other people and themselves". It is the ability to perceive and understand social information. The reaction times in POET are much longer than in the other tests, indicating the complexity of central processes governing emotional acuity.
Average Correct Reaction Time*	1169	89	23	
Omission Errors*	2	79	8	
Commission Errors*	2	107	68	
Positive Emotions				
Correct Hits	5	79	8	
Reaction Time*	1055	92	30	
Negative Emotions				
Correct Hits	5	88	21	
Reaction Time*	1282	89	23	
Reasoning Test (RT)	**Score**	**Standard**	**Percentile**	
Correct Responses	12	116	86	The NVRT measures how well a subject can perceive and understand the meaning of visual or abstract information and recognizing relationships between visual-abstract concepts. The NVRT is comprised of 15 matrices, or visual analogies. The matrices are progressively more difficult. Each is presented for 14.5 seconds. Non-verbal or visual-abstract reasoning is the process of perceiving issues and reaching conclusions through the use of symbols or generalizations rather than concrete factual information.
Average Correct Reaction Time*	5225	98	45	
Commission Errors*	2	113	81	
Omission Errors*	1	104	61	

CNS Vital Signs

CNS Vital Signs Report	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
CNSVS Duration: 38:32 (min:secs)	CNSVS Online Version 2.0.6

Four Part Continuous Performance Test	Score	Standard	Percentile
Part 1			
Average Correct Reaction Time*	416	92	30
Part 2			
Correct Responses	6	103	58
Average Correct Reaction Time*	429	90	25
Incorrect Responses*	0	104	61
Average Incorrect Reaction Time*	0		
Omission Errors*	0	103	58
Part 3			
Correct Responses	16	113	81
Average Correct Reaction Time*	604	92	30
Incorrect Responses*	0	104	61
Average Incorrect Reaction Time*	0		
Omission Errors*	0	113	81
Part 4			
Correct Responses	15	120	91
Average Correct Reaction Time*	730	95	37
Incorrect Responses*	2	97	42
Average Incorrect Reaction Time*	1112	84	14
Omission Errors*	1	120	91

The FPCPT test is a four part test that measures a subject's working memory and sustained attention. The FPCPT is a four part test: PART ONE - is a simple reaction time test, the subject must press the space bar when any stimulus is presented; PART TWO - is a variant of the continuous performance test, the subject is asked to respond to one stimulus, but not to any others. Discrimination is required, so the reaction times that are generated are "choice reaction times". PART THREE - is a "one back" CPT. The subject has to respond to a figure only if the figure immediately preceding was the same. PART FOUR - is a "two-back" CPT. It is a difficult task and is used to measure working memory. Parts two, three, and four of the tests are used to calculate sustained attention domain.

Longitudinal View	2024-03-25 14:50:35	2024-07-26 11:31:12	2024-08-19 11:19:21	2025-01-14 09:43:31	2025-08-15 09:56:53
Neurocognition Index (NCI)	91	44	96	96	99
Composite Memory	112	81	93	104	91
Verbal Memory	109	87	93	109	80
Visual Memory	109	84	97	97	106
Psychomotor Speed	80	82	86	94	103
Reaction Time*	56	69	77	70	93
Complex Attention*	105	-0	118	108	105
Cognitive Flexibility	101	-10	108	103	104
Processing Speed	92	78	94	90	92
Executive Function	100	-11	107	103	106
Social Acuity	83	104	97	90	97
Reasoning	112	101	112	115	115
Working Memory	88	104	111	88	114
Sustained Attention	96	102	113	95	115
Simple Attention	77	92	107	77	92
Motor Speed	81	91	88	98	109

Generalized Anxiety Disorder 7-item (GAD-7) Scale	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
Duration: 1:00 (min:secs)	CNSVS Online Version 2.0.6

GAD-7 Anxiety Severity:

This is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of "Not at all", "Several days", "More than half the days", and "Nearly every day" respectively. GAD-7 total score for the seven items ranges from 0 to 21.

Interpretation of Total Score:

0-4 None-Minimal anxiety

5-9 Mild anxiety

10-14 Moderate anxiety

15-21 Severe anxiety

1	Feeling nervous, anxious or on edge	2 - More than half the days
2	Not being able to stop or control worrying	1 - Several days
3	Worrying too much about different things	2 - More than half the days
4	Trouble relaxing	3 - Nearly every day
5	Being so restless that it is hard to sit still	3 - Nearly every day
6	Becoming easily annoyed or irritable	1 - Several days
7	Feeling afraid as if something awful might happen	1 - Several days
	GAD-7 Anxiety Severity	**13**

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

CNS Vital Signs

NeuroPsych Questionnaire (NPQ) SF-45	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
Duration: 3:09 (min:secs)	CNSVS Online Version 2.0.6

Domain	Score	Severity	Description
Attention	140	Mild	The Neuropsych Questionnaire Short Form asks patients (or an appropriate observer) a series of questions about their clinical state. The questions are about the symptoms of various neuropsychiatric disorders. The terminology is similar to that used in the diagnostic manuals, and in many familiar clinical questionnaires and rating scales; but it has been simplified, and all symptoms are scored on the same metric. Scores are reported on a scale of 0 (not a problem) to 300 (severe). As a rule, scores above 225 indicate a severe problem; scores from 150-224 indicate a moderate problem; and scores from 75-149, a mild problem. A high score on the Neuropsych Questionnaire Short Form means that the patient is reporting more symptoms of greater intensity. It doesn't necessarily mean that the patient has a particular condition; just that he or she (or their spouse, parent or caregiver) is saying that they have a lot of intense symptoms. Conversely, a low score simply means that the patient (or caregiver) is not reporting symptoms associated with a particular condition, at least during the period of time specified. It does not mean that the patient does not have the condition. Just as some people over-state their problems, others tend to under-state their problems. The Neuropsych Questionnaire Short Form is not a diagnostic instrument. The results it generates are only meant to be interpreted by an experienced clinician in the course of a clinical examination.
Impulsive	160	Moderate	
Memory	100	Mild	
Anxiety	200	Moderate	
Panic	200	Moderate	
Depression	140	Mild	
Mood Stability	100	Mild	
Aggression	50	Not a problem	
Fatigue	167	Moderate	
Sleep	200	Moderate	
Suicide	167	Moderate	
Pain	250	Severe	

Attention Questions		
1	Difficulty concentrating	2 - A moderate problem
2	Easily distracted	2 - A moderate problem
3	Feeling scattered, disorganized	1 - A mild problem
4	Forgetful, I need constant reminding	1 - A mild problem
5	Short attention span	1 - A mild problem
Impulsive Questions		
1	Feeling restless	3 - A severe problem
2	Fidgety, I can't sit still	2 - A moderate problem
3	Impatient	2 - A moderate problem
4	Impulsive, act without thinking	1 - A mild problem
5	Overly active	0 - Not a problem
Memory Questions		
1	Forgetful, I need constant reminding	1 - A mild problem
2	My mind goes blank	1 - A mild problem
3	Problems with memory	1 - A mild problem
4	Putting something down and then forgetting where you put it.	1 - A mild problem
Anxiety Questions		
1	Feeling anxious	2 - A moderate problem
2	Feeling nervous	2 - A moderate problem
3	Feeling restless	3 - A severe problem
4	Feeling tense	1 - A mild problem
5	Fidgety, I can't sit still	2 - A moderate problem
6	Worrying too much	2 - A moderate problem
Panic Questions		
1	Attacks of intense anxiety	2 - A moderate problem
2	Feeling so nervous it's hard to breathe	2 - A moderate problem
3	Panic attacks	2 - A moderate problem
Depression Questions		
1	Feeling depressed	2 - A moderate problem
2	Feeling discouraged about the future	2 - A moderate problem
3	Feeling irritable	1 - A mild problem
4	Feeling little or no interest in things	1 - A mild problem
5	Not enjoying things as much as before	1 - A mild problem
Mood Stability Questions		
1	Anger	1 - A mild problem
2	Easily frustrated	1 - A mild problem
3	Feeling irritable	1 - A mild problem
4	My moods change quickly	1 - A mild problem

CNS Vital Signs

NeuroPsych Questionnaire (NPQ) SF-45	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
Duration: 3:09 (min:secs)	CNSVS Online Version 2.0.6

Aggression Questions		
1	Defiant or argumentative	1 - A mild problem
2	Destructive to property or things	0 - Not a problem
3	Hostile	1 - A mild problem
4	Physical aggression toward others	0 - Not a problem
Fatigue Questions		
1	Fatigue	2 - A moderate problem
2	Feeling weak	1 - A mild problem
3	Low energy	2 - A moderate problem
Sleep Questions		
1	Hard to fall asleep	2 - A moderate problem
2	Restless or disturbed sleep	2 - A moderate problem
Suicide Questions		
1	I feel like I would be better off dead	2 - A moderate problem
2	I feel that my family would be better off if I were gone	2 - A moderate problem
3	Thoughts about ending your life	1 - A mild problem
Pain Questions		
1	A lot of aches and pains	3 - A severe problem
2	Back pain	2 - A moderate problem
3	Headache	3 - A severe problem
4	Muscle soreness	2 - A moderate problem

CNS Vital Signs

Patient Health Questionnaire (PHQ-9)	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
Duration: 1:03 (min:secs)	CNSVS Online Version 2.0.6

Major Depressive Syndrome:
If 5 more responses are shaded ("More than half the days" or "Nearly every day") and at least one the shaded questions is #1 or #2.
Other Depressive Syndrome:
If 2 to 4 responses are shaded ("More than half the days" or "Nearly every day") and at least one the shaded questions is #1 or #2.

Note: the diagnoses of Major Depressive Disorder and Other Depressive Disorder requires ruling out normal bereavement (mild symptoms, duration less than 2 months), a history of a manic episode (Bipolar Disorder) and a physical disorder, medication or other drug as the biological cause of the depressive symptoms.

PHQ-9 Depression Severity:
This is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of "Not at all", "Several days", "More than half the days", and "Nearly every day" respectively. PHQ-9 total score for the nine items ranges from 0 to 27.

Interpretation of Total Score:
1-4 Minimal depression
5-9 Mild depression
10-14 Moderate depression
15-19 Moderately severe depression
20-27 Severe depression

1	Little interest or pleasure in doing things	2 - More than half the days
2	Feeling down, depressed, or hopeless	2 - More than half the days
3	Trouble falling or staying asleep, or sleeping too much	2 - More than half the days
4	Feeling tired or having little energy	2 - More than half the days
5	Poor appetite or overeating	1 - Several days
6	Feeling bad about yourself - or that you are a failure or have let yourself or your family down	2 - More than half the days
7	Trouble concentrating on things, such as reading the newspaper or watching television	2 - More than half the days
8	Moving or speaking so slowly that other people could have noticed. Or the opposite - being so fidgety or restless that you have been moving around a lot more than usual	2 - More than half the days
9	Thoughts that you would be better off dead, or of hurting yourself in some way	1 - Several days
	PHQ-9 Score	**16**

10. How difficult have the problems you reported made it for you to do your work, take care of things at home, or get along with other people? 4 - Extremely difficult

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

CNS Vital Signs

The PTSD Checklist (PCL-5) SF-20	
Patient ID: AL19710306	Test Date: August 15, 2025 09:56:53
Age: 54	Administrator: David Burns
Total Test Time: 47:05 (min:secs)	Language: English (United States)
Duration: 2:51 (min:secs)	CNSVS Online Version 2.0.6

The PTSD Checklist (PCL-5) Is a list of problems and complaints that people sometimes have in response to stressful life experiences. The PCL-5 consists of 20 items which correspond to the DSM-5 symptom clusters for PTSD. Examinees are instructed to indicate how much they have been bothered by each symptom in the past month using a 5-point (1-5) scale. The anchors for the severity ratings are identical to those used on the SCL-90-R (Derogatis, 1983), and range from "Not at all" to "Extremely." Source: PCL-5 (9/15/2010) Weathers, Litz, Keane, Palmieri, Marx, & Schnurr -- National Center for PTSD. Used with Permission.

The event experienced: Auto accident

The event occurred on: 11/02/2023

Intrusion (Items 1 - 5)	**15**
Persistent Avoidance (Items 6 - 7)	**7**
Negative Alterations in Cognitions and Mood (Items 8 - 14)	**24**
Alterations in Arousal and Reactivity (Items 15 - 20)	**19**
Total Score (Items 1 - 20)	**65**

1	Repeated, disturbing, and unwanted memories of the stressful experience?	3 - Moderately
2	Repeated, disturbing dreams of the stressful experience?	3 - Moderately
3	Suddenly feeling or acting as if the stressful experience were actually happening again (as if you were actually back there reliving it)?	2 - A little bit
4	Feeling very upset when something reminded you of the stressful experience?	3 - Moderately
5	Having strong physical reactions when something reminded you of the stressful experience (for example, heart pounding, trouble breathing, sweating)?	4 - Quite a bit
6	Avoiding internal reminders of the stressful experience (for example, thoughts, feelings, or physical sensations)?	3 - Moderately
7	Avoiding external reminders of the stressful experience (for example, people, places, conversations, objects, activities, or situations)?	4 - Quite a bit
8	Trouble remembering important parts of the stressful experience?	3 - Moderately
9	Having strong negative beliefs about yourself, other people, or the world (for example, having thoughts such as: I am bad, there is something seriously wrong with me, no one can be trusted, the world is completely dangerous)?	4 - Quite a bit
10	Blaming yourself or someone else strongly for the stressful experience or what happened after it?	4 - Quite a bit
11	Having strong negative feelings such as fear, horror, anger, guilt, or shame?	4 - Quite a bit
12	Loss of interest in activities that you used to enjoy?	4 - Quite a bit
13	Feeling distant or cut off from other people?	3 - Moderately
14	Having trouble experiencing positive feelings (for example, being unable to have loving feelings for those close to you, or feeling emotionally numb)?	2 - A little bit
15	Feeling irritable or angry or acting aggressively?	3 - Moderately
16	Taking too many risks or doing things that cause you harm?	2 - A little bit
17	Being "superalert" or watchful or on guard?	3 - Moderately
18	Feeling jumpy or easily startled?	3 - Moderately
19	Having difficulty concentrating?	3 - Moderately
20	Trouble falling or staying asleep?	5 - Extremely