



## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/41

Paper 4 Theory (Extended)

October/November 2018

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1	The following for	rmulae	represe	nt different s	substance	es.			
		Al	Ag	CaCO <sub>3</sub>	CH₄	$Cl_2$	Cu	SO <sub>2</sub>	

Answer the following questions using only these substances. Each substance may be used once, more than once or not at all.

State which substance is:

(a)	used to make food containers	[1]
(b)	added to a blast furnace to remove impurities during the production of iron	[1]
(c)	the main constituent of natural gas	[1]
(d)	a cause of acid rain	[1]
(e)	a gas which bleaches damp litmus paper	[1]
(f)	a gas which contributes to climate change.	[1]
	[Total	: 6]


2 The table gives some information about four different particles, **A**, **B**, **C** and **D**.

particle	number of electrons	number of neutrons	number of protons	electronic structure	charge on particle
Α	11	12	11	2,8,1	0
В		14	11	2,8,1	0
С	18	20		2,8,8	0
D	18	20	17		

(a)	Complete the table. The first row has been done for you.	[4]
(b)	Give <b>two</b> particles from the table which are isotopes of each other.	
		[1]
(c)	Element <b>Z</b> is in the same group of the Periodic Table as <b>A</b> and is less reactive than <b>A</b> .	
	State the identity of element <b>Z</b> .	
		[1]
(d)	C is unreactive.	
	Use information from the table to explain why.	
		[1]
	[Total	i: 7]

**3 (a)** Copper(II) nitrate decomposes when heated. Two gases, oxygen and nitrogen dioxide, and a solid are made in the reaction.

A sample of copper(II) nitrate was decomposed using the apparatus shown.



(i) Complete the chemical equation for the reaction.

$$2Cu(NO_3)_2 \rightarrow O_2 + .....NO_2 + .....$$
 [2]

(ii) Only oxygen gas is collected at X.

Explain why.		
	[	11

(b) Nitrogen dioxide and other oxides of nitrogen are formed in car engines.

Explain how nitrogen dioxide is formed in car engines.

[2]

(c) A	teach	er he	eated 18.8g of copper	r(II) nitrate.				
(1	i) Cal	cula	te the number of mole	es of copper(II) nitrate present in the 18.8 g.				
(i			te the maximum numb II) nitrate.	per of moles of oxygen that can be made by hea	ting 18.8g of			
					mol [1]			
(iii		Calculate the maximum volume of oxygen at room temperature and pressure, in cm $^3$ , that can be made by heating 18.8 g of copper(II) nitrate.						
					cm³ [1]			
(d) A	\ samn	le of	conner(II) nitrate was	s dissolved in water to form an aqueous solution	1			
				to three portions. A separate test was done on				
	is show		o columnia opini iii	to all ou portioner, respandite test that delice on	odon pordon			
	te	st	reagent added	result				
		1	aqueous sodium hydroxide	light blue precipitate forms				
	2	2	zinc powder	solution changes from blue to colourless and a brown solid forms				
	(	3		ammonia gas is produced				
(1	i) Giv	e the	e formula of the light b	olue precipitate formed in <b>test 1</b> .				
					[1]			
(i	i) Exp	olain	the changes seen in	test 2.				
	••••							
(iii	i) Ide	ntify		t must be added to the aqueous copper(II) nitra				
	1							

[2]

(e)		oper(II) nitrate can be made by reacting $copper(II)$ carbonate with nitric acid. One of t ducts is carbon dioxide.	he
	(i)	Write a chemical equation for the reaction of copper(II) carbonate with nitric acid.	
			[2]
	(ii)	Carbon dioxide is added to the air by living things.	
		Name the chemical process by which living things add carbon dioxide to the air.	
			[1]
(	(iii)	Carbon dioxide is removed from the air by plants.	
		Name the chemical process by which plants remove carbon dioxide from the air.	
			[1]
		[Total: 1	19]

(a)	Sulf	uric	acid is made industrially by a four-step process.						
	ster ster	2 3	Sulfur is burned in air to produce sulfur dioxide.  Sulfur dioxide is converted into sulfur trioxide.  Sulfur trioxide is reacted with concentrated sulfuric acid to produce oleum.  Oleum is reacted with water to produce concentrated sulfuric acid.						
	(i)	Sor	me sulfur is obtained by mining.						
		Name <b>one</b> other major source of sulfur.							
	(ii)	 Wh	at is the name of the process by which sulfuric acid is made industrially?						
	(iii)	Des	scribe the conversion of sulfur dioxide into sulfur trioxide in <b>step 2</b> .	٠ ١٠.					
	()		our answer, include: a chemical equation for the reaction the essential reaction conditions.						
				. [5]					
(b)			oncentrated sulfuric acid is added to glucose, $C_6H_{12}O_6$ , a black solid is produced. rated sulfuric acid acts as a dehydrating agent.	The					
	(i)	Wh	at is removed from the glucose in this reaction?						
	(ii)	 Nar	me the black solid produced in this reaction.	. [1]					
				. [1]					

(c)		The gas hydrogen sulfide, $\rm H_2S$ , is produced when concentrated sulfuric acid is added to solid otassium iodide.				
	The	e reaction involves oxidation.				
	(i)	Define the term <i>oxidation</i> in terms of electron transfer.				
			[1]			
	(ii)	Complete the dot-and-cross diagram to show the electron arrangement in a molecule hydrogen sulfide. Show outer shell electrons only.	of			
		H S H	[2]			
	(iii)	Hydrogen sulfide has a simple molecular structure.				
		Explain why hydrogen sulfide has a low boiling point.				
			[2]			

(	d)	Dilute sulfuric acid	reacts with ac	lueous sodium	hydrogencarl	bonate in a ne	eutralisation i	reaction
١	u,	Dilate Saliane acia	TCacto With ac	Jacous soulaili	rryarogenican	bonate in a ne	, ati alibation i	Cachon.

$$H_2SO_4(aq) + 2NaHCO_3(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(I) + 2CO_2(g)$$

In a titration, 0.200 mol/dm³ aqueous sodium hydrogencarbonate was used to neutralise 20.0 cm³ of dilute sulfuric acid of concentration 0.150 mol/dm³.

mol [	[1]
-------	-----

- (ii) Calculate the number of moles of sodium hydrogencarbonate needed to neutralise the dilute sulfuric acid.
  - ..... mol [1]
- (iii) Calculate the volume, in cm³, of 0.200 mol/dm³ aqueous sodium hydrogencarbonate needed to neutralise the dilute sulfuric acid.

..... cm<sup>3</sup> [1]

[Total: 17]


**5** Hydrogen gas reacts with iodine gas. The equation is shown.

$$H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$$

The reaction is reversible and can reach equilibrium.

| <br> | [2 |
|------|------|------|------|------|------|------|------|------|------|------|------|----|

**(b)** The graphs show how pressure affects the yield of hydrogen iodide, HI, at two different temperatures.




(i)	Explain why	the yield a	t 500°C	does <b>not</b>	change as	s the	pressure is	increased.
-----	-------------	-------------	---------	-----------------	-----------	-------	-------------	------------

		. [1]

(ii) What can you conclude from the difference in the yield of hydrogen iodide at the **two** temperatures shown? Explain your answer.



(c) The graph shows how the concentration of hydrogen iodide, HI, changes after hydrogen gas and iodine gas are mixed together in a sealed container.



	uno
(i)	When is the rate of reaction fastest?
	[1]
(ii)	The reaction was repeated at the same temperature and pressure but in the presence of a catalyst.
	Draw a graph on the same axes to show how the concentration of hydrogen iodide changes with time in the presence of a catalyst. [2]
Αn	nixture of hydrogen gas and iodine gas is allowed to reach equilibrium.

(d)

(i)

(ii)

increasing the pressure of a gas increases its concentration.
State and explain the effect of increasing the pressure on the <b>rate</b> of the forward reaction.
[2]
State and explain the effect of increasing the temperature on the <b>rate</b> of the reverse reaction.

[Total: 13]

	hane, $\rm C_2H_6$ , is a member of the homologous series called alkanes. Hanol, $\rm C_2H_5OH$ , is a member of the homologous series called alcohols.
(i)	Alkanes are hydrocarbons.
	What is meant by the term <i>hydrocarbon</i> ?
	[2]
(ii)	All members of a homologous series can be represented by a general formula.
	State the general formula of:
	alkanes
	alcohols
(iii)	State <b>two</b> characteristics, other than having the same general formula, of members of a homologous series.
	1
	2
	[2]
(b) Eth	nane can react with chlorine in a substitution reaction.
(i)	State <b>one</b> essential reaction condition.
	[1]
(ii)	Draw the structure of the organic product formed by substitution of <b>one</b> of the hydrogen atoms in ethane with chlorine. Show all of the atoms and all of the bonds.
	[1]
(iii)	Name the product of the substitution reaction between ethane and chlorine that does <b>not</b> contain carbon.
	[1]

(c) Propan-1-ol is an alcohol.

The structure of propan-1-ol is shown.

Propan-1-ol reacts with ethanoic acid to form an ester.

Give the name of the ester formed in this reaction.

......[1]

(d) Ester Y has the structure shown.

ester Y

(i)	Give	the	molecular	formula	of	ester	Y	
-----	------	-----	-----------	---------	----	-------	---	--

.....[1]

(ii) Draw the structures of the carboxylic acid and the alcohol used to make ester Y. Show all of the atoms and all of the bonds. Give the name of the carboxylic acid and the alcohol.

structure of the carboxylic acid

name of the carboxylic acid .....

structure of the alcohol

name of the alcohol .....

(e)	Nylon is a poly	amide.								
	Complete the bonds present		structure of	nylon.	Show al	I of the	atoms	and	all o	f the
										[3]
								Γ	Total	l: 18

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

								Gro	Group								
=												≡	2	>	5	=>	₹
							- エ										2 He
Key	Key	Key	Key				hydrogen 1										helium 4
4 atomic number	atomic number	atomic number	omic number			,						2	9	7	80	6	10
Be atomic symbol	atomic symbol	atomic symbol	nic symbol	$\overline{g}$								Ω	ပ	Z	0	ш	Ne
beryllium name 9 relative atomic mass	name relative atomic mass	name relative atomic mass	name ive atomic mass	SS								boron 11	carbon 12	nitrogen 14	oxygen 16	fluorine 19	neon 20
12												13	14	15	16	17	18
Mg												Αl	S	۵	ഗ	Cl	Ā
magnesium 24												aluminium 27	silicon 28	phosphorus 31	sulfur 32	chlorine 35.5	argon 40
21 22 23	22 23	23		24		25	26	27	28	29	30	31	32	33	34	38	36
>   i	> =	>		ပ်	_	Mn	Fe	ပိ	z	Cn	Zu	Ga	Ge	As	Se	ğ	궃
calcium         scandium         titanium         vanadium         chromium           40         45         48         51         52	titanium vanadium 6	vanadium 51		chromit 52	Ę	manganese 55	iron 56	cobalt 59	nickel 59	copper 64	zinc 65	gallium 70	germanium 73	arsenic 75	selenium 79	bromine 80	krypton 84
39 40 41	40 41	41		42		43	44	45	46	47	48	49	20	51	52	53	54
Y Zr Nb	Zr Nb	Q Q		Ĭ	0	ပ	R	格	Pd	Ag	g	In	Sn	Sb	<u>Б</u>	н	Xe
strontium         yttrium         zirconium         niobium         molybdenum           88         89         91         93         96	zirconium niobium 91 93	niobium 93		molybde 96	mnu	technetium -	ruthenium 101	rhodium 103	palladium 106	silver 108	cadmium 112	indium 115	tin 119	antimony 122	tellurium 128	iodine 127	xenon 131
57–71 72 73	72 73	73		1/	<u></u>	75	92	77	78	79	80	81	82	83	84	85	98
lanthanoids Hf Ta	HTTa	Та		>	>	Re	Os	Ľ	₹	Αn	Нg	l1	Pp	<u>B</u>	Ъ	¥	R
hafnium tantalum 178 181	tantalum 181	tantalum 181		tung	gsten 84	rhenium 186	osmium 190	iridium 192	platinum 195	gold 197	mercury 201	thallium 204	lead 207	bismuth 209	polonium —	astatine -	radon
89–103 104 105	104 105	105			90	107	108	109	110	111	112		114		116		
actinoids Rf Db	Rf	Db		0,	Sg	Bh	Hs	¥	Ds		ű		F1				
radium rutherfordium dubnium seab	dubnium	dubnium		seab	seaborgium	bohrium	hassium	meitnerium -	darmstadtium	0	copernicium		flerovium		livermorium		
ı	ı	_			_ 	ı	-	ı	ı		ı		ı		I		

71	lutetium 175	103	ב	lawrencium	ı
0 <b>X</b>	ytterbium 173	102	%	nobelium	ı
69 L	thulium 169	101	Md	mendelevium	I
88 7	erbium 167	100	Fm	ferminm	1
<sup>67</sup>	holmium 165	66	Es	einsteinium	I
99	dysprosium 163	86	ŭ	californium	ı
65 T	terbium 159	97	BK	berkelium	1
<sup>64</sup>	gadolinium 157	96	Cm	curium	ı
63 <u>T</u>	europium 152	92	Am	americium	ı
62 <b>Sm</b>	samarium 150	94	Pu	plutonium	ı
61 D	promethium	93	δ	neptunium	ı
09 Z	neodymium 144	92	$\supset$	uranium	238
59 <b>7</b>	praseodymium 141	91	Ра	protactinium	231
<sub>88</sub> م	cerium 140	06	Ч	thorium	232
57	lanthanum 139	68	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is  $24\,dm^3$  at room temperature and pressure (r.t.p.).