



Cambridge Assessment International Education

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/41

Paper 4 Theory (Extended)

May/June 2019

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



BLANK PAGE

1


		the table 4_2 Mg and $^{26}_{12}$		umber of proto	ns, neutrons a	and electrons pres
		2 3 12				7
			number of protons	number of neutrons	number of electrons	
		²⁴ Mg				
		²⁶ Mg				
		12 0				
(ii) V	Vhat term	is used to	describe atom	s of the same	element, such	as ²⁴ Mg and ²⁶ Mg?
						12 - 12 -
•••						
 (iii) E	Explain wh	 ny the cher				
iii) E	Explain wh	ny the cher	mical properties			
 iii) E	Explain wh	ny the cher		s of $^{24}_{12}$ Mg and $^{26}_{12}$	ßMg are the sa	me.
 iii) E 	xplain wh	ny the cher		s of $^{24}_{12}$ Mg and $^{26}_{12}$	ßMg are the sa	
 IIII) E 	Explain wh	ny the cher	mical properties	s of ²⁴ Mg and ²¹	Mg are the sa	me.
 (iii) E 	xplain wh	ny the cher	mical properties	s of ²⁴ Mg and ²¹	Mg are the sa	me.
			mical properties	s of ²⁴ Mg and ²⁴	Mg are the sa	me.
 Comp	olete the t	able to ider	mical properties	s of ²⁴ Mg and ²⁴	Mg are the sa	me.
 Comp		able to ider	mical properties	s of ²⁴ Mg and ²⁴	Mg are the sa	me.
 Comp	olete the t	able to ider	ntify the atoms	and ions which	Mg are the sa	me.
Comp	olete the t	able to ider	mical properties	s of ²⁴ Mg and ²⁴	Mg are the sa	me.
Comp	olete the t	able to ider	ntify the atoms	and ions which	Mg are the sa	me.
 Comp	olete the t	able to ider	ntify the atoms number of protons	and ions which	have the follow	me.
 Comp	olete the t	able to ider	ntify the atoms number of protons	and ions which number of neutrons	have the followinumber of electrons	me.
 Comp	olete the t	able to ider	ntify the atoms number of protons 11	and ions which number of neutrons 12	have the following number of electrons 10	me.
 Comp	olete the t	able to ider	ntify the atoms number of protons 11	and ions which number of neutrons 12	have the following number of electrons 10	me.
Comp	plete the toons and e	able to iderelectrons.	ntify the atoms number of protons 11	and ions which number of neutrons 12 5 20	have the following number of electrons 10 4 18	me.

© UCLES 2019 0620/41/M/J/19 **[Turn over**

[Total: 13]

Z is a covalent substance. In an experiment, a sample of pure solid **Z** was continually heated for 11 minutes.

The graph shows how the temperature of the sample of pure **Z** changed during the first 9 minutes.



(a)	What is	the	melting	point	of	pure	Z ?
-----	---------	-----	---------	-------	----	------	------------

	$^{\circ}\text{C}$	[1]
--	--------------------	-----

(b) The sample of pure **Z** began to boil at 9 minutes. It was boiled for 2 minutes.

Use this information to sketch on the grid how the temperature of the sample of pure **Z** changed between 9 minutes and 11 minutes. [1]

(c) The sample of pure **Z** was continually heated between 2 minutes and 5 minutes.


Explain, in terms of attractive forces, why there was no increase in t sample of pure Z between 2 minutes and 5 minutes.	he temperature of the
sample of pare 2 between 2 minutes and 5 minutes.	

.....

(d)	Describe how the motion of particles of pure Z changed from 0 minutes to 2 minutes.
	[2]
(e)	The experiment was repeated using a solid sample of impure Z .
	Suggest the differences, if any, in the melting point and boiling point of the sample of impure ${\bf Z}$ compared to the sample of pure ${\bf Z}$.
	melting point
	boiling point
	[2]

(f) A sample of pure **Z** was allowed to cool from 120 °C to 20 °C. The total time taken was 8 minutes.

Starting from point \mathbf{x} , sketch on the grid how the temperature of the sample of pure \mathbf{Z} changed between 0 minutes and 8 minutes.



[2]

[Total: 10]

- **3** Zinc and copper are elements next to each other in the Periodic Table.
 - (a) Zinc is obtained from zinc blende in a two-step process.
 - In step 1, zinc blende is converted into zinc oxide.
 - In **step 2**, zinc oxide is converted into zinc in a blast furnace.

Outline how each of these steps are done.

In yo	our	ans	wer:
-------	-----	-----	------

- give **one** chemical equation for each step
- describe how zinc is removed from the blast furnace in step 2.

	step 1
	chemical equation
	step 2
	chemical equation
	removal of zinc in step 2
	[5]
(b)	Name the alloy formed when zinc is mixed with copper.
	[1]
(c)	Copper is a transition element. It can have variable oxidation states.
	State two other chemical properties of transition elements which make them different from Group I elements.
	1
	2
	[2]

(d)	A co	ompound of copper can be used to test for water.	
	(i)	State the full name of this compound of copper.	
			[1]
	(ii)	State the colour change that occurs when water is added to this compound of copper.	
		from to	[2]
(e)	Aqu (i)	ueous potassium iodide reacts with aqueous copper(II) sulfate to produce iodine. Balance the chemical equation for this reaction.	
	(1)	KI + CuSO ₄ \rightarrow CuI + I ₂ + K ₂ SO ₄	
		7 2 2 7	[2]
	(ii)	Deduce the charge on the copper ion in CuI.	
			[1]
	(iii)	In terms of electron transfer, explain why copper is reduced in this reaction.	
			[1]
((iv)	Identify the reducing agent.	
			[1]
		[Total: 1	16

Ethanoic acid is a weak acid and hydrochloric acid is a strong acid.

Both et	hanoic acid and hydrochloric acid dissociate in aqueous solution.
(a) (i)	Define the term acid.
	[1]
(ii)	The chemical equation shows the changes which occur when the strong acid, hydrochloric acid, is added to water.
	$HCl(aq) \rightarrow H^{+}(aq) + Cl^{-}(aq)$
	Complete the chemical equation to show the changes which occur when the weak acid ethanoic acid, is added to water.
	CH ₃ COOH(aq)[2]
av	tudent does experiments to show that hydrochloric acid is a strong acid and ethanoic acid is reak acid. The student adds an excess of hydrochloric acid and an excess of ethanoic acid separate samples of lumps of calcium carbonate.
	ly the identity of the acid is changed between the experiments. All other conditions are kept same.
(i)	State two observations which would show that hydrochloric acid is a stronger acid than ethanoic acid.
	1
	2
	[2]
(ii)	The student uses the same size container and checks that the pressure is the same for each experiment.
	State three other conditions which must be kept the same to ensure fair testing.
	1
	2
	3
	[3]

(C) H	ydrochlo	oric acid	produces	salts	called	chlorides.

Magnesium carbonate reacts with hydrochloric acid to produce magnesium chloride.

$$MgCO_3 + 2HCl \rightarrow MgCl_2 + H_2O + CO_2$$

A student used 50.00 cm³ of 2.00 mol/dm³ hydrochloric acid in an experiment to produce magnesium chloride.

Calculate the mass, in g, of magnesium carbonate needed to react exactly with 50.00 cm³ of 2.00 mol/dm³ hydrochloric acid using the following steps.

•	Calculate the number	of moles of HC1	present in 50.00 cm ³	of 2.00 mol/dm3 HC1
---	----------------------	-----------------	----------------------------------	---------------------

				 	mo	ol
D 1	 	 	 	 F0 00	3	

• Determine the number of moles of MgCO₃ which would react with 50.00 cm³ of 2.00 mol/dm³ HC*l*.

• Calculate the relative formula mass, M_r , of MgCO₃.

$$M_{\rm r}$$
 of MgCO₃ =

• Calculate the mass of MgCO $_3$ needed to react exactly with 50.00 cm 3 of 2.00 mol/dm 3 HCl.

(d)		student prepares crystals of magnesium chloride by adding an excess of gnesium carbonate to 50.00 cm³ of 2.00 mol/dm³ hydrochloric acid.	of					
	The	student filters the mixture and rinses the residue.						
	(i)	Why does the student add an excess of magnesium carbonate?						
		[1]					
	(ii)	Why does the student rinse the residue?						
		[1]					
	(iii)	Describe how the student would obtain pure crystals of magnesium chloride from th filtrate.	е					
		[3	3]					
(e)		er chloride, $AgCl$, is insoluble. It can be made by a precipitation reaction between aqueouum chloride and a suitable aqueous silver salt.	S					
	(i)	What is meant by the term precipitate?						
		[2	 21					
	(ii)	Name a suitable silver salt to use to prepare silver chloride. Complete the chemical equation to show the formation of insoluble silver chloride from						
		aqueous barium chloride and the silver salt you have named.						
		name of a suitable silver salt						
		$BaCl_2 + \dots \longrightarrow \dots + \dots $	31					
			3]					

5 The structures of five alkenes, A, B, C, D and E, are shown.

A H C=C H	H C=	CH ₃ =C H	C=C	H₂CH₃
	$\begin{array}{c} \textbf{D} \\ \textbf{H} \\ \textbf{C} = \textbf{C} \\ \textbf{H} \end{array}$	H C=	CH ₂ CH ₂ CH ₂ CH ₃	
Vhat is the gen	eral formula of alkenes?			

(a)	What is the general formula of alkenes?	
		[1]
(b)	What is the molecular formula of alkene D ?	
		[1]
(c)	Predict which alkene, A , B , C , D or E , has the highest boiling point. Explain your answer.	
	alkene	
	explanation	
		[2]

[2]

(e) A student added aqueous bromine to alkene C.

Describe the colour change seen and draw the structure of the product. Show all of the atoms and all of the bonds.

colour change from to

structure

[2]

(f) Two different alcohols can be produced from alkene **B** by an addition reaction.

(i) Draw the structures of the **two** alcohols. Show all of the atoms and all of the bonds.

[2]

[3]


(ii) State the reagent and conditions needed to produce an alcohol from alkene B.

reagent

conditions

.....

(g) Alkene C can be converted into a polymer.




(i) What type of polymerisation occurs?

.....[1]

(ii) Suggest the name of the polymer formed.

.....[1]

(iii) Complete the chemical equation to show this polymerisation.



[3]

(iv) State the empirical formula of the polymer formed.

.....[1]

[Total: 19]

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

								Gro	Group								
=												≡	2	>	I>	=>	₹
							- エ										2 He
Key	Key	Key	Key				hydrogen 1										helium 4
4 atomic number	atomic number	atomic number	tomic number									2	9	7	80	6	10
Be atomic symbol	atomic symbol	atomic symbol	mic symbol	\overline{g}								Ω	ပ	z	0	ш	Ne
beryllium name 9 relative atomic mass	name relative atomic mass	name relative atomic mass	name ive atomic mass	SS								boron 11	carbon 12	nitrogen 14	oxygen 16	fluorine 19	neon 20
12												13	14	15	16	17	18
Mg												Αl	S	۵	S	Cl	Ā
magnesium 24												aluminium 27	silicon 28	phosphorus 31	sulfur 32	chlorine 35.5	argon 40
21 22 23	22 23	23		24		25	26	27	28	29	30	31	32	33	34	35	36
> i	> =	>		ပ်	_	Mn	Fe	ပိ	Z	D C	Zn	Ga	Ge	As	Se	ă	첫
calcium scandium titanium vanadium chromium 40 45 48 51 52	titanium vanadium 6	vanadium 51		chromit 52	Ę	manganese 55	iron 56	cobalt 59	nickel 59	copper 64	zinc 65	gallium 70	germanium 73	arsenic 75	selenium 79	bromine 80	krypton 84
39 40 41	40 41	41		42		43	44	45	46	47	48	49	20	51	52	53	54
Y Zr Nb	Zr Nb	S S		Ĭ	0	ပ	R	格	Pd	Ag	g	In	Sn	Sb	Te	П	Xe
zirconium niobium 91 93	zirconium niobium 91 93	niobium 93		molybde 96	mnu	technetium -	ruthenium 101	rhodium 103	palladium 106	silver 108	cadmium 112	indium 115	tin 119	antimony 122	tellurium 128	iodine 127	xenon 131
57–71 72 73	72 73	73		1/2	Ţ	75	92	77	78	79	80	81	82	83	84	85	98
lanthanoids Hf Ta	HTTa	Та		>	>	Re	SO	'n	చ	Αn	Нg	l1	Pp	Bi	Ъо	¥	R
hafnium tantalum 178 181	tantalum 181	tantalum 181		tun	tungsten 184	rhenium 186	osmium 190	iridium 192	platinum 195	gold 197	mercury 201	thallium 204	lead 207	bismuth 209	molonium —	astatine	radon
89–103 104 105	104 105	105		10	9	107	108	109	110	111	112		114		116		
actinoids Rf Db	Rf	Dp		0)	Sg	Bh	Η̈́	Ĭ	Ds		ű		F1		_		
radium rutherfordium dubnium seabo	dubnium	dubnium		seapo	seaborgium	bohrium	hassium	meitnerium	darmstadtium	8	copernicium		flerovium		livermorium		
ı	ı	ı			_ 	ı	ı	ı	ı		ı		ı		ı		

7.1	Γn	Iutetium	175	103	۲	lawrencium	I
	Υp					_	
69	H	thulium	169	101	Md	mendelevium	1
89	ш	erbinm	167	100	Fm	ferminm	1
29	웃	holmium	165	66	Es	einsteinium	I
99	۵	dysprosium	163	86	ర్	californium	1
65	q	terbium	159	97	BK	berkelium	1
64	Gd	gadolinium	157	96	Cm	curium	1
63	En	europium	152	92	Am	americium	1
62	Sm	samarium	150	94	Pu	plutonium	I
61	Pm	promethium	I	93	ď	neptunium	ſ
09	pN	neodymium	144	92	\supset	uranium	238
29	Ā	praseodymium	141	91	Ра	protactinium	231
58	Ce	cerium	140	06	드	thorium	232
22	Гa	lanthanum	139	89	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).