

## B-splines

- A spline function of order  $m$  and knots  $\xi_1, \dots, \xi_k$  is a piecewise polynomial that can be expressed as a linear combination of the functions  $x^0, \dots, x^{m-1}, (x - \xi_1)_+^{m-1}, \dots, (x - \xi_k)_+^{m-1}$ , where

$$(x - \xi_i)_+^{m-1} = \begin{cases} (x - \xi_i)^{m-1} & \text{if } x \geq \xi_i; \\ 0 & \text{otherwise,} \end{cases}$$

for  $i = 1, \dots, k$ .

- Plot  $f(x) = (x - 0.5)_+^2$  for  $x \in [0, 1]$ .

```
f <- function(x, a=0.5){
  ans <- (x-a)^2
  ans[x<a] <- 0
  return(ans)
}
curve(f,0,1)
```

- A spline function of order  $m$  and inner knots  $\xi_1, \dots, \xi_k$  on an interval  $[a, b]$  can be expressed as a linear combination of the B-spline basis functions on  $[a, b]$  of order  $m$  with inner knots  $\xi_1, \dots, \xi_k$ .
- Each B-spline basis function of order  $m$  is characterized by a set of knots  $y_1, \dots, y_{m+1}$  (arranged in ascending order), which is denoted by  $N(\cdot|y_1, \dots, y_{m+1})$ . To compute B-spline basis functions, the following recursive formulas are used.

$$\begin{aligned} N(x|y_1, \dots, y_{m+1}) \\ = \frac{x - y_1}{y_m - y_1} N(x|y_1, \dots, y_m) + \frac{y_{m+1} - x}{y_{m+1} - y_2} N(x|y_2, \dots, y_{m+1}). \end{aligned}$$

$$N(x|y_1, y_2) = \begin{cases} 1 & \text{if } x \in [y_1, y_2); \\ 0 & \text{otherwise.} \end{cases}$$

$$N(x|\underbrace{y_1, \dots, y_1}_{m \text{ times}}, y_2) = \begin{cases} (y_2 - x)^{m-1} / (y_2 - y_1)^{m-1} & \text{if } x \in [y_1, y_2); \\ 0 & \text{otherwise.} \end{cases}$$

$$N(x|y_1, \underbrace{y_2, \dots, y_2}_{m \text{ times}}) = \begin{cases} (x - y_1)^{m-1} / (y_2 - y_1)^{m-1} & \text{if } x \in [y_1, y_2); \\ 0 & \text{otherwise.} \end{cases}$$

- B-spline basis functions on  $[a, b]$  of order  $m$  with knots  $\xi_1, \dots, \xi_k$ . Let  $y_1, \dots, y_{2m+k}$  be the sequence

$$\underbrace{a, \dots, a}_{m \text{ times}}, \xi_1, \dots, \xi_k, \underbrace{b, \dots, b}_{m \text{ times}}$$

then the  $m+k$  functions  $N(\cdot|y_1, \dots, y_{m+1}), \dots, N(\cdot|y_{m+k}, \dots, y_{2m+k})$  are the B-spline basis functions on  $[a, b]$  of order  $m$  with knots  $\xi_1, \dots, \xi_k$ .

- Note. The B-spline basis functions sum up to 1 and each one takes values in  $[0, 1]$ .
- The function `splineDesign` in R Package `splines` can be used for computing a single B-spline basis function  $N(\cdot|y_1, \dots, y_{m+1})$ .
  - Suppose that  $\mathbf{y} = (y_1, \dots, y_{m+1})$  and  $\mathbf{x} = (x_1, \dots, x_n)$  are two vectors in R, then

`splineDesign(y, x, ord=length(y)-1, outer.ok=TRUE)[,1]`

gives the vector  $(N(x_1|y_1, \dots, y_{m+1}), \dots, N(x_n|y_1, \dots, y_{m+1}))$ .

- Example 1. Check whether `splineDesign` gives

$$N(x|0, 0, 0, 1) = \begin{cases} (1-x)^{3-1}/(1-0)^{3-1} = (1-x)^2 & \text{if } x \in [0, 1]; \\ 0 & \text{otherwise.} \end{cases}$$

R commands:

```
require("splines")
y <- c(0,0,0,1)
f <- function(x){
  return(splineDesign(y, x, ord=length(y)-1, outer.ok=TRUE)[,1])
}
curve(f,0,1)
```

```
n0001 <- function(x){ (1-x)^2 }
curve(n0001, 0, 1, add=T, col=2) #Plot the N(|0,0,0,1) function on [0,1]
```

- The function `bs` in R Package `splines` can be used for computing the  $m+k$  B-spline functions on  $[a, b]$  of order  $m$  with knots  $\xi_1, \dots, \xi_k$ .

- Suppose that  $\mathbf{knots} = (\xi_1, \dots, \xi_k)$  and  $\mathbf{x} = (x_1, \dots, x_n)$  are vectors in  $\mathbb{R}$ , then B-spline basis functions on  $[a, b]$  of order  $m$  with knots  $\xi_1, \dots, \xi_k$  evaluated at  $\mathbf{x}$  form a  $n \times (m+k)$  matrix  $X$ , where the  $j$ -th column of  $X$  is the  $j$ -th B-spline basis function evaluated at  $\mathbf{x}$ . The matrix  $X$  can be computed using the R command

```
bs(x, knots=knots, deg=m-1, Boundary.knots=c(a,b), intercept=TRUE)
```

- Example 2. Compute the B-spline basis functions on  $[0, 1]$  of order three with knots 0.1 and 0.2 using `bs` and `splineDesign`. Plot the five B-spline basis functions.

```
x <- (1:1000)/1001
knotlist=c(0.1, 0.2)
ord=3
knot_all=c(rep(0,ord), knotlist, rep(1,ord))
nb = length(knotlist)+ord
plot(x,x,type="n",ylab="")
s=0
for ( i in 1:nb){
  y = knot_all[i:(i+ord)]
  f <- function(x){
    n <- length(x)
    return(splineDesign(y, x, ord=length(y)-1, outer.ok=T)[,1])
  }
  f1 <- function(x){
    bx=bs(x, deg=ord-1, knots=knotlist, Boundary.knots=c(0,1), intercept=T)
    return(bx[,i])
  }
  lines(x,f(x), type="l"); lines(x, f1(x), col=i+1)
  s=s+sum(abs(f(x)-f1(x)))
}
s
```

- Spline functions can approximate smooth functions well if the number of knots are large enough.

- Exercise 1. Let  $B_1, \dots, B_{m+k}$  be the B-spline basis functions on  $[0, 1]$  of order  $m$  with knots  $\xi_1, \dots, \xi_k$ . Check whether each of  $B_1(x), \dots, B_{m+k}(x)$  can be expressed as linear combinations of the functions  $1, x, \dots, x^{m-1}, (x - \xi_1)_+^{m-1}, \dots, (x - \xi_k)_+^{m-1}$  for  $x = (1:1000)/1001$ ,  $m = 4$ ,  $k = 3$ , and  $(\xi_1, \dots, \xi_3) = (1:3)/4$ .
- Exercise 2. Let  $f(x) = x \sin(20x)$  for  $x \in [0, 1]$ . Suppose that  $n = 1000$ ,  $(X_1, \dots, X_n) = \text{seq}(0, 1, \text{length}=n)$ , and  $Y_i = f(X_i)$  for  $i = 1, \dots, n$ .
  - (a) Find the ISE of estimating  $f$  based on  $(X_1, Y_1), \dots, (X_n, Y_n)$  by approximating  $f$  using the best linear combination of B-spline basis functions on  $[0, 1]$  of order 4 with  $k$  equally spaced knots  $((1:k)/(k+1))$  for  $k = 1, 2, \dots, 7$ . Which  $k$  gives the best ISE?
  - (b) Consider estimating  $f$  based on  $(X_1, Y_1), \dots, (X_n, Y_n)$  by approximating  $f$  using the best linear combination of  $1, x, \dots, x^m$ . Let  $m_0$  be the smallest  $m$  such that the resulting ISE is smaller than the best ISE found in Part (a). Find  $m_0$ .
- Exercise 3. Let  $f(x) = x \sin(20x)$  for  $x \in [0, 1]$ . Suppose that  $n = 1000$ ,  $(X_1, \dots, X_n) = \text{seq}(0, 1, \text{length}=n)$ , and  $Y_i = f(X_i) + \varepsilon_i$  for  $i = 1, \dots, n$ , where  $\varepsilon_i$ 's are IID  $N(0, \sigma^2)$  variables with  $\sigma = 0.2$ . Consider estimating  $f$  based on  $(X_1, Y_1), \dots, (X_n, Y_n)$  by approximating  $f$  using
  - (i) B-spline basis functions on  $[0, 1]$  of order 4 with  $k$  equally spaced knots and
  - (ii) polynomial basis functions  $1, x, \dots, x^m$ ,
 where the number of knots  $k$  is the  $k$  that gives the best ISE in Part (a) in Exercise 2, and  $m$  is  $m_0$  in Part (b) in Exercise 2. Find the IMSEs for (i) and (ii) based on 10000 simulations. Use `set.seed` to make sure that the simulation data for (i) and (ii) are the same.
- Exercise 4. Let  $f(x) = x \sin(20x)$  for  $x \in [0, 1]$ . Suppose that  $n = 1000$ ,  $(X_1, \dots, X_n) = \text{seq}(0, 1, \text{length}=n)$ , and  $Y_i = f(X_i) + \varepsilon_i$  for  $i = 1, \dots, n$ , where  $\varepsilon_i$ 's are IID  $N(0, \sigma^2)$  variables with  $\sigma = 0.2$ . Consider estimating  $f$  based on  $(X_1, Y_1), \dots, (X_n, Y_n)$  by approximating  $f$  using
  - (i) B-spline basis functions with order 4 and  $k$  equally spaced knots and

(ii) polynomial basis functions  $1, x, \dots, x^m$ ,

where  $k$  and  $m$  are chosen using cross-validation. The selection ranges for  $k$  and  $m$  are  $\{1, 2, \dots, 7\}$  and  $\{1, 2, \dots, 12\}$ . Find the IMSEs for (i) and (ii) based on 10000 simulations. Use `set.seed` to make sure that the simulation data for (i) and (ii) are the same.