

## 81

Examples of the aryl group included in the aryl sulfonate include a phenyl group, a naphthyl group, and an anthryl group. These aryl groups may have a substituent. As the substituent, for example, a linear or branched alkyl group having 1 to 6 carbon atoms or a cycloalkyl group having 3 to 6 carbon atoms is preferable. Specifically, for example, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an i-butyl group, a t-butyl group, an n-hexyl group, or a cyclohexyl group is preferable. Examples of other substituents include an alkoxy group having 1 to 6 carbon atoms, a halogen atom, a cyano group, a nitro group, an acyl group, and an acyloxy group.

The carboxylate may be an aliphatic carboxylate or an aromatic carboxylate, and examples thereof include acetate, lactate, pyruvate, trifluoroacetate, adamantane carboxylate, hydroxyadamantane carboxylate, benzoate, naphthoate, salicylate, phthalate, and phenolate, and, in particular, benzoate, naphthoate, or phenolate is preferable, and benzoate is the most preferable.

In this case, the ammonium salt is preferably tetra(n-butyl)ammonium benzoate, tetra(n-butyl)ammonium phenolate, or the like.

In a case where the ammonium salt is hydroxide, the ammonium salt is particularly preferably tetraalkylammonium hydroxide (tetraalkylammonium hydroxide such as tetramethylammonium hydroxide, tetraethylammonium hydroxide, and tetra-n-butylammonium hydroxide) having 1 to 8 carbon atoms.

(Q5) Low Molecular Compound Having Nitrogen Atom and Group Capable of Leaving by Action of Acid

The actinic ray-sensitive or radiation-sensitive resin composition used in the present invention can include a low molecular compound having a nitrogen atom and a group capable of leaving by the action of an acid. The compound preferably has basicity after the group capable of leaving by the action of an acid leaves.


With regard to the compound, reference can be made to the description in paragraphs [0324] to [0337] of JP2012-133331A, the contents of which are incorporated herein.

In the present invention, the compound may be used singly or as a mixture of two or more kinds thereof.

Other examples of the basic compound which can be used in the actinic ray-sensitive or radiation-sensitive resin composition include the compounds synthesized in Examples of JP2002-363146A and the compounds described in paragraph 0108 of JP2007-298569A.

A photosensitive basic compound may also be used as the basic compound (C'). As the photosensitive basic compound, for example, the compounds described in JP2003-524799A, J. Photopolym. Sci. & Tech., Vol. 8, P. 543 to 553 (1995), or the like can be used.


Preferred specific examples of the basic compound (C') include the following compounds, but are not limited thereto.




C-1

## 82


-continued




C-2




C-3




C-4



C-5



C-6



C-7

The molecular weight of the basic compound (C') is typically 100 to 1,500, preferably 150 to 1,300, and more preferably 200 to 1,000.

These basic compounds (C') may be used singly or in combination of thereof.

The actinic ray-sensitive or radiation-sensitive resin composition used in the present invention may or may not include the basic compound (C'). However, in a case where the basic compound (C') is included in the composition, the