

- ment effects when only a subset of patients can be expected to respond to treatment,” *Biometrics*, 44, 189-196.
- Deshpande, S.K., Hasegawa, R.B., Rabinowitz, A.R., Whyte, J., Roan, C.L., Tabatabaei, A., Baiocchi, M., Karlawish, J.H., Master, C.L., and Small, D.S. (2017), “High school football and later life cognition and mental health: An observational study,” *JAMA Neurology*, 74, 909-918. Protocol for study 2016: arXiv preprint arXiv:1607.01756.
- Fisher, R. A. (1935), *The Design of Experiments*, Edinburgh: Oliver & Boyd.
- Graves, A. B., White, E., Koepsell, T., Reier, B. V., Van Belle, G., Larson, E. B., and Raskind, M. (1990), “The association between head trauma and Alzheimer’s disease,” *American Journal of Epidemiology*, 131, 491-501.
- Hansen, B. B. (2004), “Full matching in an observational study of coaching for the SAT,” *Journal of the American Statistical Association*, 99, 609-618.
- Hansen, B. B. and Klopfer, S. O. (2006), “Optimal full matching and related designs via network flows,” *Journal of Computational and Graphical Statistics*, 15, 609-627. (R package `optmatch`)
- Hansen, B. B. (2007), “Flexible, optimal matching for observational studies,” *R News*, 7, 18-24. (R package `optmatch`)
- Hsu, J. Y. and Small, D. S. (2013), “Calibrating sensitivity analyses to observed covariates in observational studies,” *Biometrics*, 69, 803-811.
- Knopman, D. S. and Ryberg, S. (1989), “A verbal memory test with high predictive accuracy for dementia of the Alzheimer type,” *Archives of Neurology*, 46, 141-145.
- Laska, E. M. and Meisner, M. J. (1989), “Testing whether an identified treatment is best,” *Biometrics*, 45, 1139-1151.
- Lehman, E. J., Hein, M. J., Baron, S. L., and Gersic, C. M. (2012), “Neurodegenerative causes of death among retired national football league players,” *Neurology*, 79, 1970-