Hypra: Mathematical Contract and Acceptance Derivation

Leandro Almeida

Mathematical contract and exact acceptance derivation

Scope and provenance

Hypra is a **partial native Python/SymPy port** for exact regulator expansions, with three mathematically proved hypergeometric families: an integer-parameter HPL-recursion family, an even-binomial half-integer family, and a telescoping half-integer family. The native backend is the default and **requires no Mathematica installation or setup**. The derivations and limits of these three families are given below; unsupported families must fail explicitly rather than be presented as complete expansions.

HypExp [1] expands generalized hypergeometric functions around integer-valued parameters. HypExp 2 [2] adds certain half-integer classes. Hypra does not yet match their full coverage. An optional external HypExp adapter is retained as an experimental, **unverified** alternative; it is neither required by the native implementation nor evidence that a real Wolfram/HypExp run succeeded.

The calculations below independently derive the requested example. The NNLO script at `reports/infrared-consistency-review/derive_identical_two_denominator.py` was read as context only; its files were not changed and its table of logarithmic integrals is **not** imported. `scripts/validate_integral.py` derives the moments by convergent sums, a beta-function derivative and the dilogarithm reflection identity. Its offline success does not verify a Mathematica/HypExp installation.

Conventions

- The expansion variable is a dimensionless regulator ϵ, expanded at zero.
- `through=k` means every Laurent term through and **including** ϵ^k, plus a remainder $O(\epsilon^{k+1})$. Pole terms must not be discarded.
- Inputs and coefficients are exact. A binary floating-point approximation is not a substitute for a rational parameter.
- The derivation in this document uses **real** $0 < w < 1$, principal real logarithms and classical polylogarithms. Continuation outside this interval requires a branch prescription; the formulas must not be silently reused across cuts.
- An HPL object retains its complete index word and argument. Under the usual real alphabet convention, $f_0(t) = 1/t$, $f_1(t) = 1/(1-t)$ and $f_{-1}(t) = 1/(1+t)$, with $H_0(z) = \log z$, $H_1(z) = -\log(1-z)$, and $H_{-1}(z) = \log(1+z)$. Compact nonzero index m denotes $|m| - 1$ zeros followed by $\mathrm{sgn}(m)$, not a different kernel. Complex continuation, extended alphabets, and endpoint regularization are separate matters; a wrapper must not invent their conventions.

Native all-order HPL recursion for an integer family

The native implementation is based on a differential equation, not stored acceptance coefficients. Let

$$F(z,\epsilon) = {}_2F_1(1, b\epsilon; 1 + c\epsilon; z) = 1 + \sum_{k\geq 1} \epsilon^k f_k(z),$$

where b, c are regulator-independent exact constants. Its argument-series coefficient $a_n = (b\epsilon)_n/(1 + c\epsilon)_n$ satisfies $(n + c\epsilon)a_n = (n - 1 + b\epsilon)a_{n-1}$ for $n \geq 1$. Summing this identity gives

$$z(1 - z)\partial_z F + \epsilon(c - bz)F = c\epsilon, \qquad F(0,\epsilon) = 1.$$

At order one $f_1' = b/(1 - z)$, hence $f_1 = bH_1(z)$. At every higher order,

$$f_k'(z) = \left[-\frac{c}{z} + \frac{b - c}{1 - z}\right] f_{k-1}(z), \qquad f_k(0) = 0.$$

With $\mathcal{I}_a$ denoting the operation of prepending letter a to an HPL word, this is the constructive recurrence

$$\boxed{f_1 = bH_1, \qquad f_k = -c\,\mathcal{I}_0 f_{k-1} + (b - c)\,\mathcal{I}_1 f_{k-1} \quad (k \geq 2).}$$

Every word ends in 1, so its origin value is zero and these integrals require no divergent origin regularization. Equivalently, a word of length k ending in 1, with n_0 zero letters and n_1 one letters, has coefficient $b(-c)^{n_0}(b - c)^{n_1-1}$. This proves every order by induction and uniqueness of the regular solution, not only the first two acceptance orders. It covers P with $(b, c) = (-1, -2)$ and Q with $(b, c) = (-2, -2)$ and argument $1 - w$.

The derivation starts on $|z| < 1$ near $\epsilon = 0$. For real $0 < z < 1$ it uses the unambiguous real HPL branch. Analytic continuation follows only with matching HPL path/branch conventions; no branch cut crossing is implicit. The offline recurrence tests compare orders 1 through 5, at argument powers 1 through 6, against exact Pochhammer quotients for four choices of (b, c). The proof, not that finite test sample, establishes the all-order recurrence.

Two narrowly supported half-integer families

These identities do **not** establish general half-integer HypExp 2 coverage. Let $t = \sqrt{z}$ with real $0 < z < 1$.

Even-binomial family

For a independent of the regulator,

$$G(z,\epsilon) = {}_2F_1(a\epsilon, \tfrac{1}{2} + a\epsilon; \tfrac{1}{2}; z) = \frac{(1 - t)^{-2a\epsilon} + (1 + t)^{-2a\epsilon}}{2}.$$

To prove it, take the even terms of the binomial series and use $(2A)_{2n} = 2^{2n}(A)_n(A + 1/2)_n$ and $(2n)! = 2^{2n}n!(1/2)_n$, with $A = a\epsilon$. Thus

$$[\epsilon^k]G = \frac{(-2a)^k}{2k!}\left[\log^k(1-t)+\log^k(1+t)\right], \quad k \geq 0.$$

The expression is even in t and has a regular $z = 0$ limit. Its derivation from the binomial identity is exact, not a fitted expansion. No cancellation of upper and lower hypergeometric parameters at nonzero regulator was used.

Half-integer telescoping family

For $A = 1/2 + b\epsilon$,

$$T(z,\epsilon) = {}_2F_1(1, A; 1+A; z) = \sum_{n\geq 0} \frac{A}{n+A} z^n.$$

Define

$$S_k(z) = \sum_{n\geq 0} \frac{z^n}{(n+1/2)^k} = \frac{2^{k-1}}{t}\left[\mathrm{Li}_k(t) - \mathrm{Li}_k(-t)\right].$$

The equality follows by extracting odd terms of the polylogarithm series. Since $A/(n+A) = 1 - n/(n+A)$, direct geometric expansion of its denominator gives

$$T_0 = \frac{\log(1+t)-\log(1-t)}{2t}, \qquad [\epsilon^k]T = (-1)^{k-1}b^k\left[S_k - \frac{1}{2}S_{k+1}\right] \quad (k \geq 1).$$

This includes the $n = 0$ term correctly: it contributes 1 at order zero and zero at every higher order. For each fixed finite b, the expansion is local in ϵ, before the nearest denominator pole. At $z = 0$ the displayed quotients have removable limits ($T_0 = 1$, higher coefficients zero), whereas $z = 1$ is divergent. An implementation must handle those endpoints explicitly, not evaluate a spurious $0/0$ or silently return a finite value at 1.

P and Q from their defining series

Set

$$P(w,\epsilon) = {}_2F_1(1, -\epsilon; 1-2\epsilon; w), \qquad Q(w,\epsilon) = {}_2F_1(1, -2\epsilon; 1-2\epsilon; 1-w).$$

The upper parameter -2ϵ in Q is **not** the lower parameter $1-2\epsilon$; no upper/lower cancellation applies.

For $n \geq 1$, using $(1)_n/n! = 1$,

$$\frac{(-\epsilon)_n}{(1-2\epsilon)_n} = -\frac{\epsilon}{n}\left[1 + \epsilon\left(H_{n-1} + \frac{2}{n}\right) + O(\epsilon^2)\right].$$

This follows by writing the rising factorials as products and differentiating $\sum_{j=1}^{n-1}\log(1-\epsilon/j) - \sum_{j=1}^{n}\log(1-2\epsilon/j)$. On $|w| < 1$,

$$\sum_{n\geq 1}\frac{w^n}{n}=-\log(1-w),\quad \sum_{n\geq 1}\frac{w^n}{n^2}=\mathrm{Li}_2(w),\quad \sum_{n\geq 1}\frac{H_{n-1}w^n}{n}=\frac{1}{2}\log^2(1-w).$$

The last identity follows by differentiating both sides, using $\sum_{n\geq 1} H_{n-1}w^{n-1}=-\log(1-w)/(1-w)$, and the value zero at $w=0$. Consequently

$$P=1+\epsilon\log(1-w)+\epsilon^2\left[-2\,\mathrm{Li}_2(w)-\frac{1}{2}\log^2(1-w)\right]+O(\epsilon^3).$$

For Q there is an exact telescoping coefficient:

$$\frac{(-2\epsilon)_n}{(1-2\epsilon)_n}=-\frac{2\epsilon}{n-2\epsilon}.\qquad Q=1-\sum_{j\geq 1}(2\epsilon)^j\,\mathrm{Li}_j(1-w).$$

The double-series interchange is justified for fixed $0<w<1$ and $|2\epsilon|<1$ by absolute convergence. Hence

$$Q=1+2\epsilon\log w-4\epsilon^2\,\mathrm{Li}_2(1-w)+O(\epsilon^3).$$

The finite-n coefficient identities are checked exactly by the offline tests; they are not numerically fitted.

Why the J4 expansion may be integrated

Define

$$J_4(\epsilon)=\int_0^1 w^{-1-2\epsilon}(1-w)^{-\epsilon}(P-1)(Q-1)\,dw.$$

Fix a sufficiently small closed complex disk $|\epsilon|\leq r<1/4$. All hypergeometric denominator parameters avoid their singular values there. The exact Q coefficients imply $|Q-1|\leq C_r|\epsilon||\log w|$ near $w=0$ and $|Q-1|\leq C_r|\epsilon|(1-w)$ near $w=1$. For P, its coefficient can be written

$$-\epsilon\frac{\Gamma(1-2\epsilon)}{\Gamma(1-\epsilon)}\frac{\Gamma(n-\epsilon)}{\Gamma(n+1-2\epsilon)}.$$

Uniform gamma-ratio asymptotics on this compact regulator disk give $|p_n|\leq C_r|\epsilon|n^{r-1}$. Therefore $P-1=O_r(|\epsilon|w)$ near zero and $P-1=O_r(|\epsilon|(1-w)^{-r})$ near one. The full integrand is bounded by constants times

$$w^{-2r}|\log w|\quad (w\to 0),\qquad (1-w)^{1-2r}\quad (w\to 1),$$

both integrable. Dominated holomorphic parameter integration and Cauchy's formula on a smaller disk permit differentiation under this integral to any fixed finite order. Thus the Taylor coefficients computed below are legitimate integrals; the integrated remainder is $O(\epsilon^4)$. This proof is specific to J_4, **not** permission to expand each unsubtracted singular term in a larger NNLO integral separately.

Deriving all needed moments, not importing a table

All integrals in this subsection have measure dw/w on $(0,1)$. Write $L = \log w$, $M = \log(1-w)$ and $\zeta_k = \zeta(k)$. Expanding the logarithm or dilogarithm and integrating their absolutely integrable terms gives

$$\int LM = \sum_{n\geq 1} \frac{1}{n^3} = \zeta_3, \qquad \int L\,\mathrm{Li}_2(w) = -\sum_{n\geq 1} \frac{1}{n^4} = -\zeta_4,$$

$$\int L^2 M = -2\sum_{n\geq 1} \frac{1}{n^4} = -2\zeta_4.$$

Here $\int_0^1 w^{n-1}\log w\,dw = -1/n^2$ and $\int_0^1 w^{n-1}\log^2 w\,dw = 2/n^3$. The absolute sums converge.

For $B = \int LM^2$, use $\mathrm{B}(a, 1+t) - 1/a$, analytic at $(a,t) = (0,0)$, and take $\partial_a \partial_t^2$. The relevant generating function is

$$\mathrm{B}(a,1+t) = \frac{1}{a}\exp\left\{\sum_{k\geq 2} \frac{(-1)^k \zeta_k}{k}[a^k + t^k - (a+t)^k]\right\}.$$

The a^2t^2 coefficient of the exponential is $-3\zeta_4/2 + \zeta_2^2/2 = -\zeta_4/4$. After division by a and the two t derivatives, $B = -\zeta_4/2$. Differentiation is justified either after subtracting $1/a$ or directly for the mixed derivative, whose logarithmic integrand is integrable.

Finally, with $C = \int M\,\mathrm{Li}_2(1-w)$, the real reflection identity $\mathrm{Li}_2(w) + \mathrm{Li}_2(1-w) = \zeta_2 - LM$ gives

$$C = -\zeta_2^2 + \frac{1}{2}\zeta_2^2 - B = -\frac{1}{2}\zeta_2^2 - B = -\frac{3}{4}\zeta_4.$$

We used $d\,\mathrm{Li}_2(w) = -M\,dw/w$ and its endpoint values; all boundary terms are finite.

Assemble the exact coefficients

Expand the weight as $w^{-1}[1 + \epsilon(-2L - M) + O(\epsilon^2)]$. Multiplying the derived series gives

$$[\epsilon^2]J_4 = \int 2LM = 2\zeta_3,$$

$$[\epsilon^3]J_4 = \int [-4L\,\mathrm{Li}_2(w) - 3LM^2 - 4M\,\mathrm{Li}_2(1-w) - 4L^2M] = \left(4 + \frac{3}{2} + 3 + 8\right)\zeta_4 = \frac{33}{2}\zeta_4.$$

Thus the requested result is consistent:

$$\boxed{J_4 = 2\zeta(3)\epsilon^2 + \frac{11\pi^4}{60}\epsilon^3 + O(\epsilon^4).}$$

The script creates the polynomial product and integrates each monomial using the independently derived moments. It rejects any unproved monomial rather than supplying a guessed value.

Laurent prefactors and order budgeting

If a prefactor A starts at ϵ^{-p}, an analytic factor F is generally needed through order $k+p$ to obtain AF through order k. For example, $\Gamma(\epsilon)^2 F$ needs F through ϵ^{k+2}, not merely ϵ^k. Multiplication, exact cancellations and the final truncation must be performed on the **whole expression**. If a backend cannot establish the valuation/order budget or its result still contains an unexpanded hypergeometric function, a regulator derivative or a regulator-dependent coefficient, it must fail rather than label the result complete. Assumptions needed to establish a valuation or cancel a parameter are part of the mathematical input.

Validation and limits

```
uv run python scripts/validate_integral.py
uv run python scripts/validate_integral.py --latex
uv run python scripts/validate_integral.py --numeric
uv run pytest tests/test_mathematics.py
```

The optional numerical check evaluates the original integral using contiguous representations for $P-1$ and $Q-1$ to avoid catastrophic subtraction near the endpoints. It uses negative regulator samples, for which endpoint limits are zero, and introduces **no endpoint cutoff**. Convergence of the residual scaled by ϵ^4 is secondary support only, not the analytic proof and not a real-backend validation. No global continuation or arbitrary-HPL simplifier is established by these examples.

References

1. T. Huber and D. Maître, *HypExp, a Mathematica package for expanding hypergeometric functions around integer-valued parameters*, arXiv:hep-ph/0507094.
2. T. Huber and D. Maître, *HypExp 2, Expanding Hypergeometric Functions about Half-Integer Parameters*, arXiv:0708.2443.

The abstract pages were checked for these scope statements. No source code from these packages is redistributed with this derivation.