

Cheatsheet - Trigonometry

Fabio Lama – fabio.lama@pm.me

1. Intro

If $B = 90$, then AC is called the **hypotenuse**, the side **opposite** of θ is BC and the remaining side, AB , is **adjacent** to θ . Angle A is written as $\angle BAC$, $\angle A$ or just A .

Each side of a triangle is named based on the lowercase letter of the angle opposite of the side.

$$a = BC$$

$$b = AC$$

$$c = AB$$

The total degree of all angles is always 180°

$$A + B + C = 180^\circ$$

2. Convert Degree to Radian

$$360^\circ = 2\pi \text{ rads}$$

To convert x degrees to radians y :

$$x \cdot \frac{\pi}{180} = y$$

For example, 255° :

$$255 \cdot \frac{\pi}{180} \approx 3.93 \text{ rads}$$

And in reverse:

$$3.93 \cdot \frac{180}{\pi} \approx 225^\circ$$

3. Trigonometrical Ratios

Sinus, *cosine* and *tangent* are trigonometric functions that can be applied to **right-angled triangles** to determine the degree of an angle. The degree of an angle is independent from the length of each side.

$$\sin \theta = \frac{\text{side opposite to } \theta}{\text{hypotenuse}} = \frac{BC}{AC}$$

$$\cos \theta = \frac{\text{side adjacent to } \theta}{\text{hypotenuse}} = \frac{AB}{AC}$$

$$\tan \theta = \frac{\text{side opposite to } \theta}{\text{side adjacent to } \theta} = \frac{BC}{AB}$$

Sinus and *cosine* are always less than one.

$$\sin x \leq 1$$

$$\cos x \leq 1$$

The ratio of an angle can be reversed.

$$\sin 45^\circ = 0.70710678$$

$$\sin^{-1} 0.70710678 = 45^\circ$$

4. Pythagoras, Sinus and Cosine Rule

For any **right-angled** triangle, the following theorem applies:

$$a^2 + b^2 = c^2$$

$$\sqrt{a^2 + b^2} = c$$

where c is the hypotenuse. For any right-angled, acute and obtuse (non right-angled) triangles, the sinus and cosine rules apply:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

and:

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Last updated 2022-07-16 17:46:41 UTC