

# Probability Theory 1

Raazesh Sainudiin\* and Dominic Lee†,

\*Laboratory for Mathematical Statistical Experiments, Uppsala Centre, and

†Department of Mathematics, Uppsala University, Uppsala, Sweden

†School of Mathematics and Statistics, University of Canterbury, Christchurch, New Zealand

Version Date: September 27, 2019


©2007–2019 Raazesh Sainudiin. ©2008–2013 Dominic Lee.


This work is licensed under the Creative Commons Attribution-Noncommercial-Share Alike 4.0

International License. To view a copy of this license, visit

<https://creativecommons.org/licenses/by-nc-sa/4.0/>.

This work was partially supported by NSF grant DMS-03-06497 and NSF/NIGMS grant DMS-02-01037.





# Contents

<b>1 Preliminaries</b>	<b>9</b>
1.1 Elementary Set Theory . . . . .	9
1.2 Natural Numbers, Integers and Rational Numbers . . . . .	12
1.3 Real Numbers . . . . .	16
1.4 Introduction to MATLAB . . . . .	20
1.5 Permutations, Factorials and Combinations . . . . .	23
1.6 Array, Sequence, Limit, . . . . .	25
1.7 Elementary Number Theory . . . . .	31
<b>2 Probability Model</b>	<b>33</b>
2.1 Experiments . . . . .	33
2.2 Probability . . . . .	35
2.2.1 Consequences of our Definition of Probability . . . . .	37
2.2.2 Sigma Algebras of Typical Experiments* . . . . .	39
2.3 Conditional Probability . . . . .	41
2.3.1 Independence and Dependence . . . . .	44
<b>3 Random Variables</b>	<b>49</b>
3.1 Basic Definitions . . . . .	49
3.2 An Elementary Discrete Random Variable . . . . .	51
3.3 An Elementary Continuous Random Variable . . . . .	52
3.4 Transformations of random variables . . . . .	54
3.4.1 Transformation of discrete random variables . . . . .	56
3.4.2 Transformation of continuous random variables . . . . .	58
3.5 $\mathbb{R}^m$ -valued Random Variables . . . . .	62
3.5.1 $\mathbb{R}^2$ -valued Random Variables . . . . .	62
3.5.2 Conditional Random Variables . . . . .	72
3.5.3 $\mathbb{R}^m$ -valued Random Variables . . . . .	74
3.6 Expectations . . . . .	77
3.6.1 Expectations of functions of random variables . . . . .	77

<b>CONTENTS</b>	<b>4</b>
3.6.2 Properties of expectations . . . . .	80
3.7 Stochastic Processes . . . . .	83
3.8 Convergence of Random Variables . . . . .	87
3.9 Characteristic Functions . . . . .	92
3.9.1 Obtaining Moments from Characteristic Function . . . . .	93
<b>4 Statistics</b>	<b>107</b>
4.1 Data and Statistics . . . . .	107
4.2 Exploring Data and Statistics . . . . .	114
4.2.1 Univariate Data . . . . .	114
4.2.2 Bivariate Data . . . . .	116
4.2.3 Trivariate Data . . . . .	117
4.2.4 Multivariate Data . . . . .	117
4.3 Loading and Exploring Real-world Data . . . . .	119
4.3.1 Geological Data . . . . .	119
4.3.2 Metereological Data . . . . .	123
4.3.3 Textual Data . . . . .	126
4.3.4 Machine Sensor Data . . . . .	127
<b>5 Simulation</b>	<b>128</b>
5.1 Physical Random Number Generators . . . . .	128
5.2 Pseudo-Random Number Generators . . . . .	128
5.2.1 Linear Congruential Generators . . . . .	129
5.2.2 Generalized Feedback Shift Register and the “Mersenne Twister” PRNG . . . . .	132
5.3 Simulation of non-Uniform(0, 1) Random Variables . . . . .	134
5.3.1 Inversion Sampler for Continuous Random Variables . . . . .	134
5.3.2 Inversion Sampler for Discrete Random Variables . . . . .	143
5.3.3 von Neumann Rejection Sampler (RS) . . . . .	153
<b>6 Limit Laws of Statistics</b>	<b>159</b>
<b>7 Answers to Selected Exercises</b>	<b>162</b>

# List of Tables

1.1	Symbol Table: Sets and Numbers . . . . .	19
3.1	The 8 $\omega$ 's in the sample space $\Omega$ of the experiment $\mathcal{E}_\theta^3$ are given in the first row above. The RV $Y$ is the number of 'Heads' in the 3 tosses and the RV $Z$ is the number of 'Tails' in the 3 tosses. Finally, the RVs $Y'$ and $Z'$ are the indicator functions of the event that 'all three tosses were Heads' and the event that 'all three tosses were Tails', respectively. . . . .	73

# List of Figures

1.1	Union and intersection of sets shown by Venn diagrams . . . . .	10
1.2	These Venn diagram illustrate De Morgan's Laws. . . . .	11
1.3	A function $f$ ("father of") from $\mathbb{X}$ (a set of children) to $\mathbb{Y}$ (their fathers) and its inverse ("children of"). . . . .	14
1.4	A pictorial depiction of addition and its inverse. The domain is plotted in orthogonal <b>Cartesian coordinates</b> . . . . .	15
1.5	A depiction of the real line segment $[-10, 10]$ . . . . .	17
1.6	Point plot and stem plot of the finite sequence $\langle b_{1:10} \rangle$ declared as an array. . . . .	27
1.7	A plot of the sine wave over $[-2\pi, 2\pi]$ . . . . .	29
2.1	A binary tree whose leaves are all possible outcomes. . . . .	35
2.2	First ball number in 1114 NZ Lotto draws from 1987 to 2008. . . . .	38
3.1	The Indicator function of event $A \in \mathcal{F}$ is a RV $\mathbf{1}_A$ with DF $F$ . . . . .	50
3.2	A RV $X$ from a sample space $\Omega$ with 8 elements to $\mathbb{R}$ and its DF $F$ . . . . .	51
3.3	The Indicator Function $\mathbf{1}_H$ of the event 'Heads occurs', for the experiment 'Toss 1 times,' $\mathcal{E}_\theta^1$ , as the RV $X$ from the sample space $\Omega = \{H, T\}$ to $\mathbb{R}$ and its DF $F$ . The probability that 'Heads occurs' and that 'Tails occurs' are $f(1; \theta) = \mathbf{P}_\theta(X = 1) = \mathbf{P}_\theta(H) = \theta$ and $f(0; \theta) = \mathbf{P}_\theta(X = 0) = \mathbf{P}_\theta(T) = 1 - \theta$ , respectively. . . . .	53
3.4	A plot of the PDF and DF or CDF of the Uniform(0, 1) continuous RV $X$ . . . . .	54
3.5	Density and distribution functions of Exponential( $\lambda$ ) RVs, for $\lambda = 1, 10, 10^{-1}$ , in four different axes scales. . . . .	60
3.6	A plot of the PDF, DF or CDF and inverse DF of the Uniform(-1, 1) RV $X$ . . . . .	72
3.7	Mean ( $\mathbf{E}_\theta(X)$ ), variance ( $\mathbf{V}_\theta(X)$ ) and the rate of change of variance ( $\frac{d}{d\theta}\mathbf{V}_\theta(X)$ ) of a Bernoulli( $\theta$ ) RV $X$ as a function of the parameter $\theta$ . . . . .	82
3.8	PDF of $X \sim \text{Geometric}(\theta = 0.5)$ and the relative frequency histogram based on 100 and 1000 samples from $X$ according to Simulation 112 and Labwork 113. . . . .	85
3.9	Mean and variance of a Geometric( $\theta$ ) RV $X$ as a function of the parameter $\theta$ . . . . .	85
3.10	PDF of $X \sim \text{Binomial}(n = 10, \theta = 0.5)$ and the relative frequency histogram based on 100,000 samples from $X$ obtained according to Simulation 116. . . . .	87
3.11	Distribution functions of several Normal( $\mu, \sigma^2$ ) RVs for $\sigma^2 = 1, \frac{1}{10}, \frac{1}{100}, \frac{1}{1000}$ . . . . .	88
3.12	PDF of $X \sim \text{Poisson}(\lambda = 10)$ and the relative frequency histogram based on 1000 samples from $X$ according to Simulation 117. . . . .	92

4.1	Sample Space, Random Variable, Realisation, Data, and Data Space. . . . .	107
4.2	Data Spaces $\mathbb{X} = \{0, 1\}^2$ and $\mathbb{X} = \{0, 1\}^3$ for two and three Bernoulli trials, respectively.	108
4.3	Plot of the DF of $\text{Uniform}(0, 1)$ , five IID samples from it, and the ECDF $\widehat{F}_5$ for these five data points $x = (x_1, x_2, x_3, x_4, x_5) = (0.5164, 0.5707, 0.0285, 0.1715, 0.6853)$ that jumps by $1/5 = 0.20$ at each of the five samples. . . . .	113
4.4	Frequency, Relative Frequency and Density Histograms . . . . .	115
4.5	Frequency, Relative Frequency and Density Histograms . . . . .	116
4.6	2D Scatter Plot . . . . .	117
4.7	3D Scatter Plot . . . . .	118
4.8	Plot Matrix of uniformly generated data in $[0, 1]^5$ . . . . .	118
4.9	Matrix of Scatter Plots of the latitude, longitude, magnitude and depth of the 22-02-2011 earth quakes in Christchurch, New Zealand. . . . .	121
4.10	Google Earth Visualisation of the earth quakes . . . . .	122
4.11	Daily rainfalls in Christchurch since March 27 2010 . . . . .	124
4.12	Daily temperatures in Christchurch for one year since March 27 2010 . . . . .	125
4.13	Wordle of JOE 2010 . . . . .	126
4.14	Double Pendulum . . . . .	127
5.1	The linear congruential sequence of <code>LinConGen(256, 137, 0, 123, 257)</code> with non-maximal period length of 32 as a line plot over $\{0, 1, \dots, 256\}$ , scaled over $[0, 1]$ by a division by 256 and a histogram of the 256 points in $[0, 1]$ with 15 bins. . . . .	130
5.2	The LCG called <code>RANDU</code> with $(m, a, c) = (2147483648, 65539, 0)$ has strong correlation between three consecutive points as: $x_{i+2} = 6x_{k+1} - 9x_k$ . The two plots are showing $(x_i, x_{i+1}, x_{i+2})$ from two different view points. . . . .	132
5.3	Triplet point clouds from the “Mersenne Twister” with two different seeds (see Lab-work 96). . . . .	134
5.4	A plot of the PDF, DF or CDF and inverse DF of the $\text{Uniform}(-1, 1)$ RV $X$ . . . . .	135
5.5	Visual Cognitive Tool GUI: Inversion Sampling from $X \sim \text{Uniform}(-5, 5)$ . . . . .	136
5.6	The PDF $f$ , DF $F$ , and inverse DF $F^{[-1]}$ of the the $\text{Exponential}(\lambda = 1.0)$ RV. . . . .	137
5.7	Visual Cognitive Tool GUI: Inversion Sampling from $X \sim \text{Exponential}(0.5)$ . . . . .	138
5.8	Visual Cognitive Tool GUI: Inversion Sampling from $X \sim \text{Laplace}(5)$ . . . . .	140
5.9	Visual Cognitive Tool GUI: Inversion Sampling from $X \sim \text{Cauchy}$ . . . . .	142
5.10	Unending fluctuations of the running means based on $n$ IID samples from the Standard Cauchy RV $X$ in each of five replicate simulations (blue lines). The running means, based on $n$ IID samples from the $\text{Uniform}(0, 10)$ RV, for each of five replicate simulations (magenta lines). . . . .	143
5.11	The DF $F(x; 0.3, 0.7)$ of the de Moivre(0.3, 0.7) RV and its inverse $F^{[-1]}(u; 0.3, 0.7)$ . . . . .	146
5.12	Visual Cognitive Tool GUI: Rejection Sampling from $X \sim \text{Normal}(0, 1)$ with PDF $f$ based on proposals from $Y \sim \text{Laplace}(1)$ with PDF $g$ . . . . .	155
5.13	Rejection Sampling from $X \sim \text{Normal}(0, 1)$ with PDF $f$ based on 100 proposals from $Y \sim \text{Laplace}(1)$ with PDF $g$ . . . . .	156

# Chapter 1

## Preliminaries

### 1.1 Elementary Set Theory

A **set** is a collection of distinct objects. We write a set by enclosing its elements with curly braces. For example, we denote a set of the two objects  $\circ$  and  $\bullet$  by:

$$\{\circ, \bullet\}.$$

Sometimes, we give names to sets. For instance, we might call the first example set  $A$  and write:

$$A = \{\circ, \bullet\}.$$

We do not care about the order of elements within a set, i.e.  $A = \{\circ, \bullet\} = \{\bullet, \circ\}$ . We do not allow a set to contain multiple copies of any of its elements unless the copies are distinguishable, say by labels. So,  $B = \{\circ, \bullet, \bullet\}$  is not a set unless the two copies of  $\bullet$  in  $B$  are labelled or marked to make them distinct, e.g.  $B = \{\circ, \tilde{\bullet}, \bullet'\}$ . Names for sets that arise in a mathematical discourse are given upper-case letters ( $A, B, C, D, \dots$ ). Special symbols are reserved for commonly encountered sets.

Here is the set  $\mathcal{G}$  of twenty two Greek lower-case alphabets that we may encounter later:

$$\mathcal{G} = \{ \alpha, \beta, \gamma, \delta, \epsilon, \zeta, \eta, \theta, \kappa, \lambda, \mu, \nu, \xi, \pi, \rho, \sigma, \tau, \upsilon, \varphi, \chi, \psi, \omega \}.$$

They are respectively named alpha, beta, gamma, delta, epsilon, zeta, eta, theta, kappa, lambda, mu, nu, xi, pi, rho, sigma, tau, upsilon, phi, chi, psi and omega. *LHS* and *RHS* are abbreviations for objects on the Left and Right Hand Sides, respectively, of some binary relation. By the notation:

$$LHS := RHS,$$

we mean that *LHS* is equal, by definition, to *RHS*.

The set which does not contain any element (the collection of nothing) is called the **empty set**:

$$\emptyset := \{ \}.$$

We say an element  $b$  **belongs to** a set  $B$ , or simply that  $b$  belongs to  $B$  or that  $b$  is an element of  $B$ , if  $b$  is one of the elements that make up the set  $B$ , and write:

$$b \in B.$$

When  $b$  **does not belong to**  $B$ , we write:

$$b \notin B.$$

For our example set  $A = \{\circ, \bullet\}$ ,  $\star \notin A$  but  $\bullet \in A$ .

We say that a set  $C$  is a **subset** of another set  $D$  and write:

$$C \subset D$$

if every element of  $C$  is also an element of  $D$ . By this definition, any set is a subset of itself.

We say that two sets  $C$  and  $D$  are **equal** (as sets) and write  $C = D$  ‘if and only if’ ( $\iff$ ) every element of  $C$  is also an element of  $D$ , and every element of  $D$  is also an element of  $C$ . This definition of set equality is notationally summarised as follows:

$$C = D \iff C \subset D, D \subset C .$$

When two sets  $C$  and  $D$  are not equal by the above definition, we say that  $C$  is **not equal** to  $D$  and write:

$$C \neq D .$$

The **union** of two sets  $C$  and  $D$ , written as  $C \cup D$ , is the set of elements that belong to  $C$  or  $D$ . We can formally express our definition of set union as:

$$C \cup D := \{x : x \in C \text{ or } x \in D\} .$$

When a colon (:) appears inside a set, it stands for ‘such that’. Thus, the above expression is read as ‘ $C$  union  $D$  is equal by definition to the set of all elements  $x$ , such that  $x$  belongs to  $C$  or  $x$  belongs to  $D$ .’

Similarly, the **intersection** of two sets  $C$  and  $D$ , written as  $C \cap D$ , is the set of elements that belong to both  $C$  and  $D$ . Formally:

$$C \cap D := \{x : x \in C \text{ and } x \in D\} .$$

**Venn diagrams** are visual aids for set operations as in the diagrams below.

Figure 1.1: Union and intersection of sets shown by Venn diagrams

The set-difference or **difference** of two sets  $C$  and  $D$ , written as  $C \setminus D$ , is the set of elements in  $C$  that do not belong to  $D$ . Formally:

$$C \setminus D := \{x : x \in C \text{ and } x \notin D\} .$$

When a universal set, e.g.  $U$  is well-defined, the **complement** of a given set  $B$  denoted by  $B^c$  is the set of all elements of  $U$  that don’t belong to  $B$ , i.e.:

$$B^c := U \setminus B .$$

We say two sets  $C$  and  $D$  are **disjoint** if they have no elements in common, i.e.  $C \cap D = \emptyset$ .

By drawing Venn diagrams, let us check **De Morgan’s Laws**:

$$(A \cup B)^c = A^c \cap B^c \text{ and } (A \cap B)^c = A^c \cup B^c$$

**Classwork 1 (Fruits and colours)** Consider a set of fruits  $F = \{\text{orange, banana, apple}\}$  and a set of colours  $C = \{\text{red, green, blue, orange}\}$ . Then,




Figure 1.2: These Venn diagram illustrate De Morgan's Laws.

1.  $F \cap C =$

2.  $F \cup C =$

3.  $F \setminus C =$

4.  $C \setminus F =$

**Classwork 2 (Subsets of a universal set)** Suppose we are given a universal set  $U$ , and three of its subsets,  $A$ ,  $B$  and  $C$ . Also suppose that  $A \subset B \subset C$ . Find the circumstances, if any, under which each of the following statements is true (T) and justify your answer:

- |                           |                                |                           |                        |
|---------------------------|--------------------------------|---------------------------|------------------------|
| (1) $C \subset B$         | T when $B = C$                 | (2) $A \subset C$         | T by assumption        |
| (3) $C \subset \emptyset$ | T when $A = B = C = \emptyset$ | (4) $\emptyset \subset A$ | T always               |
| (5) $C \subset U$         | T by assumption                | (6) $U \subset A$         | T when $A = B = C = U$ |

## Exercises

**Ex. 1.1** — Let  $\Omega$  be the universal set of students, lecturers and tutors involved in a course. Now consider the following subsets:

- The set of 50 students,  $S = \{S_1, S_2, S_3, \dots, S_{50}\}$ .
- The set of 3 lecturers,  $L = \{L_1, L_2, L_3\}$ .
- The set of 4 tutors,  $T = \{T_1, T_2, T_3, L_3\}$ .

Note that one of the lecturers also tutors in the course. Find the following sets:

- |                       |                  |
|-----------------------|------------------|
| (a) $T \cap L$        | (f) $S \cap L$   |
| (b) $T \cap S$        | (g) $S^c \cap L$ |
| (c) $T \cup L$        | (h) $T^c$        |
| (d) $T \cup L \cup S$ | (i) $T^c \cap L$ |
| (e) $S^c$             | (j) $T^c \cap T$ |

**Ex. 1.2** — Using Venn diagram, sketch and check the rule:  

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

**Ex. 1.3** — Using Venn diagram, sketch and check the rule:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

**Ex. 1.4** — Using a Venn diagram, illustrate the idea that  $A \subseteq B$  if and only if  $A \cup B = B$ .

### SET SUMMARY

$\{a_1, a_2, \dots, a_n\}$	— a set containing the elements, $a_1, a_2, \dots, a_n$ .
$a \in A$	— $a$ is an element of the set $A$ .
$A \subseteq B$	— the set $A$ is a subset of $B$ .
$A \cup B$	— “union”, meaning the set of all elements which are in $A$ or $B$ , or both.
$A \cap B$	— “intersection”, meaning the set of all elements in both $A$ and $B$ .
$\{\} \text{ or } \emptyset$	— empty set.
$\Omega$	— universal set.
$A^c$	— the complement of $A$ , meaning the set of all elements in $\Omega$ , the universal set, which are not in $A$ .

## 1.2 Natural Numbers, Integers and Rational Numbers

We denote the number of elements in a set named  $B$  by:

$$\#B := \text{Number of elements in the set } B .$$

In fact, the Hindu-Arab numerals we have inherited are based on this intuition of the size of a collection. The elements of the set of **natural numbers**:

$$\mathbb{N} := \{1, 2, 3, 4, \dots\} , \text{ may be defined using } \# \text{ as follows:}$$

$$\begin{aligned} 1 &:= \#\{\star\} = \#\{\bullet\} = \#\{\alpha\} = \#\{\{\bullet\}\} = \#\{\{\bullet, \bullet'\}\} = \dots, \\ 2 &:= \#\{\star', \star\} = \#\{\bullet, \circ\} = \#\{\alpha, \omega\} = \#\{\{\circ\}, \{\alpha, \star, \bullet\}\} = \dots, \\ &\vdots \end{aligned}$$

For our example sets,  $A = \{\circ, \bullet\}$  and the set of Greek alphabets  $\mathcal{G}$ ,  $\#A = 2$  and  $\#\mathcal{G} = 22$ . The number zero may be defined as the size of an empty set:

$$0 := \#\emptyset = \#\{\}$$

The set of **non-negative integers** is:

$$\mathbb{Z}_+ := \mathbb{N} \cup \{0\} = \{0, 1, 2, 3, \dots\} .$$

A **product set** is the **Cartesian product** ( $\times$ ) of two or more possibly distinct sets:

$$A \times B := \{(a, b) : a \in A \text{ and } b \in B\}$$

For example, if  $A = \{\circ, \bullet\}$  and  $B = \{\star\}$ , then  $A \times B = \{(\circ, \star), (\bullet, \star)\}$ . Elements of  $A \times B$  are called **ordered pairs**.

The binary arithmetic operation of **addition** (+) between a pair of non-negative integers  $c, d \in \mathbb{Z}_+$  can be defined via sizes of disjoint sets. Suppose,  $c = \#C$ ,  $d = \#D$  and  $C \cap D = \emptyset$ , then:

$$c + d = \#C + \#D := \#(C \cup D) .$$

For example, if  $A = \{\circ, \bullet\}$  and  $B = \{\star\}$ , then  $A \cap B = \emptyset$  and  $\#A + \#B = \#(A \cup B) \iff 2 + 1 = 3$ .

The binary arithmetic operation of **multiplication** ( $\cdot$ ) between a pair of non-negative integers  $c, d \in \mathbb{Z}_+$  can be defined via sizes of product sets. Suppose,  $c = \#C$ ,  $d = \#D$ , then:

$$c \cdot d = \#C \cdot \#D := \#(C \times D) .$$

For example, if  $A = \{\circ, \bullet\}$  and  $B = \{\star\}$ , then  $\#A \cdot \#B = \#(A \times B) \iff 2 \cdot 1 = 2$ .

More generally, a product set of  $A_1, A_2, \dots, A_m$  is:

$$A_1 \times A_2 \times \cdots \times A_m := \{(a_1, a_2, \dots, a_m) : a_1 \in A_1, a_2 \in A_2, \dots, a_m \in A_m\}$$

Elements of an  $m$ -product set are called **ordered  $m$ -tuples**. When we take the product of the same set we abbreviate as follows:

$$A^m := \underbrace{A \times A \times \cdots \times A}_{m \text{ times}} := \{(a_1, a_2, \dots, a_m) : a_1 \in A, a_2 \in A, \dots, a_m \in A\}$$

**Classwork 3 (Cartesian product of sets)** 1. Let  $A = \{\circ, \bullet\}$ . What are the elements of  $A^2$ ? 2. Suppose  $\#A = 2$  and  $\#B = 3$ . What is  $\#(A \times B)$ ? 3. Suppose  $\#A_1 = s_1, \#A_2 = s_2, \dots, \#A_m = s_m$ . What is  $\#(A_1 \times A_2 \times \cdots \times A_m)$ ?

Now, let's recall the definition of a function. A **function** is a “mapping” that associates each element in some set  $\mathbb{X}$  (the domain) to exactly one element in some set  $\mathbb{Y}$  (the range). Two different elements in  $\mathbb{X}$  can be mapped to or associated with the same element in  $\mathbb{Y}$ , and not every element in  $\mathbb{Y}$  needs to be mapped. Suppose  $x \in \mathbb{X}$ . Then we say  $f(x) = y \in \mathbb{Y}$  is the **image** of  $x$ . To emphasise that  $f$  is a **function** from  $\mathbb{X} \ni x$  to  $\mathbb{Y} \ni y$ , we write:

$$f(x) = y : \mathbb{X} \rightarrow \mathbb{Y} .$$

And for some  $y \in \mathbb{Y}$ , we call the set:

$$f^{[-1]}(y) := \{x \in \mathbb{X} : f(x) = y\} \subset \mathbb{X} ,$$

the **pre-image** or **inverse image** of  $y$ , and

$$f^{[-1]} := f^{[-1]}(y \in \mathbb{Y}) = X \subset \mathbb{X} ,$$

Figure 1.3: A function  $f$  (“father of”) from  $\mathbb{X}$  (a set of children) to  $\mathbb{Y}$  (their fathers) and its inverse (“children of”).

as the **inverse** of  $f$ .

We motivated the non-negative integers  $\mathbb{Z}_+$  via the size of a set. With the notion of two directions (+ and -) and the magnitude of the current position from the origin zero (0) of a dynamic entity, we can motivate the set of **integers**:

$$\mathbb{Z} := \{\dots, -3, -2, -1, 0, +1, +2, +3, \dots\} .$$

The integers with a **minus** or **negative sign** (-) before them are called negative integers and those with a **plus** or **positive sign** (+) before them are called positive integers. Conventionally, + signs are dropped. Some examples of functions you may have encountered are **arithmetic operations** such as **addition** (+), **subtraction** (-), **multiplication** ( $\cdot$ ) and **division** (/) of ordered pairs of integers. The reader is assumed to be familiar with such arithmetic operations with pairs of integers. Every integer is either positive, negative, or zero. In terms of this we define the notion of **order**. We say an integer  $a$  is **less than** an integer  $b$  and write  $a < b$  if  $b - a$  is positive. We say an integer  $a$  is **less than or equal to** an integer  $b$  and write  $a \leq b$  if  $b - a$  is positive or zero. Finally, we say that  $a$  is greater than  $b$  and write  $a > b$  if  $b < a$ . Similarly,  $a$  is greater than equal to  $b$ , i.e.  $a \geq b$ , if  $b \leq a$ . The set of integers are **well-ordered**, i.e., for every integer  $a$  there is a next largest integer  $a + 1$ .

**Classwork 4 (Addition over integers)** Consider the set of integers  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ . Try to set up the arithmetic operation of addition as a function. The domain for addition is the Cartesian product of  $\mathbb{Z}$ :

$$\mathbb{Z}^2 := \mathbb{Z} \times \mathbb{Z} := \{(a, b) : a \in \mathbb{Z}, b \in \mathbb{Z}\}$$

What is its range ?

$$+ : \mathbb{Z} \times \mathbb{Z} \rightarrow$$

If the magnitude of the entity’s position is measured in units (e.g. meters) that can be rationally divided into  $q$  pieces with  $q \in \mathbb{N}$ , then we have the set of rational numbers:

$$\mathbb{Q} := \{p/q : p \in \mathbb{Z}, q \in \mathbb{Z} \setminus \{0\}\}$$

The expressions  $p/q$  and  $p'/q'$  denote the same rational number if and only if  $p \cdot q' = p' \cdot q$ . Every rational number has a unique irreducible expression  $p/q$ , where  $q$  is positive and as small as possible. For example,  $1/2$ ,  $2/4$ ,  $3/6$ , and  $1001/2002$  are different expressions for the same rational number whose irreducible unique expression is  $1/2$ .

Figure 1.4: A pictorial depiction of addition and its inverse. The domain is plotted in orthogonal Cartesian coordinates.

Addition and multiplication are defined for rational numbers by:

$$\frac{p}{q} + \frac{p'}{q'} = \frac{p \cdot q' + p' \cdot q}{q \cdot q'} \quad \text{and} \quad \frac{p}{q} \cdot \frac{p'}{q'} = \frac{p \cdot p'}{q \cdot q'} .$$

The rational numbers form a **field** under the operations of addition and multiplication defined above in terms of addition and multiplication over integers. This means that the following properties are satisfied:

1. Addition and multiplication are each **commutative**

$$a + b = b + a, \quad a \cdot b = b \cdot a ,$$

and associative

$$a + (b + c) = (a + b) + c, \quad a \cdot (b \cdot c) = (a \cdot b) \cdot c .$$

2. Multiplication **distributes** over addition

$$a \cdot (b + c) = (a \cdot b) + (a \cdot c) .$$

3. 0 is the **additive identity** and 1 is the multiplicative identity

$$0 + a = a \quad \text{and} \quad 1 \cdot a = a .$$

4. Every rational number  $a$  has a negative,  $a + (-a) = 0$  and every non-zero rational number  $a$  has a reciprocal,  $a \cdot 1/a = 1$ .

The field axioms imply the usual laws of arithmetic and allow subtraction and division to be defined in terms of addition and multiplication as follows:

$$\frac{p}{q} - \frac{p'}{q'} := \frac{p}{q} + \frac{-p'}{q'} \quad \text{and} \quad \frac{p}{q} / \frac{p'}{q'} := \frac{p}{q} \cdot \frac{q'}{p'}, \quad \text{provided } p' \neq 0 .$$

We will see later that the theory of finite fields is necessary for the study of pseudo-random number generators (PRNGs) and PRNGs are the heart-beat of randomness and statistics with computers.

### 1.3 Real Numbers

Unlike rational numbers which are expressible in their reduced forms by  $p/q$ , it is fairly tricky to define or express real numbers. It is possible to define real numbers formally and constructively via equivalence classes of Cauchy sequence of rational numbers. For this all we need are notions of (1) infinity, (2) sequence of rational numbers and (3) distance between any two rational numbers in an infinite sequence of them. These are topics usually covered in an introductory course in real analysis and are necessary for a firm foundation in computational statistics. Instead of a formal constructive definition of real numbers, we give a more concrete one via decimal expansions. See Donald E. Knuth's treatment [*Art of Computer Programming, Vol. I, Fundamental Algorithms*, 3rd Ed., 1997, pp. 21-25] for a fuller story. A **real number** is a numerical quantity  $x$  that has a decimal expansion:

$$x = n + 0.d_1d_2d_3 \dots , \text{ where, each } d_i \in \{0, 1, \dots, 9\}, n \in \mathbb{Z} ,$$

and the sequence  $0.d_1d_2d_3 \dots$  does not terminate with infinitely many consecutive 9s. By the above decimal representation, the following arbitrarily accurate enclosure of the real number  $x$  by rational numbers is implied:

$$n + \frac{d_1}{10} + \frac{d_2}{100} + \dots + \frac{d_k}{10^k} =: \underline{x}_k \leq x < \bar{x}_k := n + \frac{d_1}{10} + \frac{d_2}{100} + \dots + \frac{d_k}{10^k} + \frac{1}{10^{k+1}}$$

for every  $k \in \mathbb{N}$ . Thus, rational arithmetic  $(+, -, \cdot, /)$  can be extended with arbitrary precision to any ordered pair of real numbers  $x$  and  $y$  by operations on their rational enclosures  $\underline{x}, \bar{x}$  and  $\underline{y}, \bar{y}$ .

Some examples of real numbers that are not rational (**irrational numbers**) are:

$$\sqrt{2} = 1.41421356237309 \dots \text{the side length of a square with area of 2 units}$$

$$\pi = 3.14159265358979 \dots \text{the ratio of the circumference to diameter of a circle}$$

$$e = 2.71828182845904 \dots \text{Euler's constant}$$

We can think of  $\pi$  as being enclosed by the following pairs of rational numbers:

$$\begin{aligned} 3 + \frac{1}{10} &=: \underline{\pi}_1 \leq \pi < \bar{\pi}_1 := 3 + \frac{1}{10} + \frac{1}{10^1} \\ 3 + \frac{1}{10} + \frac{4}{100} &=: \underline{\pi}_2 \leq \pi < \bar{\pi}_2 := 3 + \frac{1}{10} + \frac{4}{100} + \frac{1}{100} \\ 3 + \frac{1}{10} + \frac{4}{100} + \frac{1}{10^3} &=: \underline{\pi}_3 \leq \pi < \bar{\pi}_3 := 3 + \frac{1}{10} + \frac{4}{100} + \frac{1}{10^3} + \frac{1}{10^3} \\ &\vdots \\ 3.14159265358979 &=: \underline{\pi}_{14} \leq \pi < \bar{\pi}_{14} := 3.14159265358979 + \frac{1}{10^{14}} \\ &\vdots \end{aligned}$$

Think of the real number system as the continuum of points that make up a line, as shown in Figure 1.5.

Let  $y$  and  $z$  be two real numbers such that  $y \leq z$ . Then, the **closed interval**  $[y, z]$  is the set of real numbers  $x$  such that  $y \leq x \leq z$ :

$$[y, z] := \{x : y \leq x \leq z\} .$$

Figure 1.5: A depiction of the real line segment  $[-10, 10]$ .

The **half-open interval**  $(y, z]$  or  $[y, z)$  and the **open interval**  $(y, z)$  are defined analogously:

$$\begin{aligned}(y, z] &:= \{x : y < x \leq z\} , \\ [y, z) &:= \{x : y \leq x < z\} , \\ (y, z) &:= \{x : y < x < z\} .\end{aligned}$$

We also allow  $y$  to be **minus infinity** (denoted  $-\infty$ ) or  $z$  to be **infinity** (denoted  $\infty$ ) at an open endpoint of an interval, meaning that there is no lower or upper bound. With this allowance we get the set of **real numbers**  $\mathbb{R} := (-\infty, \infty)$ , the **non-negative real numbers**  $\mathbb{R}_+ := [0, \infty)$  and the **positive real numbers**  $\mathbb{R}_{>0}(0, \infty)$  as follows:

$$\begin{aligned}\mathbb{R} &:= (-\infty, \infty) = \{x : -\infty < x < \infty\} , \\ \mathbb{R}_+ &:= [0, \infty) = \{x : 0 \leq x < \infty\} , \\ \mathbb{R}_{>0} &:= (0, \infty) = \{x : 0 < x < \infty\} .\end{aligned}$$

For a positive real number  $b \in \mathbb{R}_{>0}$  and an integer  $n \in \mathbb{Z}$ , the  $n$ -th **power** or **exponent** of  $b$  is:

$$b^0 = 1, \quad b^n = b^{n-1} \cdot b \quad \text{if } n > 0, \quad b^n = b^{n+1}/b \quad \text{if } n < 0 .$$

The following **laws of exponents** hold by mathematical induction when  $m, n \in \mathbb{Z}$ :

$$b^{m+n} = b^m \cdot b^n, \quad (b^m)^n = b^{m \cdot n} .$$

If  $y \in \mathbb{R}$  and  $m \in \mathbb{N}$ , the unique positive real number  $z \in \mathbb{R}_{>0}$  such that  $z^m = y$  is called the  $m$ -th **root of  $y$**  and denoted by  $\sqrt[m]{y}$ , i.e.,

$$z^m = y \implies z = \sqrt[m]{y} .$$

For a rational number  $r = p/q \in \mathbb{Q}$ , we define the  $r$ -th power of  $b \in \mathbb{R}$  as follows:

$$b^r = b^{p/q} := \sqrt[q]{b^p}$$

The laws of exponents hold for this definition and different expressions for the same rational number  $r = ap/aq$  yield the same power, i.e.,  $b^{p/q} = b^{ap/aq}$ . Recall that a real number  $x = n + 0.d_1d_2d_3\dots \in \mathbb{R}$  can be arbitrarily precisely enclosed by the rational numbers  $\underline{x}_k := n + \frac{d_1}{10} + \frac{d_2}{100} + \dots + \frac{d_k}{10^k}$  and  $\bar{x}_k := n + \frac{d_1}{10} + \frac{d_2}{100} + \dots + \frac{d_k}{10^k} + \frac{1}{10^k}$  by increasing  $k$ . Suppose first that  $b > 1$ . Then, using rational powers, we can enclose  $b^x$ ,

$$b^{n+\frac{d_1}{10}+\frac{d_2}{100}+\dots+\frac{d_k}{10^k}} =: b^{\underline{x}_k} \leq b^x < b^{\bar{x}_k} =: b^{n+\frac{d_1}{10}+\frac{d_2}{100}+\dots+\frac{d_k}{10^k}+\frac{1}{10^k}} ,$$

within an interval of width  $b^{n+\frac{d_1}{10}+\frac{d_2}{100}+\dots+\frac{d_k}{10^k}} \left( b^{\frac{1}{10^k}} - 1 \right) < b^{n+1}(b-1)/10^k$ . By taking a large enough  $k$  we can evaluate  $b^x$  to any accuracy. Finally, when  $b < 1$  we define  $b^x := (1/b)^{-x}$  and when  $b = 0$ ,  $b^x := 1$ .

Suppose  $y \in \mathbb{R}_{>0}$  and  $b \in \mathbb{R} \setminus \{1\}$  then the real number  $x$  such that  $y = b^x$  is called the **logarithm of  $y$  to the base  $b$**  and we write this as:

$$y = b^x \iff x = \log_b y$$

The definition implies:

$$x = \log_b(b^x) = b^{\log_b x},$$

and the laws of exponents imply:

$$\begin{aligned}\log_b(xy) &= \log_b x + \log_b y, \quad \text{if } x > 0, y > 0 \text{ and} \\ \log_b(c^y) &= y \log_b c, \quad \text{if } c > 0.\end{aligned}$$

The **common logarithm** is  $\log_{10}(y)$ , the **binary logarithm** is  $\log_2(y)$  and the **natural logarithm** is  $\log_e(y)$ , where  $e$  is the Euler's constant. Since we will mostly work with  $\log_e(y)$  we use  $\log(y)$  to mean  $\log_e(y)$ . You are assumed to be familiar with trigonometric functions ( $\sin(x)$ ,  $\cos(x)$ ,  $\tan(x)$ , ...). We sometimes denote the special power function  $e^y$  by  $\exp(y)$ .

Familiar extremal elements of a set of real numbers, say  $A$ , are the following:

$$\boxed{\max A := \text{greatest element in } A}$$

For example,  $\max\{1, 4, -9, 345\} = 345$ ,  $\max[-93.8889, 1002.786] = 1002.786$ .

$$\boxed{\min A := \text{least element in } A}$$

For example,  $\min\{1, 4, -9, 345\} = -9$ ,  $\min[-93.8889, 1002.786] = -93.8889$ . We need a slightly more sophisticated notion for the extremal elements of a set  $A$  that may not belong to  $A$ . We say that a real number  $x$  is a **lower bound** for a non-empty set of real numbers  $A$ , provided  $x \leq a$  for every  $a \in A$ . We say that the set  $A$  is **bounded below** if it has at least one lower bound. A lower bound is the **greatest lower bound** if it is at least as large as any other lower bound. The greatest lower bound of a set of real numbers  $A$  is called the **infimum** of  $A$  and is denoted by:

$$\boxed{\inf A := \text{greatest lower bound of } A}$$

For example,  $\inf(0, 1) = 0$  and  $\inf\{10.333 \cup [-99, 1001.33]\} = -99$ . We similarly define the **least upper bound** of a non-empty set of real numbers  $A$  to be the **supremum** of  $A$  and denote it as:

$$\boxed{\sup A := \text{least upper bound of } A}$$

For example,  $\sup(0, 1) = 1$  and  $\sup\{10.333 \cup [-99, 1001.33]\} = 1001.33$ . By convention, we define  $\inf \emptyset := \infty$ ,  $\sup \emptyset := -\infty$ . Finally, if a set  $A$  is not bounded below then  $\inf A := -\infty$  and if a set  $A$  is not bounded above then  $\sup A := \infty$ .

Symbol	Meaning
$A = \{\star, \circ, \bullet\}$	$A$ is a set containing the elements $\star, \circ$ and $\bullet$
$\circ \in A$	$\circ$ belongs to $A$ or $\circ$ is an element of $A$
$A \ni \circ$	$\circ$ belongs to $A$ or $\circ$ is an element of $A$
$\circ \notin A$	$\circ$ does not belong to $A$
$\#A$	Size of the set $A$ , for e.g. $\#\{\star, \circ, \bullet, \odot\} = 4$
$\mathbb{N}$	The set of natural numbers $\{1, 2, 3, \dots\}$
$\mathbb{Z}$	The set of integers $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
$\mathbb{D}_+$	The set of non-negative integers $\{0, 1, 2, 3, \dots\}$
$\emptyset$	Empty set or the collection of nothing or $\{\}$
$A \subset B$	$A$ is a subset of $B$ or $A$ is contained by $B$ , e.g. $A = \{\circ\}, B = \{\bullet\}$
$A \supset B$	$A$ is a superset of $B$ or $A$ contains $B$ e.g. $A = \{\circ, \star, \bullet\}, B = \{\circ, \bullet\}$
$A = B$	$A$ equals $B$ , i.e. $A \subset B$ and $B \subset A$
$Q \implies R$	Statement $Q$ implies statement $R$ or If $Q$ then $R$
$Q \iff R$	$Q \implies R$ and $R \implies Q$
$\{x : x \text{ satisfies property } R\}$	The set of all $x$ such that $x$ satisfies property $R$
$A \cup B$	$A$ union $B$ , i.e. $\{x : x \in A \text{ or } x \in B\}$
$A \cap B$	$A$ intersection $B$ , i.e. $\{x : x \in A \text{ and } x \in B\}$
$A \setminus B$	$A$ minus $B$ , i.e. $\{x : x \in A \text{ and } x \notin B\}$
$A := B$	$A$ is equal to $B$ by definition
$A =: B$	$B$ is equal to $A$ by definition
$A^c$	$A$ complement, i.e. $\{x : x \in U, \text{ the universal set, but } x \notin A\}$
$A_1 \times A_2 \times \dots \times A_m$	The $m$ -product set $\{(a_1, a_2, \dots, a_m) : a_1 \in A_1, a_2 \in A_2, \dots, a_m \in A_m\}$
$A^m$	The $m$ -product set $\{(a_1, a_2, \dots, a_m) : a_1 \in A, a_2 \in A, \dots, a_m \in A\}$
$f := f(x) = y : \mathbb{X} \rightarrow \mathbb{Y}$	A function $f$ from domain $\mathbb{X}$ to range $\mathbb{Y}$
$f^{[-1]}(y)$	Inverse image of $y$
$f^{[-1]} := f^{[-1]}(y \in \mathbb{Y}) = X \subset \mathbb{X}$	Inverse of $f$
$a < b$ or $a \leq b$	$a$ is less than $b$ or $a$ is less than or equal to $b$
$a > b$ or $a \geq b$	$a$ is greater than $b$ or $a$ is greater than or equal to $b$
$\mathbb{Q}$	Rational numbers
$(x, y)$	the open interval $(x, y)$ , i.e. $\{r : x < r < y\}$
$[x, y]$	the closed interval $(x, y)$ , i.e. $\{r : x \leq r \leq y\}$
$(x, y]$	the half-open interval $(x, y]$ , i.e. $\{r : x < r \leq y\}$
$[x, y)$	the half-open interval $[x, y)$ , i.e. $\{r : x \leq r < y\}$
$\mathbb{R} := (-\infty, \infty)$	Real numbers, i.e. $\{r : -\infty < r < \infty\}$
$\mathbb{R}_+ := [0, \infty)$	Real numbers, i.e. $\{r : 0 \leq r < \infty\}$
$\mathbb{R}_{>0} := (0, \infty)$	Real numbers, i.e. $\{r : 0 < r < \infty\}$

Table 1.1: Symbol Table: Sets and Numbers

## 1.4 Introduction to MATLAB

We use MATLAB to perform computations and visualisations. MATLAB is a numerical computing environment and programming language that is optimised for vector and matrix processing. STAT 218/313 students will have access to Maths & Stats Department's computers that are licensed to run MATLAB . You can remotely connect to these machines from home by following instructions at <http://www.math.canterbury.ac.nz/php/resources/comdocs/remote>.

**Labwork 5 (Basics of MATLAB )** Let us familiarize ourselves with MATLAB in this session. First, you need to launch MATLAB from your terminal. Since this is system dependent, ask your tutor for help. The command window within the MATLAB window is where you need to type commands. Here is a minimal set of commands you need to familiarize yourself with in this session.

1. Type the following command to add 2 numbers in the command window right after the command prompt `>>` .

```
>> 13+24
```

Upon hitting **Enter** or **Return** on your keyboard, you should see:

```
ans =
37
```

The summand 37 of 13 and 24 is stored in the default variable called `ans` which is short for answer.

2. We can write **comments** in MATLAB following the % character. All the characters in a given line that follow the percent character % are ignored by MATLAB . It is very helpful to comment what is being done in the code. You won't get full credit without sufficient comments in your coding assignments. For example we could have added the comment to the previous addition. To save space in these notes, we suppress the blank lines and excessive line breaks present in MATLAB 's command window.

```
>> 13+24 % adding 13 to 24 using the binary arithmetic operator +
ans =      37
```

3. You can **create or reopen a diary file** in MATLAB to record your work. Everything you typed or input and the corresponding output in the command window will be recorded in the diary file. You can create or reopen a diary file by typing `diary filename.txt` in the command window. When you have finished recording, simply type `diary off` in the command window **to turn off the diary file**. The diary file with .txt extension is simply a text-file. It can be edited in different editors after the diary is turned off in MATLAB . You need to type `diary LabWeek1.txt` to start recording your work for electronic submission if needed.

```
>> diary blah.txt % start a diary file named blah.txt
>> 3+56
ans =      59
>> diary off % turn off the current diary file blah.txt
```

```
>> type blah.txt % this allows you to see the contents of blah.txt
3+56
ans =      59
diary off
>> diary blah.txt % reopen the existing diary file blah.txt
>> 45-54
ans =      -9
>> diary off % turn off the current diary file blah.txt again
>> type blah.txt % see its contents
3+56
ans =      59
diary off
45-54
ans =      -9
diary off
```

4. Let's learn to store values in variables of our choice. Type the following at the command prompt :

```
>> VariableCalledX = 12
```

Upon hitting enter you should see that the number 12 has been assigned to the variable named **VariableCalledX** :

```
VariableCalledX =      12
```

5. MATLAB stores default value for some variables, such as **pi** ( $\pi$ ), **i** and **j** (complex numbers).

```
>> pi
ans =      3.1416
>> i
ans =      0 + 1.0000i
>> j
ans =      0 + 1.0000i
```

All predefined symbols (variables, constants, function names, operators, etc) in MATLAB are written in lower-case. Therefore, it is a good practice to name the variable you define using upper and mixed case letters in order to prevent an unintended overwrite of some predefined MATLAB symbol.

6. We could have stored the sum of 13 and 24 in the variable **X**, by entering:

```
>> X = 13 + 24
X =      37
```

7. Similarly, you can store the outcome of multiplication (via operation **\*** ), subtraction (via operation **-** ), division (via **/** ) and exponentiation (via **^** )of any two numbers of your choice in a variable name of your choice. Evaluate the following expressions in MATLAB :

$$\begin{aligned} p &= 45.89 * 1.00009 \\ m &= 5376.0 - 6.00 \end{aligned}$$

$$\begin{aligned} d &= 89.0 / 23.3454 \\ p &= 2^{0.5} \end{aligned}$$

8. You may compose the elementary operations to obtain rational expressions by using parenthesis to specify the order of the operations. To obtain  $\sqrt{2}$ , you can type the following into MATLAB 's command window.

```
>> 2^(1/2)
ans =      1.4142
```

The omission of parenthesis about  $1/2$  means something else and you get the following output:

```
>> 2^1/2
ans =      1
```

MATLAB first takes the 1st power of 2 and then divides it by 2 using its default precedence rules for binary operators in the absence of parenthesis. The order of operations or default precedence rule for arithmetic operations is 1. brackets or parentheses; 2. exponents (powers and roots); 3. division and multiplication; 4. addition and subtraction. The mnemonic **bedmas** can be handy. When in doubt, use parenthesis to force the intended order of operations.

9. When you try to divide by 0, MATLAB returns **Inf** for infinity.

```
>> 10/0
ans =    Inf
```

10. We can clear the value we have assigned to a particular variable and reuse it. We demonstrate it by the following commands and their output:

```
>> X
X =      37
>> clear X
>> X
??? Undefined function or variable 'X'.
```

Entering **X** after **clearing** it gives the above self-explanatory error message preceded by **???**.

11. We can suppress the output on the screen by ending the command with a semi-colon. Take a look at the simple command that sets **X** to  $\sin(3.145678)$  with and without the ‘;**;**’ at the end:

```
>> X = sin(3.145678)
X =    -0.0041
>> X = sin(3.145678);
```

12. If you do not understand a MATLAB function or command then type **help** or **doc** followed by the function or command. For example:

```
>> help sin
SIN    Sine of argument in radians.
SIN(X) is the sine of the elements of X.
See also asin, sind.
Overloaded methods:
darray/sin
Reference page in Help browser
    doc sin
>> doc sin
```

It is a good idea to use the help files before you ask your tutor.

13. Set the variable `x` to equal 17.13 and evaluate  $\cos(x)$ ,  $\log(x)$ ,  $\exp(x)$ ,  $\arccos(x)$ ,  $\text{abs}(x)$ ,  $\text{sign}(x)$  using the MATLAB commands `cos`, `log`, `exp`, `acos`, `abs`, `sign`, respectively. Read the help files to understand what each function does.
14. When we work with real numbers (floating-point numbers) or really large numbers, we might want the output to be displayed in concise notation. This can be controlled in MATLAB using the `format` command with the `short` or `long` options with/without `e` for scientific notation. `format compact` is used for getting compacted output and `format` returns the default format. For example:

```
>> format compact
>> Y=15;
>> Y = Y + acos(-1)
Y = 18.1416
>> format short
>> Y
Y = 18.1416
>> format short e
>> Y
Y = 1.8142e+001
>> format long
>> Y
Y = 18.141592653589793
>> format long e
>> Y
Y = 1.814159265358979e+001
>> format
>> Y
Y = 18.1416
```

15. Finally, to quit from MATLAB just type `quit` or `exit` at the prompt.

```
>> quit
```

16. An **M-file** is a special text file with a `.m` extension that contains a set of code or instructions in MATLAB . In this course we will be using two types of M-files: **script** and **function** files. A script file is simply a list of commands that we want executed and saves us from retyping code modules we are pleased with. A function file allows us to write specific tasks as functions with input and output. These functions can be called from other script files, function files or command window. We will see such examples shortly.

By now, you are expected to be familiar with arithmetic operations, simple function evaluations, format control, starting and stopping a diary file and launching and quitting MATLAB .

## 1.5 Permutations, Factorials and Combinations

**Definition 1 (Permutations and Factorials)** A **permutation** of  $n$  objects is an arrangement of  $n$  distinct objects in a row. For example, there are 2 permutations of the two objects  $\{1, 2\}$ :

$$12, \quad 21,$$

and 6 permutations of the three objects  $\{a, b, c\}$ :

$$abc, \quad acb, \quad bac, \quad bca, \quad cab, \quad cba.$$

Let the number of ways to choose  $k$  objects out of  $n$  and to arrange them in a row be denoted by  $p_{n,k}$ . For example, we can choose two ( $k = 2$ ) objects out of three ( $n = 3$ ) objects,  $\{a, b, c\}$ , and arrange them in a row in six ways ( $p_{3,2}$ ):

$$ab, \quad ac, \quad ba, \quad bc, \quad ca, \quad cb.$$

Given  $n$  objects, there are  $n$  ways to choose the left-most object, and once this choice has been made there are  $n - 1$  ways to select a different object to place next to the left-most one. Thus, there are  $n(n - 1)$  possible choices for the first two positions. Similarly, when  $n > 2$ , there are  $n - 2$  choices for the third object that is distinct from the first two. Thus, there are  $n(n - 1)(n - 2)$  possible ways to choose three distinct objects from a set of  $n$  objects and arrange them in a row. In general,

$$p_{n,k} = n(n - 1)(n - 2) \dots (n - k + 1)$$

and the total number of permutations called ‘ $n$  factorial’ and denoted by  $n!$  is

$$n! := p_{n,n} = n(n - 1)(n - 2) \dots (n - n + 1) = n(n - 1)(n - 2) \dots (3)(2)(1) =: \prod_{i=1}^n i.$$

Some factorials to bear in mind

$$0! := 1 \quad 1! = 1, \quad 2! = 2, \quad 3! = 6, \quad 4! = 24, \quad 5! = 120 \quad 10! = 3,628,800.$$

When  $n$  is large we can get a good idea of  $n!$  without laboriously carrying out the  $n - 1$  multiplications via Stirling’s approximation (*Methodus Differentialis* (1730), p. 137) :

$$n! \cong \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

**Definition 2 (Combinations)** The combinations of  $n$  objects taken  $k$  at a time are the possible choices of  $k$  different elements from a collection of  $n$  objects, disregarding order. They are called the  $k$ -combinations of the collection. The combinations of the three objects  $\{a, b, c\}$  taken two at a time, called the 2-combinations of  $\{a, b, c\}$ , are

$$ab, \quad ac, \quad bc,$$

and the combinations of the five objects  $\{1, 2, 3, 4, 5\}$  taken three at a time, called the 3-combinations of  $\{1, 2, 3, 4, 5\}$  are

$$123, \quad 124, \quad 125, \quad 134, \quad 135, \quad 145, \quad 234, \quad 235, \quad 245, \quad 345.$$

The total number of  $k$ -combination of  $n$  objects, called a **binomial coefficient**, denoted  $\binom{n}{k}$  and read “ $n$  choose  $k$ ,” can be obtained from  $p_{n,k} = n(n - 1)(n - 2) \dots (n - k + 1)$  and  $k! := p_{k,k}$ . Recall that  $p_{n,k}$  is the number of ways to choose the first  $k$  objects from the set of  $n$  objects and arrange them in a row with regard to order. Since we want to disregard order and each  $k$ -combination appears exactly  $p_{k,k}$  or  $k!$  times among the  $p_{n,k}$  many permutations, we perform a division:

$$\binom{n}{k} := \frac{p_{n,k}}{p_{k,k}} = \frac{n(n - 1)(n - 2) \dots (n - k + 1)}{k(k - 1)(k - 2) \dots 2 \ 1}.$$

Binomial coefficients are often called “Pascal’s Triangle” and attributed to Blaise Pascal’s *Traité du Triangle Arithmétique* from 1653, but they have many “fathers”. There are earlier treatises of the binomial coefficients including Szu-yüan Yü-chien (“The Precious Mirror of the Four Elements”) by the Chinese mathematician Chu Shih-Chieh in 1303, and in an ancient Hindu classic, *Pingala’s Chandasāstra*, due to Halāyudha (10-th century AD).

## 1.6 Array, Sequence, Limit, ...

In this section we will study a basic data structure in MATLAB called an **array** of numbers. Arrays are finite sequences and they can be processed easily in MATLAB . The notion of infinite sequences lead to **limits**, one of the most fundamental concepts in mathematics.

For any natural number  $n$ , we write

$$\langle x_{1:n} \rangle := x_1, x_2, \dots, x_{n-1}, x_n$$

to represent the **finite sequence** of real numbers  $x_1, x_2, \dots, x_{n-1}, x_n$ . For two integers  $m$  and  $n$  such that  $m \leq n$ , we write

$$\langle x_{m:n} \rangle := x_m, x_{m+1}, \dots, x_{n-1}, x_n$$

to represent the **finite sequence** of real numbers  $x_m, x_{m+1}, \dots, x_{n-1}, x_n$ . In mathematical analysis, finite sequences and their countably infinite counterparts play a fundamental role in limiting processes. Given an integer  $m$ , we denote an **infinite sequence** or simply a sequence as:

$$\langle x_{m:\infty} \rangle := x_m, x_{m+1}, x_{m+2}, x_{m+3}, \dots$$

Given index set  $\mathcal{I}$  which may be finite or infinite in size, a sequence can either be seen as a set of ordered pairs:

$$\{(i, x_i) : i \in \mathcal{I}\},$$

or as a function that maps the index set to the set of real numbers:

$$x(i) = x_i : \mathcal{I} \rightarrow \{x_i : i \in \mathcal{I}\},$$

The finite sequence  $\langle x_{m:n} \rangle$  has  $\mathcal{I} = \{m, m+1, m+2, m+3, \dots, n\}$  as its index set while an infinite sequence  $\langle x_{m:\infty} \rangle$  has  $\mathcal{I} = \{m, m+1, m+2, m+3, \dots\}$  as its index set. A **sub-sequence**  $\langle x_{j:k} \rangle$  of a finite sequence  $\langle x_{m:n} \rangle$  or an infinite sequence  $\langle x_{m:\infty} \rangle$  is:

$$\langle x_{j:k} \rangle = x_j, x_{j+1}, \dots, x_{k-1}, x_k \quad \text{where, } \quad m \leq j \leq k \leq n < \infty.$$

A rectangular arrangement of  $m \cdot n$  real numbers in  $m$  rows and  $n$  columns is called an  $m \times n$  **matrix**. The ' $m \times n$ ' represents the **size** of the matrix. We use bold upper-case letters to denote matrices, for e.g:

$$\mathbf{X} = \begin{bmatrix} x_{1,1} & x_{1,2} & \dots & x_{1,n-1} & x_{1,n} \\ x_{2,1} & x_{2,2} & \dots & x_{2,n-1} & x_{2,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{m-1,1} & x_{m-1,2} & \dots & x_{m-1,n-1} & x_{m-1,n} \\ x_{m,1} & x_{m,2} & \dots & x_{m,n-1} & x_{m,n} \end{bmatrix}$$

Matrices with only one row or only one column are called **vectors**. An  $1 \times n$  matrix is called a **row vector** since there is only one row and an  $m \times 1$  matrix is called a **column vector** since there is only one column. We use bold-face lowercase letters to denote row and column vectors.

$$\text{A row vector } \mathbf{x} = [x_1 \ x_2 \ \dots \ x_n] = (x_1, x_2, \dots, x_n)$$

$$\text{and a column vector } \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{m-1} \\ y_m \end{bmatrix} = [y_1 \ y_2 \ \dots \ y_m]' = (y_1, y_2, \dots, y_m)'.$$

The superscripting by  $'$  is the transpose operation and simply means that the rows and columns are exchanged. Thus the transpose of the matrix  $\mathbf{X}$  is:

$$\mathbf{X}' = \begin{bmatrix} x_{1,1} & x_{2,1} & \dots & x_{m-1,1} & x_{m,1} \\ x_{1,2} & x_{2,2} & \dots & x_{m-1,2} & x_{m,2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{1,n-1} & x_{2,n-1} & \dots & x_{m-1,n-1} & x_{m,n-1} \\ x_{1,n} & x_{2,n} & \dots & x_{m-1,n} & x_{m,n} \end{bmatrix}$$

In linear algebra and calculus, it is natural to think of vectors and matrices as points (ordered  $m$ -tuples) and ordered collection of points in Cartesian co-ordinates. We assume that the reader has heard of operations with matrices and vectors such as matrix multiplication, determinants, transposes, etc. Such concepts will be introduced as they are needed in the sequel.

Finite sequences, vectors and matrices can be represented in a computer by an elementary data structure called an **array**.


**Labwork 6 (Sequences as arrays)** Let us learn to represent, visualise and operate finite sequences as MATLAB arrays. Try out the commands and read the comments for clarification.

```
>> a = [17] % Declare the sequence of one element 17 in array a
a =
17
>> % Declare the sequence of 10 numbers in array b
>> b=[-1.4508 0.6636 -1.4768 -1.2455 -0.8235 1.1254 -0.4093 0.1199 0.2043 -0.8236]
b =
-1.4508    0.6636   -1.4768   -1.2455   -0.8235    1.1254   -0.4093    0.1199    0.2043   -0.8236
>> c = [1 2 3] % Declare the sequence of 3 consecutive numbers 1,2,3
c =
1     2     3
>> % linspace(x1, x2, n) generates n points linearly spaced between x1 and x2
>> r = linspace(1, 3, 3) % Declare sequence r = c using linspace
r =
1     2     3
>> s1 = 1:10 % declare an array s1 starting at 1, ending by 10, in increments of 1
s =
1     2     3     4     5     6     7     8     9     10
>> s2 = 1:2:10 % declare an array s2 starting at 1, ending by 10, in increments of 2
s =
1     3     5     7     9
>> s2(3) % obtain the third element of the finite sequence s2
ans =
5
>> s2(2:4) % obtain the subsequence from second to fourth elements of the finite sequence s2
ans =
3     5     7
```

We may visualise (as per Figure 1.6) the finite sequences  $\langle b_{1:n} \rangle$  stored in the array **b** as the set of ordered pairs  $\{(1, b_1), (2, b_2), \dots, (10, b_n)\}$  representing the function  $b(i) = b_i : \{1, 2, \dots, n\} \rightarrow \{b_1, b_2, \dots, b_n\}$  via **point plot** and **stem plot** using Matlab's **plot** and **stem** commands, respectively.

```
>> display(b) % display the array b in memory
b =
-1.4508    0.6636   -1.4768   -1.2455   -0.8235    1.1254   -0.4093    0.1199    0.2043   -0.8236
>> plot(b,'o') % point plot of ordered pairs (1,b(1)), (2,b(2)), ..., (10,b(10))
>> stem(b) % stem plot of ordered pairs (1,b(1)), (2,b(2)), ..., (10,b(10))
>> plot(b,'-o') % point plot of ordered pairs (1,b(1)), (2,b(2)), ..., (10,b(10)) connected by lines
```

**Labwork 7 (Vectors and matrices as arrays)** Let us learn to represent, visualise and operate vectors as MATLAB arrays. Syntactically, a vector is stored in an array exactly in the same way we stored a finite sequence. However, mathematically, we think of a vector as an ordered  $m$ -tuple that can be visualised as a point in Cartesian co-ordinates. Try out the commands and read the comments for clarification.

Figure 1.6: Point plot and stem plot of the finite sequence  $\langle b_{1:10} \rangle$  declared as an array.

```

>> a = [1 2]          % an 1 X 2 row vector
>> z = [1 2 3]        % Declare an 1 X 3 row vector z with three numbers
z =
    1      2      3
>> % linspace(x1, x2, n) generates n points linearly spaced between x1 and x2
>> r = linspace(1, 3, 3)           % Declare an 1 X 3 row vector r = z using linspace
r =
    1      2      3
>> c = [1; 2; 3]                % Declare a 3 X 1 column vector c with three numbers. Semicolons delineate columns
c =
    1
    2
    3
>> rT = r'                    % The column vector (1,2,3)' by taking the transpose of r via r'
rT =
    1
    2
    3
>> y = [1 1 1]                  % y is a sequence or row vector of 3 1's
y =
    1      1      1
>> ones(1,10)                  % ones(m,n) is an m X n matrix of ones. Useful when m or n is large.
ans =
    1      1      1      1      1      1      1      1      1      1

```

We can use two dimensional arrays to represent matrices. Some useful built-in commands to generate standard matrices are:

```

>> Z=zeros(2,10) % the 2 X 10 matrix of zeros
Z =
    0      0      0      0      0      0      0      0      0      0
    0      0      0      0      0      0      0      0      0      0
>> O=ones(4,5) % the 4 X 5 matrix of ones
O =
    1      1      1      1      1
    1      1      1      1      1
    1      1      1      1      1
    1      1      1      1      1
>> E=eye(4) % the 4 X 4 identity matrix
E =
    1      0      0      0
    0      1      0      0
    0      0      1      0
    0      0      0      1

```

We can also perform operations with arrays representing vectors, finite sequences, or matrices.

```

>> y % the array y is
y =      1      1      1
>> z % the array z is
z =      1      2      3
>> x = y + z
x =      2      3      4
                                % x is the sum of vectors y and z (with same size 1 X 3)
>> y = y * 2
y =      2      2      2
                                % y is updated to 2 * y (each term of y is multiplied by 2)
>> p = z .* y
p =      2      4      6
                                % p is the vector obtained by term-by-term product of z and y
>> d = z ./ y
d = 0.5000    1.0000    1.5000
                                % d is the vector obtained by term-by-term division of z and y
>> t=linspace(-10,10,4)
t = -10.0000   -3.3333   3.3333   10.0000
                                % t has 4 numbers equally-spaced between -10 and 10
>> s = sin(t)
s = 0.5440    0.1906   -0.1906   -0.5440
                                % s is a vector obtained from the term-wise sin of the vector t
>> sSq = sin(t) .^ 2
sSq = 0.2960    0.0363    0.0363    0.2960
                                % sSq is an array obtained from term-wise squaring (. ^ 2) of the sin(t) array
>> cSq = cos(t) .^ 2
cSq = 0.7040    0.9637    0.9637    0.7040
                                % cSq is an array obtained from term-wise squaring (. ^ 2) of the cos(t) array
>> sSq + cSq % we can add the two arrays sSq and cSq to get the array of 1's
ans =      1      1      1      1
>> n = sin(t) .^ 2 + cos(t) .^ 2
n =      1      1      1      1
                                % we can directly do term-wise operation sin^2(t) + cos^2(t) of t as well
>> t2 = (-10:6.666665:10)
t2 = -10.0000   -3.3333   3.3333   10.0000
                                % t2 is similar to t above but with ':' syntax of (start:increment:stop)

```

Similarly, operations can be performed with matrices.

```

>> (0+0) .^ (1/2) % term-by-term square root of the matrix obtained by adding 0=ones(4,5) to itself
ans =
  1.4142    1.4142    1.4142    1.4142    1.4142
  1.4142    1.4142    1.4142    1.4142    1.4142
  1.4142    1.4142    1.4142    1.4142    1.4142
  1.4142    1.4142    1.4142    1.4142    1.4142

```

We can access specific rows or columns of a matrix as follows:

```

>> % declare a 3 X 3 array A of row vectors
>> A = [0.2760    0.4984    0.7513; 0.6797    0.9597    0.2551; 0.1626    0.5853    0.6991]
A =
  0.2760    0.4984    0.7513
  0.6797    0.9597    0.2551
  0.1626    0.5853    0.6991
>> A(2,:) % access the second row of A
ans =
  0.6797    0.9597    0.2551
>> B = A(2:3,:)
B =
  0.6797    0.9597    0.2551
  0.1626    0.5853    0.6991
>> C = A(:,[1 3]) % store the first and third columns of A in matrix C
C =
  0.2760    0.7513
  0.6797    0.2551

```

**Labwork 8 (Plotting a function as points of ordered pairs in two arrays)** Next we plot the function  $\sin(x)$  from several ordered pairs  $(x_i, \sin(x_i))$ . Here  $x_i$ 's are from the domain  $[-2\pi, 2\pi]$ .

We use the `plot` function in MATLAB . Create an M-file called `MySineWave.m` and copy the following commands in it. By entering `MySineWave` in the command window you should be able to run the script and see the figure in the figure window.

---

```

x = linspace(-2*pi,2*pi,100);           % x has 100 points equally spaced in [-2*pi, 2*pi]
y = sin(x);                            % y is the term-wise sin of x, ie sin of every number in x is in y, resp.
plot(x,y,'.');                         % plot x versus y as dots should appear in the Figure window
xlabel('x');                           % label x-axis with the single quote enclosed string x
ylabel('sin(x)', 'FontSize',16);        % label y-axis with the single quote enclosed string
title('Sine Wave in [-2 pi, 2 pi]', 'FontSize',16);    % give a title; click Figure window to see changes
set(gca,'XTick',-8:1:8,'FontSize',16) % change the range and size of X-axis ticks
% you can go to the Figure window's File menu to print/save the plot

```

---

The plot was saved as an encapsulated postscript file from the File menu of the Figure window and is displayed below.

Figure 1.7: A plot of the sine wave over  $[-2\pi, 2\pi]$ .

Let us first recall some elementary ideas from real analysis.

**Definition 3 (Convergent sequence of real numbers)** A sequence of real numbers  $\langle x_i \rangle_{i=1}^{\infty} := x_1, x_2, \dots$  is said to converge to a limit  $a \in \mathbb{R}$  and denoted by:

$$\lim_{i \rightarrow \infty} x_i = a ,$$

if for every natural number  $m \in \mathbb{N}$ , a natural number  $N_m \in \mathbb{N}$  exists such that for every  $j \geq N_m$ ,  $|x_j - a| \leq \frac{1}{m}$ .

**Example 9 (Limit of a sequence of 17s)** Let  $\langle x_i \rangle_{i=1}^{\infty} = 17, 17, 17, \dots$ . Then  $\lim_{i \rightarrow \infty} x_i = 17$ . This is because for every  $m \in \mathbb{N}$ , we can take  $N_m = 1$  and satisfy the definition of the limit, i.e.:

$$\text{for every } j \geq N_m = 1, |x_j - 17| = |17 - 17| = 0 \leq \frac{1}{m} .$$

**Example 10 (Limit of  $1/i$ )** Let  $\langle x_i \rangle_{i=1}^{\infty} = \frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \dots$ , i.e.  $x_i = \frac{1}{i}$ , then  $\lim_{i \rightarrow \infty} x_i = 0$ . This is because for every  $m \in \mathbb{N}$ , we can take  $N_m = m$  and satisfy the definition of the limit, i.e.:

$$\text{for every } j \geq N_m = m, |x_j - 0| = \left| \frac{1}{j} - 0 \right| = \frac{1}{j} \leq \frac{1}{m} .$$

However, several other sequences also approach the limit 0. Some such sequences that approach the limit 0 from the right are:

$$\langle x_{1:\infty} \rangle = \frac{1}{1}, \frac{1}{4}, \frac{1}{9}, \dots \quad \text{and} \quad \langle x_{1:\infty} \rangle = \frac{1}{1}, \frac{1}{8}, \frac{1}{27}, \dots ,$$

and some that approach the limit 0 from the left are:

$$\langle x_{1:\infty} \rangle = -\frac{1}{1}, -\frac{1}{2}, -\frac{1}{3}, \dots \quad \text{and} \quad \langle x_{1:\infty} \rangle = -\frac{1}{1}, -\frac{1}{4}, -\frac{1}{9}, \dots ,$$

and finally some that approach 0 from either side are:

$$\langle x_{1:\infty} \rangle = -\frac{1}{1}, +\frac{1}{2}, -\frac{1}{3}, \dots \quad \text{and} \quad \langle x_{1:\infty} \rangle = -\frac{1}{1}, +\frac{1}{4}, -\frac{1}{9}, \dots .$$

When we do not particularly care about the specifics of a sequence of real numbers  $\langle x_{1:\infty} \rangle$ , in terms of the exact values it takes for each  $i$ , but we are only interested that it converges to a limit  $a$  we write:

$$x \rightarrow a$$

and say that  $x$  approaches  $a$ . If we are only interested in those sequences that converge to the limit  $a$  from the right or left, we write:

$$x \rightarrow a^+ \quad \text{or} \quad x \rightarrow a^-$$

and say  $x$  approaches  $a$  from the right or left, respectively.

**Definition 4 (Limits of Functions)** We say a function  $f(x) : \mathbb{R} \rightarrow \mathbb{R}$  has a **limit**  $L \in \mathbb{R}$  as  $x$  approaches  $a$  and write:

$$\lim_{x \rightarrow a} f(x) = L ,$$

provided  $f(x)$  is arbitrarily close to  $L$  for all values of  $x$  that are sufficiently close to, but not equal to,  $a$ . We say that  $f$  has a **right limit**  $L_R$  or **left limit**  $L_L$  as  $x$  approaches  $a$  from the left or right, and write:

$$\lim_{x \rightarrow a^+} f(x) = L_R \quad \text{or} \quad \lim_{x \rightarrow a^-} f(x) = L_L ,$$

provided  $f(x)$  is arbitrarily close to  $L_R$  or  $L_L$  for all values of  $x$  that are sufficiently close to, but not equal to,  $a$  from the right of  $a$  or the left of  $a$ , respectively. When the limit is not an element of  $\mathbb{R}$  or when the left and right limits are distinct, we say that the limit does not exist.

**Example 11 (Limit of  $1/x^2$ )** Consider the function  $f(x) = \frac{1}{x^2}$ . Then

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{1}{x^2} = 1$$

exists since the limit  $1 \in \mathbb{R}$ , and the right and left limits are the same:

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} \frac{1}{x^2} = 1 \quad \text{and} \quad \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} \frac{1}{x^2} = 1 .$$

However, the following limit does not exist:

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$$

since  $\infty \notin \mathbb{R}$ .

**Example 12 (Limit of  $(1+x)^{\frac{1}{x}}$ )** The limit of  $f(x) = (1+x)^{\frac{1}{x}}$  as  $x$  approaches 0 exists and it is the Euler's constant  $e$ :

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \approx 2.71828 .$$

Notice that the above limit exists despite the fact that  $f(0) = (1+0)^{\frac{1}{0}}$  itself is undefined and does not exist.

**Example 13 (Limit of  $\frac{x^3-1}{x-1}$ )** For  $f(x) = \frac{x^3-1}{x-1}$ , this limit exists:

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{x^3 - 1}{x - 1} = \lim_{x \rightarrow 1} \frac{(x-1)(x^2 + x + 1)}{(x-1)} = \lim_{x \rightarrow 1} x^2 + x + 1 = 3$$

despite the fact that  $f(1) = \frac{1^3-1}{1-1} = \frac{0}{0}$  itself is undefined and does not exist.

Next we look at some examples of limits at infinity.

**Example 14 (Limit of  $(1 - \frac{\lambda}{n})^n$ )** The limit of  $f(n) = (1 - \frac{\lambda}{n})^n$  as  $n$  approaches  $\infty$  exists and it is  $e^{-\lambda}$ :

$$\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n = e^{-\lambda}.$$

**Example 15 (Limit of  $(1 - \frac{\lambda}{n})^{-\alpha}$ )** The limit of  $f(n) = (1 - \frac{\lambda}{n})^{-\alpha}$ , for some  $\alpha > 0$ , as  $n$  approaches  $\infty$  exists and it is 1 :

$$\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^{-\alpha} = 1.$$

**Definition 5 (Continuity of a function)** We say a real-valued function  $f(x) : D \rightarrow \mathbb{R}$  with the domain  $D \subset \mathbb{R}$  is **right continuous** or **left continuous** at a point  $a \in D$ , provided:

$$\lim_{x \rightarrow a^+} f(x) = f(a) \quad \text{or} \quad \lim_{x \rightarrow a^-} f(x) = f(a),$$

respectively. We say  $f$  is **continuous** at  $a \in D$ , provided:

$$\lim_{x \rightarrow a^+} f(x) = f(a) = \lim_{x \rightarrow a^-} f(x).$$

Finally,  $f$  is said to be continuous if  $f$  is continuous at every  $a \in D$ .

**Example 16 (Discontinuity of  $f(x) = (1+x)^{\frac{1}{x}}$  at 0)** Let us reconsider the function  $f(x) = (1+x)^{\frac{1}{x}} : \mathbb{R} \rightarrow \mathbb{R}$ . Clearly,  $f(x)$  is continuous at 1, since:

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} (1+x)^{\frac{1}{x}} = 2 = f(1) = (1+1)^{\frac{1}{1}},$$

but it is not continuous at 0, since:

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \approx 2.71828 \neq f(0) = (1+0)^{\frac{1}{0}}.$$

Thus,  $f(x)$  is not a continuous function over  $\mathbb{R}$ .

## 1.7 Elementary Number Theory

We introduce basic notions that we need from elementary number theory here. These notions include integer functions and modular arithmetic as they will be needed later on.

For any real number  $x$ :

$\lfloor x \rfloor := \max\{y : y \in \mathbb{Z} \text{ and } y \leq x\}$ , i.e., the greatest integer less than or equal to  $x$  (the **floor** of  $x$ ),  
 $\lceil x \rceil := \min\{y : y \in \mathbb{Z} \text{ and } y \geq x\}$ , i.e., the least integer greater than or equal to  $x$  (the **ceiling** of  $x$ ).

**Example 17 (Floors and ceilings)**

$$\lfloor 1 \rfloor = 1, \quad \lceil 1 \rceil = 1, \quad \lfloor 17.8 \rfloor = 17, \quad \lceil -17.8 \rceil = -18, \quad \lceil \sqrt{2} \rceil = 2, \quad \lfloor \pi \rfloor = 3, \quad \lceil \frac{1}{10^{100}} \rceil = 1.$$

**Labwork 18 (Floors and ceilings in MATLAB )** We can use MATLAB functions `floor` and `ceil` to compute  $\lfloor x \rfloor$  and  $\lceil x \rceil$ , respectively. Also, the argument  $x$  to these functions can be an array.

```
>> sqrt(2) % the square root of 2 is
ans =      1.4142
>> ceil(sqrt(2)) % ceiling of square root of 2
ans =      2
>> floor(-17.8) % floor of -17.8
ans =     -18
>> ceil([1 sqrt(2) pi -17.8 1/(10^100)]) %the ceiling of each element of an array
ans =      1      2      4     -17      1
>> floor([1 sqrt(2) pi -17.8 1/(10^100)]) % the floor of each element of an array
ans =      1      1      3    -18      0
```

**Classwork 19 (Relations between floors and ceilings)** Convince yourself of the following formulae. Use examples, plots and/or formal arguments.

$$\begin{aligned}\lceil x \rceil &= \lfloor x \rfloor \iff x \in \mathbb{Z} \\ \lceil x \rceil &= \lfloor x \rfloor + 1 \iff x \notin \mathbb{Z} \\ \lfloor -x \rfloor &= -\lceil x \rceil \\ x - 1 < \lfloor x \rfloor &\leq x \leq \lceil x \rceil < x + 1\end{aligned}$$

Let us define modular arithmetic next. Suppose  $x$  and  $y$  are any real numbers, i.e.  $x, y \in \mathbb{R}$ , we define the binary operation called “ $x \bmod y$ ” as:

$$x \bmod y := \begin{cases} x - y\lfloor x/y \rfloor & \text{if } y \neq 0 \\ x & \text{if } y = 0 \end{cases}$$

# Chapter 2

## Probability Model

### 2.1 Experiments

Ideas about chance events and random behaviour arose out of thousands of years of game playing, long before any attempt was made to use mathematical reasoning about them. Board and dice games were well known in Egyptian times, and Augustus Caesar gambled with dice. Calculations of odds for gamblers were put on a proper theoretical basis by Fermat and Pascal in the early 17th century.

**Definition 6** An **experiment** is an activity or procedure that produces distinct, well-defined possibilities called **outcomes**. The set of all outcomes is called the **sample space**, and is denoted by  $\Omega$ .

The subsets of  $\Omega$  are called **events**. A single outcome,  $\omega$ , when seen as a subset of  $\Omega$ , as in  $\{\omega\}$ , is called a **simple event**.

Events,  $E_1, E_2 \dots E_n$ , that cannot occur at the same time are called **mutually exclusive** events, or **pair-wise disjoint** events. This means that  $E_i \cap E_j = \emptyset$  where  $i \neq j$ .

**Example 20** Some standard examples of experiments are the following:

- $\Omega = \{\text{Defective, Non-defective}\}$  if our experiment is to inspect a light bulb.

There are only two outcomes here, so  $\Omega = \{\omega_1, \omega_2\}$  where  $\omega_1 = \text{Defective}$  and  $\omega_2 = \text{Non-defective}$ .

- $\Omega = \{\text{Heads, Tails}\}$  if our experiment is to note the outcome of a coin toss.

This time,  $\Omega = \{\omega_1, \omega_2\}$  where  $\omega_1 = \text{Heads}$  and  $\omega_2 = \text{Tails}$ .

- If our experiment is to roll a die then there are six outcomes corresponding to the number that shows on the top. For this experiment,  $\Omega = \{1, 2, 3, 4, 5, 6\}$ .

Some examples of events are the set of odd numbered outcomes  $A = \{1, 3, 5\}$ , and the set of even numbered outcomes  $B = \{2, 4, 6\}$ .

The simple events of  $\Omega$  are  $\{1\}, \{2\}, \{3\}, \{4\}, \{5\}$ , and  $\{6\}$ .

The outcome of a random experiment is uncertain until it is performed and observed. Note that sample spaces need to reflect the problem in hand. The example below is to convince you that an experiment's sample space is merely a collection of distinct elements called outcomes and these outcomes have to be *discernible in some well-specified sense* to the experimenter!

**Example 21** Consider a generic die-tossing experiment by a human experimenter. Here  $\Omega = \{\omega_1, \omega_2, \omega_3, \dots, \omega_6\}$ , but the experiment might correspond to rolling a die whose faces are:

1. sprayed with six different scents (nose!), or
2. studded with six distinctly flavoured candies (tongue!), or
3. contoured with six distinct bumps and pits (touch!), or
4. acoustically discernible at six different frequencies (ears!), or
5. painted with six different colours (eyes!), or
6. marked with six different numbers 1, 2, 3, 4, 5, 6 (eyes!), or , ...

These six experiments are equivalent as far as probability goes.

**Definition 7** A **trial** is a single performance of an experiment and it results in an outcome.

**Example 22** Some standard examples of a trial are:

- A roll of a die.
- A toss of a coin.
- A release of a chaotic double pendulum.

An experimenter often performs more than one trial. Repeated trials of an experiment forms the basis of science and engineering as the experimenter learns about the phenomenon by repeatedly performing the same mother experiment with possibly different outcomes. This repetition of trials in fact provides the very motivation for the definition of probability.

**Definition 8** An **n-product experiment** is obtained by repeatedly performing  $n$  trials of some experiment. The experiment that is repeated is called the “mother” experiment.

**Example 23 (Toss a coin  $n$  times)** Suppose our experiment entails tossing a coin  $n$  times and recording H for Heads and T for Tails. When  $n = 3$ , one possible outcome of this experiment is HHT, ie. a Head followed by another Head and then a Tail. Seven other outcomes are possible.

The sample space for “toss a coin three times” experiment is:

$$\Omega = \{H, T\}^3 = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\},$$

with a particular sample point or outcome  $\omega = HTH$ , and another distinct outcome  $\omega' = HHH$ . An event, say  $A$ , that ‘at least two Heads occur’ is the following subset of  $\Omega$ :

$$A = \{HHH, HHT, HTH, THH\}.$$

Another event, say  $B$ , that ‘no Heads occur’ is:

$$B = \{TTT\}$$

Note that the event  $B$  is also an outcome or sample point. Another interesting event is the empty set  $\emptyset \subset \Omega$ . The event that ‘nothing in the sample space occurs’ is  $\emptyset$ .

Figure 2.1: A binary tree whose leaves are all possible outcomes.

**Classwork 24 (A thrice-bifurcating tree of outcomes)** Can you think of a graphical way to enumerate the outcomes of the Experiment 23? Draw a diagram of this under the caption of Figure 2.1, using the caption as a hint (in other words, draw your own Figure 2.1).

#### EXPERIMENT SUMMARY

Experiment	–	an activity producing distinct outcomes.
$\Omega$	–	set of all outcomes of the experiment.
$\omega$	–	an individual outcome in $\Omega$ , called a simple event.
$A \subseteq \Omega$	–	a subset $A$ of $\Omega$ is an event.
Trial	–	one performance of an experiment resulting in 1 outcome.

## 2.2 Probability

The mathematical model for probability or the probability model is an axiomatic system that may be motivated by the intuitive idea of ‘long-term relative frequency’. If the axioms and definitions are intuitively motivated, the probability model simply follows from the application of logic to these axioms and definitions. No attempt to define probability in the real world is made. However, the application of probability models to real-world problems through statistical experiments has a fruitful track record. In fact, you are here for exactly this reason.

**Idea 9 (The long-term relative frequency (LTRF) idea)** Suppose we are interested in the fairness of a coin, i.e. if landing Heads has the same “probability” as landing Tails. We can toss it  $n$  times and call  $N(H, n)$  the fraction of times we observed Heads out of  $n$  tosses. Suppose that after conducting the tossing experiment 1000 times, we rarely observed Heads, e.g. 9 out of the 1000 tosses, then  $N(H, 1000) = 9/1000 = 0.009$ . Suppose we continued the number of tosses to a million and found that this number approached closer to 0.1, or, more generally,  $N(H, n) \rightarrow 0.1$  as  $n \rightarrow \infty$ . We might, at least intuitively, think that the coin is unfair and has a lower “probability” of 0.1 of landing Heads. We might think that it is fair had we observed  $N(H, n) \rightarrow 0.5$  as  $n \rightarrow \infty$ . Other crucial assumptions that we have made here are:

1. **Something Happens:** Each time we toss a coin, we are certain to observe Heads **or** Tails, denoted by  $H \cup T$ . The probability that “something happens” is 1. More formally:

$$N(H \cup T, n) = \frac{n}{n} = 1.$$

This is an intuitively reasonable assumption that simply says that one of the possible outcomes is certain to occur, provided the coin is not so thick that it can land on or even roll along its circumference.

2. **Addition Rule:** Heads and Tails are mutually exclusive events in any given toss of a coin, i.e. they cannot occur simultaneously. The intersection of mutually exclusive events is the empty set and is denoted by  $H \cap T = \emptyset$ . The event  $H \cup T$ , namely that the event that “coin lands Heads **or** coin lands Tails” satisfies:

$$N(H \cup T, n) = N(H, n) + N(T, n).$$

3. The coin-tossing experiment is repeatedly performed in an **independent** manner, i.e. the outcome of any individual coin-toss does not affect that of another. This is an intuitively reasonable assumption since the coin has no memory and the coin is tossed identically each time.

We will use the LTRF idea more generally to motivate a mathematical model of probability called probability model. Suppose  $A$  is an event associated with some experiment  $\mathcal{E}$ , so that  $A$  either does or does not occur when the experiment is performed. We want the probability that event  $A$  occurs in a specific performance of  $\mathcal{E}$ , denoted by  $\mathbf{P}(A)$ , to intuitively mean the following: if one were to perform a super-experiment  $\mathcal{E}^\infty$  by independently repeating the experiment  $\mathcal{E}$  and recording  $N(A, n)$ , the fraction of times  $A$  occurs in the first  $n$  performances of  $\mathcal{E}$  within the super-experiment  $\mathcal{E}^\infty$ . Then the LTRF idea suggests:

$$N(A, n) := \frac{\text{Number of times } A \text{ occurs}}{n = \text{Number of performances of } \mathcal{E}} \rightarrow \mathbf{P}(A), \text{ as } n \rightarrow \infty \quad (2.1)$$

Now, we are finally ready to define probability.

**Definition 10 (Probability)** Let  $\mathcal{E}$  be an experiment with sample space  $\Omega$ . Let  $\mathcal{F}$  denote a suitable collection of events in  $\Omega$  that satisfy the following conditions:

1. It (the collection) contains the sample space:  $\boxed{\Omega \in \mathcal{F}}$ .
2. It is closed under complementation:  $\boxed{A \in \mathcal{F} \implies A^c \in \mathcal{F}}$ .
3. It is closed under countable unions:  $\boxed{A_1, A_2, \dots \in \mathcal{F} \implies \bigcup_i A_i := A_1 \cup A_2 \cup \dots \in \mathcal{F}}$ .

Formally, this collection of events is called a **sigma field** or a **sigma algebra**. Our experiment  $\mathcal{E}$  has a sample space  $\Omega$  and a collection of events  $\mathcal{F}$  that satisfy the three condition.

Given a double, e.g.  $(\Omega, \mathcal{F})$ , **probability** is just a function  $\mathbf{P}$  which assigns each event  $A \in \mathcal{F}$  a number  $\mathbf{P}(A)$  in the real interval  $[0, 1]$ , i.e.  $\boxed{\mathbf{P} : \mathcal{F} \rightarrow [0, 1]}$ , such that:

1. The ‘Something Happens’ axiom holds, i.e.  $\boxed{\mathbf{P}(\Omega) = 1}$ .
2. The ‘Addition Rule’ axiom holds, i.e. for events  $A$  and  $B$ :

$$\boxed{A \cap B = \emptyset \implies \mathbf{P}(A \cup B) = \mathbf{P}(A) + \mathbf{P}(B)}.$$

### 2.2.1 Consequences of our Definition of Probability

It is important to realize that we accept the ‘addition rule’ as an axiom in our mathematical definition of probability (or our probability model) and we do **not** prove this rule. However, the facts which are stated (with proofs) below, are logical consequences of our definition of probability:

1. For any event  $A$ ,  $\boxed{\mathbf{P}(A^c) = 1 - \mathbf{P}(A)}.$

**Proof:** One line proof.

$$\overbrace{\mathbf{P}(A) + \mathbf{P}(A^c)}^{LHS} = \underbrace{\mathbf{P}(A \cup A^c)}_{+ \text{ rule } \because A \cap A^c = \emptyset} = \underbrace{\mathbf{P}(\Omega)}_{A \cup A^c = \Omega} = \underbrace{1}_{\because \mathbf{P}(\Omega) = 1} \stackrel{RHS}{\Rightarrow} \mathbf{P}(A^c) = 1 - \mathbf{P}(A)$$

- If  $A = \Omega$  then  $A^c = \Omega^c = \emptyset$  and  $\boxed{\mathbf{P}(\emptyset) = 1 - \mathbf{P}(\Omega) = 1 - 1 = 0}.$

2. For any two events  $A$  and  $B$ , we have the **inclusion-exclusion principle**:

$$\boxed{\mathbf{P}(A \cup B) = \mathbf{P}(A) + \mathbf{P}(B) - \mathbf{P}(A \cap B)}.$$

**Proof:** Since:

$$\begin{aligned} A &= (A \setminus B) \cup (A \cap B) && \text{and} && (A \setminus B) \cap (A \cap B) = \emptyset, \\ A \cup B &= (A \setminus B) \cup B && \text{and} && (A \setminus B) \cap B = \emptyset \end{aligned}$$

the addition rule implies that:

$$\begin{aligned} \mathbf{P}(A) &= \mathbf{P}(A \setminus B) + \mathbf{P}(A \cap B) \\ \mathbf{P}(A \cup B) &= \mathbf{P}(A \setminus B) + \mathbf{P}(B) \end{aligned}$$

Substituting the first equality above into the second, we get:

$$\mathbf{P}(A \cup B) = \mathbf{P}(A \setminus B) + \mathbf{P}(B) = \mathbf{P}(A) - \mathbf{P}(A \cap B) + \mathbf{P}(B)$$

3. For a sequence of mutually disjoint events  $A_1, A_2, A_3, \dots, A_n$ :

$$\boxed{A_i \cap A_j = \emptyset \text{ for any } i, j \implies \mathbf{P}(A_1 \cup A_2 \cup \dots \cup A_n) = \mathbf{P}(A_1) + \mathbf{P}(A_2) + \dots + \mathbf{P}(A_n).}$$

**Proof:** If  $A_1, A_2, A_3$  are mutually disjoint events, then  $A_1 \cup A_2$  is disjoint from  $A_3$ . Thus, two applications of the addition rule for disjoint events yields:

$$\mathbf{P}(A_1 \cup A_2 \cup A_3) = \mathbf{P}((A_1 \cup A_2) \cup A_3) \underset{+ \text{ rule}}{=} \mathbf{P}(A_1 \cup A_2) + \mathbf{P}(A_3) \underset{+ \text{ rule}}{=} \mathbf{P}(A_1) + \mathbf{P}(A_2) + \mathbf{P}(A_3)$$

The  $n$ -event case follows by mathematical induction.

We have formally defined the **probability model** specified by the **probability triple**  $(\Omega, \mathcal{F}, \mathbf{P})$  that can be used to model an **experiment**  $\mathcal{E}$ .

**Example 25 (First Ball out of NZ Lotto)** Let us observe the number on *the first ball that pops out in a New Zealand Lotto trial*. There are forty balls labelled 1 through 40 for this experiment and so the sample space is

$$\Omega = \{1, 2, 3, \dots, 39, 40\}.$$

Because the balls are vigorously whirled around inside the Lotto machine before the first one pops out, we can model each ball to pop out first with the same probability. So, we assign each outcome  $\omega \in \Omega$  the same probability of  $\frac{1}{40}$ , i.e., our probability model for this experiment is:

$$\mathbf{P}(\omega) = \frac{1}{40}, \text{ for each } \omega \in \Omega = \{1, 2, 3, \dots, 39, 40\} .$$

(Note: We sometimes abuse notation and write  $\mathbf{P}(\omega)$  instead of the more accurate but cumbersome  $\mathbf{P}(\{\omega\})$  when writing down probabilities of simple events.)

Figure 2.2 (a) shows the frequency of the first ball number in 1114 NZ Lotto draws. Figure 2.2 (b) shows the relative frequency, i.e., the frequency divided by 1114, the number of draws. Figure 2.2 (b) also shows the equal probabilities under our model.




Figure 2.2: First ball number in 1114 NZ Lotto draws from 1987 to 2008.

Next, let us take a detour into how one might interpret it in the real world. The following is an adaptation from Williams D, *Weighing the Odds: A Course in Probability and Statistics*, Cambridge University Press, 2001, which henceforth is abbreviated as WD2001.

### Probability Model

Sample space  $\Omega$

Sample point  $\omega$

(No counterpart)

Event A, a (suitable) subset of  $\Omega$

$\mathbf{P}(A)$ , a number between 0 and 1

### Real-world Interpretation

Set of all outcomes of an experiment

Possible outcome of an experiment

Actual outcome  $\omega^*$  of an experiment

The real-world event corresponding to A

occurs if and only if  $\omega^* \in A$

Probability that A will occur for an experiment yet to be performed

### Events in Probability Model

Sample space  $\Omega$

The  $\emptyset$  of  $\Omega$

The intersection  $A \cap B$

$A_1 \cap A_2 \cap \dots \cap A_n$

The union  $A \cup B$

$A_1 \cup A_2 \cup \dots \cup A_n$

$A^c$ , the complement of A

$A \setminus B$

$A \subset B$

### Real-world Interpretation

The certain even ‘something happens’

The impossible event ‘nothing happens’

‘Both A and B occur’

‘All of the events  $A_1, A_2, \dots, A_n$  occur simultaneously’

‘At least one of A and B occurs’

‘At least one of the events  $A_1, A_2, \dots, A_n$  occurs’

‘A does not occur’

‘A occurs, but B does not occur’

‘If A occurs, then B must occur’

In the probability model of Example 25, show that for any event  $E \subset \Omega$ ,

$$\mathbf{P}(E) = \frac{1}{40} \times \text{number of elements in } E .$$

Let  $E = \{\omega_1, \omega_2, \dots, \omega_k\}$  be an event with  $k$  outcomes (simple events). Then by the addition rule for mutually exclusive events we get:

$$\mathbf{P}(E) = \mathbf{P}(\{\omega_1, \omega_2, \dots, \omega_k\}) = \mathbf{P}\left(\bigcup_{i=1}^k \{\omega_i\}\right) = \sum_{i=1}^k \mathbf{P}(\{\omega_i\}) = \sum_{i=1}^k \frac{1}{40} = \frac{k}{40} .$$

### 2.2.2 Sigma Algebras of Typical Experiments\*

**Example 26 ('Toss a fair coin once')** Consider the 'Toss a fair coin once' experiment. What is its sample space  $\Omega$  and a reasonable collection of events  $\mathcal{F}$  that underpin this experiment?

$$\Omega = \{\text{H}, \text{T}\}, \quad \mathcal{F} = \{\Omega, \emptyset, \{\text{H}\}, \{\text{T}\}\} ,$$

A function that will satisfy the definition of probability for this collection of events  $\mathcal{F}$  and assign  $\mathbf{P}(\text{H}) = \frac{1}{2}$  is summarized below. First check that the above  $\mathcal{F}$  is a sigma-algebra. Draw a picture for  $\mathbf{P}$  with arrows that map elements in the domain  $\mathcal{F}$  given above to elements in its range.

Event $A \in \mathcal{F}$	$\mathbf{P} : \mathcal{F} \rightarrow [0, 1]$	$\mathbf{P}(A) \in [0, 1]$
$\Omega = \{\text{H}, \text{T}\} \bullet$	→	1
$\text{T} \bullet$	→	$1 - \frac{1}{2}$
$\text{H} \bullet$	→	$\frac{1}{2}$
$\emptyset \bullet$	→	0

**Classwork 27 (The trivial sigma algebra)** Note that  $\mathcal{F}' = \{\Omega, \emptyset\}$  is also a sigma algebra of the sample space  $\Omega = \{\text{H}, \text{T}\}$ . Can you think of a probability for the collection  $\mathcal{F}'$ ?

Event $A \in \mathcal{F}'$	$\mathbf{P} : \mathcal{F}' \rightarrow [0, 1]$	$\mathbf{P}(A) \in [0, 1]$
$\Omega = \{\text{H}, \text{T}\} \bullet$	→	
$\emptyset \bullet$	→	

Thus,  $\mathcal{F}$  and  $\mathcal{F}'$  are two distinct sigma algebras over our  $\Omega = \{\text{H}, \text{T}\}$ . Moreover,  $\mathcal{F}' \subset \mathcal{F}$  and is called a sub sigma algebra. Try to show that  $\{\Omega, \emptyset\}$  is the smallest possible sigma algebra over all possible sigma algebras over any given sample space  $\Omega$  (think of intersecting an arbitrary family of sigma algebras)?

Generally one encounters four types of sigma algebras and they are:

- When the sample space  $\Omega = \{\omega_1, \omega_2, \dots, \omega_k\}$  is a finite set with  $k$  outcomes and  $\mathbf{P}(\omega_i)$ , the probability for each outcome  $\omega_i \in \Omega$  is known, then one typically takes the sigma-algebra  $\mathcal{F}$  to be the set of all subsets of  $\Omega$  called the **power set** and denoted by  $2^\Omega$ . The probability of each event  $A \in 2^\Omega$  can be obtained by adding the probabilities of the outcomes in  $A$ , i.e.,  $\mathbf{P}(A) = \sum_{\omega_i \in A} \mathbf{P}(\omega_i)$ . Clearly,  $2^\Omega$  is indeed a sigma-algebra and it contains  $2^{\#\Omega}$  events in it.
- When the sample space  $\Omega = \{\omega_1, \omega_2, \dots\}$  is a countable set then one typically takes the sigma-algebra  $\mathcal{F}$  to be the set of all subsets of  $\Omega$ . Note that this is very similar to the case with finite  $\Omega$  except now  $\mathcal{F} = 2^\Omega$  could have uncountably many events in it.

3. If  $\Omega = \mathbb{R}^d$  for finite  $d \in \{1, 2, 3, \dots\}$  then the **Borel sigma-algebra** is the smallest sigma-algebra containing all **half-spaces**, i.e., sets of the form

$$\{x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : x_1 \leq c_1, x_2 \leq c_2, \dots, x_d \leq c_d\}, \quad \text{for any } c = (c_1, c_2, \dots, c_d) \in \mathbb{R}^d ,$$

When  $d = 1$  the half-spaces are the half-lines  $\{(-\infty, c] : c \in \mathbb{R}\}$  and when  $d = 2$  the half-spaces are the south-west quadrants  $\{(-\infty, c_1] \times (-\infty, c_2] : (c_1, c_2) \in \mathbb{R}^2\}$ , etc. (Equivalently, the Borel sigma-algebra is the smallest sigma-algebra containing all open sets in  $\mathbb{R}^d$ ).

4. Given a finite set  $\mathbb{S} = \{s_1, s_2, \dots, s_k\}$ , let  $\Omega$  be the sequence space  $\mathbb{S}^\infty := \mathbb{S} \times \mathbb{S} \times \mathbb{S} \times \dots$ , i.e., the set of sequences of infinite length that are made up of elements from  $\mathbb{S}$ . A set of the form

$$A_1 \times A_2 \times \dots \times A_n \times \mathbb{S} \times \mathbb{S} \times \dots, \quad A_k \subset \mathbb{S} \text{ for all } k \in \{1, 2, \dots, n\} ,$$

is called a **cylinder set**. The set of events in  $\mathbb{S}^\infty$  is the smallest sigma-algebra containing the cylinder sets.

## Exercises

**Ex. 2.1** — In English language text, the twenty six letters in the alphabet occur with the following frequencies:

E	13%	R	7.7%	A	7.3%	H	3.5%	F	2.8%	M	2.5%	W	1.6%	X	0.5%	J	0.2%
T	9.3%	O	7.4%	S	6.3%	L	3.5%	P	2.7%	Y	1.9%	V	1.3%	K	0.3%	Z	0.1%
N	7.8%	I	7.4%	D	4.4%	C	3%	U	2.7%	G	1.6%	B	0.9%	Q	0.3%		

Suppose you pick one letter at random from a randomly chosen English book from our central library with  $\Omega = \{A, B, C, \dots, Z\}$  (ignoring upper/lower cases), then what is the probability of these events?

- (a)  $\mathbf{P}(\{Z\})$
- (b)  $\mathbf{P}(\text{'picking any letter'})$
- (c)  $\mathbf{P}(\{E, Z\})$
- (d)  $\mathbf{P}(\text{'picking a vowel'})$
- (e)  $\mathbf{P}(\text{'picking any letter in the word WAZZZUP'})$
- (f)  $\mathbf{P}(\text{'picking any letter in the word WAZZZUP or a vowel'})$ .

**Ex. 2.2** — Find the sample spaces for the following experiments:

1. Tossing 2 coins whose faces are sprayed with black paint denoted by  $B$  and white paint denoted by  $W$ .
2. Drawing 4 screws from a bucket of left-handed and right-handed screws denoted by  $L$  and  $R$ , respectively.
3. Rolling a die and recording the number on the upturned face until the first 6 appears.

**Ex. 2.3** — Suppose we pick a letter at random from the word WAIMAKARIRI.

1. What is the sample space  $\Omega$ ?
2. What probabilities should be assigned to the outcomes?
3. What is the probability of *not* choosing the letter R?

**Ex. 2.4** — There are seventy five balls in total inside the Bingo Machine. Each ball is labelled by one of the following five letters: B, I, N, G, and O. There are fifteen balls labelled by each letter. The letter on the first ball that comes out of a BINGO machine after it has been well-mixed is the outcome of our experiment.

- (a) Write down the sample space of this experiment.
- (b) Find the probabilities of each simple event.
- (c) Show that  $\mathbf{P}(\Omega)$  is indeed 1.
- (d) Check that the addition rule for mutually exclusive events holds for the simple events  $\{B\}$  and  $\{I\}$ .
- (e) Consider the following events:  $C = \{B, I, G\}$  and  $D = \{G, I, N\}$ . Using the addition rule for two arbitrary events, find  $\mathbf{P}(C \cup D)$ .

### PROBABILITY SUMMARY

Axioms:

1. If  $A \subseteq \Omega$  then  $0 \leq \mathbf{P}(A) \leq 1$  and  $\mathbf{P}(\Omega) = 1$ .
2. If  $A, B$  are disjoint events, then  $\mathbf{P}(A \cup B) = \mathbf{P}(A) + \mathbf{P}(B)$ .  
[This is true only when  $A$  and  $B$  are disjoint.]
3. If  $A_1, A_2, \dots$  are disjoint then  $\mathbf{P}(A_1 \cup A_2 \cup \dots) = \mathbf{P}(A_1) + \mathbf{P}(A_2) + \dots$

Rules:

$$\begin{aligned}\mathbf{P}(A^c) &= 1 - \mathbf{P}(A) \\ \mathbf{P}(A \cup B) &= \mathbf{P}(A) + \mathbf{P}(B) - \mathbf{P}(A \cap B) \quad [\text{always true}]\end{aligned}$$

## 2.3 Conditional Probability

Next, we define conditional probability and the notion of independence of events. We use the LTRF idea to motivate the definition.

**Idea 11 (LTRF intuition for conditional probability)** Let  $A$  and  $B$  be any two events associated with our experiment  $\mathcal{E}$  with  $\mathbf{P}(A) \neq 0$ . The ‘conditional probability that  $B$  occurs given that  $A$  occurs’ denoted by  $\mathbf{P}(B|A)$  is again intuitively underpinned by the super-experiment  $\mathcal{E}^\infty$  which is the ‘independent’ repetition of our original experiment  $\mathcal{E}$  ‘infinitely’ often. The LTRF idea is that  $\mathbf{P}(B|A)$  is the long-term proportion of those experiments on which  $A$  occurs that  $B$  also occurs.

Recall that  $N(A, n)$  as defined in (2.1) is the fraction of times  $A$  occurs out of  $n$  independent repetitions of our experiment  $\mathcal{E}$  (ie. the experiment  $\mathcal{E}^n$ ). If  $A \cap B$  is the event that ‘ $A$  and  $B$  occur simultaneously’, then we intuitively want

$$\mathbf{P}(B|A) \quad “\rightarrow” \quad \frac{N(A \cap B, n)}{N(A, n)} = \frac{N(A \cap B, n)/n}{N(A, n)/n} = \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(A)}$$

as our  $\mathcal{E}^n \rightarrow \mathcal{E}^\infty$ . So, we **define** conditional probability as we want.

**Definition 12 (Conditional Probability)** Suppose we are given an experiment  $\mathcal{E}$  with a triple  $(\Omega, \mathcal{F}, \mathbf{P})$ . Let  $A$  and  $B$  be events, ie.  $A, B \in \mathcal{F}$ , such that  $\mathbf{P}(A) \neq 0$ . Then, we define the **conditional probability** of  $B$  given  $A$  by,

$$\mathbf{P}(B|A) := \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(A)} . \quad (2.2)$$

Note that for a **fixed** event  $A \in \mathcal{F}$  with  $\mathbf{P}(A) > 0$  and **any** event  $B \in \mathcal{F}$ , the conditional probability  $\mathbf{P}(B|A)$  is a probability as in Definition 10, ie. a function:

$$\mathbf{P}(B|A) : \mathcal{F} \rightarrow [0, 1]$$

that assigns to each  $B \in \mathcal{F}$  a number in the interval  $[0, 1]$ , such that,

1.  $\mathbf{P}(\Omega|A) = 1$  Meaning ‘Something Happens given the event A happens’
2. The ‘Addition Rule’ axiom holds, ie. for events  $B_1, B_2 \in \mathcal{F}$ ,

$$B_1 \cap B_2 = \emptyset \text{ implies } \mathbf{P}(B_1 \cup B_2|A) = \mathbf{P}(B_1|A) + \mathbf{P}(B_2|A) .$$

3. For mutually exclusive or pairwise-disjoint events,  $B_1, B_2, \dots$ ,

$$\mathbf{P}(B_1 \cup B_2 \cup \dots |A) = \mathbf{P}(B_1|A) + \mathbf{P}(B_2|A) + \dots .$$

From the definition of conditional probability we get the following rules:

1. Complementation rule:  $\mathbf{P}(B|A) = 1 - \mathbf{P}(B^c|A)$  .
2. Addition rule for two arbitrary events  $B_1$  and  $B_2$ :

$$\mathbf{P}(B_1 \cup B_2|A) = \mathbf{P}(B_1|A) + \mathbf{P}(B_2|A) - \mathbf{P}(B_1 \cap B_2|A) .$$

3. Multiplication rule for two likely events:

If  $A$  and  $B$  are events, and if  $\mathbf{P}(A) \neq 0$  and  $\mathbf{P}(B) \neq 0$ , then

$$\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B|A) = \mathbf{P}(B)\mathbf{P}(A|B) .$$

**Example 28 (Wasserman03, p. 11)** A medical test for a disease  $D$  has outcomes + and -. the probabilities are:

	Have Disease ( $D$ )	Don't have disease ( $D^c$ )
Test positive (+)	0.009	0.099
Test negative (-)	0.001	0.891

Using the definition of conditional probability, we can compute the conditional probability that you test positive given that you have the disease:

$$\mathbf{P}(+|D) = \frac{\mathbf{P}(+ \cap D)}{\mathbf{P}(D)} = \frac{0.009}{0.009 + 0.001} = 0.9 ,$$

and the conditional probability that you test negative given that you don't have the disease:

$$\mathbf{P}(-|D^c) = \frac{\mathbf{P}(- \cap D^c)}{\mathbf{P}(D^c)} = \frac{0.891}{0.099 + 0.891} \approx 0.9 .$$

Thus, the test is quite accurate since sick people test positive 90% of the time and healthy people test negative 90% of the time.

Now, suppose you go for a test and test positive. What is the probability that you have the disease ?

$$\mathbf{P}(D|+) = \frac{\mathbf{P}(D \cap +)}{\mathbf{P}(+)} = \frac{0.009}{0.009 + 0.099} \approx 0.08$$

Most people who are not used to the definition of conditional probability would intuitively associate a number much bigger than 0.08 for the answer. Interpret conditional probability in terms of the meaning of the numbers that appear in the numerator and denominator of the above calculations.

Next we look at one of the most elegant applications of the definition of conditional probability along with the addition rule for a partition of  $\Omega$ .

**Proposition 13 (Bayes' Theorem, 1763)** Suppose the events  $A_1, A_2, \dots, A_k \in \mathcal{F}$ , with  $\mathbf{P}(A_h) > 0$  for each  $h \in \{1, 2, \dots, k\}$ , partition the sample space  $\Omega$ , ie. they are mutually exclusive (disjoint) and exhaustive events with positive probability:

$$A_i \cap A_j = \emptyset, \text{ for any distinct } i, j \in \{1, 2, \dots, k\}, \quad \bigcup_{h=1}^k A_h = \Omega, \quad \mathbf{P}(A_h) > 0$$

Thus, precisely one of the  $A_h$ 's will occur on any performance of our experiment  $\mathcal{E}$ .

Let  $B \in \mathcal{F}$  be some event with  $\mathbf{P}(B) > 0$ , then

$$\mathbf{P}(A_h|B) = \frac{\mathbf{P}(B|A_h)\mathbf{P}(A_h)}{\sum_{h=1}^k \mathbf{P}(B|A_h)\mathbf{P}(A_h)} \quad (2.3)$$

**Proof:** We apply elementary set theory, the definition of conditional probability  $k + 2$  times and the addition rule once:

$$\begin{aligned} \mathbf{P}(A_h|B) &= \frac{\mathbf{P}(A_h \cap B)}{\mathbf{P}(B)} = \frac{\mathbf{P}(B \cap A_h)}{\mathbf{P}(B)} = \frac{\mathbf{P}(B|A_h)\mathbf{P}(A_h)}{\mathbf{P}(B)} \\ &= \frac{\mathbf{P}(B|A_h)\mathbf{P}(A_h)}{\mathbf{P}\left(\bigcup_{h=1}^k (B \cap A_h)\right)} = \frac{\mathbf{P}(B|A_h)\mathbf{P}(A_h)}{\sum_{h=1}^k \mathbf{P}(B \cap A_h)} \\ &= \frac{\mathbf{P}(B|A_h)\mathbf{P}(A_h)}{\sum_{h=1}^k \mathbf{P}(B|A_h)\mathbf{P}(A_h)} \end{aligned}$$

The operations done to the denominator in the proof above:

$$\mathbf{P}(B) = \sum_{h=1}^k \mathbf{P}(B|A_h)\mathbf{P}(A_h) \quad (2.4)$$

is also called ‘the law of total probability’ or ‘the total probability theorem’. We call  $\mathbf{P}(A_h)$  the **prior probability of  $A_h$**  and  $\mathbf{P}(A_h|B)$  the **posterior probability of  $A_h$** .

**Example 29 (Wasserman2003 p.12)** Suppose Larry divides his email into three categories:  $A_1$  = “spam”,  $A_2$  = “low priority”, and  $A_3$  = “high priority”. From previous experience, he finds that  $\mathbf{P}(A_1) = 0.7$ ,  $\mathbf{P}(A_2) = 0.2$  and  $\mathbf{P}(A_3) = 0.1$ . Note that  $\mathbf{P}(A_1 \cup A_2 \cup A_3) = \mathbf{P}(\Omega) = 0.7 + 0.2 + 0.1 = 1$ . Let  $B$  be the event that the email contains the word “free.” From previous experience,  $\mathbf{P}(B|A_1) = 0.9$ ,  $\mathbf{P}(B|A_2) = 0.01$  and  $\mathbf{P}(B|A_3) = 0.01$ . Note that  $\mathbf{P}(B|A_1) + \mathbf{P}(B|A_2) + \mathbf{P}(B|A_3) = 0.9 + 0.01 + 0.01 \neq 1$ . Now, suppose Larry receives an email with the word “free.” What is the probability that it is “spam,” “low priority,” and “high priority” ?

$$\mathbf{P}(A_1|B) = \frac{\mathbf{P}(B|A_1)\mathbf{P}(A_1)}{\mathbf{P}(B|A_1)\mathbf{P}(A_1) + \mathbf{P}(B|A_2)\mathbf{P}(A_2) + \mathbf{P}(B|A_3)\mathbf{P}(A_3)} = \frac{0.9 \times 0.7}{(0.9 \times 0.7) + (0.01 \times 0.2) + (0.01 \times 0.1)} = \frac{0.63}{0.633} \approx 0.995$$

$$\mathbf{P}(A_2|B) = \frac{\mathbf{P}(B|A_2)\mathbf{P}(A_2)}{\mathbf{P}(B|A_1)\mathbf{P}(A_1) + \mathbf{P}(B|A_2)\mathbf{P}(A_2) + \mathbf{P}(B|A_3)\mathbf{P}(A_3)} = \frac{0.01 \times 0.2}{(0.9 \times 0.7) + (0.01 \times 0.2) + (0.01 \times 0.1)} = \frac{0.002}{0.633} \approx 0.003$$


$$\mathbf{P}(A_3|B) = \frac{\mathbf{P}(B|A_3)\mathbf{P}(A_3)}{\mathbf{P}(B|A_1)\mathbf{P}(A_1) + \mathbf{P}(B|A_2)\mathbf{P}(A_2) + \mathbf{P}(B|A_3)\mathbf{P}(A_3)} = \frac{0.01 \times 0.1}{(0.9 \times 0.7) + (0.01 \times 0.2) + (0.01 \times 0.1)} = \frac{0.001}{0.633} \approx 0.002$$

Note that  $\mathbf{P}(A_1|B) + \mathbf{P}(A_2|B) + \mathbf{P}(A_3|B) = 0.995 + 0.003 + 0.002 = 1$ .

**Example 30 (Urn with red and black balls)** A well-mixed urn contains five red and ten black balls. We draw two balls from the urn without replacement. What is the probability that the second ball drawn is red?

This is easy to see if we draw a probability tree diagram. The first split in the tree is based on the outcome of the first draw and the second on the outcome of the last draw. The outcome of the first draw dictates the probabilities for the second one since we are sampling without replacement. We multiply the probabilities on the edges to get probabilities of the four endpoints, and then sum the ones that correspond to red in the second draw, that is

$$P(\text{second ball is red}) = 4/42 + 10/42 = 1/3 .$$



Alternatively, use the total probability theorem to break the problem down into manageable pieces. Let  $R_1 = \{(red, red), (red, black)\}$  and  $R_2 = \{(red, red), (black, red)\}$  be the events corresponding to a red ball in the 1st and 2nd draws, respectively, and let  $B_1 = \{(black, red), (black, black)\}$  be the event of a black ball on the first draw.

Now  $R_1$  and  $B_1$  partition  $\Omega$  so we can write:

$$\begin{aligned} P(R_2) &= \mathbf{P}(R_2 \cap R_1) + \mathbf{P}(R_2 \cap B_1) \\ &= \mathbf{P}(R_2|R_1)\mathbf{P}(R_1) + \mathbf{P}(R_2|B_1)\mathbf{P}(B_1) \\ &= (4/14)(1/3) + (5/14)(2/3) = 1/3 . \end{aligned}$$

### 2.3.1 Independence and Dependence

**Definition 14 (Independence of two events)** Any two events  $A$  and  $B$  are said to be **independent** if and only if

$$\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B) . \quad (2.5)$$

Let us make sense of this definition in terms of our previous definitions. When  $\mathbf{P}(A) = 0$  or  $\mathbf{P}(B) = 0$ , both sides of the above equality are 0. If  $\mathbf{P}(A) \neq 0$ , then rearranging the above equation we get:

$$\frac{\mathbf{P}(A \cap B)}{\mathbf{P}(A)} = \mathbf{P}(B) .$$

But, the LHS is  $\mathbf{P}(B|A)$  by definition 2.2, and thus for independent events  $A$  and  $B$ , we get:

$$\mathbf{P}(B|A) = \mathbf{P}(B) .$$

This says that information about the occurrence of  $A$  does not affect the occurrence of  $B$ . If  $\mathbf{P}(B) \neq 0$ , then an analogous argument:

$$\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B) \iff \mathbf{P}(B \cap A) = \mathbf{P}(A)\mathbf{P}(B) \iff \frac{\mathbf{P}(B \cap A)}{\mathbf{P}(B)} = \mathbf{P}(A) \iff \mathbf{P}(A|B) = \mathbf{P}(A) ,$$

says that information about the occurrence of  $B$  does not affect the occurrence of  $A$ . Therefore, the probability of their joint occurrence  $\mathbf{P}(A \cap B)$  is simply the product of their individual probabilities  $\mathbf{P}(A)\mathbf{P}(B)$ .

**Definition 15 (Independence of a sequence of events)** We say that a finite or infinite sequence of events  $A_1, A_2, \dots$  are independent if whenever  $i_1, i_2, \dots, i_k$  are distinct elements from the set of indices  $\mathbb{N}$ , such that  $A_{i_1}, A_{i_2}, \dots, A_{i_k}$  are defined (elements of  $\mathcal{F}$ ), then

$$\mathbf{P}(A_{i_1} \cap A_{i_2} \dots \cap A_{i_k}) = \mathbf{P}(A_{i_1})\mathbf{P}(A_{i_2}) \cdots \mathbf{P}(A_{i_k})$$

**Example 31 (Some Standard Examples)** A sequence of events in a sequence of independent trials is independent.

(a) Suppose you toss a fair coin twice such that the first toss is independent of the second. Then,

$$\mathbf{P}(\text{Heads on the first toss} \cap \text{Tails on the second toss}) = \mathbf{P}(\mathsf{H})\mathbf{P}(\mathsf{T}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} .$$

(b) Suppose you independently toss a fair die three times. Let  $E_i$  be the event that the outcome is an even number on the  $i$ -th trial. The probability of getting an even number in all three trials is:

$$\begin{aligned} \mathbf{P}(E_1 \cap E_2 \cap E_3) &= \mathbf{P}(E_1)\mathbf{P}(E_2)\mathbf{P}(E_3) \\ &= (\mathbf{P}(\{2, 4, 6\}))^3 \\ &= (\mathbf{P}(\{2\}) \cup \{4\} \cup \{6\}))^3 \\ &= (\mathbf{P}(\{2\}) + \mathbf{P}(\{4\}) + \mathbf{P}(\{6\}))^3 \\ &= \left(\frac{1}{6} + \frac{1}{6} + \frac{1}{6}\right)^3 = \left(\frac{1}{2}\right)^3 = \frac{1}{8} . \end{aligned}$$

(c) Suppose you toss a fair coin independently  $m$  times. Then each of the  $2^m$  possible outcomes in the sample space  $\Omega$  has equal probability of  $\frac{1}{2^m}$  due to independence.

**Example 32 (dependence and independence)** Suppose we toss two fair dice. Let  $A$  denote the event that the sum of the dice is six and  $B$  denote the event that the first die equals four. The sample space encoding the thirty six ordered pairs of outcomes for the two dice is  $\Omega =$

$\{(1,1), (1,2), \dots, (1,6), (2,1), \dots, (2,6), \dots, (5,6), (6,6)\}$  and due to independence  $\mathbf{P}(\omega) = 1/36$  for each  $\omega \in \Omega$ . Then

$$\mathbf{P}(A \cap B) = \mathbf{P}(\{(4,2)\}) = \frac{1}{36},$$

but

$$\begin{aligned}\mathbf{P}(A)\mathbf{P}(B) &= \mathbf{P}(\{(1,5), (2,4), (3,3), (4,2), (5,1)\})\mathbf{P}(\{(4,1), (4,2), (4,3), (4,4), (4,5), (4,6)\}) \\ &= \frac{5}{36} \times \frac{6}{36} = \frac{5}{36} \times \frac{1}{6} = \frac{5}{216},\end{aligned}$$

and therefore  $A$  and  $B$  are not independent. The reason for the events  $A$  and  $B$  being dependent is clear because the chance of getting a total of six depends on the outcome of the first die (not being six).

Now, let  $C$  be the event that the sum of the two dice equals seven. Then

$$\mathbf{P}(C \cap B) = \mathbf{P}(\{(4,3)\}) = \frac{1}{36},$$

while

$$\begin{aligned}\mathbf{P}(C \cap B) &= \mathbf{P}(\{(1,6), (2,5), (3,4), (4,3), (5,2), (6,1)\})\mathbf{P}(\{(4,1), (4,2), (4,3), (4,4), (4,5), (4,6)\}) \\ &= \frac{6}{36} \times \frac{6}{36} = \frac{1}{36},\end{aligned}$$

and therefore  $C$  and  $B$  are independent events. Once again this is clear because the chance of getting a total of seven does not depend any more on the outcome of the first die (it is allowed to be any one of the six possible outcomes).

**Example 33 (Pairwise independent events that are not jointly independent)** Let a ball be drawn from an well-stirred urn containing four balls labelled 1,2,3,4. Consider the events  $A = \{1,2\}$ ,  $B = \{1,3\}$  and  $C = \{1,4\}$ . Then,

$$\begin{aligned}\mathbf{P}(A \cap B) &= \mathbf{P}(A)\mathbf{P}(B) = \frac{2}{4} \times \frac{2}{4} = \frac{1}{4}, \\ \mathbf{P}(A \cap C) &= \mathbf{P}(A)\mathbf{P}(C) = \frac{2}{4} \times \frac{2}{4} = \frac{1}{4}, \\ \mathbf{P}(B \cap C) &= \mathbf{P}(B)\mathbf{P}(C) = \frac{2}{4} \times \frac{2}{4} = \frac{1}{4},\end{aligned}$$

but,

$$\frac{1}{4} = \mathbf{P}(\{1\}) = \mathbf{P}(A \cap B \cap C) \neq \mathbf{P}(A)\mathbf{P}(B)\mathbf{P}(C) = \frac{2}{4} \times \frac{2}{4} \times \frac{2}{4} = \frac{1}{8}.$$

Therefore, inspite of being pairwise independent, the events  $A$ ,  $B$  and  $C$  are not jointly independent.

## Exercises

**Ex. 2.5 —** What gives the greater probability of hitting some target at least once:

- 1.hitting in a shot with probability  $\frac{1}{2}$  and firing 1 shot, or
- 2.hitting in a shot with probability  $\frac{1}{3}$  and firing 2 shots?

First guess. Then calculate.

**Ex. 2.6** — Suppose we independently roll two fair dice each of whose faces are marked by numbers 1,2,3,4, 5 and 6.

- 1.List the sample space for the experiment if we note the numbers on the 2 upturned faces.
- 2.What is the probability of obtaining a sum greater than 4 but less than 7?

**Ex. 2.7** — Based on past experience, 70% of students in a certain course pass the midterm test. The final exam is passed by 80% of those who passed the midterm test, but only by 40% of those who fail the midterm test. What fraction of students pass the final exam?

**Ex. 2.8** — A small brewery has two bottling machines. Machine 1 produces 75% of the bottles and machine 2 produces 25%. One out of every 20 bottles filled by machine 1 is rejected for some reason, while one out of every 30 bottles filled by machine 2 is rejected. What is the probability that a randomly selected bottle comes from machine 1 given that it is accepted?

**Ex. 2.9** — A process producing micro-chips, produces 5% defective, at random. Each micro-chip is tested, and the test will correctly detect a defective one  $4/5$  of the time, and if a good micro-chip is tested the test will declare it is defective with probability  $1/10$ .

- (a)If a micro-chip is chosen at random, and tested to be good, what was the probability that it was defective anyway?
- (b)If a micro-chip is chosen at random, and tested to be defective, what was the probability that it was good anyway?
- (c)If 2 micro-chips are tested and determined to be good, what is the probability that at least one is in fact defective?

**Ex. 2.10** — Suppose that  $\frac{2}{3}$  of all gales are force 1,  $\frac{1}{4}$  are force 2 and  $\frac{1}{12}$  are force 3. Furthermore, the probability that force 1 gales cause damage is  $\frac{1}{4}$ , the probability that force 2 gales cause damage is  $\frac{2}{3}$  and the probability that force 3 gales cause damage is  $\frac{5}{6}$ .

- (a)If a gale is reported, what is the probability of it causing damage?
- (b)If the gale caused damage, find the probabilities that it was of: force 1; force 2; force 3.
- (c)If the gale did NOT cause damage, find the probabilities that it was of: force 1; force 2; force 3.

**Ex. 2.11** — \*\*The sensitivity and specificity of a medical diagnostic test for a disease are defined as follows:

$$\begin{aligned}\text{sensitivity} &= \mathbf{P}(\text{test is positive} \mid \text{patient has the disease}) , \\ \text{specificity} &= \mathbf{P}(\text{test is negative} \mid \text{patient does not have the disease}) .\end{aligned}$$

Suppose that a medical test has a sensitivity of 0.7 and a specificity of 0.95. If the prevalence of the disease in the general population is 1%, find

- (a)the probability that a patient who tests positive actually has the disease,
- (b)the probability that a patient who tests negative is free from the disease.

**Ex. 2.12** — \*\*The detection rate and false alarm rate of an intrusion sensor are defined as

$$\begin{aligned}\text{detection rate} &= \mathbf{P}(\text{detection declared} \mid \text{intrusion}) , \\ \text{false alarm rate} &= \mathbf{P}(\text{detection declared} \mid \text{no intrusion}) .\end{aligned}$$

If the detection rate is 0.999 and the false alarm rate is 0.001, and the probability of an intrusion occurring is 0.01, find

- (a)the probability that there is an intrusion when a detection is declared,
- (b)the probability that there is no intrusion when no detection is declared.

**Ex. 2.13** — \*\*Let  $A$  and  $B$  be events such that  $\mathbf{P}(A) \neq 0$  and  $\mathbf{P}(B) \neq 0$ . When  $A$  and  $B$  are disjoint, are they also independent? Explain clearly why or why not.

#### CONDITIONAL PROBABILITY SUMMARY

$\mathbf{P}(A|B)$  means the probability that  $A$  occurs given that  $B$  has occurred.

$$\mathbf{P}(A|B) = \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(B)} = \frac{\mathbf{P}(A)\mathbf{P}(B|A)}{\mathbf{P}(B)} \quad \text{if } \mathbf{P}(B) \neq 0$$

$$\mathbf{P}(B|A) = \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(A)} = \frac{\mathbf{P}(B)\mathbf{P}(A|B)}{\mathbf{P}(A)} \quad \text{if } \mathbf{P}(A) \neq 0$$

Conditional probabilities obey the axioms and rules of probability.

# Chapter 3

## Random Variables

It can be inconvenient to work with a set of outcomes  $\Omega$  upon which arithmetic is not possible. We are often measuring our outcomes with subsets of real numbers. Some examples include:

Experiment	Possible measured outcomes
Counting the number of typos up to now	$\mathbb{Z}_+ := \{0, 1, 2, \dots\} \subset \mathbb{R}$
Length in centi-meters of some shells on New Brighton beach	$(0, +\infty) \subset \mathbb{R}$
Waiting time in minutes for the next Orbiter bus to arrive	$\mathbb{R}_+ := [0, \infty) \subset \mathbb{R}$
Vertical displacement from current position of a pollen on water	$\mathbb{R}$

### 3.1 Basic Definitions

To take advantage of our measurements over the real numbers, in terms of its metric structure and arithmetic, we need to formally define this measurement process using the notion of a random variable.

**Definition 16 (Random Variable)** Let  $(\Omega, \mathcal{F}, P)$  be some probability triple. Then, a **Random Variable (RV)**, say  $X$ , is a function from the sample space  $\Omega$  to the set of real numbers  $\mathbb{R}$

$$X : \Omega \rightarrow \mathbb{R}$$

such that for every  $x \in \mathbb{R}$ , the inverse image of the half-open real interval  $(-\infty, x]$  is an element of the collection of events  $\mathcal{F}$ , i.e.:

$$\text{for every } x \in \mathbb{R}, \quad X^{[-1]}((-\infty, x]) := \{\omega : X(\omega) \leq x\} \in \mathcal{F}.$$

This definition can be summarised by the statement that a RV is an  $\mathcal{F}$ -measurable map. We assign probability to the RV  $X$  as follows:

$$\mathbf{P}(X \leq x) = \mathbf{P}(X^{[-1]}((-\infty, x])) := \mathbf{P}(\{\omega : X(\omega) \leq x\}). \quad (3.1)$$

**Definition 17 (Distribution Function)** The **Distribution Function (DF)** or **Cumulative Distribution Function (CDF)** of any RV  $X$ , over a probability triple  $(\Omega, \mathcal{F}, P)$ , denoted by  $F$  is:

$$F(x) := \mathbf{P}(X \leq x) = \mathbf{P}(\{\omega : X(\omega) \leq x\}), \quad \text{for any } x \in \mathbb{R}. \quad (3.2)$$


Thus,  $F(x)$  or simply  $F$  is a non-decreasing, right continuous,  $[0, 1]$ -valued function over  $\mathbb{R}$ . When a RV  $X$  has DF  $F$  we write  $X \sim F$ .

A special RV that often plays the role of ‘building-block’ in Probability and Statistics is the indicator function of an event  $A$  that tells us whether the event  $A$  has occurred or not. Recall that an event belongs to the collection of possible events  $\mathcal{F}$  for our experiment.

**Definition 18 (Indicator Function)** The **Indicator Function** of an event  $A$  denoted  $\mathbb{1}_A$  is defined as follows:

$$\mathbb{1}_A(\omega) := \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases} \quad (3.3)$$

Figure 3.1: The Indicator function of event  $A \in \mathcal{F}$  is a RV  $\mathbb{1}_A$  with DF  $F$



**Classwork 34 (Indicator function is a random variable)** Let us convince ourselves that  $\mathbb{1}_A$  is really a RV. For  $\mathbb{1}_A$  to be a RV, we need to verify that for any real number  $x \in \mathbb{R}$ , the inverse image  $\mathbb{1}_A^{[-1]}( (-\infty, x] )$  is an event, ie :

$$\mathbb{1}_A^{[-1]}( (-\infty, x] ) := \{\omega : \mathbb{1}_A(\omega) \leq x\} \in \mathcal{F} .$$

All we can assume about the collection of events  $\mathcal{F}$  is that it contains the event  $A$  and that it is a sigma algebra. A careful look at the Figure 3.1 yields:


$$\mathbb{1}_A^{[-1]}( (-\infty, x] ) := \{\omega : \mathbb{1}_A(\omega) \leq x\} = \begin{cases} \emptyset & \text{if } x < 0 \\ A^c & \text{if } 0 \leq x < 1 \\ A \cup A^c = \Omega & \text{if } 1 \leq x \end{cases}$$

Thus,  $\mathbb{1}_A^{[-1]}( (-\infty, x] )$  is one of the following three sets that belong to  $\mathcal{F}$ ; (1)  $\emptyset$ , (2)  $A^c$  and (3)  $\Omega$  depending on the value taken by  $x$  relative to the interval  $[0, 1]$ . We have proved that  $\mathbb{1}_A$  is indeed a RV.

Some useful properties of the Indicator Function are:

$$\mathbb{1}_{A^c} = 1 - \mathbb{1}_A, \quad \mathbb{1}_{A \cap B} = \mathbb{1}_A \mathbb{1}_B, \quad \mathbb{1}_{A \cup B} = \mathbb{1}_A + \mathbb{1}_B - \mathbb{1}_A \mathbb{1}_B$$

We slightly abuse notation when  $A$  is a single element set by ignoring the curly braces.

Figure 3.2: A RV  $X$  from a sample space  $\Omega$  with 8 elements to  $\mathbb{R}$  and its DF  $F$ 

**Classwork 35 (A random variable with three values and eight sample points)** Consider the RV  $X$  of Figure 3.2. Let the events  $A = \{\omega_1, \omega_2\}$ ,  $B = \{\omega_3, \omega_4, \omega_5\}$  and  $C = \{\omega_6, \omega_7, \omega_8\}$ . Define the RV  $X$  formally. What sets should  $\mathcal{F}$  minimally include? What do you need to do to make sure that  $\mathcal{F}$  is a sigma algebra?

## 3.2 An Elementary Discrete Random Variable

When a RV takes at most countably many values from a discrete set  $\mathbb{D} \subset \mathbb{R}$ , we call it a **discrete** RV. Often,  $\mathbb{D}$  is the set of integers  $\mathbb{Z}$ .

**Definition 19 (probability mass function (PMF))** Let  $X$  be a discrete RV over a probability triple  $(\Omega, \mathcal{F}, P)$ . We define the **probability mass function** (PMF)  $f$  of  $X$  to be the function  $f : \mathbb{D} \rightarrow [0, 1]$  defined as follows:

$$f(x) := \mathbf{P}(X = x) = \mathbf{P}(\{\omega : X(\omega) = x\}), \quad \text{where } x \in \mathbb{D}.$$

The DF  $F$  and PMF  $f$  for a discrete RV  $X$  satisfy the following:

1. For any  $x \in \mathbb{R}$ ,

$$\mathbf{P}(X \leq x) = F(x) = \sum_{\mathbb{D} \ni y \leq x} f(y) := \sum_{y \in \mathbb{D} \cap (-\infty, x]} f(y).$$

2. For any  $a, b \in \mathbb{D}$  with  $a < b$ ,

$$\mathbf{P}(a < X \leq b) = F(b) - F(a) = \sum_{y \in \mathbb{D} \cap (a, b]} f(y) .$$

In particular, when  $\mathbb{D} = \mathbb{Z}$  and  $a = b - 1$ ,

$$\mathbf{P}(b - 1 < X \leq b) = F(b) - F(b - 1) = f(b) = \mathbf{P}(\{\omega : X(\omega) = b\}) .$$

3. And of course

$$\sum_{x \in \mathbb{D}} f(x) = 1$$

The Indicator Function  $\mathbf{1}_A$  of the event that ‘ $A$  occurs’ for the  $\theta$ -specific experiment  $\mathcal{E}$  over some probability triple  $(\Omega, \mathcal{F}, \mathbf{P}_\theta)$ , with  $A \in \mathcal{F}$ , is the Bernoulli( $\theta$ ) RV. The parameter  $\theta$  denotes the probability that ‘ $A$  occurs’ (see Figure 3.3 when  $A$  is the event that ‘H occurs’). This is our first example of a discrete RV.

**Model 1** (Bernoulli( $\theta$ )) Given a parameter  $\theta \in [0, 1]$ , the probability mass function (PMF) for the Bernoulli( $\theta$ ) RV  $X$  is:

$$f(x; \theta) = \theta^x (1 - \theta)^{1-x} \mathbf{1}_{\{0,1\}}(x) = \begin{cases} \theta & \text{if } x = 1, \\ 1 - \theta & \text{if } x = 0, \\ 0 & \text{otherwise} \end{cases} \quad (3.4)$$

and its DF is:

$$F(x; \theta) = \begin{cases} 1 & \text{if } 1 \leq x, \\ 1 - \theta & \text{if } 0 \leq x < 1, \\ 0 & \text{otherwise} \end{cases} \quad (3.5)$$

We emphasise the dependence of the probabilities on the parameter  $\theta$  by specifying it following the semicolon in the argument for  $f$  and  $F$  and by subscripting the probabilities, i.e.  $\mathbf{P}_\theta(X = 1) = \theta$  and  $\mathbf{P}_\theta(X = 0) = 1 - \theta$ .

### 3.3 An Elementary Continuous Random Variable


When a RV takes values in the continuum we call it a **continuous** RV. An example of such a RV is the vertical position (in micro meters) since the original release of a pollen grain on water. Another example of a continuous RV is the volume of water (in cubic meters) that fell on the southern Alps last year.

**Definition 20 (probability density function (PDF))** A RV  $X$  is said to be ‘continuous’ if there exists a piecewise-continuous function  $f$ , called the probability density function (PDF) of  $X$ , such that for any  $a, b \in \mathbb{R}$  with  $a < b$ ,

$$\mathbf{P}(a < X \leq b) = F(b) - F(a) = \int_a^b f(x) dx .$$

The following hold for a continuous RV  $X$  with PDF  $f$ :

Figure 3.3: The Indicator Function  $\mathbb{1}_H$  of the event ‘Heads occurs’, for the experiment ‘Toss 1 times,’  $\mathcal{E}_\theta^1$ , as the RV  $X$  from the sample space  $\Omega = \{H, T\}$  to  $\mathbb{R}$  and its DF  $F$ . The probability that ‘Heads occurs’ and that ‘Tails occurs’ are  $f(1; \theta) = \mathbf{P}_\theta(X = 1) = \mathbf{P}_\theta(H) = \theta$  and  $f(0; \theta) = \mathbf{P}_\theta(X = 0) = \mathbf{P}_\theta(T) = 1 - \theta$ , respectively.



1. For any  $x \in \mathbb{R}$ ,  $\mathbf{P}(X = x) = 0$ .
2. Consequentially, for any  $a, b \in \mathbb{R}$  with  $a \leq b$ ,

$$\mathbf{P}(a < X < b) = \mathbf{P}(a < X \leq b) = \mathbf{P}(a \leq X \leq b) = \mathbf{P}(a \leq X < b) .$$

3. By the fundamental theorem of calculus, except possibly at finitely many points (where the continuous pieces come together in the piecewise-continuous  $f$ ):

$$f(x) = \frac{d}{dx} F(x)$$

4. And of course  $f$  must satisfy:

$$\int_{-\infty}^{\infty} f(x) dx = \mathbf{P}(-\infty < X < \infty) = 1 .$$

An elementary and fundamental example of a continuous RV is the Uniform(0, 1) RV of Model 2. It forms the foundation for random variate generation and simulation. In fact, it is appropriate to call this the fundamental model since every other experiment can be obtained from this one.


**Model 2 (The Fundamental Model)** The probability density function (PDF) of the fundamental model or the Uniform(0, 1) RV is

$$f(x) = \mathbb{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1, \\ 0 & \text{otherwise} \end{cases} \quad (3.6)$$

and its distribution function (DF) or cumulative distribution function (CDF) is:

$$F(x) := \int_{-\infty}^x f(y) dy = \begin{cases} 0 & \text{if } x < 0, \\ x & \text{if } 0 \leq x \leq 1, \\ 1 & \text{if } x > 1 \end{cases} \quad (3.7)$$

Note that the DF is the identity map in  $[0, 1]$ . The PDF and DF are depicted in Figure 3.4.

Figure 3.4: A plot of the PDF and DF or CDF of the Uniform(0, 1) continuous RV  $X$ .**\*\*tossing a fair coin infinitely often and the fundamental model**

- The fundamental model is equivalent to infinite tosses of a fair coin (see using binary expansion of any  $x \in (0, 1)$ )
- The fundamental model has infinitely many copies of itself within it!

**\*\*universality of the fundamental model**

- one can obtain any other random variable from the fundamental model!

### 3.4 Transformations of random variables

Suppose we know the distribution of a random variable  $X$ . How do we find the distribution of a transformation of  $X$ , say  $g(X)$ ? Before we answer this question let us ask a motivational question. Why are we interested in functions of random variables?

**Example 36** Consider a simple financial example where an individual sells  $X$  items per day, the profit per item is \$5 and the overhead costs are \$500 per day. The original random variable is  $X$ , but the random variable  $Y$  which gives the daily profit is of more interest, where

$$Y = 5X - 500 .$$

**Example 37** In a cell-phone system a mobile signal may have a signal-to-noise-ratio of  $X$ , but engineers prefer to express such ratios in decibels, i.e.,

$$Y = 10 \log_{10}(X) .$$

Hence in a great many situations we are more interested in functions of random variables. Let us return to our original question of determining the distribution of a transformation or function of  $X$ . First note that this transformation of  $X$  is itself another random variable, say  $Y = g(X)$ , where  $g$  is a function from a subset  $\mathbb{X}$  of  $\mathbb{R}$  to a subset  $\mathbb{Y}$  of  $\mathbb{R}$ , i.e.,  $g : \mathbb{X} \rightarrow \mathbb{Y}$ ,  $\mathbb{X} \subset \mathbb{R}$  and  $\mathbb{Y} \subset \mathbb{R}$ .

The **inverse image** of a set  $A$  is the set of all real numbers in  $\mathbb{X}$  whose image is in  $A$ , i.e.,

$$g^{[-1]}(A) = \{x \in \mathbb{X} : g(x) \in A\} .$$

In other words,

$$x \in g^{[-1]}(A) \text{ if and only if } g(x) \in A .$$

For example,

- if  $g(x) = 2x$  then  $g^{[-1]}([4, 6]) = [2, 3]$
- if  $g(x) = 2x + 1$  then  $g^{[-1]}([5, 7]) = [2, 3]$
- if  $g(x) = x^3$  then  $g^{[-1]}([1, 8]) = [1, 2]$
- if  $g(x) = x^2$  then  $g^{[-1]}([1, 4]) = [-2, -1] \cup [1, 2]$
- if  $g(x) = \sin(x)$  then  $g^{[-1]}([-1, 1]) = \mathbb{R}$
- if ...

For the singleton set  $A = \{y\}$ , we write  $g^{[-1]}(y)$  instead of  $g^{[-1]}(\{y\})$ . For example,

- if  $g(x) = 2x$  then  $g^{[-1]}(4) = \{2\}$
- if  $g(x) = 2x + 1$  then  $g^{[-1]}(7) = \{3\}$
- if  $g(x) = x^3$  then  $g^{[-1]}(8) = \{2\}$
- if  $g(x) = x^2$  then  $g^{[-1]}(4) = \{-2, 2\}$
- if  $g(x) = \sin(x)$  then  $g^{[-1]}(0) = \{k\pi : k \in \mathbb{Z}\} = \{\dots, -3\pi, -2\pi, -\pi, 0, \pi, 2\pi, 3\pi, \dots\}$
- if ...

If  $g : \mathbb{X} \rightarrow \mathbb{Y}$  is one-to-one (injective) and onto (surjective), then the inverse image of a singleton set is itself a singleton set. Thus, the inverse image of such a function  $g$  becomes itself a function and is called the **inverse function**. One can find the inverse function, if it exists by the following steps:

Step 1; write  $y = g(x)$

Step 2; solve for  $x$  in terms of  $y$

Step 3; set  $g^{-1}(y)$  to be this solution

We write  $g^{-1}$  whenever the inverse image  $g^{[-1]}$  exists as an inverse function of  $g$ . Thus, the inverse function  $g^{-1}$  is a specific type of inverse image  $g^{[-1]}$ . For example,

- if  $g(x) = 2x$  then  $g : \mathbb{R} \rightarrow \mathbb{R}$  is injective and surjective and therefore its inverse function is:  
Step 1;  $y = 2x$ , Step 2;  $x = \frac{y}{2}$ , Step 3;  $g^{-1}(y) = \frac{y}{2}$
- if  $g(x) = 2x + 1$  then  $g : \mathbb{R} \rightarrow \mathbb{R}$  is injective and surjective and therefore its inverse function is:  
Step 1;  $y = 2x + 1$ , Step 2;  $x = \frac{y-1}{2}$ , Step 3;  $g^{-1}(y) = \frac{y-1}{2}$
- if  $g(x) = x^3$  then  $g : \mathbb{R} \rightarrow \mathbb{R}$  is injective and surjective and therefore its inverse function is:  
Step 1;  $y = x^3$ , Step 2;  $x = y^{\frac{1}{3}}$ , Step 3;  $g^{-1}(y) = y^{\frac{1}{3}}$

However, you need to be careful by limiting the domain to obtain the inverse function for the following examples:

- if  $g(x) = x^2$  and domain of  $g$  is  $[0, +\infty)$  then its inverse function is  $g^{-1}(y) = \sqrt{y}$ , i.e., if  $g(x) = x^2 : [0, +\infty) \rightarrow [0, +\infty)$  then the inverse image  $g^{[-1]}(y)$  for  $y \in [0, +\infty)$  is given by the inverse function  $g^{-1}(y) = \sqrt{y} : [0, +\infty) \rightarrow [0, +\infty)$ .
- if  $g(x) = x^2$  and domain of  $g$  is  $(-\infty, 0]$  then its inverse function is  $g^{-1}(y) = -\sqrt{y}$ , i.e., if  $g(x) = x^2 : (-\infty, 0] \rightarrow [0, +\infty)$  then the inverse image  $g^{[-1]}(y)$  for  $y \in [0, +\infty)$  is given by the inverse function  $g^{-1}(y) = -\sqrt{y} : [0, +\infty) \rightarrow (-\infty, 0]$ .
- if  $g(x) = \sin(x)$  and domain of  $g$  is  $[0, \frac{\pi}{2}]$  then its inverse function  $g^{-1}(y) = \arcsin(y)$ , i.e., if  $g(x) = \sin(x) : [0, \frac{\pi}{2}] \rightarrow [0, 1]$  then the inverse image  $g^{[-1]}(y)$  for  $y \in [0, 1]$  is given by the inverse function  $g^{-1}(y) = \arcsin(y) : [0, 1] \rightarrow [0, \frac{\pi}{2}]$ .
- if  $g(x) = \sin(x)$  and domain of  $g$  is  $[-\frac{\pi}{2}, \frac{\pi}{2}]$  then its inverse function  $g^{-1}(y) = \arcsin(y)$ , i.e., if  $g(x) = \sin(x) : [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow [-1, 1]$  then the inverse image  $g^{[-1]}(y)$  for  $y \in [-1, 1]$  is given by the inverse function  $g^{-1}(y) = \arcsin(y) : [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- if ...

Now, let us return to our question of determining the distribution of the transformation  $g(X)$ . To answer this question we must first observe that the inverse image  $g^{[-1]}$  satisfies the following properties:

- $g^{[-1]}(\mathbb{Y}) = \mathbb{X}$
- For any set  $A$ ,  $g^{[-1]}(A^c) = (g^{[-1]}(A))^c$
- For any collection of sets  $\{A_1, A_2, \dots\}$ ,

$$g^{[-1]}(A_1 \cup A_2 \cup \dots) = g^{[-1]}(A_1) \cup g^{[-1]}(A_2) \cup \dots .$$

Consequentially,

$$P(g(X) \in A) = P\left(X \in g^{[-1]}(A)\right) \quad (3.8)$$

satisfies the axioms of probability and gives the desired probability of the event  $A$  from the transformation  $Y = g(X)$  in terms of the probability of the event given by the inverse image of  $A$  underpinned by the random variable  $X$ . It is crucial to understand this from the sample space  $\Omega$  of the underlying experiment in the sense that Equation (3.8) is just short-hand for its actual meaning:

$$P(\{\omega \in \Omega : g(X(\omega)) \in A\}) = P\left(\left\{\omega \in \Omega : X(\omega) \in g^{[-1]}(A)\right\}\right) .$$

Because we have more than one random variable to consider, namely,  $X$  and its transformation  $Y = g(X)$  we will subscript the probability density or mass function and the distribution function by the random variable itself. For example we denote the distribution function of  $X$  by  $F_X(x)$  and that of  $Y$  by  $F_Y(y)$ .

### 3.4.1 Transformation of discrete random variables

For a discrete random variable  $X$  with probability mass function  $f_X$  we can obtain the probability mass function  $f_Y$  of  $Y = g(X)$  using Equation (3.8) as follows:

$$\begin{aligned} f_Y(y) &= \mathbf{P}(Y = y) = \mathbf{P}(Y \in \{y\}) \\ &= P(g(X) \in \{y\}) = P\left(X \in g^{[-1]}\{y\}\right) \\ &= P\left(X \in g^{[-1]}(y)\right) = \sum_{x \in g^{[-1]}(y)} f_X(x) = \sum_{x \in \{x : g(x)=y\}} f_X(x) . \end{aligned}$$

This gives the formula:

$$\boxed{f_Y(y) = \mathbf{P}(Y = y) = \sum_{x \in g^{[-1]}(y)} f_X(x) = \sum_{x \in \{x: g(x) = y\}} f_X(x) .} \quad (3.9)$$

**Example 38** Let  $X$  be the discrete random variable with probability mass function  $f_X$  as tabulated below:

$x$	-1    0    1
$f_X(x) = \mathbf{P}(X = x)$	$\frac{1}{4}$ $\frac{1}{2}$ $\frac{1}{4}$

If  $Y = 2X$  then the transformation  $g(X) = 2X$  has inverse image  $g^{[-1]}(y) = \{y/2\}$ . Then, by Equation (3.9) the probability mass function of  $Y$  is expressed in terms of the known probabilities of  $X$  as:

$$f_Y(y) = \mathbf{P}(Y = y) = \sum_{x \in g^{[-1]}(y)} f_X(x) = \sum_{x \in \{y/2\}} f_X(x) = f_X(y/2) ,$$

and tabulated below:

$y$	-2    0    2
$f_Y(y)$	$\frac{1}{4}$ $\frac{1}{2}$ $\frac{1}{4}$

**Example 39** If  $X$  is the random variable in the previous Example then what is the probability mass function of  $Y = 2X + 1$ ? Once again,

$$f_Y(y) = \mathbf{P}(Y = y) = \sum_{x \in g^{[-1]}(y)} f_X(x) = \sum_{x \in \{(y-1)/2\}} f_X(x) = f_X((y-1)/2) ,$$

and tabulated below:

$y$	-1    1    3
$f_Y(y)$	$\frac{1}{4}$ $\frac{1}{2}$ $\frac{1}{4}$

In fact, obtaining the probability of a one-to-one transformation of a discrete random variable as in Examples 38 and 39 is merely a matter of looking up the probability at the image of the inverse function. This is because there is only one term in the sum that appears in Equation (3.9). When the transformation is not one-to-one the number of terms in the sum can be more than one as shown in the next Example.

**Example 40** Reconsider the random variable  $X$  of the last two Examples and let  $Y = X^2$ . Recall that  $g(x) = x^2$  does not have an inverse function unless the domain is restricted to the positive or the negative parts of the real line. Since our random variable  $X$  takes values on both sides of the real line, namely  $\{-1, 0, 1\}$ , let us note that the transformation  $g(X) = X^2$  is no longer a one-to-one function. Then, by Equation (3.9) the probability mass function of  $Y$  is expressed in

terms of the known probabilities of  $X$  as:

$$f_Y(y) = \mathbf{P}(Y = y) = \sum_{x \in g^{[-1]}(y)} f_X(x) = \sum_{\{x: g(x)=y\}} f_X(x) = \sum_{\{x: x^2=y\}} f_X(x) ,$$

computed for each  $y \in \{0, 1\}$  as follows:

$$\begin{aligned} f_Y(0) &= \sum_{\{x: x^2=0\}} f_X(x) = f_X(0) = \frac{1}{2} , \\ f_Y(1) &= \sum_{\{x: x^2=1\}} f_X(x) = f_X(-1) + f_X(1) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2} , \end{aligned}$$

and finally tabulated below:

$y$	0	1
$f_Y(y)$	$\frac{1}{2}$	$\frac{1}{2}$

### 3.4.2 Transformation of continuous random variables

Suppose we know  $F_X$  and/or  $f_X$  of a continuous random variable  $X$ . Let  $Y = g(X)$  be a transformation of  $X$ . Our objective is to obtain  $F_Y$  and/or  $f_Y$  of  $Y$  from  $F_X$  and/or  $f_X$ .

#### One-to-one transformations

The easiest case for transformations of continuous random variables is when  $g$  is **one-to-one and monotone**.

- First, let us consider the case when  $g$  is **monotone and increasing** on the range of the random variable  $X$ . In this case  $g^{-1}$  is also an increasing function and we can obtain the distribution function of  $Y = g(X)$  in terms of the distribution function of  $X$  as

$$F_Y(y) = P(Y \leq y) = P(g(X) \leq y) = P(X \leq g^{-1}(y)) = F_X(g^{-1}(y)) .$$

Now, let us use a form of chainrule to compute the density of  $Y$  as follows:

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dy} F_X(g^{-1}(y)) = f_X(g^{-1}(y)) \frac{d}{dy}(g^{-1}(y)) .$$

- Second, let us consider the case when  $g$  is **monotone and decreasing** on the range of the random variable  $X$ . In this case  $g^{-1}$  is also a decreasing function and we can obtain the distribution function of  $Y = g(X)$  in terms of the distribution function of  $X$  as

$$F_Y(y) = P(Y \leq y) = P(g(X) \leq y) = P(X \geq g^{-1}(y)) = 1 - F_X(g^{-1}(y)) ,$$

and the density of  $Y$  as

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dy} (1 - F_X(g^{-1}(y))) = -f_X(g^{-1}(y)) \frac{d}{dy}(g^{-1}(y)) .$$

For a monotonic and decreasing  $g$ , its inverse function  $g^{-1}$  is also decreasing and consequently the density  $f_Y$  is indeed positive because  $\frac{d}{dy}(g^{-1}(y))$  is negative.

We can combine the above two cases and obtain the following **change of variable formula** for the probability density of  $Y = g(X)$  when  $g$  is one-to-one and monotone on the range of  $X$ .

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| . \quad (3.10)$$

The steps involved in finding the density of  $Y = g(X)$  for a one-to-one and monotone  $g$  are:

1. Write  $y = g(x)$  for  $x$  in range of  $x$  and check that  $g(x)$  is monotone over the required range to apply the change of variable formula.
2. Write  $x = g^{-1}(y)$  for  $y$  in range of  $y$ .
3. Obtain  $\left| \frac{d}{dy} g^{-1}(y) \right|$  for  $y$  in range of  $y$ .
4. Finally, from Equation (3.10) get  $f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|$  for  $y$  in range of  $y$ .

Let us use these four steps to obtain the density of monotone transformations of continuous random variables.

**Example 41** Let  $X$  be Uniform(0, 1) random variable and let  $Y = g(X) = 1 - X$ . We are interested in the density of the tranformed random variable  $Y$ . Let us follow the four steps and use the change of variable formula to obtain  $f_Y$  from  $f_X$  and  $g$ .

1.  $y = g(x) = 1 - x$  is a monotone decreasing function over  $0 \leq x \leq 1$ , the range of  $X$ . So, we can apply the change of variable formula.
  2.  $x = g^{-1}(y) = 1 - y$  is a monotone decreasing function over  $1 - 0 \geq 1 - x \geq 1 - 1$ , i.e.,  $0 \leq y \leq 1$ .
  3. For  $0 \leq y \leq 1$ ,
- $$\left| \frac{d}{dy} g^{-1}(y) \right| = \left| \frac{d}{dy} (1 - y) \right| = |-1| = 1 .$$
4. we can use Equation (3.10) to find the density of  $Y$  as follows:

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| = f_X(1 - y) 1 = 1 ,$$

for  $0 \leq y \leq 1$

Thus, we have shown that if  $X$  is a Uniform(0, 1) random variable then  $Y = 1 - X$  is also a Uniform(0, 1) random variable.

**Example 42** Let  $X$  be a Uniform(0, 1) random variable and let  $Y = g(X) = -\log(X)$ . We are interested in the density of the tranformed random variable  $Y$ . Once again, since  $g$  is a one-to-one monotone function let us follow the four steps and use the change of variable formula to obtain  $f_Y$  from  $f_X$  and  $g$ .

1.  $y = g(x) = -\log(x)$  is a monotone decreasing function over  $0 < x < 1$ , the range of  $X$ . So, we can apply the change of variable formula.

2.  $x = g^{-1}(y) = \exp(-y)$  is a monotone decreasing function over  $0 < y < \infty$ .

3. For  $0 < y < \infty$ ,

$$\left| \frac{d}{dy} g^{-1}(y) \right| = \left| \frac{d}{dy} (\exp(-y)) \right| = |-\exp(-y)| = \exp(-y) .$$

4. We can use Equation (3.10) to find the density of  $Y$  as follows:

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| = f_X(\exp(-y)) \exp(-y) = 1 \exp(-y) = \exp(-y) .$$

Note that  $0 < \exp(-y) < 1$  for  $0 < y < \infty$ .


Thus, we have shown that if  $X$  is a Uniform(0, 1) random variable then  $Y = -\log(X)$  is a random variable with PDF  $f_Y(y) = \mathbb{1}_{(0,\infty)}(y) \exp(-y)$ . We can similarly show that for a parameter  $\lambda > 0$ , if  $X \sim \text{Uniform}(0, 1)$  then  $Y = -\lambda^{-1} \log(X)$  yields a probability model of RVs that are parameterized by  $\lambda$  and extremely useful in applications. We introduce it next.

**Model 3** (Exponential( $\lambda$ )) For a given  $\lambda > 0$ , an Exponential( $\lambda$ ) RV has the following PDF  $f$  and DF  $F$ :

$$f(x; \lambda) = \lambda e^{-\lambda x} \quad F(x; \lambda) = 1 - e^{-\lambda x} . \quad (3.11)$$

This distribution is fundamental because of its property of **memorylessness** and plays a fundamental role in continuous time processes as we will see later.

Figure 3.5: Density and distribution functions of Exponential( $\lambda$ ) RVs, for  $\lambda = 1, 10, 10^{-1}$ , in four different axes scales.



### Direct method

If the transformation  $g$  in  $Y = g(X)$  is not necessarily one-to-one then special care is needed to obtain the distribution function or density of  $Y$ . For a continuous random variable  $X$  with a

known distribution function  $F_X$  we can obtain the distribution function  $F_Y$  of  $Y = g(X)$  using Equation (3.8) as follows:

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(Y \in (-\infty, y]) \\ &= P(g(X) \in (-\infty, y]) = P\left(X \in g^{[-1]}((-\infty, y])\right) = P(X \in \{x : g(x) \in (-\infty, y]\}) . \end{aligned}$$

In words, the above equalities just mean that the probability that  $Y \leq y$  is the probability that  $X$  takes a value  $x$  that satisfies  $g(x) \leq y$ . We can use this approach if it is reasonably easy to find the set  $g^{[-1]}((-\infty, y]) = \{x : g(x) = (-\infty, y]\}$ .

**Example 43** Let  $X$  be any random variable with distribution function  $F_X$ . Let  $Y = g(X) = X^2$ . Then we can find  $F_Y$ , the distribution function of  $Y$  from  $F_X$  as follows:

- Since  $Y = X^2 \geq 0$ , if  $y < 0$  then  $F_Y(y) = P(X \in \{x : x^2 < y\}) = \mathbf{P}(X \in \emptyset) = 0$ .
- If  $y \geq 0$  then

$$\begin{aligned} F_Y(y) = P(Y \leq y) &= P(X^2 \leq y) \\ &= P(-\sqrt{y} \leq X \leq \sqrt{y}) \\ &= F_X(\sqrt{y}) - F_X(-\sqrt{y}) . \end{aligned}$$

By differentiation we get:

- If  $y < 0$  then  $f_Y(y) = \frac{d}{dy}(F_Y(y)) = \frac{d}{dy}0 = 0$ .
- If  $y \geq 0$  then

$$\begin{aligned} f_Y(y) = \frac{d}{dy}(F_Y(y)) &= \frac{d}{dy}(F_X(\sqrt{y}) - F_X(-\sqrt{y})) \\ &= \frac{d}{dy}(F_X(\sqrt{y})) - \frac{d}{dy}(F_X(-\sqrt{y})) \\ &= \frac{1}{2}y^{-\frac{1}{2}}f_X(\sqrt{y}) - \left(-\frac{1}{2}y^{-\frac{1}{2}}f_X(-\sqrt{y})\right) \\ &= \frac{1}{2\sqrt{y}}(f_X(\sqrt{y}) + f_X(-\sqrt{y})) . \end{aligned}$$

Therefore, the distribution function of  $Y = X^2$  is:

$$F_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ F_X(\sqrt{y}) - F_X(-\sqrt{y}) & \text{if } y \geq 0 \end{cases} . \quad (3.12)$$

and the probability density function of  $Y = X^2$  is:

$$f_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ \frac{1}{2\sqrt{y}}(f_X(\sqrt{y}) + f_X(-\sqrt{y})) & \text{if } y \geq 0 \end{cases} . \quad (3.13)$$

### 3.5 $\mathbb{R}^m$ -valued Random Variables

Often, in experiments we are measuring two or more aspects simultaneously. For example, we may be measuring the diameters and lengths of cylindrical shafts manufactured in a plant or heights, weights and blood-sugar levels of individuals in a clinical trial. Thus, the underlying outcome  $\omega \in \Omega$  needs to be mapped to measurements as realizations of random vectors in the real plane  $\mathbb{R}^2 = (-\infty, \infty) \times (-\infty, \infty)$  or the real space  $\mathbb{R}^3 = (-\infty, \infty) \times (-\infty, \infty) \times (-\infty, \infty)$ :

$$\omega \mapsto (X(\omega), Y(\omega)) : \Omega \rightarrow \mathbb{R}^2 \quad \omega \mapsto (X(\omega), Y(\omega), Z(\omega)) : \Omega \rightarrow \mathbb{R}^3$$

More generally, we may be interested in heights, weights, blood-sugar levels, family medical history, known allergies, etc. of individuals in the clinical trial and thus need to make  $m$  measurements of the outcome in  $\mathbb{R}^m$  using a “measurable mapping” from  $\Omega \rightarrow \mathbb{R}^m$ . To deal with such multivariate measurements we need the notion of **random vectors** ( $\vec{RV}$ s), i.e. ordered pairs of random variables  $(X, Y)$ , ordered triples of random variables  $(X, Y, Z)$ , or more generally ordered  $m$ -tuples of random variables  $(X_1, X_2, \dots, X_m)$ .

#### 3.5.1 $\mathbb{R}^2$ -valued Random Variables

We first focus on understanding  $(X, Y)$ , a bivariate  $\vec{RV}$  or  $\mathbb{R}^2$ -valued RV that is obtained from a pair of discrete or continuous RVs. We then generalize to  $\mathbb{R}^m$ -valued RVs with  $m > 2$  in the next section.

**Definition 21 (JDF)** The **joint distribution function (JDF)** or **joint cumulative distribution function (JCDF)**,  $F_{X,Y}(x, y) : \mathbb{R}^2 \rightarrow [0, 1]$ , of the bivariate random vector  $(X, Y)$  is

$$\begin{aligned} F_{X,Y}(x, y) &= \mathbf{P}(X \leq x \cap Y \leq y) = \mathbf{P}(X \leq x, Y \leq y) \\ &= P(\{\omega : X(\omega) \leq x, Y(\omega) \leq y\}), \text{ for any } (x, y) \in \mathbb{R}^2, \end{aligned} \quad (3.14)$$

where the right-hand side represents the probability that the random vector  $(X, Y)$  takes on a value in  $\{(x', y') : x' \leq x, y' \leq y\}$ , the set of points in the plane that are south-west of the point  $(x, y)$ .

The JDF  $F_{X,Y}(x, y) : \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfies the following conditions to remain a probability:

1.  $0 \leq F_{X,Y}(x, y) \leq 1$
2.  $F_{X,Y}(x, y)$  is a non-decreasing function of both  $x$  and  $y$
3.  $F_{X,Y}(x, y) \rightarrow 1$  as  $x \rightarrow \infty$  and  $y \rightarrow \infty$
4.  $F_{X,Y}(x, y) \rightarrow 0$  as  $x \rightarrow -\infty$  and  $y \rightarrow -\infty$

**Definition 22 (JPMF)** If  $(X, Y)$  is a **discrete random vector** that takes values in a discrete support set  $\mathcal{S}_{X,Y} = \{(x_i, y_j) : i = 1, 2, \dots, j = 1, 2, \dots\} \subset \mathbb{R}^2$  with probabilities  $p_{i,j} = \mathbf{P}(X = x_i, Y = y_j) > 0$ , then its **joint probability mass function** (or JPMF) is:

$$f_{X,Y}(x_i, y_j) = \mathbf{P}(X = x_i, Y = y_j) = \begin{cases} p_{i,j} & \text{if } (x_i, y_j) \in \mathcal{S}_{X,Y} \\ 0 & \text{otherwise} \end{cases}. \quad (3.15)$$


Since  $\mathbf{P}(\Omega) = 1$ ,  $\sum_{(x_i, y_j) \in \mathcal{S}_{X,Y}} f_{X,Y}(x_i, y_j) = 1$ .

From JPMF  $f_{X,Y}$  we can get the values of the JDF  $F_{X,Y}(x, y)$  and the probability of any event  $B$  by simply taking sums,

$$\boxed{F_{X,Y}(x, y) = \sum_{x_i \leq x, y_j \leq y} f_{X,Y}(x_i, y_j) , \quad \mathbf{P}(B) = \sum_{(x_i, y_j) \in B \cap \mathcal{S}_{X,Y}} f_{X,Y}(x_i, y_j)} , \quad (3.16)$$

**Example 44** Let  $(X, Y)$  be a discrete bivariate R $\vec{V}$  with the following joint probability mass function (JPMF):

$$f_{X,Y}(x, y) := P(X = x, Y = y) = \begin{cases} 0.1 & \text{if } (x, y) = (0, 0) \\ 0.3 & \text{if } (x, y) = (0, 1) \\ 0.2 & \text{if } (x, y) = (1, 0) \\ 0.4 & \text{if } (x, y) = (1, 1) \\ 0.0 & \text{otherwise.} \end{cases}$$



It is helpful to write down the JPMF  $f_{X,Y}(x, y)$  in a tabular form:

	$Y = 0$	$Y = 1$
$X = 0$	0.1	0.3
$X = 1$	0.2	0.4

From the above Table we can read for instance that the joint probability  $f_{X,Y}(0,0) = 0.1$ .

Find  $\mathbf{P}(B)$  for the event  $B = \{(0,0), (1,1)\}, F_{X,Y}(1/2, 1/2), F_{X,Y}(3/2, 1/2), F_{X,Y}(4, 5)$  and  $F_{X,Y}(-4, -1)$ .

1.  $\mathbf{P}(B) = \sum_{(x,y) \in \{(0,0), (1,1)\}} f_{X,Y}(x, y) = f_{X,Y}(0, 0) + f_{X,Y}(1, 1) = 0.1 + 0.4$
2.  $F_{X,Y}(1/2, 1/2) = \sum_{\{(x,y): x \leq 1/2, y \leq 1/2\}} f_{X,Y}(x, y) = f_{X,Y}(0, 0) = 0.1$
3.  $F_{X,Y}(3/2, 1/2) = \sum_{\{(x,y): x \leq 3/2, y \leq 1/2\}} f_{X,Y}(x, y) = f_{X,Y}(0, 0) + f_{X,Y}(1, 0) = 0.1 + 0.2 = 0.3$
4.  $F_{X,Y}(4, 5) = \sum_{\{(x,y): x \leq 4, y \leq 5\}} f_{X,Y}(x, y) = f_{X,Y}(0, 0) + f_{X,Y}(0, 1) + f_{X,Y}(1, 0) + f_{X,Y}(1, 1) = 1$
5.  $F_{X,Y}(-4, -1) = \sum_{\{(x,y): x \leq -4, y \leq -1\}} f_{X,Y}(x, y) = 0$

**Definition 23 (JPDF)** We say  $(X, Y)$  is a **continuous  $R^2$ -valued random variable** if its JDF  $F_{X,Y}(x, y)$  is differentiable and its **joint probability density function (JPDF)** is given by:

$$f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y) .$$

For notational convenience, we sometimes suppress the subscripting when the random variables are clear from the context and write  $f(x, y)$  and  $F(x, y)$  instead of  $f_{X,Y}(x, y)$  and  $F_{X,Y}(x, y)$ , respectively.

From JPDF  $f_{X,Y}$  we can compute the JDF  $F_{X,Y}$  at any point  $(x, y) \in \mathbb{R}^2$  and more generally we can compute the probability of any event  $B$ , that can be cast as a region in  $\mathbb{R}^2$ , by simply taking two-dimensional integrals:

$$\boxed{F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) du dv}, \quad (3.17)$$

and

$$\boxed{\mathbf{P}(B) = \int \int_B f_{X,Y}(x, y) dx dy}. \quad (3.18)$$

In particular, if  $\mathbb{B}_\delta(x, y)$  denotes a square of a small area  $\delta > 0$  that is centered at  $(x, y)$ , then the following approximate equality holds and improves as  $\delta \rightarrow 0$ :

$$\mathbf{P}((X, Y) \in \mathbb{B}_\delta(x, y)) \approx \delta f_{X,Y}(x, y). \quad (3.19)$$


The JPDF satisfies the following two properties:

1. integrates to 1, i.e.,  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$
2. is a non-negative function, i.e.,  $f_{X,Y}(x, y) \geq 0$  for every  $(x, y) \in \mathbb{R}^2$ .

**Example 45** Let  $(X, Y)$  be a continuous RV that is uniformly distributed on the unit square  $[0, 1]^2 := [0, 1] \times [0, 1]$  with following JPDF:

$$f(x, y) = \mathbf{1}_{[0,1]^2}(x) \begin{cases} 1 & \text{if } (x, y) \in [0, 1]^2 \\ 0 & \text{otherwise.} \end{cases}$$

Find explicit expressions for the following: (1) DF  $F(x, y)$  for any  $(x, y) \in [0, 1]^2$ , (2)  $\mathbf{P}(X \leq 1/3, Y \leq 1/2)$ , (3)  $P((X, Y) \in [1/4, 1/2] \times [1/3, 2/3])$ .



Let us begin to find the needed expressions.

1. Let  $(x, y) \in [0, 1]^2$  then by Equation (3.17):

$$F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) du dv = \int_0^y \int_0^x 1 du dv = \int_0^y [u]_{u=0}^x dv = \int_0^y x dv = [xv]_{v=0}^y = xy$$

2. We can obtain  $\mathbf{P}(X \leq 1/3, Y \leq 1/2)$  by evaluating  $F_{X,Y}$  at  $(1/3, 1/2)$ :

$$\mathbf{P}(X \leq 1/3, Y \leq 1/2) = F_{X,Y}(1/3, 1/2) = \frac{1}{3} \frac{1}{2} = \frac{1}{6}$$

We can also find  $\mathbf{P}(X \leq 1/3, Y \leq 1/2)$  by integrating the JPDF over the rectangular event  $A = \{X < 1/3, Y < 1/2\} \subset [0, 1]^2$  according to Equation (3.18). This amounts here to finding the area of  $A$ , we compute  $\mathbf{P}(A) = (1/3)(1/2) = 1/6$ .

3. We can find  $P((X, Y) \in [1/4, 1/2] \times [1/3, 2/3])$  by integrating the JPDF over the rectangular event  $B = [1/4, 1/2] \times [1/3, 2/3]$  according to Equation (3.18):

$$\begin{aligned} P((X, Y) \in [1/4, 1/2] \times [1/3, 2/3]) &= \int \int_B f_{X,Y}(x, y) dx dy = \int_{1/3}^{2/3} \int_{1/4}^{1/2} 1 dx dy \\ &= \int_{1/3}^{2/3} [x]_{1/4}^{1/2} dy = \int_{1/3}^{2/3} \left[ \frac{1}{2} - \frac{1}{4} \right] dy = \left( \frac{1}{2} - \frac{1}{4} \right) [y]_{1/3}^{2/3} \\ &= \left( \frac{1}{2} - \frac{1}{4} \right) \left( \frac{2}{3} - \frac{1}{3} \right) = \frac{1}{4} \left( \frac{1}{3} \right) = \frac{1}{12} \end{aligned}$$


In general, for a bivariate uniform RV  $\vec{V}$  on the unit square the  $\mathbf{P}([a, b] \times [c, d]) = (b-a)(d-c)$  for any event given by the rectangular region  $[a, b] \times [c, d]$  inside the unit square  $[0, 1] \times [0, 1]$ . Thus any two events with the same rectangular area have the same probability (imagine sliding a small rectangle inside the unit square... no matter where you slide this rectangle to while remaining in the unit square, the probability of  $\omega \mapsto (X(\omega), Y(\omega)) = (x, y)$  falling inside this “slidable” rectangle is the same...).

**Example 46** Let the RV  $X$  denote the time until a web server connects to your computer, and let the RV  $Y$  denote the time until the server authorizes you as a valid user. Each of these RVs measures the waiting time from a common starting time (in milliseconds) and  $X < Y$ . From past response times of the web server we know that a good approximation for the JPDF of the RV  $(X, Y)$  is

$$f_{X,Y}(x, y) = \begin{cases} \frac{6}{10^6} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) & \text{if } x > 0, y > 0, x < y \\ 0 & \text{otherwise.} \end{cases}$$

Answer the following:

1. identify the support of  $(X, Y)$ , i.e., the region in the plane where  $f_{X,Y}$  takes positive values
2. check that  $f_{X,Y}$  indeed integrates to 1 as it should
3. Find  $\mathbf{P}(X \leq 400, Y \leq 800)$
4. It is known that humans prefer a response time of under 1/10 seconds ( $10^2$  milliseconds) from the web server before they get impatient. What is  $\mathbf{P}(X + Y < 10^2)$ ?



Let us answer the questions.

1. The support is the intersection of the positive quadrant with the  $y > x$  half-plane.
- 2.

$$\begin{aligned}
 \int_{y=-\infty}^{\infty} \int_{x=-\infty}^{\infty} f_{X,Y}(x,y) dx dy &= \int_{x=0}^{\infty} \int_{y=x}^{\infty} f_{X,Y}(x,y) dy dx \\
 &= \int_{x=0}^{\infty} \int_{y=x}^{\infty} \frac{6}{10^6} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) dy dx \\
 &= \frac{6}{10^6} \int_{x=0}^{\infty} \left( \int_{y=x}^{\infty} \exp\left(-\frac{2}{1000}y\right) dy \right) \exp\left(-\frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \int_{x=0}^{\infty} \left[ -\frac{1000}{2} \exp\left(-\frac{2}{1000}y\right) \right]_{y=x}^{\infty} \exp\left(-\frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \int_{x=0}^{\infty} \left[ 0 + \frac{1000}{2} \exp\left(-\frac{2}{1000}x\right) \right] \exp\left(-\frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \int_{x=0}^{\infty} \frac{1000}{2} \exp\left(-\frac{2}{1000}x - \frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \frac{1000}{2} \left[ -\frac{1000}{3} \exp\left(-\frac{3}{1000}x\right) \right]_{x=0}^{\infty} \\
 &= \frac{6}{10^6} \frac{1000}{2} \left[ 0 + \frac{1000}{3} \right] \\
 &= 1
 \end{aligned}$$

3. First, identify the region with positive JPDF for the event  $(X \leq 400, Y \leq 800)$

$$\begin{aligned}
 \mathbf{P}(X \leq 400, Y \leq 800) &= \int_{x=0}^{400} \int_{y=x}^{800} f_{X,Y}(x,y) dy dx \\
 &= \int_{x=0}^{400} \int_{y=x}^{800} \frac{6}{10^6} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) dy dx \\
 &= \frac{6}{10^6} \int_{x=0}^{400} \left[ -\frac{1000}{2} \exp\left(-\frac{2}{1000}y\right) \right]_{y=x}^{800} \exp\left(-\frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \frac{1000}{2} \int_{x=0}^{400} \left( -\exp\left(-\frac{1600}{1000}\right) + \exp\left(-\frac{2}{1000}x\right) \right) \exp\left(-\frac{1}{1000}x\right) dx \\
 &= \frac{6}{10^6} \frac{1000}{2} \int_{x=0}^{400} \left( \exp\left(-\frac{3}{1000}x\right) - e^{-8/5} \exp\left(-\frac{1}{1000}x\right) \right) dx \\
 &= \frac{6}{10^6} \frac{1000}{2} \left( \left( -\frac{1000}{3} \exp\left(-\frac{3}{1000}x\right) \right)_{x=0}^{400} - e^{-8/5} \left( -1000 \exp\left(-\frac{1}{1000}x\right) \right)_{x=0}^{400} \right) \\
 &= \frac{6}{10^6} \frac{1000}{2} 1000 \left( \frac{1}{3} \left( 1 - e^{-6/5} \right) - e^{-8/5} \left( 1 - e^{-2/5} \right) \right) \\
 &= 3 \left( \frac{1}{3} \left( 1 - e^{-6/5} \right) - e^{-8/5} \left( 1 - e^{-2/5} \right) \right) \\
 &\approx 0.499 .
 \end{aligned}$$

4. First, identify the region with positive JPDF for the event  $(X + Y \leq c)$ , say  $c = 500$  (but generally  $c$  can be any positive number). This is the triangular region at the intersection of the four half-planes:  $x > 0$ ,  $x < c$ ,  $y > x$  and  $y < c - x$ . (*Draw picture here*) Let's integrate

the JPDF over our triangular event as follows:

$$\begin{aligned}
\mathbf{P}(X + Y \leq c) &= \int_{x=0}^{c/2} \int_{y=x}^{c-x} f_{X,Y}(x,y) dy dx \\
&= \int_{x=0}^{c/2} \int_{y=x}^{c-x} \frac{6}{10^6} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) dy dx \\
&= \frac{6}{10^6} \int_{x=0}^{c/2} \int_{y=x}^{c-x} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) dy dx \\
&= \frac{6}{10^6} \frac{1000}{2} \int_{x=0}^{c/2} \left[-\exp\left(-\frac{2}{1000}y\right)\right]_{y=x}^{c-x} \exp\left(-\frac{1}{1000}x\right) dx \\
&= \frac{3}{10^3} \int_{x=0}^{c/2} \left[-\exp\left(-\frac{2c-2x}{1000}\right) + \exp\left(-\frac{2x}{1000}\right)\right] \exp\left(-\frac{x}{1000}\right) dx \\
&= \frac{3}{10^3} \int_{x=0}^{c/2} \left(\exp\left(-\frac{3x}{1000}\right) - \exp\left(\frac{x-2c}{1000}\right)\right) dx \\
&= 3 \left( \left[ -\frac{1}{3} \exp\left(-\frac{3x}{1000}\right) \right]_{x=0}^{c/2} - \left[ e^{-2c/1000} \exp\left(\frac{x}{1000}\right) \right]_{x=0}^{c/2} \right) \\
&= 3 \left( \frac{1}{3} (1 - e^{-3c/2000}) - e^{-2c/1000} (e^{c/2000} - 1) \right) \\
&= 1 - e^{-3c/2000} + 3e^{-2c/1000} - 3e^{-3c/2000} \\
&= 1 - 4e^{-3c/2000} + 3e^{-c/500}
\end{aligned}$$

5.  $\mathbf{P}(X + Y < 100) = 1 - 4e^{-300/2000} + 3e^{-100/500} \approx 0.134$ . This means only about one in one hundred requests to this server will be processed within 100 milliseconds.

We can obtain  $\mathbf{P}(X + Y < c)$  for several values of  $c$  using MATLAB and note that about 96% of requests are processed in less than 3000 milliseconds or 3 seconds.

```

>> c = [100 1000 2000 3000 4000]
c = 100      1000      2000      3000      4000

>> p = 1 - 4 * exp(-3*c/2000) + 3 * exp(-c/500)

p = 0.0134    0.5135    0.8558    0.9630    0.9911

```

**Definition 24 (Marginal PDF or PMF)** If the R $\vec{V}$   $(X, Y)$  has  $f_{X,Y}(x, y)$  as its joint PDF or joint PMF, then the **marginal PDF or PMF** of a random vector  $(X, Y)$  is defined by :

$$f_X(x) = \begin{cases} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy & \text{if } (X, Y) \text{ is a continuous R}\vec{V} \\ \sum_y f_{X,Y}(x,y) & \text{if } (X, Y) \text{ is a discrete R}\vec{V} \end{cases}$$

and the **marginal PDF or PMF** of  $Y$  is defined by:

$$f_Y(y) = \begin{cases} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx & \text{if } (X, Y) \text{ is a continuous R}\vec{V} \\ \sum_x f_{X,Y}(x,y) & \text{if } (X, Y) \text{ is a discrete R}\vec{V} \end{cases}$$

**Example 47** Obtain the marginal PMFs  $f_Y(y)$  and  $f_X(x)$  from the joint PMF  $f_{X,Y}(x, y)$  of the discrete R $\vec{V}$  in Example 44. Just sum  $f_{X,Y}(x, y)$  over  $x$ 's and  $y$ 's (reported in a tabular form):

	$Y = 0$	$Y = 1$
$X = 0$	0.1	0.3
$X = 1$	0.2	0.4

From the above Table we can find:

$$\begin{aligned} f_X(x) &= \mathbf{P}(X = x) = \sum_y f_{X,Y}(x, y) \\ &= f_{X,Y}(x, 0) + f_{X,Y}(x, 1) = \begin{cases} f_{X,Y}(0, 0) + f_{X,Y}(0, 1) = 0.1 + 0.3 = 0.4 & \text{if } x = 0 \\ f_{X,Y}(1, 0) + f_{X,Y}(1, 1) = 0.2 + 0.4 = 0.6 & \text{if } x = 1 \end{cases} \end{aligned}$$


Similarly,

$$\begin{aligned} f_Y(y) &= \mathbf{P}(Y = y) = \sum_x f_{X,Y}(x, y) \\ &= f_{X,Y}(0, y) + f_{X,Y}(1, y) = \begin{cases} f_{X,Y}(0, 0) + f_{X,Y}(1, 0) = 0.1 + 0.2 = 0.3 & \text{if } y = 0 \\ f_{X,Y}(0, 1) + f_{X,Y}(1, 1) = 0.3 + 0.4 = 0.7 & \text{if } y = 1 \end{cases} \end{aligned}$$

Just report the marginal probabilities as row and column sums of the JPDF table.

Thus marginal PMF gives us the probability of a specific RV, within a R $\vec{V}$ , taking a value irrespective of the value taken by the other RV in this R $\vec{V}$ .

**Example 48** Obtain the marginal PDFs  $f_Y(y)$  and  $f_X(x)$  from the joint PDF  $f_{X,Y}(x, y)$  of the continuous R $\vec{V}$  in Example 45 (the bivariate uniform R $\vec{V}$  on  $[0, 1]^2$ ).



Let us suppose  $(x, y) \in [0, 1]^2$  and note that  $f_{X,Y} = 0$  if  $(x, y) \notin [0, 1]^2$ . We can obtain marginal PMFs  $f_X(x)$  and  $f_Y(y)$  by integrating the JPDF  $f_{X,Y} = 1$  along  $y$  and  $x$ , respectively.

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy = \int_0^1 f_{X,Y}(x, y) dy = \int_0^1 1 dy = [y]_0^1 = 1 - 0 = 1$$

Similarly,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx = \int_0^1 f_{X,Y}(x, y) dx = \int_0^1 1 dx = [x]_0^1 = 1 - 0 = 1$$

We are seeing a histogram of the **marginal samples** and their marginal PDFs in the Figure.

Thus marginal PDF gives us the probability density of a specific RV in a R $\vec{V}$ , irrespective of the value taken by the other RV in this R $\vec{V}$ .

**Example 49** Obtain the marginal PDF  $f_Y(y)$  from the joint PDF  $f_{X,Y}(x,y)$  of the continuous R $\vec{V}$  in Example 46 that gave the response times of a web server.

$$f_{X,Y}(x,y) = \begin{cases} \frac{6}{10^6} \exp\left(-\frac{1}{1000}x - \frac{2}{1000}y\right) & \text{if } x > 0, y > 0, x < y \\ 0 & \text{otherwise.} \end{cases}$$

Use  $f_Y(y)$  to compute the probability that  $Y$  exceeds 2000 milliseconds.

First we need to obtain an expression for  $f_Y(y)$ . For  $y > 0$ ,

$$\begin{aligned} f_Y(y) &= \int_{x=-\infty}^{\infty} f_{X,Y}(x,y) dx \\ &= \int_{x=-\infty}^{\infty} 6 \times 10^{-6} e^{-0.001x - 0.002y} dx \\ &= 6 \times 10^{-6} \int_{x=0}^y e^{-0.001x - 0.002y} dx \\ &= 6 \times 10^{-6} e^{-0.002y} \int_{x=0}^y e^{-0.001x} dx \\ &= 6 \times 10^{-6} e^{-0.002y} \left[ \frac{e^{-0.001x}}{-0.001} \right]_{x=0}^{x=y} \\ &= 6 \times 10^{-6} e^{-0.002y} \left( \frac{e^{-0.001y}}{-0.001} - \frac{e^{-0.001 \times 0}}{-0.001} \right) \\ &= 6 \times 10^{-6} e^{-0.002y} \left( \frac{1 - e^{-0.001y}}{0.001} \right) \\ &= 6 \times 10^{-3} e^{-0.002y} (1 - e^{-0.001y}) \end{aligned}$$

We have the marginal PDF of  $Y$  and from this we can obtain

$$\begin{aligned} \mathbf{P}(Y > 2000) &= \int_{2000}^{\infty} f_Y(y) dy \\ &= \int_{2000}^{\infty} 6 \times 10^{-3} e^{-0.002y} (1 - e^{-0.001y}) dy \\ &= 6 \times 10^{-3} \int_{2000}^{\infty} e^{-0.002y} dy - \int_{2000}^{\infty} e^{-0.003y} dy \\ &= 6 \times 10^{-3} \left( \left[ \frac{e^{-0.002y}}{-0.002} \right]_{2000}^{\infty} - \left( \left[ \frac{e^{-0.003y}}{-0.003} \right]_{2000}^{\infty} \right) \right) \\ &= 6 \times 10^{-3} \left( \frac{e^{-4}}{0.002} - \frac{e^{-6}}{0.003} \right) \\ &= 0.05 \end{aligned}$$

Alternatively, you can obtain  $\mathbf{P}(Y > 2000)$  by directly integrating the joint PDF  $f_{X,Y}(x,y)$  over the appropriate region (but you may now have to integrate two pieces: rectangular infinite strip  $(x,y) : 0 < x < 2000, y > 2000$  and a triangular infinite piece  $\{(x,y) : y > x, y > 2000, x > 2000\}$ )... more involved but we get the same answer.

$$\begin{aligned} \mathbf{P}(Y > 2000) &= \int_{x=0}^{2000} \left( \int_{y=2000}^{\infty} 6 \times 10^{-6} e^{-0.001x - 0.002y} dy \right) dx + \\ &\quad \int_{x=2000}^{\infty} \left( \int_{y=x}^{\infty} 6 \times 10^{-6} e^{-0.001x - 0.002y} dy \right) dx \\ &\quad \vdots \text{(try as a tutorial problem)} \end{aligned}$$

$$\mathbf{P}(Y > 2000) = 0.0475 + 0.0025 = 0.05$$

We have seen the notion of independence of two events in Definition 14 or of a sequence of events in Definition 15. Recall that independence amounts to having the probability of the joint occurrence of the events to be given by the product of the probabilities of each of the events.

We can use the definition of independence of two events to define the independence of two random variables using their distribution functions.

**Definition 25 (Independence of Two RVs)** Consider an  $\mathbb{R}^2$ -valued RV  $X := (X_1, X_2)$ . Then the  $\mathbb{R}$ -valued RVs  $X_1$  and  $X_2$  are said to be independent or independently distributed if and only if

$$\mathbf{P}(X_1 \leq x_1, X_2 \leq x_2) = \mathbf{P}(X_1 \leq x_1)\mathbf{P}(X_2 \leq x_2)$$

or equivalently,

$$F_{X_1, X_2}(x_1, x_2) = F_{X_1}(x_1)F_{X_2}(x_2) ,$$

for any pair of real numbers  $(x_1, x_2) \in \mathbb{R}^2$ .

By the above definition, for **discrete** RVs  $X_1, X_2$  that are independent, the following equality is satisfied between the joint and marginal PMFs:

$$f_{X_1, X_2}(x_1, x_2) = \mathbf{P}(X_1 = x_1, X_2 = x_2) = \mathbf{P}(X_1 = x_1)\mathbf{P}(X_2 = x_2) = f_{X_1}(x_1)f_{X_2}(x_2) \text{ for any } (x_1, x_2) \in \mathbb{R}^2 ,$$

and for **continuous** RVs  $X_1, X_2$  that are independent, the following equality is satisfied between the joint and marginal PDFs:

$$f_{X_1, X_2}(x_1, x_2) = f_{X_1}(x_1)f_{X_2}(x_2) \text{ for any } (x_1, x_2) \in \mathbb{R}^2 .$$

In summary, two RVs  $X$  and  $Y$  are said to be **independent** if and only if for every  $(x, y)$

$F_{X,Y}(x, y) = F_X(x) \times F_Y(y)$	or	$f_{X,Y}(x, y) = f_X(x) \times f_Y(y)$
--	----	--

Let us confirm that our familiar experiment of tossing a fair coin twice independently when encoded by a pair of independent Bernoulli(1/2) RVs satisfies the above definition.

**Example 50 (Pair of independent Bernoulli(1/2) RVs)** Let  $X_1$  and  $X_2$  be a pair of independent Bernoulli(1/2) RVs each taking values in the set  $\{0, 1\}$  with the following tabulated probabilities. Verify that the JPMF  $f_{X_1, X_2}(x_1, x_2) = 1/4$  for each  $(x_1, x_2) \in \{0, 1\}^2$  is indeed given by the marginal PMF  $f_{X_i}(x_i) = 1/2$  for each  $i \in \{1, 2\}$  and each  $x_i \in \{0, 1\}$ .

	$X_2 = 0$	$X_2 = 1$	
$X_1 = 0$	1/4	1/4	1/2
$X_1 = 1$	1/4	1/4	1/2
	1/2	1/2	1

From the above Table we can read for instance that the *joint probability* that  $\mathbb{R}^2$ -valued RV  $(X_1, X_2)$  takes the value or realization  $(0, 0)$  is 1/4 from the first entry of the inner-most tabulated rectangle, i.e.,  $\mathbf{P}((X_1, X_2) = (0, 0)) = 1/4$ , and that the *marginal probability* that the RV  $X_1$  takes the value or realization 0 is 1/2, i.e.,  $\mathbf{P}(X_1 = 0) = 1/2$ . Clearly,  $1/4 = 1/2 \times 1/2$ , and so our familiar experiment when seen as an  $\mathbb{R}^2$ -valued RV is indeed composed of two independent  $\mathbb{R}$ -valued Bernoulli(1/2) RVs.

**Example 51** Recall the  $\mathbb{R}^2$ -valued continuous RV  $(X, Y)$  of Example 45 that is uniformly distributed on the unit square  $[0, 1]^2$ . First show that  $X$  and  $Y$  independent. Then show that both  $X$  and  $Y$  are identically distributed according to the  $\text{Uniform}(0, 1)$  RV.

This can be shown by checking that the joint PDF is indeed equal to the product of the marginal PDFs of  $\text{Uniform}(0, 1)$  RVs as follows:

$$\begin{cases} 1 = f_{X,Y}(x, y) = f_X(x) \times f_Y(y) = 1 \times 1 = 1 & \text{if } (x, y) \in [0, 1]^2 \\ 0 = f_{X,Y}(x, y) = f_X(x) \times f_Y(y) = 0 \times 0 = 0 & \text{if } (x, y) \notin [0, 1]^2 \end{cases}$$

Are  $X$  and  $Y$  independent in the server times  $\vec{RV}$  from Example 46?

We can compute  $f_X(x)$  and use the already computed  $f_Y(y)$  to mechanically check if the JPDF is the product of the marginal PDFs. But intuitively, we know that these RVs (connection time and authentication time) are dependent – one is strictly greater than the other. Also the JPDF has zero density when  $x > y$ , but the product of the marginal densities won't.

Now, let us take advantage of independent random variables and solve some problems.

**Example 52 (distance between random faults in a manufactured line)** Suppose two points are tossed independently and uniformly at random onto a line segment of unit length. What is the probability that the distance between the two points does not exceed a given length  $l$ ?

done in lectures...

Let us introduce parameters for the lower and upper bounds of the interval upon which a continuous RV is uniformly distributed using the following probability model.


**Model 4 (Uniform( $\theta_1, \theta_2$ ))** Given two real parameters  $\theta_1, \theta_2 \in \mathbb{R}$ , such that  $\theta_1 < \theta_2$ , the PDF of the  $\text{Uniform}(\theta_1, \theta_2)$  RV  $X$  is:

$$f(x; \theta_1, \theta_2) = \begin{cases} \frac{1}{\theta_2 - \theta_1} & \text{if } \theta_1 \leq x \leq \theta_2, \\ 0 & \text{otherwise} \end{cases} \quad (3.20)$$

and its DF given by  $F(x; \theta_1, \theta_2) = \int_{-\infty}^x f(y; \theta_1, \theta_2) dy$  is:

$$F(x; \theta_1, \theta_2) = \begin{cases} 0 & \text{if } x < \theta_1 \\ \frac{x - \theta_1}{\theta_2 - \theta_1} & \text{if } \theta_1 \leq x \leq \theta_2, \\ 1 & \text{if } x > \theta_2 \end{cases} \quad (3.21)$$

Recall that we emphasise the dependence of the probabilities on the two parameters  $\theta_1$  and  $\theta_2$  by specifying them following the semicolon in the argument for  $f$  and  $F$ .

Figure 3.6: A plot of the PDF, DF or CDF and inverse DF of the Uniform( $-1, 1$ ) RV  $X$ .**Example 53 (Buffon's Needle Experiment to Physically Estimate  $\pi$ )**

done in lectures...

### 3.5.2 Conditional Random Variables

Often we will have a condition where one of the two random variables that make up a random vector  $(X_1, X_2)$  already occurs and takes a value. And we might want to compute the probability of the occurrence of the other random variable given this conditional information. For this all we need to do is extend the idea of conditional probabilities to  $\mathbb{R}^2$ -valued random variables as defined below.

**Definition 26 (Conditional PDF or PMF)** Let  $(X_1, X_2)$  be a discrete bivariate RV. The conditional PMF of  $X_1|X_2 = x_2$ , where  $f_{X_2}(x_2) := \mathbf{P}(X_2 = x_2) > 0$  is:

$$f_{X_1|X_2}(x_1|x_2) := \mathbf{P}(X_1 = x_1|X_2 = x_2) = \frac{\mathbf{P}(X_1 = x_1, X_2 = x_2)}{\mathbf{P}(X_2 = x_2)} = \frac{f_{X_1, X_2}(x_1, x_2)}{f_{X_2}(x_2)} .$$

Similarly, if  $f_{X_1}(x_1) := \mathbf{P}(X_1 = x_1) > 0$ , then the conditional PMF of  $X_2|X_1 = x_1$  is:

$$f_{X_2|X_1}(x_2|x_1) := \mathbf{P}(X_2 = x_2|X_1 = x_1) = \frac{\mathbf{P}(X_1 = x_1, X_2 = x_2)}{\mathbf{P}(X_1 = x_1)} = \frac{f_{X_1, X_2}(x_1, x_2)}{f_{X_1}(x_1)} .$$

If  $(X_1, X_2)$  are continuous RVs such that the marginal PDF  $f_{X_2}(x_2) > 0$ , then the conditional PDF of  $X_1|X_2 = x_2$  is:

$$f_{X_1|X_2}(x_1|x_2) = \frac{f_{X_1, X_2}(x_1, x_2)}{f_{X_2}(x_2)}, \quad \mathbf{P}(X_1 \in A|X_2 = x_2) = \int_A f_{X_1|X_2}(x_1|x_2) dx_1 .$$

Similarly, if  $f_{X_1}(x_1) > 0$ , then the conditional PDF of  $X_2|X_1 = x_1$  is:

$$f_{X_2|X_1}(x_2|x_1) = \frac{f_{X_1,X_2}(x_1, x_2)}{f_{X_1}(x_1)}, \quad \mathbf{P}(X_2 \in A|X_1 = x_1) = \int_A f_{X_2|X_1}(x_2|x_1) dx_2.$$

Let us consider a few discrete RVs for the simple coin tossing experiment  $\mathcal{E}_\theta^3$  that build on the Bernoulli( $\theta$ ) RV  $X_i$  for the  $i$ -th toss in an **independent and identically distributed (IID.)** manner.

Table 3.1: The 8  $\omega$ 's in the sample space  $\Omega$  of the experiment  $\mathcal{E}_\theta^3$  are given in the first row above. The RV  $Y$  is the number of ‘Heads’ in the 3 tosses and the RV  $Z$  is the number of ‘Tails’ in the 3 tosses. Finally, the RVs  $Y'$  and  $Z'$  are the indicator functions of the event that ‘all three tosses were Heads’ and the event that ‘all three tosses were Tails’, respectively.

$\omega:$	HHH	HHT	HTH	HTT	THH	THT	TTH	TTT	RV Definitions / Model
$\mathbf{P}(\omega):$	$\frac{1}{8}$	$X_i \stackrel{\text{IID}}{\sim} \text{Bernoulli}(\frac{1}{2})$							
$Y(\omega):$	3	2	2	1	2	1	1	0	$Y := X_1 + X_2 + X_3$
$Z(\omega):$	0	1	1	2	1	2	2	3	$Z := (1 - X_1) + (1 - X_2) + (1 - X_3)$
$Y'(\omega):$	1	0	0	0	0	0	0	0	$Y' := X_1 X_2 X_3$
$Z'(\omega):$	0	0	0	0	0	0	0	1	$Z' := (1 - X_1)(1 - X_2)(1 - X_3)$

**Classwork 54 (Two random variables of ‘toss a coin thrice’ experiment)** Describe the probability of the RV  $Y$  and  $Y'$  of Table 3.1 in terms of its PMF. Repeat the process for the RV  $Z$  in your spare time.

$$\mathbf{P}(Y = y) = \left\{ \begin{array}{l} \dots \\ \mathbf{P}(Y' = y') = \left\{ \begin{array}{l} \dots \end{array} \right. \end{array} \right.$$

**Classwork 55 (The number of ‘Heads’ given there is at least one ‘Tails’)** Consider the following two questions.

1. What is conditional probability  $\mathbf{P}(Y|Y' = 0)$  ?

$$\mathbf{P}(Y = y|Y' = 0) = \frac{\mathbf{P}(Y=y, Y'=0)}{\mathbf{P}(Y'=0)} = \frac{\mathbf{P}(\{\omega: Y(\omega)=y \cap Y'(\omega)=0\})}{\mathbf{P}(\{\omega: Y'(\omega)=0\})} = ?$$

$\mathbf{P}(Y = 0 Y' = 0)$	$\frac{\mathbf{P}(Y=0, Y'=0)}{\mathbf{P}(Y'=0)}$	$\frac{\frac{1}{8}}{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}$	$\frac{1}{7}$
$\mathbf{P}(Y = 1 Y' = 0)$	$\frac{\mathbf{P}(Y=1, Y'=0)}{\mathbf{P}(Y'=0)}$	$\frac{\frac{1}{8} + \frac{1}{8} + \frac{1}{8}}{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}$	$\frac{3}{7}$
$\mathbf{P}(Y = 2 Y' = 0)$	$\frac{\mathbf{P}(Y=2, Y'=0)}{\mathbf{P}(Y'=0)}$	$\frac{\frac{1}{8} + \frac{1}{8} + \frac{1}{8}}{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}$	$\frac{3}{7}$
$\mathbf{P}(Y = 3 Y' = 0)$	$\frac{\mathbf{P}(Y=3, Y'=0)}{\mathbf{P}(Y'=0)}$	$\frac{\mathbf{P}(\emptyset)}{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}$	0
$\mathbf{P}(Y \in \{0, 1, 2, 3\} Y' = 0)$	$\frac{\sum_{y=0}^3 \mathbf{P}(Y=y, Y'=0)}{\mathbf{P}(Y'=0)}$	$\frac{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}{\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}}$	1

2. What is  $\mathbf{P}(Y|Y' = 1)$  ?

$$\mathbf{P}(Y = y|Y' = 1) = \begin{cases} 1 & \text{if } y = 3 \\ 0 & \text{otherwise} \end{cases}$$

### 3.5.3 $\mathbb{R}^m$ -valued Random Variables

Consider the R $\vec{V}$   $X$  whose components are the RVs  $X_1, X_2, \dots, X_m$ , i.e.,  $X := (X_1, X_2, \dots, X_m)$ , where  $m \geq 2$ . A particular realization of this RV is a point  $(x_1, x_2, \dots, x_m)$  in  $\mathbb{R}^m$ . Now, let us extend the notions of JCDF, JPMF and JPDF to  $\mathbb{R}^m$ .

**Definition 27 (multivariate JDF)** The **joint distribution function (JDF)** or **joint cumulative distribution function (JCDF)**,  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) : \mathbb{R}^m \rightarrow [0, 1]$ , of the multivariate random vector  $(X_1, X_2, \dots, X_m)$  is

$$\begin{aligned} F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) &= \mathbf{P}(X \leq x_1 \cap X_2 \leq x_2 \cap \dots \cap X_m \leq x_m) \\ &= \mathbf{P}(X_1 \leq x_1, X_2 \leq x_2, \dots, X_m \leq x_m) \\ &= P(\{\omega : X_1(\omega) \leq x_1, X_2(\omega) \leq x_2, \dots, X_m(\omega) \leq x_m\}), \end{aligned} \quad (3.22)$$

for any  $(x_1, x_2, \dots, x_m) \in \mathbb{R}^m$ , where the right-hand side represents the probability that the random vector  $(X_1, X_2, \dots, X_m)$  takes on a value in  $\{(x'_1, x'_2, \dots, x'_m) : x'_1 \leq x_1, x'_2 \leq x_2, \dots, x'_m \leq x_m\}$ , the set of points in  $\mathbb{R}^m$  that are less than the point  $(x_1, x_2, \dots, x_m)$  in each coordinate  $1, 2, \dots, m$ .

The JDF  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) : \mathbb{R}^m \rightarrow \mathbb{R}$  satisfies the following conditions to remain a probability:

1.  $0 \leq F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) \leq 1$
2.  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m)$  is an increasing function of  $x_1, x_2, \dots$  and  $x_m$
3.  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) \rightarrow 1$  as  $x_1 \rightarrow \infty, x_2 \rightarrow \infty, \dots$  and  $x_m \rightarrow \infty$

$$4. F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) \rightarrow 0 \text{ as } x_1 \rightarrow -\infty, x_2 \rightarrow -\infty, \dots \text{ and } x_m \rightarrow -\infty$$

**Definition 28 (Multivariate JPMF)** If  $(X_1, X_2, \dots, X_m)$  is a **discrete random vector** that takes values in a discrete support set  $\mathcal{S}_{X_1, X_2, \dots, X_m}$ , then its **joint probability mass function** (or JPMF) is:

$$f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) = \mathbf{P}(X_1 = x_1, X_2 = x_2, \dots, X_m = x_m) . \quad (3.23)$$

Since  $\mathbf{P}(\Omega) = 1$ ,  $\sum_{(x_1, x_2, \dots, x_m) \in \mathcal{S}_{X_1, X_2, \dots, X_m}} f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) = 1$ .

From JPMF  $f_{X_1, X_2, \dots, X_m}$  we can get the JCDF  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m)$  and the probability of any event  $B$  by simply taking sums as in Equation (3.16) but now over all  $m$  coordinates.

**Definition 29 (Multivariate JPFD)**  $(X_1, X_2, \dots, X_m)$  is a **continuous random vector** if its JDF  $F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m)$  is differentiable and the **joint probability density function (JPDF)** is given by:

$$f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) = \frac{\partial^m}{\partial x_1 \partial x_2 \dots \partial x_m} F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) ,$$

From JPFD  $f_{X_1, X_2, \dots, X_m}$  we can compute the JDF  $F_{X_1, X_2, \dots, X_m}$  at any point  $(x_1, x_2, \dots, x_m) \in \mathbb{R}^m$  and more generally we can compute the probability of any event  $B$ , that can be cast as a region in  $\mathbb{R}^m$ , by “simply” taking  $m$ -dimensional integrals (you have done such iterated integrals when  $m = 3$ ):

$$\boxed{F_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) = \int_{-\infty}^{x_m} \dots \int_{-\infty}^{x_2} \int_{-\infty}^{x_1} f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) dx_1 dx_2 \dots dx_m} , \quad (3.24)$$

and

$$\boxed{\mathbf{P}(B) = \int \dots \int \int_B f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) dx_1 dx_2 \dots dx_m} . \quad (3.25)$$

The JPFD satisfies the following two properties:

1. integrates to 1, i.e.,  $\int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) dx_1 dx_2 \dots dx_m = 1$
2. is a non-negative function, i.e.,  $f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) \geq 0$ .

The marginal PDF (marginal PMF) is obtained by integrating (summing) the JPFD (JPMF) over all other random variables. For example, the marginal PDF of  $X_1$  is

$$f_{X_1}(x_1) = \int_{x_2=-\infty}^{\infty} \dots \int_{x_m=-\infty}^{\infty} f_{X_1, X_2, \dots, X_m}(x_1, x_2, \dots, x_m) dx_2 \dots dx_m$$

**Definition 30 (Independence of Sequence of RVs)** A finite or infinite sequence of RVs  $X_1, X_2, \dots$  is said to be independent or independently distributed if and only if

$$\mathbf{P}(X_{i_1} \leq x_{i_1}, X_{i_2} \leq x_{i_2}, \dots, X_{i_k} \leq x_{i_k}) = \mathbf{P}(X_{i_1} \leq x_{i_1}) \mathbf{P}(X_{i_2} \leq x_{i_2}) \dots \mathbf{P}(X_{i_k} \leq x_{i_k})$$

or equivalently,

$$F_{X_{i_1}, X_{i_2}, \dots, X_{i_m}}(x_{i_1}, x_{i_2}, \dots, x_{i_m}) = F_{X_{i_1}}(x_{i_1}) F_{X_{i_2}}(x_{i_2}) \dots F_{X_{i_m}}(x_{i_m}) ,$$

for any distinct subset of indices  $\{i_1, i_2, \dots, i_m\}$  of  $\{1, 2, \dots\}$ , the index set of the sequence of RVs and any sequence of real numbers  $x_{i_1}, x_{i_2}, \dots, x_{i_m}$ .

By the above definition, the sequence of **discrete** RVs  $X_1, X_2, \dots$  taking values in an at most countable set  $\mathbb{D}$  are said to be independently distributed if for any distinct subset of indices  $\{i_1, i_2, \dots, i_k\}$  such that the corresponding RVs  $X_{i_1}, X_{i_2}, \dots, X_{i_k}$  exists as a distinct subset of our original sequence of RVs  $X_1, X_2, \dots$  and for any elements  $x_{i_1}, x_{i_2}, \dots, x_{i_k}$  in  $\mathbb{D}$ , the following equality is satisfied:

$$\mathbf{P}(X_{i_1} = x_{i_1}, X_{i_2} = x_{i_2}, \dots, X_{i_k} = x_{i_k}) = \mathbf{P}(X_{i_1} = x_{i_1})\mathbf{P}(X_{i_2} = x_{i_2}) \cdots \mathbf{P}(X_{i_k} = x_{i_k})$$

For an independent sequence of RVs  $\{X_1, X_2, \dots\}$ , we have

$$\begin{aligned} & \mathbf{P}(X_{i+1} \leq x_{i+1} | X_i \leq x_i, X_{i-1} \leq x_{i-1}, \dots, X_1 \leq x_1) \\ &= \frac{\mathbf{P}(X_{i+1} \leq x_{i+1}, X_i \leq x_i, X_{i-1} \leq x_{i-1}, \dots, X_1 \leq x_1)}{\mathbf{P}(X_i \leq x_i, X_{i-1} \leq x_{i-1}, \dots, X_1 \leq x_1)} \\ &= \frac{\mathbf{P}(X_{i+1} \leq x_{i+1})\mathbf{P}(X_i \leq x_i)\mathbf{P}(X_{i-1} \leq x_{i-1}) \cdots \mathbf{P}(X_1 \leq x_1)}{\mathbf{P}(X_i \leq x_i)\mathbf{P}(X_{i-1} \leq x_{i-1}) \cdots \mathbf{P}(X_1 \leq x_1)} \\ &= \mathbf{P}(X_{i+1} \leq x_{i+1}) \end{aligned}$$

The above equality that

$$\mathbf{P}(X_{i+1} \leq x_{i+1} | X_i \leq x_i, X_{i-1} \leq x_{i-1}, \dots, X_1 \leq x_1) = \mathbf{P}(X_{i+1} \leq x_{i+1})$$

simply says that the conditional distribution of the RV  $X_{i+1}$  given all previous RVs  $X_i, X_{i-1}, \dots, X_1$  is simply determined by the distribution of  $X_{i+1}$ .

When a sequence of RVs are not independent they are said to be **dependent**.

## 3.6 Expectations

Expectation is perhaps the most fundamental concept in probability theory. In fact, probability is itself an expectation as you will soon see!

Expectation is one of the fundamental concepts in probability. The expected value of a real-valued random variable gives the population mean, a measure of the centre of the distribution of the variable in some sense. Its variance measures its spread and so on.

**Definition 31 (Expectation of a RV)** The **expectation**, or **expected value**, or **mean**, or **first moment**, of a random variable  $X$ , with distribution function  $F$  and density  $f$ , is defined to be

$$\mathbf{E}(X) := \int x dF(x) = \begin{cases} \sum_x x f(x) & \text{if } X \text{ is discrete} \\ \int x f(x) dx & \text{if } X \text{ is continuous,} \end{cases} \quad (3.26)$$

provided the sum or integral is well-defined. We say the expectation exists if

$$\int |x| dF(x) < \infty. \quad (3.27)$$

Sometimes, we denote  $\mathbf{E}(X)$  by  $\mathbf{E}X$  for brevity. Thus, the expectation is a single-number summary of the RV  $X$  and may be thought of as the average. We subscript  $E$  to specify the parameter  $\theta \in \Theta$  with respect to which the integration is undertaken.

$$\mathbf{E}_\theta X := \int x dF(x; \theta)$$

**Definition 32 (Variance of a RV)** Let  $X$  be a RV with mean or expectation  $\mathbf{E}(X)$ . Variance of  $X$  denoted by  $\mathbf{V}(X)$  or  $VX$  is

$$\mathbf{V}(X) := \mathbf{E}((X - \mathbf{E}(X))^2) = \int (x - \mathbf{E}(X))^2 dF(x),$$

provided this expectation exists. The **standard deviation** denoted by  $\text{sd}(X) := \sqrt{\mathbf{V}(X)}$ . Thus variance is a measure of “spread” of a distribution.

**Definition 33 ( $k$ -th moment of a RV)** We call

$$\mathbf{E}(X^k) = \int x^k dF(x)$$

as the  $k$ -th moment of the RV  $X$  and say that the  $k$ -th moment exists when  $\mathbf{E}(|X|^k) < \infty$ . We call the following expectation as the  $k$ -th central moment:

$$\mathbf{E}((X - \mathbf{E}(X))^k).$$

### 3.6.1 Expectations of functions of random variables

More generally, by taking the expected value of various functions of a random variable, we can measure many interesting features of its distribution, including spread and correlation.

**Definition 34 (Expectation of a function of a RV)** The **Expectation** of a function  $g(X)$  of a random variable  $X$  is defined as:

$$\mathbf{E}(g(X)) := \int g(x)dF(x) = \begin{cases} \sum_x g(x)f(x) & \text{if } X \text{ is a discrete RV} \\ \int_{-\infty}^{\infty} g(x)f(x)dx & \text{if } X \text{ is a continuous RV} \end{cases}$$

provided  $\mathbf{E}(g(X))$  exists, i.e.,  $\int |g(x)|dF(x) < \infty$ .

The **mean** which characterises the central location of the random variable  $X$  is merely the expectation of the identity function  $g(x) = x$ :

$$\mathbf{E}(X) = \begin{cases} \sum_x xf(x) & \text{if } X \text{ is a discrete RV} \\ \int_{-\infty}^{\infty} xf(x)dx & \text{if } X \text{ is a continuous RV} \end{cases}$$

Often, mean is denoted by  $\mu$ .

The **variance** which characterises the spread or the variability of the random variable  $X$  is also the expectation of the function  $g(x) = (x - \mathbf{E}(X))^2$ :

$$\mathbf{V}(X) = \mathbf{E}((X - \mathbf{E}(X))^2) = \begin{cases} \sum_x (x - \mathbf{E}(X))^2 f(x) & \text{if } X \text{ is a discrete RV} \\ \int_{-\infty}^{\infty} (x - \mathbf{E}(X))^2 f(x) dx & \text{if } X \text{ is a continuous RV} \end{cases}$$

Often, variance is denoted by  $\sigma^2$ .

### INTUITIVELY, WHAT IS EXPECTATION?

Definition 34 gives expectation as a “weighted average” of the possible values. This is true but some intuitive idea of expectation is also helpful.

- Expectation is what you expect.

Consider tossing a fair coin. If it is heads you lose \$10. If it is tails you win \$10. What do you expect to win? Nothing. If  $X$  is the amount you win then

$$\mathbf{E}(X) = -10 \times \frac{1}{2} + 10 \times \frac{1}{2} = 0.$$

So what you expect (nothing) and the weighted average ( $\mathbf{E}(X) = 0$ ) agree.

- Expectation is a long run average.

Suppose you are able to repeat an experiment independently, over and over again. Each experiment produces one value  $x$  of a random variable  $X$ . If you take the average of the  $x$  values for a large number of trials, then this average converges to  $\mathbf{E}(X)$  as the number of trials grows. In fact, this is called the **law of large numbers**.

We can concretize the above two intuitive insights by the following two examples.

**Example 56 (Winnings on Average)** Let  $Y = r(X)$ . Then

$$\mathbf{E}(Y) = \mathbf{E}(r(X)) = \int r(x) dF(x) .$$

Think of playing a game where we draw  $x \sim X$  and then I pay you  $y = r(x)$ . Then your average income is  $r(x)$  times the chance that  $X = x$ , summed (or integrated) over all values of  $x$ .

**Example 57 (Probability is an Expectation)** Let  $A$  be an event and let  $r(X) = \mathbb{1}_A(x)$ . Recall  $\mathbb{1}_A(x)$  is 1 if  $x \in A$  and  $\mathbb{1}_A(x) = 0$  if  $x \notin A$ . Then

$$\mathbf{E}(\mathbb{1}_A(X)) = \int \mathbb{1}_A(x) dF(x) = \int_A dF(x) = \mathbf{P}(X \in A) = \mathbf{P}(A) \quad (3.28)$$

Thus, probability is a special case of expectation. Recall our LTRF motivation for the definition of probability and make the connection.

### Expectations of functions of $\mathbb{R}^2$ -valued random variables

In the case of a single random variable we saw that its expectation gives the population mean, a measure of the center of the distribution of the variable in some sense. Similarly, by taking the expected value of various functions of a  $\mathbb{R}^2$ -valued random variable, we can measure many interesting features of its joint distribution.

**Definition 35** The **Expectation** of a function  $g(X, Y)$  of the  $\mathbb{R}^2$ -valued RV  $(X, Y)$  is defined as:

$$\mathbf{E}(g(X, Y)) = \begin{cases} \sum_{(x,y)} g(x, y) f_{X,Y}(x, y) & \text{if } (X, Y) \text{ is a discrete RV} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy & \text{if } (X, Y) \text{ is a continuous RV} \end{cases}$$

Some typical expectations for  $\mathbb{R}^2$ -valued random variables are:

1. Joint Moments

$$\mathbf{E}(X^r Y^s)$$

When  $r = s = 1$ , we have  $\mathbf{E}(XY)$ , the expectation of the product of two RVs.

2. We need a new notion for the variance of two RVs.

If  $\mathbf{E}(X^2) < \infty$  and  $\mathbf{E}(Y^2) < \infty$  then  $\mathbf{E}(|XY|) < \infty$  and  $\mathbf{E}(|(X - \mathbf{E}(X))(Y - \mathbf{E}(Y))|) < \infty$ . This allows the definition of **covariance** of  $X$  and  $Y$  as

$$\mathbf{Cov}(X, Y) := \mathbf{E}((X - \mathbf{E}(X))(Y - \mathbf{E}(Y))) = \mathbf{E}(XY) - \mathbf{E}(X)\mathbf{E}(Y)$$

The same ideas naturally extend, via multiple sums and integrals, to define the expectation of functions of  $\mathbb{R}^k$ -valued random variables with  $k > 2$ .

### Viewing a deterministic real variable as a random variable

Consider the class of discrete RVs with distributions that place all probability mass on a single real number. This is the probability model for the deterministic real variable, which is often thought of as an unknown constant  $\theta \in \mathbb{R}$ .

**Model 5** (Point Mass( $\theta$ )) Given a specific point  $\theta \in \mathbb{R}$ , we say an RV  $X$  has point mass at  $\theta$  or is Point Mass( $\theta$ ) distributed if the DF is:

$$F(x; \theta) = \begin{cases} 0 & \text{if } x < \theta \\ 1 & \text{if } x \geq \theta \end{cases} \quad (3.29)$$

and the PMF is:

$$f(x; \theta) = \begin{cases} 0 & \text{if } x \neq \theta \\ 1 & \text{if } x = \theta \end{cases} \quad (3.30)$$

Thus, Point Mass( $\theta$ ) RV  $X$  is deterministic in the sense that every realisation of  $X$  is exactly equal to  $\theta \in \mathbb{R}$ . We will see that this distribution plays a central limiting role in asymptotic statistics.

**Example 58 (Mean and variance of Point Mass( $\theta$ ) RV)** Let  $X \sim \text{Point Mass}(\theta)$ . Then:

$$\mathbf{E}(X) = \sum_x x f(x) = \theta \times 1 = \theta , \quad \mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = \theta^2 - \theta^2 = 0 .$$

### 3.6.2 Properties of expectations

The following results, where  $a$  is a constant, may easily be proved using the properties of summations and integrals:

$$\boxed{\mathbf{E}(a) = a}$$

$$\boxed{\mathbf{E}(a g(X)) = a \mathbf{E}(g(X))}$$

$$\boxed{\mathbf{E}(g(X) + h(X)) = \mathbf{E}(g(X)) + \mathbf{E}(h(X))}$$

Note that here  $g(X)$  and  $h(X)$  are functions of the random variable  $X$ : e.g.  $g(X) = X^2$ .

Using these results we can obtain the following useful formula for variance:

$$\begin{aligned} V(X) &= E((X - \mathbf{E}(X))^2) \\ &= E(X^2 - 2X\mathbf{E}(X) + (\mathbf{E}(X))^2) \\ &= E(X^2) - E(2X\mathbf{E}(X)) + E((\mathbf{E}(X))^2) \\ &= E(X^2) - 2\mathbf{E}(X)\mathbf{E}(X) + (\mathbf{E}(X))^2 \\ &= E(X^2) - 2(\mathbf{E}(X))^2 + (\mathbf{E}(X))^2 \\ &= \mathbf{E}(X^2) - (\mathbf{E}(X))^2 . \end{aligned}$$

That is,

$$\boxed{V(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2} .$$

The above properties of expectations imply that for constants  $a$  and  $b$ ,

$$\boxed{\mathbf{V}(aX + b) = a^2\mathbf{V}(X)} \quad (3.31)$$

More generally, for random variables  $X_1, X_2, \dots, X_n$  and constants  $a_1, a_2, \dots, a_n$

- $\mathbf{E}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i \mathbf{E}(X_i)$  . (3.32)

- $\mathbf{V}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i^2 \mathbf{V}(X_i)$ , provided  $X_1, X_2, \dots, X_n$  are independent . (3.33)

- Let  $X_1, X_2, \dots, X_n$  be independent RVs, then

$$\mathbf{E}\left(\prod_{i=1}^n X_i\right) = \prod_{i=1}^n \mathbf{E}(X_i), \text{ provided } X_1, X_2, \dots, X_n \text{ are independent .} \quad (3.34)$$

Next, let us compute the mean and variance of our familiar RVs.

**Example 59 (Mean and variance of Bernoulli( $\theta$ ) RV)** Let  $X \sim \text{Bernoulli}(\theta)$ . Then,

$$\mathbf{E}(X) = \sum_{x=0}^1 xf(x) = (0 \times (1 - \theta)) + (1 \times \theta) = 0 + \theta = \theta ,$$

$$\mathbf{E}(X^2) = \sum_{x=0}^1 x^2 f(x) = (0^2 \times (1 - \theta)) + (1^2 \times \theta) = 0 + \theta = \theta ,$$

$$\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = \theta - \theta^2 = \theta(1 - \theta) .$$

Parameter specifically,

$$\mathbf{E}_\theta(X) = \theta \quad \text{and} \quad \mathbf{V}_\theta(X) = \theta(1 - \theta) .$$


Maximum of the variance  $\mathbf{V}_\theta(X)$  is found by setting the derivative to zero, solving for  $\theta$  and showing the second derivative is locally negative, i.e.  $\mathbf{V}_\theta(X)$  is concave down:

$$\mathbf{V}'_\theta(X) := \frac{d}{d\theta} \mathbf{V}_\theta(X) = 1 - 2\theta = 0 \iff \theta = \frac{1}{2} , \quad \mathbf{V}''_\theta(X) := \frac{d}{d\theta} \left( \frac{d}{d\theta} \mathbf{V}_\theta(X) \right) = -2 < 0 ,$$

$$\max_{\theta \in [0,1]} \mathbf{V}_\theta(X) = \frac{1}{2} \left( 1 - \frac{1}{2} \right) = \frac{1}{4} , \text{ since } \mathbf{V}_\theta(X) \text{ is maximized at } \theta = \frac{1}{2}$$

The plot depicting these expectations as well as the rate of change of the variance are depicted in Figure 3.7. Note from this Figure that  $\mathbf{V}_\theta(X)$  attains its maximum value of  $1/4$  at  $\theta = 0.5$  where  $\frac{d}{d\theta} \mathbf{V}_\theta(X) = 0$ . Furthermore, we know that we don't have a minimum at  $\theta = 0.5$  since the second derivative  $\mathbf{V}''_\theta(X) = -2$  is negative for any  $\theta \in [0, 1]$ . This confirms that  $\mathbf{V}_\theta(X)$  is concave down and therefore we have a maximum of  $\mathbf{V}_\theta(X)$  at  $\theta = 0.5$ . We will revisit this example when we employ a numerical approach called Newton-Raphson method to solve for the maximum of a differentiable function by setting its derivative equal to zero.

Figure 3.7: Mean ( $\mathbf{E}_\theta(X)$ ), variance ( $\mathbf{V}_\theta(X)$ ) and the rate of change of variance ( $\frac{d}{d\theta}\mathbf{V}_\theta(X)$ ) of a Bernoulli( $\theta$ ) RV  $X$  as a function of the parameter  $\theta$ .



**Example 60 (Mean and variance of Uniform(0, 1) RV)** Let  $X \sim \text{Uniform}(0, 1)$ . Then,

$$\mathbf{E}(X) = \int_{x=0}^1 xf(x) dx = \int_{x=0}^1 x 1 dx = \frac{1}{2} (x^2)_{x=0}^{x=1} = \frac{1}{2} (1 - 0) = \frac{1}{2},$$

$$\mathbf{E}(X^2) = \int_{x=0}^1 x^2 f(x) dx = \int_{x=0}^1 x^2 1 dx = \frac{1}{3} (x^3)_{x=0}^{x=1} = \frac{1}{3} (1 - 0) = \frac{1}{3},$$

$$\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = \frac{1}{3} - \left(\frac{1}{2}\right)^2 = \frac{1}{3} - \frac{1}{4} = \frac{1}{12}.$$

**Example 61 (Expected Exponential of the Uniform(0, 1) RV)** Let  $X \sim \text{Uniform}(0, 1)$  and  $Y = r(X) = e^X$ . Compute  $\mathbf{E}(Y)$ .

We can simply apply the definition of  $\mathbf{E}(r(X))$ , since  $Y = r(X)$ , is just a function of  $X$ , as follows:

$$\mathbf{E}(Y) = \int_0^1 e^x f(x) dx = \int_0^1 e^x 1 dx = e - 1.$$

**Mean and Variance of Exponential( $\lambda$ ):** Show that the mean of an Exponential( $\lambda$ ) RV  $X$  is:

$$\mathbf{E}_\lambda(X) = \int_0^\infty xf(x; \lambda) dx = \int_0^\infty x\lambda e^{-\lambda x} dx = \frac{1}{\lambda},$$

and the variance is:

$$\mathbf{V}_\lambda(X) = \left(\frac{1}{\lambda}\right)^2.$$

### 3.7 Stochastic Processes

**Definition 36 (Stochastic Process)** A collection of RVs

$$(X_\alpha)_{\alpha \in \mathbb{A}} := (X_\alpha : \alpha \in \mathbb{A})$$

is called a **stochastic process**. Thus, for every  $\alpha \in \mathbb{A}$ , the index set of the stochastic process,  $X_\alpha$  is a RV. If the index set  $\mathbb{A} \subset \mathbb{Z}$  then we have a **discrete time stochastic process**, typically denoted by

$$(X_i)_{i \in \mathbb{Z}} := \dots, X_{-2}, X_{-1}, X_0, X_1, X_2, \dots, \text{ or}$$

$$(X_i)_{i \in \mathbb{N}} := X_1, X_2, \dots, \text{ or}$$

$$(X_i)_{i \in [n]} := X_1, X_2, \dots, X_n, \text{ where, } [n] := \{1, 2, \dots, n\}.$$

If  $\mathbb{A} \subset \mathbb{R}$  then we have a **continuous time stochastic process**, typically denoted by  $\{X_t\}_{t \in \mathbb{R}}$ , etc.

Of course the above process is quite general and can allow for arbitrary dependence among the RVs. Generally, we cannot produce useful models without making simplifying assumptions. The absolutely simplest but extremely useful assumption is that of the **Independent and Identically Distributed or IID Process** or merely **IID Sequence of RVs** when the index set is a subset of  $\mathbb{N}$ .

This is exactly the sequence of RVs associated with our product experiment  $\mathcal{E}^{\otimes \infty} := (\Omega, \mathcal{F}_\mathcal{X}, P_\theta)^{\otimes \infty}$ .

**Definition 37 (Independent and Identically Distributed (IID) Process)** The finite or infinite sequence of RVs or the stochastic process  $X_1, X_2, \dots$  is said to be independent and identically distributed or IID if :

- they are independently distributed according to Definition 30, and
- $F(X_1) = F(X_2) = \dots$ , ie. all the  $X_i$ 's have the same DF  $F(X_1)$ .

This is perhaps the most elementary class of stochastic processes and we succinctly denote it by

$$(X_i)_{i \in [n]} := X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} F, \quad \text{or} \quad (X_i)_{i \in \mathbb{N}} := X_1, X_2, \dots \stackrel{\text{IID}}{\sim} F.$$

We sometimes replace the DF  $F$  above by the name of the RV.

**Definition 38 (Independently Distributed)** The sequence of RVs or the stochastic process  $(X_i)_{i \in \mathbb{N}} := X_1, X_2, \dots$  is said to be independently distributed if :

- $X_1, X_2, \dots$  is independently distributed according to Definition 30.

This is a class of stochastic processes that is more general than the IID class.

Let us consider a RV that arises from an IID stochastic process of Bernoulli( $\theta$ ) RVs  $\{X_i\}_{i \in \mathbb{N}}$ , ie.

$$\{X_i\}_{i \in \mathbb{N}} := \{X_1, X_2, \dots\} \stackrel{\text{IID}}{\sim} \text{Bernoulli}(\theta).$$

When we consider the number of IID Bernoulli( $\theta$ ) trials before the first ‘Head’ occurs we get the following discrete RV.

**Model 6** (Geometric( $\theta$ ) RV) Given a parameter  $\theta \in (0, 1)$ , the PMF of the Geometric( $\theta$ ) RV  $X$  is

$$f(x; \theta) = \begin{cases} \theta(1 - \theta)^x & \text{if } x \in \mathbb{Z}_+ := \{0, 1, 2, \dots\} \\ 0 & \text{otherwise} \end{cases} \quad (3.35)$$

It is straightforward to verify that  $f(x; \theta)$  is indeed a PDF :

$$\sum_{x=0}^{\infty} f(x; \theta) = \sum_{x=0}^{\infty} \theta(1 - \theta)^x = \theta \left( \frac{1}{1 - (1 - \theta)} \right) = \theta \left( \frac{1}{\theta} \right) = 1$$

The above equality is a consequence of the geometric series identity (3.36) with  $a = \theta$  and  $\vartheta := 1 - \theta$ :

$$\sum_{x=0}^{\infty} a\vartheta^x = a \left( \frac{1}{1 - \vartheta} \right), \text{ provided, } 0 < \vartheta < 1. \quad (3.36)$$

**Proof:**

$$a + a\vartheta + a\vartheta^2 + \dots + a\vartheta^n = \sum_{0 \leq x \leq n} a\vartheta^x = a + \sum_{1 \leq x \leq n} a\vartheta^x = a + \vartheta \sum_{1 \leq x \leq n} a\vartheta^{x-1} = a + \vartheta \sum_{0 \leq x \leq n-1} a\vartheta^x = a + \vartheta \sum_{0 \leq x \leq n} a\vartheta^x - a\vartheta^{n+1}$$

Therefore,

$$\begin{aligned} \sum_{0 \leq x \leq n} a\vartheta^x &= a + \vartheta \sum_{0 \leq x \leq n} a\vartheta^x - a\vartheta^{n+1} \\ \left( \sum_{0 \leq x \leq n} a\vartheta^x \right) - \left( \vartheta \sum_{0 \leq x \leq n} a\vartheta^x \right) &= a - a\vartheta^{n+1} \\ \left( \sum_{0 \leq x \leq n} a\vartheta^x \right) (1 - \vartheta) &= a(1 - \vartheta^{n+1}) \\ \sum_{0 \leq x \leq n} a\vartheta^x &= a \left( \frac{1 - \vartheta^{n+1}}{1 - \vartheta} \right) \\ \sum_{x=0}^{\infty} a\vartheta^x := \lim_{n \rightarrow \infty} \sum_{0 \leq x \leq n} a\vartheta^x &= a \left( \frac{1}{1 - \vartheta} \right), \text{ provided, } 0 < \vartheta < 1 \end{aligned}$$

The outcome of a Geometric( $\theta$ ) RV can be thought of as “the number of tosses needed before the appearance of the first ‘Head’ when tossing a coin with probability of ‘Heads’ equal to  $\theta$  in a independent and identical manner.”

**Example 62 (Mean and variance of Geometric( $\theta$ ) RV)** Let  $X \sim \text{Geometric}(\theta)$  RV. Then,

$$\mathbf{E}(X) = \sum_{x=0}^{\infty} x\theta(1 - \theta)^x = \theta \sum_{x=0}^{\infty} x(1 - \theta)^x$$

In order to simplify the RHS above, let us employ differentiation with respect to  $\theta$ :

$$\frac{-1}{\theta^2} = \frac{d}{d\theta} \left( \frac{1}{\theta} \right) = \frac{d}{d\theta} \sum_{x=0}^{\infty} (1 - \theta)^x = \sum_{x=0}^{\infty} -x(1 - \theta)^{x-1}$$

Multiplying the LHS and RHS above by  $-(1 - \theta)$  and substituting in  $\mathbf{E}(X) = \theta \sum_{x=0}^{\infty} x(1 - \theta)^x$ , we get a much simpler expression for  $\mathbf{E}(X)$  :

$$\frac{1 - \theta}{\theta^2} = \sum_{x=0}^{\infty} x(1 - \theta)^x \implies \mathbf{E}(X) = \theta \left( \frac{1 - \theta}{\theta^2} \right) = \frac{1 - \theta}{\theta}.$$

Similarly, it can be shown that

$$\mathbf{V}(X) = \frac{1 - \theta}{\theta^2}.$$

Figure 3.8: PDF of  $X \sim \text{Geometric}(\theta = 0.5)$  and the relative frequency histogram based on 100 and 1000 samples from  $X$  according to Simulation 112 and Labwork 113.





Figure 3.9: Mean and variance of a Geometric( $\theta$ ) RV  $X$  as a function of the parameter  $\theta$ .



**Example 63 (Coupon Collectors Problem)** Recall the Coupon Collector's Problem from lectures.

The RV  $Y$  in Table 3.1 may be generalized to an experiment  $\mathcal{E}_\theta^n$  with  $n$  coin tosses. Let  $X_i$  be the Indicator function of the event ‘Heads on the  $i$ -th toss’ as before. Then  $Y$  defined by,

$$Y := \sum_{i=1}^n X_i := X_1 + X_2 + \cdots + X_n ,$$

is the number of ‘Heads’ in  $n$  tosses. Akin to the second row of Table 3.1, for the ‘Toss  $n$  times’ experiment  $\mathcal{E}_\theta^n$  the RV  $Y$  as defined above will take values in  $\{0, 1, 2, \dots, n\}$  and is therefore a discrete RV. This is called the Binomial RV as defined next. But, first we remind ourselves of some elementary definitions involving arrangements of objects from a collection (recall Section 1.5).

**Model 7 (Binomial( $n, \theta$ ) RV)** Let the RV  $X = \sum_{i=1}^n X_i$  be the sum of  $n$  independent and identically distributed Bernoulli( $\theta$ ) RVs, i.e.:

$$X = \sum_{i=1}^n X_i, \quad X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} \text{Bernoulli}(\theta) .$$

Given two parameters  $n$  and  $\theta$ , the PMF of the Binomial( $n, \theta$ ) RV  $X$  is:

$$f(x; n, \theta) = \begin{cases} \binom{n}{x} \theta^x (1 - \theta)^{n-x} & \text{if } x \in \{0, 1, 2, 3, \dots, n\} , \\ 0 & \text{otherwise} \end{cases} \quad (3.37)$$

where,  $\binom{n}{x}$  is:

$$\binom{n}{x} = \frac{n(n-1)(n-2)\dots(n-x+1)}{x(x-1)(x-2)\dots(2)(1)} = \frac{n!}{x!(n-x)!} .$$

$\binom{n}{x}$  is read as “ $n$  choose  $x$ .”

**Proof:** This is only a sketch. A formal proof should start with the mathematical induction for the very formula for the binomial coefficient.

Observe that for the Binomial( $n, \theta$ ) RV  $X$ ,  $\mathbf{P}(X = x) = f(x; n, \theta)$  is the probability that  $x$  of the  $n$  Bernoulli( $\theta$ ) trials result in an outcome of 1’s. Next note that if all  $n$   $X_i$ ’s are 0’s, then  $X = 0$ , and if all  $n$   $X_i$ ’s are 1’s, then  $X = n$ . In general, if some of the  $n$   $X_i$ ’s are 1’s and the others are 0, then  $X$  can only take values in  $\{0, 1, 2, \dots, n\}$  and therefore  $f(x; n, \theta) = 0$  if  $x \notin \{0, 1, 2, \dots, n\}$ .

Now, let us compute  $f(x; n, \theta)$  when  $x \in \{0, 1, 2, \dots, n\}$ . Consider the set of indices  $\{1, 2, 3, \dots, n\}$  for the  $n$  IID Bernoulli( $\theta$ ) RVs  $\{X_1, X_2, \dots, X_n\}$ . Now choose  $x$  indices from  $\{1, 2, \dots, n\}$  to mark

those trials in a particular realization of  $\{x_1, x_2, \dots, x_n\}$  with the Bernoulli outcome of 1. The probability of each such event is  $\theta^x(1 - \theta)^{n-x}$  due to the IID assumption. For each realization  $\{x_1, x_2, \dots, x_n\} \in \{0, 1\}^n := \{\text{all binary } (0-1) \text{ strings of length } n\}$ , specified by a choice of  $x$  trial indices with Bernoulli outcome 1, the binomial RV  $X = \sum_{i=1}^n X_i$  takes the value  $x$ . Since there are exactly  $\binom{n}{x}$  many ways in which we can choose  $x$  trial indices (with outcome 1) from the set of  $n$  trial indices  $\{1, 2, \dots, n\}$ , we get the desired product for  $f(x; n, \theta) = \binom{n}{x} \theta^x (1 - \theta)^{n-x}$  when  $x \in \{0, 1, \dots, n\}$ .


**Mean and variance of  $\text{Binomial}(n, \theta)$  RV:** Let  $X \sim \text{Binomial}(n, \theta)$ . Based on the definition of expectation:

$$\mathbf{E}(X) = \int x dF(x; n, \theta) = \sum_x x f(x; n, \theta) = \sum_{x=0}^n x \binom{n}{x} \theta^x (1 - \theta)^{n-x} .$$

However, this is a nontrivial sum to evaluate. Instead, we may use (3.32) and (3.33) by noting that  $X = \sum_{i=1}^n X_i$ , where the  $\{X_1, X_2, \dots, X_n\} \stackrel{\text{IID}}{\sim} \text{Bernoulli}(\theta)$ ,  $\mathbf{E}(X_i) = \theta$  and  $\mathbf{V}(X_i) = \theta(1 - \theta)$ :

$$\begin{aligned} \mathbf{E}(X) &= \mathbf{E}(X_1 + X_2 + \dots + X_n) = \mathbf{E}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \mathbf{E}(X_i) = n\theta , \\ \mathbf{V}(X) &= \mathbf{V}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \mathbf{V}(X_i) = \sum_{i=1}^n \theta(1 - \theta) = n\theta(1 - \theta) . \end{aligned}$$

Figure 3.10: PDF of  $X \sim \text{Binomial}(n = 10, \theta = 0.5)$  and the relative frequency histogram based on 100,000 samples from  $X$  obtained according to Simulation 116.



### 3.8 Convergence of Random Variables

This important topic is concerned with the limiting behavior of sequences of RVs


$$\{X_i\}_{i=1}^n := X_1, X_2, X_3, \dots, X_{n-1}, X_n \quad \text{as } n \rightarrow \infty .$$

From a statistical viewpoint  $n \rightarrow \infty$  is associated with the amount of data or information  $\rightarrow \infty$ . Refresh yourself with notions of convergence, limits and continuity in the real line (**S 1.6**) before proceeding further.

**Classwork 64 (Convergence of  $X_i \sim \text{Normal}(0, 1/i)$ )** Suppose you are given an independent sequence of RVs  $\{X_i\}_{i=1}^n$ , where  $X_i \sim \text{Normal}(0, 1/i)$ . How would you talk about the convergence of  $X_n \sim \text{Normal}(0, 1/n)$  as  $n$  approaches  $\infty$ ? Take a look at Figure 3.11 for insight. The probability mass of  $X_n$  increasingly concentrates about 0 as  $n$  approaches  $\infty$  and the variance  $1/n$  approaches 0, as depicted in Figure 3.11. Based on this observation, can we expect  $\lim_{n \rightarrow \infty} X_n = X$ , where the limiting RV  $X \sim \text{Point Mass}(0)$ ?

The answer is **no**. This is because  $\mathbf{P}(X_n = X) = 0$  for any  $n$ , since  $X \sim \text{Point Mass}(0)$  is a discrete RV with exactly one outcome 0 and  $X_n \sim \text{Normal}(0, 1/n)$  is a continuous RV for every  $n$ , however large. In other words, a continuous RV, such as  $X_n$ , has 0 probability of realizing any single real number in its support, such as 0.

Figure 3.11: Distribution functions of several  $\text{Normal}(\mu, \sigma^2)$  RVs for  $\sigma^2 = 1, \frac{1}{10}, \frac{1}{100}, \frac{1}{1000}$ .



Thus, we need more sophisticated notions of convergence for sequences of RVs. Two such notions are formalized next as they are minimal prerequisites for a clear understanding of three basic propositions in Statistics :

1. Weak Law of Large Numbers,
2. Central Limit Theorem,
3. Gilvenko-Cantelli Theorem.

**Definition 39 (Convergence in Distribution)** Let  $X_1, X_2, \dots$ , be a sequence of RVs and let  $X$  be another RV. Let  $F_n$  denote the DF of  $X_n$  and  $F$  denote the DF of  $X$ . Then we say that  $X_n$  converges to  $X$  in distribution, and write:

$$X_n \rightsquigarrow X$$

if for any real number  $t$  at which  $F$  is continuous,

$$\lim_{n \rightarrow \infty} F_n(t) = F(t) \quad [\text{in the sense of Definition 4}].$$

The above limit, by (3.2) in our Definition 17 of a DF, can be equivalently expressed as follows:

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbf{P}(\{\omega : X_n(\omega) \leq t\}) &= \mathbf{P}(\{\omega : X(\omega) \leq t\}), \\ \text{i.e. } \mathbf{P}(\{\omega : X_n(\omega) \leq t\}) &\rightarrow \mathbf{P}(\{\omega : X(\omega) \leq t\}), \quad \text{as } n \rightarrow \infty. \end{aligned}$$

**Definition 40 (Convergence in Probability)** Let  $X_1, X_2, \dots$ , be a sequence of RVs and let  $X$  be another RV. Let  $F_n$  denote the DF of  $X_n$  and  $F$  denote the DF of  $X$ . Then we say that  $X_n$  converges to  $X$  in probability, and write:

$$X_n \xrightarrow{P} X$$

if for every real number  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} \mathbf{P}(|X_n - X| > \epsilon) = 0 \quad [\text{in the sense of Definition 4}].$$

Once again, the above limit, by (3.1) in our Definition 16 of a RV, can be equivalently expressed as follows:

$$\lim_{n \rightarrow \infty} \mathbf{P}(\{\omega : |X_n(\omega) - X(\omega)| > \epsilon\}) = 0, \quad \text{ie,} \quad \mathbf{P}(\{\omega : |X_n(\omega) - X(\omega)| > \epsilon\}) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Let us revisit the problem of convergence in Classwork 64 armed with our new notions of convergence.

**Example 65 (Convergence in distribution)** Suppose you are given an independent sequence of RVs  $\{X_i\}_{i=1}^n$ , where  $X_i \sim \text{Normal}(0, 1/i)$  with DF  $F_n$  and let  $X \sim \text{Point Mass}(0)$  with DF  $F$ . We can formalize our observation in Classwork 64 that  $X_n$  is concentrating about 0 as  $n \rightarrow \infty$  by the statement:

$$X_n \text{ is converging in distribution to } X, \text{ ie,} \quad X_n \rightsquigarrow X.$$

**Proof:** To check that the above statement is true we need to verify that the definition of convergence in distribution is satisfied for our sequence of RVs  $X_1, X_2, \dots$  and the limiting RV  $X$ . Thus, we need to verify that for any continuity point  $t$  of the Point Mass(0) DF  $F$ ,  $\lim_{n \rightarrow \infty} F_n(t) = F(t)$ . First note that

$$X_n \sim \text{Normal}(0, 1/n) \implies Z := \sqrt{n}X_n \sim \text{Normal}(0, 1),$$

and thus

$$F_n(t) = \mathbf{P}(X_n < t) = \mathbf{P}(\sqrt{n}X_n < \sqrt{nt}) = \mathbf{P}(Z < \sqrt{nt}).$$

The only discontinuous point of  $F$  is 0 where  $F$  jumps from 0 to 1.

When  $t < 0$ ,  $F(t)$ , being the constant 0 function over the interval  $(-\infty, 0)$ , is continuous at  $t$ . Since  $\sqrt{nt} \rightarrow -\infty$ , as  $n \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} F_n(t) = \lim_{n \rightarrow \infty} \mathbf{P}(Z < \sqrt{nt}) = 0 = F(t).$$

And, when  $t > 0$ ,  $F(t)$ , being the constant 1 function over the interval  $(0, \infty)$ , is again continuous at  $t$ . Since  $\sqrt{nt} \rightarrow \infty$ , as  $n \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} F_n(t) = \lim_{n \rightarrow \infty} \mathbf{P}(Z < \sqrt{nt}) = 1 = F(t).$$

Thus, we have proved that  $X_n \rightsquigarrow X$  by verifying that for any  $t$  at which the Point Mass(0) DF  $F$  is continuous, we also have the desired equality:  $\lim_{n \rightarrow \infty} F_n(t) = F(t)$ .

However, note that

$$F_n(0) = \frac{1}{2} \neq F(0) = 1,$$

and so convergence fails at 0, i.e.  $\lim_{n \rightarrow \infty} F_n(t) \neq F(t)$  at  $t = 0$ . But,  $t = 0$  is not a continuity point of  $F$  and the definition of convergence in distribution only requires the convergence to hold at continuity points of  $F$ .

For the same sequence of RVs in Classwork 64 and Example 65 we are tempted to ask whether  $X_n \sim \text{Normal}(0, 1/n)$  converges in probability to  $X \sim \text{Point Mass}(0)$ , i.e. whether  $X_n \xrightarrow{P} X$ . We need some elementary inequalities in Probability to help us answer this question. We visit these inequalities next.

**Proposition 41 (Markov's Inequality)** Let  $(\Omega, \mathcal{F}, P)$  be a probability triple and let  $X = X(\omega)$  be a non-negative RV. Then,

$$\mathbf{P}(X \geq \epsilon) \leq \frac{\mathbf{E}(X)}{\epsilon}, \quad \text{for any } \epsilon > 0. \quad (3.38)$$

**Proof:**

$$\begin{aligned} X &= X\mathbf{1}_{\{y:y \geq \epsilon\}}(x) + X\mathbf{1}_{\{y:y < \epsilon\}}(x) \\ &\geq X\mathbf{1}_{\{y:y \geq \epsilon\}}(x) \\ &\geq \epsilon\mathbf{1}_{\{y:y \geq \epsilon\}}(x) \end{aligned} \quad (3.39)$$

Finally, taking expectations on both sides of the above inequality and then using the fact that the expectation of an indicator function of an event is simply the probability of that event (3.28), we get the desired result:

$$\mathbf{E}(X) \geq \epsilon\mathbf{E}(\mathbf{1}_{\{y:y \geq \epsilon\}}(x)) = \epsilon\mathbf{P}(X \geq \epsilon).$$

Let us look at some immediate consequences of Markov's inequality.

**Proposition 42 (Chebychev's Inequality)** For any RV  $X$  and any  $\epsilon > 0$ ,

$$\mathbf{P}(|X| > \epsilon) \leq \frac{\mathbf{E}(|X|)}{\epsilon} \quad (3.40)$$

$$\mathbf{P}(|X| > \epsilon) = \mathbf{P}(X^2 \geq \epsilon^2) \leq \frac{\mathbf{E}(X^2)}{\epsilon^2} \quad (3.41)$$

$$\mathbf{P}(|X - \mathbf{E}(X)| \geq \epsilon) = \mathbf{P}((X - \mathbf{E}(X))^2 \geq \epsilon^2) \leq \frac{\mathbf{E}(X - \mathbf{E}(X))^2}{\epsilon^2} = \frac{\mathbf{V}(X)}{\epsilon^2} \quad (3.42)$$

**Proof:** All three forms of Chebychev's inequality are mere corollaries (careful reapplications) of Markov's inequality.

Armed with Markov's inequality we next enquire the convergence in probability for the sequence of RVs in Classwork 64 and Example 65.

**Example 66 (Convergence in probability)** Does the the sequence of RVs  $\{X_n\}_{n=1}^{\infty}$ , where  $X_n \sim \text{Normal}(0, 1/n)$ , converge in probability to  $X \sim \text{Point Mass}(0)$ , i.e. does  $X_n \xrightarrow{P} X$ ?

To find out if  $X_n \xrightarrow{P} X$ , we need to show that for any  $\epsilon > 0$ ,  $\lim_{n \rightarrow \infty} \mathbf{P}(|X_n - X| > \epsilon) = 0$ .

Let  $\epsilon$  be any real number greater than 0, then

$$\begin{aligned} \mathbf{P}(|X_n| > \epsilon) &= \mathbf{P}(|X_n|^2 > \epsilon^2) \\ &= \frac{\mathbf{E}(X_n^2)}{\epsilon^2} \quad [\text{by Markov's Inequality (3.38)}] \\ &= \frac{\frac{1}{n}}{\epsilon^2} \rightarrow 0, \quad \text{as } n \rightarrow \infty \quad [\text{in the sense of Definition 4}]. \end{aligned}$$

Hence, we have shown that for any  $\epsilon > 0$ ,  $\lim_{n \rightarrow \infty} \mathbf{P}(|X_n - X| > \epsilon) = 0$  and therefore by Definition 40,  $X_n \xrightarrow{P} X$  or  $X_n \xrightarrow{P} 0$ .

**Convention:** When  $X$  has a Point Mass( $\theta$ ) distribution and  $X_n \xrightarrow{P} X$ , we simply write  $X_n \xrightarrow{P} \theta$ .

Now that we have been introduced to two notions of convergence for sequences of RVs we can begin to appreciate the construction of limiting random variables. In several situations it becomes cumbersome to model the events using the Binomial( $n, \theta$ ) RV, especially when the parameter

$\theta \propto 1/n$  and the events become rare. However, for some real parameter  $\lambda > 0$ , the Binomial( $n, \lambda/n$ ) RV with probability of the number of successes in  $n$  trials, with per-trial success probability  $\lambda/n$ , approaches the Poisson distribution with expectation  $\lambda$ , as  $n$  approaches  $\infty$  (actually, it converges in distribution as defined later). The Poisson( $\lambda$ ) RV is much simpler to work with than the combinatorially laden Binomial( $n, \theta = \lambda/n$ ) RV. We sketch the details of this next.

Let  $X \sim \text{Binomial}(n, \theta = \lambda/n)$ , then for any  $x \in \{0, 1, 2, 3, \dots, n\}$ ,

$$\begin{aligned} \mathbf{P}(X = x) &= \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x} \\ &= \frac{n(n-1)(n-2)\cdots(n-x+1)}{x(x-1)(x-2)\cdots(2)(1)} \left(\frac{\lambda^x}{n^x}\right) \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-x} \\ &= \underbrace{\left(\frac{n}{n}\right) \left(\frac{n-1}{n}\right) \left(\frac{n-2}{n}\right) \cdots \left(\frac{n-x+1}{n}\right)}_{\text{underbrace}} \underbrace{\left(\frac{\lambda^x}{x!}\right)}_{\text{underbrace}} \underbrace{\left(1 - \frac{\lambda}{n}\right)^n}_{\text{underbrace}} \underbrace{\left(1 - \frac{\lambda}{n}\right)^{-x}}_{\text{underbrace}} \end{aligned} \quad (3.43)$$

As  $n \rightarrow \infty$ , the expression below the first overbrace  $\rightarrow 1$ , while that below the second overbrace, being independent of  $n$  remains the same. By the elementary examples of limits 14 and 15, as  $n \rightarrow \infty$ , the expression over the first underbrace approaches  $e^{-\lambda}$  while that over the second underbrace approaches 1. Finally, we get the desired limit:

$$\lim_{n \rightarrow \infty} \mathbf{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!} .$$

**Model 8 (Poisson( $\lambda$ ) RV)** Given a real parameter  $\lambda > 0$ , the discrete RV  $X$  is said to be Poisson( $\lambda$ ) distributed if  $X$  has PDF:

$$f(x; \lambda) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!} & \text{if } x \in \mathbb{Z}_+ := \{0, 1, 2, \dots\} , \\ 0 & \text{otherwise .} \end{cases} \quad (3.44)$$

Note that the PDF integrates to 1:

$$\sum_{x=0}^{\infty} f(x; \lambda) = \sum_{x=0}^{\infty} \frac{e^{-\lambda} \lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^{-\lambda} e^{\lambda} = 1 ,$$

where we exploit the Taylor series of  $e^\lambda$  to obtain the second-last equality above.

**Mean and variance of Poisson( $\lambda$ ) RV:** Let  $X \sim \text{Poisson}(\lambda)$ . Then:

$$\mathbf{E}(X) = \sum_{x=0}^{\infty} x f(x; \lambda) = \sum_{x=0}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} x \frac{\lambda^x}{x!} = e^{-\lambda} \sum_{x-1=0}^{\infty} \frac{\lambda \lambda^{x-1}}{(x-1)!} = e^{-\lambda} \lambda e^{\lambda} = \lambda .$$

Similarly,


$$\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = \lambda + \lambda^2 - \lambda^2 = \lambda .$$

since

$$\begin{aligned}
 \mathbf{E}(X^2) &= \sum_{x=0}^{\infty} x^2 \frac{e^{-\lambda} \lambda^x}{x!} = \lambda e^{-\lambda} \sum_{x=1}^{\infty} \frac{x \lambda^{x-1}}{(x-1)!} = \lambda e^{-\lambda} \left( 1 + \frac{2\lambda}{1} + \frac{3\lambda^2}{2!} + \frac{4\lambda^3}{3!} + \dots \right) \\
 &= \lambda e^{-\lambda} \left( \left( 1 + \frac{\lambda}{1} + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \dots \right) + \left[ \frac{\lambda}{1} + \frac{2\lambda^2}{2!} + \frac{3\lambda^3}{3!} + \dots \right] \right) \\
 &= \lambda e^{-\lambda} \left( (e^\lambda) + \lambda \left( 1 + \frac{2\lambda}{2!} + \frac{3\lambda^2}{3!} + \dots \right) \right) = \lambda e^{-\lambda} \left( e^\lambda + \lambda \left( 1 + \lambda + \frac{\lambda^2}{2!} + \dots \right) \right) \\
 &= \lambda e^{-\lambda} (e^\lambda + \lambda e^\lambda) = \lambda e^{-\lambda} (e^\lambda + \lambda e^\lambda) = \lambda(1 + \lambda) = \lambda + \lambda^2
 \end{aligned}$$

Note that Poisson( $\lambda$ ) distribution is one whose mean and variance are the same, namely  $\lambda$ .

Figure 3.12: PDF of  $X \sim \text{Poisson}(\lambda = 10)$  and the relative frequency histogram based on 1000 samples from  $X$  according to Simulation 117.



The Poisson( $\lambda$ ) RV  $X$  is also related to the IID Exponential( $\lambda$ ) RV  $Y_1, Y_2, \dots$ :  $X$  is the number of occurrences, per unit time, of an instantaneous event whose inter-occurrence time is the IID Exponential( $\lambda$ ) RV. For example, the number of buses arriving at our bus-stop in the next minute, with exponentially distributed inter-arrival times, has a Poisson distribution.

### 3.9 Characteristic Functions

The characteristic function (CF) of a random variable gives another way to specify its distribution. Thus CF is a powerful tool for analytical results involving random variables (more).

**Definition 43 (Characteristic Function (CF))** Let  $X$  be a RV and  $i = \sqrt{-1}$ . The function  $\varphi_X(t) : \mathbb{R} \rightarrow \mathbb{C}$  defined by

$$\varphi_X(t) := E(\exp(itX)) = \begin{cases} \sum_x \exp(itx) f_X(x) & \text{if } X \text{ is discrete RV} \\ \int_{-\infty}^{\infty} \exp(itx) f_X(x) dx & \text{if } X \text{ is continuous RV} \end{cases} \quad (3.45)$$

is called the **characteristic function** of  $X$ .

NOTE:  $\varphi_X(t)$  exists for any  $t \in \mathbb{R}$ , because

$$\begin{aligned}\varphi_X(t) &= E(\exp(\imath t X)) \\ &= E(\cos(tX) + \imath \sin(tX)) \\ &= E(\cos(tX)) + \imath E(\sin(tX))\end{aligned}$$

and the last two expected values are well-defined, because the sine and cosine functions are bounded by  $[-1, 1]$ .

For a continuous RV,  $\int_{-\infty}^{\infty} \exp(-\imath tx) f_X(x) dx$  is called the *Fourier transform* of  $f_X$ . This is the CF but with  $t$  replaced by  $-t$ . You will also encounter Fourier transforms when solving differential equations.

### 3.9.1 Obtaining Moments from Characteristic Function

Recall that the  $k$ -th moment of  $X$  is  $E(X^k)$  for any  $k \in \mathbb{N} := \{1, 2, 3, \dots\}$  is

$$E(X^k) = \begin{cases} \sum_x x^k f_X(x) & \text{if } X \text{ is a discrete RV} \\ \int_{-\infty}^{\infty} x^k f_X(x) dx & \text{if } X \text{ is a continuous RV} \end{cases}$$

The characteristic function can be used to derive the moments of  $X$  due to the following nice relationship between the the  $k$ -th moment of  $X$  and the  $k$ -th derivative of the CF of  $X$ .

**Proposition 44 (Moment & CF.)** Let  $X$  be a random variable and  $\varphi_X(t)$  be its CF. If  $E(X^k)$  exists and is finite, then  $\varphi_X(t)$  is  $k$  times continuously differentiable and

$$E(X^k) = \frac{1}{\imath^k} \left[ \frac{d^k \varphi_X(t)}{dt^k} \right]_{t=0} .$$

where  $\left[ \frac{d^k \varphi_X(t)}{dt^k} \right]_{t=0}$  is the  $k$ -th derivative of  $\varphi_X(t)$  with respect to  $t$ , evaluated at the point  $t = 0$ .

**Proof:** The proper proof is very messy so we just give a sketch of the ideas in the proof. Due to the linearity of the expectation (integral) and the derivative operators, we can change the order of operations:

$$\frac{d^k \varphi_X(t)}{dt^k} = \frac{d^k}{dt^k} E(\exp(\imath t X)) = E \left( \frac{d^k}{dt^k} \exp(\imath t X) \right) = E \left( (\imath X)^k \exp(\imath t X) \right) = \imath^k E \left( X^k \exp(\imath t X) \right)$$

The RHS evaluated at  $t = 0$  is

$$\left[ \frac{d^k \varphi_X(t)}{dt^k} \right]_{t=0} = \left[ \imath^k E \left( X^k \exp(\imath t X) \right) \right]_{t=0} = \imath^k E \left( X^k \right)$$

This completes the sketch of the proof.

The above Theorem gives us the relationship between the moments and the derivatives of the CF if we already know that the moment exists. When one wants to compute a moment of a random variable, what we need is the following Theorem.

**Proposition 45 (Moments from CF.)** Let  $X$  be a random variable and  $\varphi_X(t)$  be its CF. If  $\varphi_X(t)$  is  $k$  times differentiable at the point  $t = 0$ , then

1. if  $k$  is even, the  $n$ -th moment of  $X$  exists and is finite for any  $0 \leq n \leq k$ ;
2. if  $k$  is odd, the  $n$ -th moment of  $X$  exists and is finite for any  $0 \leq n \leq k - 1$ .

In both cases,

$$E(X^k) = \frac{1}{i^k} \left[ \frac{d^k \varphi_X(t)}{dt^k} \right]_{t=0}. \quad (3.46)$$

where  $\left[ \frac{d^k \varphi_X(t)}{dt^k} \right]_{t=0}$  is the  $k$ -th derivative of  $\varphi_X(t)$  with respect to  $t$ , evaluated at the point  $t = 0$ .

**Proof:** For proof see e.g., Ushakov, N. G. (1999) Selected topics in characteristic functions, VSP (p. 39).

**Example 67** Let  $X$  be the Bernoulli( $\theta$ ) RV. Find the CF of  $X$ . Then use CF to find  $E(X)$ ,  $E(X^2)$  and from this obtain the variance  $V(X) = E(X^2) - (E(X))^2$ .

Solution:

### Part 1

Recall the PMF for this discrete RV with parameter  $\theta \in (0, 1)$  is

$$f_X(x; \theta) = \begin{cases} \theta & \text{if } x = 1 \\ 1 - \theta & \text{if } x = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Let's first find the CF of  $X$

$$\begin{aligned} \varphi_X(t) &= E(\exp(itX)) = \sum_x \exp(itx)f_X(x; \theta) \quad \text{By Defn. in Equation (3.45)} \\ &= \exp(it \times 0)(1 - \theta) + \exp(it \times 1)\theta = \exp(0)(1 - \theta) + \exp(it)\theta = 1 - \theta + \theta \exp(it) \end{aligned}$$

### Part 2:

Let's differentiate CF

$$\frac{d}{dt} \varphi_X(t) = \frac{d}{dt} (1 - \theta + \theta e^{it}) = \theta i \exp(it)$$

We get  $E(X)$  by evaluating  $\frac{d}{dt} \varphi_X(t)$  at  $t = 0$  and dividing by  $i$  according to Equation (3.46) as follows:

$$E(X) = \frac{1}{i} \left[ \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{i} [\theta i \exp(it)]_{t=0} = \frac{1}{i} (\theta i \exp(i0)) = \theta .$$

Similarly from Equation (3.46) we can get  $E(X^2)$  as follows:

$$\begin{aligned} E(X^2) &= \frac{1}{\iota^2} \left[ \frac{d^2}{dt^2} \varphi_X(t) \right]_{t=0} = \frac{1}{\iota^2} \left[ \frac{d}{dt} \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{\iota^2} \left[ \frac{d}{dt} \theta \exp(\iota t) \right]_{t=0} \\ &= \frac{1}{\iota^2} [\theta \iota^2 \exp(\iota t)]_{t=0} = \frac{1}{\iota^2} (\theta \iota^2 \exp(\iota 0)) = \theta . \end{aligned}$$

Finally, from the first and second moments we can get the variance as follows:

$$V(X) = E(X^2) - (E(X))^2 = \theta - \theta^2 = \theta(1 - \theta) .$$

Let's check that this is what we have as variance for the Bernoulli( $\theta$ ) RV if we directly computed it using weighted sums in the definition of expectations:  $E(X) = 1 \times \theta + 0 \times (1 - \theta) = \theta$ ,  $E(X^2) = 1^2 \times \theta + 0^2 \times (1 - \theta) = \theta$  and thus giving the same  $V(X) = E(X^2) - (E(X))^2 = \theta - \theta^2 = \theta(1 - \theta)$ .

**Example 68** Let  $X$  be an Exponential( $\lambda$ ) RV. First show that its CF is  $\lambda/(\lambda - \iota t)$ . Then use CF to find  $E(X)$ ,  $E(X^2)$  and from this obtain the variance  $V(X) = E(X^2) - (E(X))^2$ .

Solution:

Recall that the PDF of an Exponential( $\lambda$ ) RV for a given parameter  $\lambda \in (0, \infty)$  is  $\lambda e^{-\lambda x}$  if  $x \in [0, \infty)$  and 0 if  $x \notin [0, \infty)$ .

**Part 1:** Find the CF.

We will use the fact that

$$\int_0^\infty \alpha e^{-\alpha x} dx = [-e^{-\alpha x}]_0^\infty = 1$$

$$\begin{aligned} \varphi_X(t) &= E(\exp(\iota X t)) = E(e^{\iota X t}) = \int_{-\infty}^\infty e^{\iota x t} \lambda e^{-\lambda x} dx = \lambda \int_0^\infty e^{-(\lambda - \iota t)x} dx \\ &= \frac{\lambda}{\lambda - \iota t} \int_0^\infty (\lambda - \iota t) e^{-(\lambda - \iota t)x} dx = \frac{\lambda}{\lambda - \iota t} \int_0^\infty \alpha e^{-\alpha x} dx = \frac{\lambda}{\lambda - \iota t} , \end{aligned}$$

where  $\alpha = \lambda - \iota t$  with  $\lambda > 0$ .

Alternatively, you can use  $e^{\iota t x} = \cos(tx) + \iota \sin(tx)$  and do integration by parts to arrive at the same answer starting from:

$$\varphi_X(t) = \int_{-\infty}^\infty e^{\iota x t} \lambda e^{-\lambda x} dx = \int_{-\infty}^\infty \cos(tx) e^{-\lambda x} dx + \iota \int_{-\infty}^\infty \sin(tx) e^{-\lambda x} dx = \frac{\lambda}{\lambda - \iota t} .$$

**Part 2:**

Let us differentiate the CF to get moments using Equation (3.46) (CF has to be once and twice differentiable at  $t = 0$  to get the first and second moments).

$$\begin{aligned} \frac{d}{dt} \varphi_X(t) &= \frac{d}{dt} \left( \frac{\lambda}{\lambda - \iota t} \right) = \lambda \left( -1 \times (\lambda - \iota t)^{-2} \times \frac{d}{dt} (\lambda - \iota t) \right) \\ &= \lambda \left( \frac{-1}{(\lambda - \iota t)^2} \times (-\iota) \right) = \frac{\lambda \iota}{(\lambda - \iota t)^2} \end{aligned}$$

We get  $E(X)$  by evaluating  $\frac{d}{dt} \varphi_X(t)$  at  $t = 0$  and dividing by  $\iota$  according to Equation (3.46) as follows:

$$E(X) = \frac{1}{\iota} \left[ \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{\iota} \left[ \frac{\lambda \iota}{(\lambda - \iota t)^2} \right]_{t=0} = \frac{1}{\iota} \left( \frac{\lambda \iota}{\lambda^2} \right) = \frac{1}{\iota} \left( \frac{\iota}{\lambda} \right) = \frac{1}{\lambda}$$

Let's pause and see if this makes sense.... Yes, because the expected value of  $\text{Exponential}(\lambda)$  RV is indeed  $1/\lambda$  (recall from when we introduced this RV).

Similarly from Equation (3.46) we can get  $E(X^2)$  as follows:

$$\begin{aligned} E(X^2) &= \frac{1}{\iota^2} \left[ \frac{d^2}{dt^2} \varphi_X(t) \right]_{t=0} = \frac{1}{\iota^2} \left[ \frac{d}{dt} \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{\iota^2} \left[ \frac{d}{dt} \frac{\lambda \iota}{(\lambda - \iota t)^2} \right]_{t=0} \\ &= \frac{1}{\iota^2} \left[ \lambda \iota \times \frac{d}{dt} (\lambda - \iota t)^{-2} \right]_{t=0} = \frac{1}{\iota^2} \left[ \lambda \iota \left( -2(\lambda - \iota t)^{-3} \frac{d}{dt} (\lambda - \iota t) \right) \right]_{t=0} \\ &= \frac{1}{\iota^2} [\lambda \iota (-2(\lambda - \iota t)^{-3} \times (-\iota))]_{t=0} = \frac{1}{\iota^2} \left[ \frac{2\lambda \iota^2}{(\lambda - \iota t)^3} \right]_{t=0} = \frac{1}{\iota^2} \left( \frac{2\lambda \iota^2}{\lambda^3} \right) = \frac{2}{\lambda^2}. \end{aligned}$$

Finally, from the first and second moments we can get the variance as follows:

$$V(X) = E(X^2) - (E(X))^2 = \frac{2}{\lambda^2} - \left( \frac{1}{\lambda} \right)^2 = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{2-1}{\lambda^2} = \frac{1}{\lambda^2}.$$

Let's check that this is what we had as variance for the  $\text{Exponential}(\lambda)$  RV when we first introduced it and directly computed using integrals for definition of expectation.

Characteristic functions can be used to characterize the distribution of a random variable.

Two RVs  $X$  and  $Y$  have the same DFs, i.e.,  $F_X(x) = F_Y(x)$  for all  $x \in \mathbb{R}$ , if and only if they have the same characteristic functions, i.e.  $\varphi_X(t) = \varphi_Y(t)$  for all  $t \in \mathbb{R}$  (for proof see Resnick, S. I. (1999) A Probability Path, Birkhauser).

Thus, if we can show that two RVs have the same CF then we know they are the same. This can be much more challenging or impossible to do directly with their DFs.

Let  $Z$  be  $\text{Normal}(0, 1)$ , the standard normal RV. We can find the CF for  $Z$  using couple of tricks as follows

$$\begin{aligned} \varphi_Z(t) &= E(e^{\iota t Z}) \\ &= \int_{-\infty}^{\infty} e^{\iota t z} f_Z(z) dz = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\iota t z} e^{-z^2/2} dz = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\iota t z - z^2/2} dz \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-(t^2 + (z - \iota t)^2)/2} dz = e^{-t^2/2} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-(z - \iota t)^2/2} dz \\ &= e^{-t^2/2} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-y^2/2} dy \quad \text{substituting } y = z - \iota t, dy = dz \\ &= e^{-t^2/2} \frac{1}{\sqrt{2\pi}} \sqrt{2\pi} \quad \text{using the normalizing constant in PDF of } \text{Normal}(0, 1) \text{ RV} \\ &= e^{-t^2/2} \end{aligned}$$

Thus the CF of the standard normal RV  $Z$  is

$$\boxed{\varphi_Z(t) = e^{-t^2/2}} \tag{3.47}$$

Let  $X$  be a RV with CF  $\varphi_X(t)$ . Let  $Y$  be a linear transformation of  $X$

$$Y = a + bX$$

where  $a$  and  $b$  are two constant real numbers and  $b \neq 0$ . Then the CF of  $Y$  is

$$\boxed{\varphi_Y(t) = \exp(\iota at)\varphi_X(bt)} \tag{3.48}$$

**Proof:** This is easy to prove using the definition of CF as follows:

$$\begin{aligned}\varphi_Y(t) &= E(\exp(itY)) = E(\exp(it(a+bX))) = E(\exp(ita+itbX)) \\ &= E(\exp(ita)\exp(itbX)) = \exp(ita)E(\exp(itbX)) = \exp(ita)\varphi_X(bt)\end{aligned}$$

**Example 69** Let  $Y$  be a  $\text{Normal}(\mu, \sigma^2)$  RV. Recall that  $Y$  is a linear transformation of  $Z$ , i.e.,  $Y = \mu + \sigma Z$  where  $Z$  is a  $\text{Normal}(0, 1)$  RV. Using Equations (3.47) and (3.48) find the CF of  $Y$ .

Solution:

$$\begin{aligned}\varphi_Y(t) &= \exp(i\mu t)\varphi_Z(\sigma t), \quad \text{since } Y = \mu + \sigma Z \\ &= e^{i\mu t}e^{(-\sigma^2 t^2)/2}, \quad \text{since } \varphi_Z(t) = e^{-t^2/2} \\ &= e^{i\mu t - (\sigma^2 t^2)/2}\end{aligned}$$

A generalization of (3.48) is the following. If  $X_1, X_2, \dots, X_n$  are independent RVs and  $a_1, a_2, \dots, a_n$  are some constants, then the CF of the linear combination  $Y = \sum_{i=1}^n a_i X_i$  is

$$\varphi_Y(t) = \varphi_{X_1}(a_1 t) \times \varphi_{X_2}(a_2 t) \times \cdots \times \varphi_{X_n}(a_n t) = \prod_{i=1}^n \varphi_{X_i}(a_i t) . \quad (3.49)$$

**Example 70** Using the following three facts:

- Eqn. (3.49)
- the Binomial( $n, \theta$ ) RV  $Y$  is the sum of  $n$  independent Bernoulli( $\theta$ ) RVs (from Probability Course)
- the CF of Bernoulli( $\theta$ ) RV (from lecture notes for Inference Course)

find the CF of the Binomial( $n, \theta$ ) RV  $Y$ .

Solution:

Let  $X_1, X_2, \dots, X_n$  be independent Bernoulli( $\theta$ ) RVs with CF  $(1 - \theta + \theta e^{it})$  then  $Y = \sum_{i=1}^n X_i$  is the Binomial( $n, \theta$ ) RV and by Eqn. (3.49) with  $a_1 = a_2 = \cdots = 1$ , we get

$$\varphi_Y(t) = \varphi_{X_1}(t) \times \varphi_{X_2}(t) \cdots \varphi_{X_n}(t) = \prod_{i=1}^n \varphi_{X_i}(t) = \prod_{i=1}^n (1 - \theta + \theta e^{it}) = (1 - \theta + \theta e^{it})^n .$$

**Example 71** Let  $Z_1$  and  $Z_2$  be independent  $\text{Normal}(0, 1)$  RVs.

1. Use Eqn. (3.49) to find the CF of  $Z_1 + Z_2$ .
2. From the CF of  $Z_1 + Z_2$  identify what RV it is.
3. Use Eqn. (3.49) to find the CF of  $2Z_1$ .
4. From the CF of  $2Z_1$  identify what RV it is.

5. Try to understand the difference between the distributions of  $Z_1 + Z_2$  and  $2Z_1$  inspite of  $Z_1$  and  $Z_2$  having the same distribution.

Hint: from lectures we know that  $\varphi_X(t) = e^{\mu t - (\sigma^2 t^2)/2}$  for a  $\text{Normal}(\mu, \sigma^2)$  RV  $X$ .

Solution:

1. By Eqn. (3.49) we just multiply the characteristic functions of  $Z_1$  and  $Z_2$ , both of which are  $e^{-t^2/2}$ ,

$$\varphi_{Z_1+Z_2}(t) = \varphi_{Z_1}(t) \times \varphi_{Z_2}(t) = e^{-t^2/2} \times e^{-t^2/2} = e^{-2t^2/2} = e^{-t^2} .$$

2. The CF of  $Z_1 + Z_2$  is that of the  $\text{Normal}(\mu, \sigma^2)$  RV with  $\mu = 0$  and  $\sigma^2 = 2$ . Thus  $Z_1 + Z_2$  is the  $\text{Normal}(0, 2)$  RV with mean parameter  $\mu = 0$  and variance parameter  $\sigma^2 = 2$ .

3. We can again use Eqn. (3.49) to find the CF of  $2Z_1$  as follows

$$\varphi_{2Z_1} = \varphi_{Z_1}(2t) = e^{-2^2 t^2/2} .$$

4. The CF of  $2Z_1$  is that of the  $\text{Normal}(\mu, \sigma^2)$  RV with  $\mu = 0$  and  $\sigma^2 = 2^2 = 4$ . Thus  $2Z_1$  is the  $\text{Normal}(0, 4)$  RV with mean parameter  $\mu = 0$  and variance parameter  $\sigma^2 = 4$ .

5.  $2Z_1$  has a bigger variance from multiplying the standard normal RV by 2 while  $Z_1 + Z_2$  has a smaller variance from adding two independent standard normal RVs. Thus, the result of adding the same RV twice does not have the same distribution as that of multiplying it by 2. In other words  $2 \times Z$  is not equal to  $Z + Z$  in terms of its probability distribution!

## Exercises

**Ex. 3.1** — One number in the following table for the probability function of a random variable  $X$  is incorrect. Which is it, and what should the correct value be?

$x$	1	2	3	4	5
$\mathbf{P}(X = x)$	0.07	0.10	1.10	0.32	0.40

**Ex. 3.2** — Let  $X$  be the number of years before a particular type of machine will need replacement. Assume that  $X$  has the probability function  $f(1) = 0.1$ ,  $f(2) = 0.2$ ,  $f(3) = 0.2$ ,  $f(4) = 0.2$ ,  $f(5) = 0.3$ .

1. Find the distribution function,  $F$ , for  $X$ , and graph both  $f$  and  $F$ .
2. Find the probability that the machine needs to be replaced during the first 3 years.
3. Find the probability that the machine needs no replacement during the first 3 years.

**Ex. 3.3** — Of 200 adults, 176 own one TV set, 22 own two TV sets, and 2 own three TV sets. A person is chosen at random. What is the probability mass function of  $X$ , the number of TV sets owned by that person?

**Ex. 3.4** — Suppose a discrete random variable  $X$  has probability function give by

$x$	3	4	5	6	7	8	9	10	11	12	13
$\mathbf{P}(X = x)$	0.07	0.01	0.09	0.01	0.16	0.25	0.20	0.03	0.02	0.11	0.05

- (a) Construct a row of cumulative probabilities for this table, that is, find the distribution function of  $X$ .  
(b) Find the following probabilities.

(i) $\mathbf{P}(X \leq 5)$	(iii) $\mathbf{P}(X > 9)$	(v) $\mathbf{P}(4 < X \leq 9)$
(ii) $\mathbf{P}(X < 12)$	(iv) $\mathbf{P}(X \geq 9)$	(vi) $\mathbf{P}(4 < X < 11)$

**Ex. 3.5** — A box contains 4 right-handed and 6 left-handed screws. Two screws are drawn at random without replacement. Let  $X$  be the number of left-handed screws drawn. Find the probability mass function for  $X$ , and then calculate the following probabilities:

1.  $\mathbf{P}(X \leq 1)$
2.  $\mathbf{P}(X \geq 1)$
3.  $\mathbf{P}(X > 1)$

**Ex. 3.6** — Let  $X$  be the outcome of a fair die roll with probability mass function given by

$$f_X(x) = \begin{cases} \frac{1}{6} & \text{if } x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise.} \end{cases}$$

If  $Y = (X - 3)^2$  then find the probability mass function of  $Y$ ,  $f_Y(y)$ .

**Ex. 3.7** — Given a natural number  $n$  as a parameter, i.e., given a parameter  $n \in \{1, 2, 3, \dots\}$ , let  $X$  be a discrete uniform random variable on the finite set

$$\mathbb{X} = \{-n, -n + 1, \dots, -1, 0, 1, \dots, n - 1, n\}$$

i.e. the probability mass function of  $X$  is:

$$f_X(x; n) = \begin{cases} \frac{1}{2^{n+1}} & \text{if } x \in \mathbb{X} \\ 0 & \text{otherwise.} \end{cases}$$

Find the probability mass function  $f_Y(y; n)$  for  $Y = |X|$ , the absolute value of  $X$ .

**Ex. 3.8** — Suppose that a random variable  $X$  has geometric probability mass function,

$$f(x) = \frac{k}{2^x} \quad (x = 0, 1, 2, \dots).$$

1. Find the value of  $k$ .

2. What is  $\mathbf{P}(X \geq 4)$ ?

**Ex. 3.9** — If  $X$  is a Geometric( $\theta$ ) random variable and  $Y = (\frac{1}{2})^X$  then find an expression for  $f_Y(y)$ .

**Ex. 3.10** — If  $X$  is a continuous random variable with probability density function

$$f_X(x) = \begin{cases} xe^{-x} & x \geq 0 \\ 0 & x < 0 \end{cases},$$

find the probability density function of  $Y = e^X$ .

**Ex. 3.11** — Four fair coins are tossed simultaneously. If we count the number of heads that appear then we have a binomial random variable,  $X = \text{the number of heads}$ .

1. Find the probability mass function of  $X$ .

2. Compute the probabilities of obtaining no heads, precisely 1 head, at least 1 head, not more than 3 heads.

**Ex. 3.12** — The distribution of blood types in a certain population is as follows:

Blood type	Type O	Type A	Type B	Type AB
Proportion	0.45	0.40	0.10	0.05

A random sample of 15 blood donors is observed from this population. Find the probabilities of the following events.

1. Only one type  $AB$  donor is included.

2. At least three of the donors are type  $B$ .

3. More than ten of the donors are either type  $O$  or type  $A$ .

4. Fewer than five of the donors are not type  $A$ .

**Ex. 3.13** — If the probability of hitting a target in a single shot is 10% and 10 shots are fired independently, what is the probability that the target will be hit at least once?

**Ex. 3.14** — Consider the probability density function

$$f(x) = \begin{cases} k & -4 \leq x \leq 4 \\ 0 & \text{otherwise} \end{cases}.$$

1. Find the value of  $k$ .
2. Find the distribution function,  $F$ .
3. Graph  $f$  and  $F$ .

**Ex. 3.15** — Assume that a new light bulb will burn out at time  $t$  hours according to the probability density function given by

$$f(t) = \begin{cases} \lambda e^{-\lambda t} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases}.$$

In this context,  $\lambda$  is often called the failure rate of the bulb.

- (a) Assume that  $\lambda = 0.01$ , and find the probability that the bulb will not burn out before  $\tau$  hours. This  $\tau$ -specific probability is often called the reliability of the bulb.  
Hint: Use the distribution function for an Exponential( $\lambda$ ) random variable (recall,  $F(\tau; \lambda) = \int_{-\infty}^{\tau} f(t)dt$ )!
- (b) For what value of  $\tau$  is the reliability of the bulb exactly  $\frac{1}{2}$ ?

**Ex. 3.16** — Feller discusses the probability and statistics of flying bomb hits in an area of southern London during II world war. The area in question was partitioned into  $24 \times 24 = 576$  small squares. The total number of hits was 537. There were 229 squares with 0 hits, 211 with 1 hit, 93 with 2 hits, 35 with 3 hits, 7 with 4 hits and 1 with 5 or more hits. Assuming the hits were purely random, use the Poisson approximation to find the probability that a particular square would have exactly  $k$  hits. Compute the expected number of squares that would have 0, 1, 2, 3, 4, and 5 or more hits and compare this with the observed results (Snell 9.2.14).

**Ex. 3.17** — Suppose that a certain type of magnetic tape contains, on the average, 2 defects per 100 meters. What is the probability that a roll of tape 300 meters long will contain no defects?

**Ex. 3.18** — In 1910, E. Rutherford and H. Geiger showed experimentally that the number of alpha particles emitted per second in a radioactive process is a random variable  $X$  having a Poisson distribution. If the average number of particles emitted per second is 0.5, what is the probability of observing two or more particles during any given second?

**Ex. 3.19** — The number of lacunae (surface pits) on specimens of steel, polished and examined in a metallurgical laboratory, is known to have Poisson distribution.

1. Write down the formula for the probability that a specimen has  $x$  defects, explaining the meanings of the symbols you use.
2. Simplify the formula in the case  $x = 0$ .
3. In a large homogeneous collection of specimens, 10% have one or more lacunae. Find (approximately) the percentage having exactly two.
4. Why might the Poisson distribution not apply in this situation?

**Ex. 3.20** — If  $X$  is a Poisson( $\lambda$ ) random variable find the probability mass function,  $f_Y(y)$ , of

$$Y = \frac{1}{(X + 1)^2}.$$

**Ex. 3.21** — Find the probability that none of the three bulbs in a traffic signal need to be replaced during the first 1200 hours of operation if the length of time before a single bulb needs to be replaced is a continuous random variable  $X$  with density

$$f(x) = \begin{cases} 6(0.25 - (x - 1.5)^2) & 1 < x < 2 \\ 0 & \text{otherwise} \end{cases}.$$

Note:  $X$  is measured in multiples of 1000 hours.

**Ex. 3.22** — Let the random variable  $X$  be the time after which certain ball bearings wear out, with density

$$f(x) = \begin{cases} ke^{-x} & 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}.$$

Note:  $X$  is measured in years.

1. Find  $k$ .
2. Find the probability that a bearing will last at least 1 year.

**Ex. 3.23** — If  $X$ , the received power at an antenna is an Exponential( $\lambda$ ) random variable then find the probability density function of the amplitude  $Y = \sqrt{X}$ .

**Ex. 3.24** — If  $X$  is a Uniform( $a, b$ ) random variable where  $0 < a < b$ , find the probability density function,  $f_Y(y)$ , of

$$Y = \log_e(X).$$

**Ex. 3.25** — Let  $X$  be the number of air conditioners a store sells each day, and assume that  $X$  has probability mass function  $f(10) = 0.1$ ,  $f(11) = 0.3$ ,  $f(12) = 0.4$ ,  $f(13) = 0.2$ .

1. Find the expected number of conditioners that the store sells each day.
2. If the profit per conditioner is \$55, what is the expected daily profit?

**Ex. 3.26** — A small petrol station is supplied with fuel every Saturday afternoon. Assume that its volume of sales  $X$ , in ten thousands of litres, has density

$$f(x) = \begin{cases} 6x(1-x) & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}.$$

Determine the mean and variance of  $X$ .

**Ex. 3.27** — Starting from the definition of the variance of a random variable (Definition 32) show that

$$\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2.$$

**Ex. 3.28** — Show that  $V(aX + b) = a^2V(X)$  for constants  $a$  and  $b$  and a random variable  $X$ .

**Ex. 3.29** — \*\*Let  $X$  be a discrete random variable with PMF given by

$$f(x) = \begin{cases} \frac{x}{10} & \text{if } x \in \{1, 2, 3, 4\}, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find:

- (i)  $\mathbf{P}(X = 0)$
  - (ii)  $\mathbf{P}(2.5 < X < 5)$
  - (iii)  $\mathbf{E}(X)$
  - (iv)  $\mathbf{V}(X)$
- (b) Write down the DF (or CDF) of  $X$ .
- (c) Plot the PMF and CDF of  $X$ .

**Ex. 3.30** — Find the mean and the variance of the following random variables.

1.  $X$  a discrete uniform random variable on  $\{1, 2, 3, 4, 5, 6\}$ , i.e., *the number a fair die turns up*.
2.  $X$  is a Uniform(0, 8) random variable, i.e., *a continuous uniform random variable from the interval [0, 8]*.
3.  $X$  has a density function

$$f(x) = \begin{cases} 2e^{-2x} & \text{if } x \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

**Ex. 3.31** — Let  $(X, Y)$  be a continuous R $\vec{V}$  with joint probability density function (JPDF)

$$f_{X,Y}(x, y) = \begin{cases} a(x^2 + y) & \text{if } 0 < x < 1 \text{ and } 0 < y < 1 \\ 0 & \text{otherwise.} \end{cases}$$

Find the following:

1. the normalizing constant  $a$  which will ensure  $\mathbf{P}(\Omega) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$
2.  $f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$  called the marginal probability density function (MPDF) of  $X$
3.  $f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx$  called the marginal probability density function (MPDF) of  $Y$
4. Check if  $f_X(x)f_Y(y) = f_{X,Y}(x, y)$  for every  $(x, y)$  and decide whether  $X$  and  $Y$  are independent random variables. Hint:  $X$  and  $Y$  are said to be independent if  $f_X(x)f_Y(y) = f_{X,Y}(x, y)$  for every  $(x, y)$ .
5.  $F_{X,Y}(x, y)$ , the joint cumulative distribution function (JCDF) of  $(X, Y)$  for any  $(x, y) \in (0, 1) \times (0, 1)$
6. the probability that  $X > 0.5$  and  $Y < 0.6$ , i.e.,  $\mathbf{P}(X > 0.5, Y < 0.6)$
7.  $E(X)$ , the expectation of  $X$  or the first moment of  $X$
8.  $E(Y)$ , the expectation of  $Y$  or the first moment of  $Y$
9.  $E(XY)$ , the expectation of  $XY$
10.  $\mathbf{Cov}(X, Y) = E(XY) - E(X)E(Y)$ , the covariance of  $X$  and  $Y$ .

**Ex. 3.32** — Logs are milled to have a width of  $\mu$ . The actual width of a randomly selected item is  $X$ . If  $X$  is a  $\text{Normal}(\mu, \sigma^2)$  random variable then find the probability density function of the *squared-error* of the milling process,

$$Y = (X - \mu)^2.$$

**Ex. 3.33** — Let  $(X, Y)$  be a discrete random vector (R $\vec{V}$ ) with support:

$$\mathcal{S}_{X,Y} = \{(0, 0), (0, 1), (1, 0), (1, 1)\} .$$

Let its joint probability mass function (JPMF) be:

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{4} & \text{if } (x,y) = (0,0) \\ \frac{1}{4} & \text{if } (x,y) = (0,1) \\ \frac{1}{4} & \text{if } (x,y) = (1,0) \\ \frac{1}{4} & \text{if } (x,y) = (1,1) \\ 0 & \text{otherwise .} \end{cases}$$

Are  $X$  and  $Y$  independent?

**Ex. 3.34** — A semiconductor product consists of three layers that are fabricated independently. If the variances in thickness of the first, second and third layers are 25, 40 and 30 nanometers squared, what is the variance of the thickness of the final product?

**Ex. 3.35** — Find the covariance for the discrete R $\vec{V}$  ( $X, Y$ ) with joint probability mass function

$$f_{X,Y}(x,y) = \begin{cases} 0.2 & \text{if } (x,y) = (0,0) \\ 0.1 & \text{if } (x,y) = (1,1) \\ 0.1 & \text{if } (x,y) = (1,2) \\ 0.1 & \text{if } (x,y) = (2,1) \\ 0.1 & \text{if } (x,y) = (2,2) \\ 0.4 & \text{if } (x,y) = (3,3) \\ 0 & \text{otherwise .} \end{cases}$$

[Hint: Recall that  $\mathbf{Cov}(X, Y) = E(XY) - E(X)E(Y)$ ]

**Ex. 3.36** — Consider two random variables (RVs)  $X$  and  $Y$  having marginal distribution functions

$$F_X(x) = \begin{cases} 0 & \text{if } x < 1 \\ \frac{1}{2} & \text{if } 1 \leq x < 2 \\ 1 & \text{if } x \geq 2 \end{cases}$$

$$F_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ 1 - \frac{1}{2}e^{-y} - \frac{1}{2}e^{-2y} & \text{if } y \geq 0 \end{cases}$$

If  $X$  and  $Y$  are independent, what is their joint distribution function  $F_{X,Y}(x,y)$ ? [Hint: you need to express  $F_{X,Y}(x,y)$  for any  $(x,y) \in \mathbb{R}^2$ .]

**Ex. 3.37** — Let  $(X, Y)$  be a continuous R $\vec{V}$  with joint probability density function (JPDF):

$$f_{X,Y}(x,y) = \begin{cases} e^{-x} & \text{if } x \in [0, \infty) \text{ and } y \in [2, 3] \\ 0 & \text{otherwise .} \end{cases}$$

Are  $X$  and  $Y$  independent?

**Ex. 3.38** — In an electronic assembly, let the RVs  $X_1, X_2, X_3, X_4$  denote the lifetimes of four components in hours. Suppose that the JPDF of these variables is

$$f_{X_1,X_2,X_3,X_4}(x_1, x_2, x_3, x_4) \\ = \begin{cases} 9 \times 10^{-12} e^{-0.001x_1 - 0.002x_2 - 0.0015x_3 - 0.003x_4} & \text{if } x_1 \geq 0, x_2 \geq 0, x_3 \geq 0, x_4 \geq 0 \\ 0 & \text{otherwise .} \end{cases}$$

What is the probability that the device operates for more than 1000 hours without any failures? [Hint: The requested probability is  $\mathbf{P}(X_1 > 1000, X_2 > 1000, X_3 > 1000, X_4 > 1000)$  since each one of the four components of the device must not fail before 1000 hours.]

**Ex. 3.39** — Suppose the RVs  $Y_1$ ,  $Y_2$  and  $Y_3$  represent the thickness in micrometers of a substrate, an active layer, and a coating layer of a chemical product. Assume  $Y_1$ ,  $Y_2$  and  $Y_3$  are  $\text{Normal}(10000, 250^2)$ ,  $\text{Normal}(1000, 20^2)$  and  $\text{Normal}(80, 4^2)$  RVs, respectively. Further suppose that they are independent. The required specifications for the thickness of the substrate, active layer and coating layer are  $[9500, 10500]$ ,  $[950, 1050]$  and  $[75, 85]$ , respectively. What proportion of chemical products meets all thickness specifications? [Hint: this is just  $\mathbf{P}(9500 < Y_1 < 10500, 950 < Y_2 < 1050, 75 < Y_3 < 85)$ ] Which one of the three thicknesses has the least probability of meeting specifications?

**Ex. 3.40** — Soft drink cans are filled by an automated filling machine. Assume the fill volumes of the cans are independent  $\text{Normal}(12.1, 0.01)$  RVs. What is the probability that the average volume of ten cans selected from this process is less than 12.01 fluid ounces?

**Ex. 3.41** — Let  $X_1, X_2, X_3, X_4$  be RVs that denote the number of bits received in a digital channel that are classified as *excellent*, *good*, *fair* and *poor*, respectively. In a transmission of 10 bits, what is the probability that 6 of the bits received are *excellent*, 2 are *good*, 2 are *fair* and none are *poor* under the assumption that the classification of bits are independent events and that the probabilities of each bit being *excellent*, *good*, *fair* and *poor* are 0.6, 0.3, 0.08 and 0.02, respectively. [Hint: Think of  $\text{Multinomial}(n = 10, \theta_1 = 0.6, \theta_2 = 0.3, \theta_3 = 0.08, \theta_4 = 0.02)$  as a model for bit classification in this digital channel.]

**Ex. 3.42** — Let  $X$  be a discrete random variable (RV) with probability mass function (PMF)

$$f_X(x) = \begin{cases} \frac{1}{3} & \text{if } x = 0 \\ \frac{1}{3} & \text{if } x = 1 \\ \frac{1}{3} & \text{if } x = 2 \\ 0 & \text{otherwise} . \end{cases}$$

1. Find the characteristic function (CF) of  $X$
2. Using the CF find  $V(X)$ , the variance of  $X$ . Hint:  $V(X) = E(X^2) - (E(X))^2$

**Ex. 3.43** — Recall that the  $\text{Geometric}(\theta)$  RV  $X$  has the following PMF

$$f_X(x; \theta) = \begin{cases} \theta(1 - \theta)^x & \text{if } x \in \{0, 1, 2, 3, \dots\} \\ 0 & \text{otherwise} \end{cases}$$

1. Find the CF of  $X$ . (Hint: the sum of the infinite geometric series  $\sum_{x=0}^{\infty} ar^x = \frac{a}{1-r}$ .)
2. Using the CF find  $E(X)$ .

**Ex. 3.44** — Let  $X$  be the  $\text{Uniform}(a, b)$  RV with the following probability density function (PDF)

$$f_X(x; a, b) = \begin{cases} \frac{1}{b-a} & \text{if } x \in [a, b] \\ 0 & \text{otherwise} . \end{cases}$$

Find the CF of  $X$ .

**Ex. 3.45** — Recall that the Poisson( $\lambda$ ) RV has the following PMF

$$f_X(x; \lambda) = \begin{cases} \frac{\lambda^x}{x!} e^{-\lambda} & \text{if } x \in \{0, 1, 2, 3, \dots\} \\ 0 & \text{otherwise} . \end{cases}$$

Hint: the power series of  $e^\alpha = \sum_{x=0}^{\infty} \frac{\alpha^x}{x!}$ .

1. Find the CF of  $X$ .
2. Find the variance of  $X$  using its CF.

**Ex. 3.46** — Let  $X$  be a Poisson( $\lambda$ ) RV and  $Y$  be another Poisson( $\mu$ ) RV. Suppose  $X$  and  $Y$  are independent. Use Eqn. (3.49) to first find the CF of the RV  $W = X + Y$ . From the CF of  $W$  try to identify what RV it is.

**Ex. 3.47** — Recall from lecture that if  $Y = a + bX$  for some constants  $a$  and  $b$  with  $b \neq 0$  then  $\varphi_Y(t) = e^{iat} \varphi_X(bt)$  and that  $\varphi_Z(t) = e^{-t^2/2}$  if  $Z$  is the Normal( $0, 1$ ) RV. Using these facts find the CF of  $-Z$ , the RV obtained from  $Z$  by simply switching its sign. From the CF of  $-Z$  identify what RV it is.

# Chapter 4

## Statistics

### 4.1 Data and Statistics


**Definition 46 (Data)** The function  $X$  measures the outcome  $\omega$  of an experiment with sample space  $\Omega$  [Often, the sample space is also denoted by  $S$ ]. Formally,  $X$  is a random variable [or a random vector  $X = (X_1, X_2, \dots, X_n)$ , i.e. a vector of random variables] taking values in the **data space**  $\mathbb{X}$ :

$$X(\omega) : \Omega \rightarrow \mathbb{X}.$$

The realisation of the RV  $X$  when an experiment is performed is the observation or data  $x \in \mathbb{X}$ . That is, when the experiment is performed once and it yields a specific  $\omega \in \Omega$ , the data  $X(\omega) = x \in \mathbb{X}$  is the corresponding realisation of the RV  $X$ .

Figure 4.1: Sample Space, Random Variable, Realisation, Data, and Data Space.

**Example 72 (Tossing a coin  $n$  times)** For some given parameter  $\theta \in \Theta := [0, 1]$ , consider  $n$  IID Bernoulli( $\theta$ ) trials, i.e.  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} \text{Bernoulli}(\theta)$ . Then the random vector  $X = (X_1, X_2, \dots, X_n)$ , which takes values in the data space  $\mathbb{X} = \{0, 1\}^n := \{(x_1, x_2, \dots, x_n) : x_i \in \{0, 1\}, i = 1, 2, \dots, n\}$ , made up of vertices of the  $n$ -dimensional hyper-cube, measures the outcomes of this experiment. A particular realisation of  $X$ , upon performance of this experiment, is the observation, data or data vector  $(x_1, x_2, \dots, x_n)$ . For instance, if we observed  $n - 1$  tails and 1 heads, in that order, then our data vector  $(x_1, x_2, \dots, x_{n-1}, x_n) = (0, 0, \dots, 0, 1)$ .

Figure 4.2: Data Spaces  $\mathbb{X} = \{0, 1\}^2$  and  $\mathbb{X} = \{0, 1\}^3$  for two and three Bernoulli trials, respectively.

**Definition 47 (Statistic)** A **statistic**  $T$  is any function of the data:

$$T(x) : \mathbb{X} \rightarrow \mathbb{T} .$$

Thus, a statistic  $T$  is also an RV that takes values in the space  $\mathbb{T}$ . When  $x \in \mathbb{X}$  is the realisation of an experiment, we let  $T(x) = t$  denote the corresponding realisation of the statistic  $T$ . Sometimes we use  $T_n(X)$  and  $\mathbb{T}_n$  to emphasise that  $X$  is an  $n$ -dimensional random vector, i.e.  $\mathbb{X} \subset \mathbb{R}^n$ .

**Classwork 73 (Is data a statistic?)** Is the RV  $X$ , for which the realisation is the observed data  $X(\omega) = x$ , a statistic? In other words, is the data a statistic? [Hint: consider the identity map  $T(x) = x : \mathbb{X} \rightarrow \mathbb{T} = \mathbb{X}$ .]

Next, we define two important statistics called the **sample mean** and **sample variance**. Since they are obtained from the sample data, they are called **sample moments**, as opposed to the **population moments**. The corresponding population moments are  $\mathbf{E}(X_1)$  and  $\mathbf{V}(X_1)$ , respectively.

**Definition 48 (Sample Mean)** From a given a sequence of RVs  $X_1, X_2, \dots, X_n$ , we may obtain another RV called the  $n$ -samples mean or simply the sample mean:

$$T_n( (X_1, X_2, \dots, X_n) ) = \bar{X}_n( (X_1, X_2, \dots, X_n) ) := \frac{1}{n} \sum_{i=1}^n X_i . \quad (4.1)$$

For brevity, we write

$$\bar{X}_n( (X_1, X_2, \dots, X_n) ) \quad \text{as} \quad \bar{X}_n ,$$

and its realisation

$$\bar{X}_n( (x_1, x_2, \dots, x_n) ) \quad \text{as} \quad \bar{x}_n .$$

Note that the expectation and variance of  $\bar{X}_n$  are:

$$\begin{aligned} \mathbf{E}(\bar{X}_n) &= \mathbf{E}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) && [\text{by definition (4.1)}] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbf{E}(X_i) && [\text{by property (3.32)}] \end{aligned}$$

Furthermore, if every  $X_i$  in the original sequence of RVs  $X_1, X_2, \dots$  is **identically** distributed with the same expectation, by convention  $\mathbf{E}(X_1)$ , then:

$$\mathbf{E}(\bar{X}_n) = \frac{1}{n} \sum_{i=1}^n \mathbf{E}(X_i) = \frac{1}{n} \sum_{i=1}^n \mathbf{E}(X_1) = \frac{1}{n} n \mathbf{E}(X_1) = \mathbf{E}(X_1) . \quad (4.2)$$

Similarly, we can show that:

$$\begin{aligned} \mathbf{V}(\bar{X}_n) &= \mathbf{V}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) && [\text{by definition (4.1)}] \\ &= \left(\frac{1}{n}\right)^2 \mathbf{V}\left(\sum_{i=1}^n X_i\right) && [\text{by property (3.31)}] \end{aligned}$$

Furthermore, if the original sequence of RVs  $X_1, X_2, \dots$  is **independently** distributed then:

$$\mathbf{V}(\bar{X}_n) = \left(\frac{1}{n}\right)^2 \mathbf{V}\left(\sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \mathbf{V}(X_i) \quad [\text{by property (3.33)}]$$

Finally, if the original sequence of RVs  $X_1, X_2, \dots$  is **independently and identically** distributed with the same variance ( $\mathbf{V}(X_1)$  by convention) then:

$$\mathbf{V}(\bar{X}_n) = \frac{1}{n^2} \sum_{i=1}^n \mathbf{V}(X_i) = \frac{1}{n^2} \sum_{i=1}^n \mathbf{V}(X_1) = \frac{1}{n^2} n \mathbf{V}(X_1) = \frac{1}{n} \mathbf{V}(X_1) . \quad (4.3)$$

**Labwork 74 (Sample mean)** After initializing the fundamental sampler, we draw five samples and then obtain the sample mean using the MATLAB function `mean`. In the following, we will reuse the samples stored in the array `XsFromUni01Twstr101`.

```
>> rand('twister',101); % initialise the fundamental Uniform(0,1) sampler
>> XsFromUni01Twstr101=rand(1,5); % simulate n=5 IID samples from Uniform(0,1) RV
>> SampleMean=mean(XsFromUni01Twstr101);% find sample mean
>> disp(XsFromUni01Twstr101); % The data-points x_1,x_2,x_3,x_4,x_5 are:
    0.5164    0.5707    0.0285    0.1715    0.6853
>> disp(SampleMean); % The Sample mean is :
    0.3945
```

We can thus use `mean` to obtain the sample mean  $\bar{x}_n$  of  $n$  sample points  $x_1, x_2, \dots, x_n$ .

We may also obtain the sample mean using the `sum` function and a division by sample size:

```
>> sum(XsFromUni01Twstr101) % take the sum of the elements of the XsFromUni01Twstr101 array
ans =      1.9723
>> sum(XsFromUni01Twstr101) / 5 % divide the sum by the sample size 5
ans =      0.3945
```

We can also obtain the sample mean via matrix product or multiplication as follows:

```
>> size(XsFromUni01Twstr101) % size(SomeArray) gives the size or dimensions of the arrar SomeArray
ans =      1      5
>> ones(5,1) % here ones(5,1) is an array of 1's with size or dimension 5 X 1
ans =
    1
    1
```

```

1
1
1
>> XsFromUni01Twstr101 * ones(5,1) % multiplying an 1 X 5 matrix with a 5 X 1 matrix of Ones
ans = 1.9723
>> XsFromUni01Twstr101 * ( ones(5,1) * 1/5) % multiplying an 1 X 5 matrix with a 5 X 1 matrix of 1/5 's
ans = 0.3945

```

**Definition 49 (Sample Variance & Standard Deviation)** From a given a sequence of random variables  $X_1, X_2, \dots, X_n$ , we may obtain another statistic called the  $n$ -samples variance or simply the sample variance :

$$T_n( (X_1, X_2, \dots, X_n) ) = S_n^2( (X_1, X_2, \dots, X_n) ) := \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2 . \quad (4.4)$$

For brevity, we write  $S_n^2( (X_1, X_2, \dots, X_n) )$  as  $S_n^2$  and its realisation  $S_n^2( (x_1, x_2, \dots, x_n) )$  as  $s_n^2$ .

Sample standard deviation is simply the square root of sample variance:

$$S_n( (X_1, X_2, \dots, X_n) ) = \sqrt{S_n^2( (X_1, X_2, \dots, X_n) )} \quad (4.5)$$

For brevity, we write  $S_n( (X_1, X_2, \dots, X_n) )$  as  $S_n$  and its realisation  $S_n( (x_1, x_2, \dots, x_n) )$  as  $s_n$ .

Once again, if  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} X_1$ , the expectation of the sample variance is:

$$\mathbf{E}(S_n^2) = \mathbf{V}(X_1) .$$

**Labwork 75 (Sample variance and sample standard deviation)** We can compute the sample variance and sample standard deviation for the five samples stored in the array `XsFromUni01Twstr101` from Labwork 74 using MATLAB's functions `var` and `std`, respectively.

```

>> disp(XsFromUni01Twstr101); % The data-points x_1,x_2,x_3,x_4,x_5 are :
    0.5164    0.5707    0.0285    0.1715    0.6853
>> SampleVar=var(XsFromUni01Twstr101);% find sample variance
>> SampleStd=std(XsFromUni01Twstr101);% find sample standard deviation
>> disp(SampleVar) % The sample variance is:
    0.0785
>> disp(SampleStd) % The sample standard deviation is:
    0.2802

```

It is important to bear in mind that the statistics such as sample mean and sample variance are random variables and have an underlying distribution.

**Definition 50 (Order Statistics)** Suppose  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} F$ , where  $F$  is the DF from the set of all DFs over the real line. Then, the  $n$ -sample **order statistics**  $X_{([n])}$  is:

$$X_{([n])}( (X_1, X_2, \dots, X_n) ) := (X_{(1)}, X_{(2)}, \dots, X_{(n)}) , \text{ such that, } X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(n)} . \quad (4.6)$$

For brevity, we write  $X_{([n])}( (X_1, X_2, \dots, X_n) )$  as  $X_{([n])}$  and its realisation  $X_{([n])}( (x_1, x_2, \dots, x_n) )$  as  $x_{([n])} = (x_{(1)}, x_{(2)}, \dots, x_{(n)})$ .

Without going into the details of how to sort the data in ascending order to obtain the order statistics (an elementary topic of an Introductory Computer Science course), we simply use MATLAB's function `sort` to obtain the order statistics, as illustrated in the following example.

**Labwork 76 (Order statistics and sorting)** The order statistics for the five samples stored in `XsFromUni01Twstr101` from Labwork 74 can be computed using `sort` as follows:

```
>> disp(XsFromUni01Twstr101); % display the sample points
    0.5164    0.5707    0.0285    0.1715    0.6853
>> SortedXsFromUni01Twstr101=sort(XsFromUni01Twstr101); % sort data
>> disp(SortedXsFromUni01Twstr101); % display the order statistics
    0.0285    0.1715    0.5164    0.5707    0.6853
```

Therefore, we can use `sort` to obtain our order statistics  $x_{(1)}, x_{(2)}, \dots, x_{(n)}$  from  $n$  sample points  $x_1, x_2, \dots, x_n$ .

Next, we will introduce a family of common statistics, called the  $q^{\text{th}}$  quantile, by first defining the function:

**Definition 51 (Inverse DF or Inverse CDF or Quantile Function)** Let  $X$  be an RV with DF  $F$ . The **inverse DF** or **inverse CDF** or **quantile function** is:

$$F^{[-1]}(q) := \inf \{x : F(x) > q\}, \quad \text{for some } q \in [0, 1] . \quad (4.7)$$

If  $F$  is strictly increasing and continuous then  $F^{[-1]}(q)$  is the unique  $x \in \mathbb{R}$  such that  $F(x) = q$ .

A **functional** is merely a function of another function. Thus,  $T(F) : \{\text{All DFs}\} \rightarrow \mathbb{T}$ , being a map or function from the space of DFs to its range  $\mathbb{T}$ , is a functional. Some specific examples of functionals we have already seen include:

1. The **mean** of RV  $X \sim F$  is a function of the DF  $F$ :

$$T(F) = \mathbf{E}(X) = \int x dF(x) .$$

2. The **variance** of RV  $X \sim F$  is a function of the DF  $F$ :

$$T(F) = \mathbf{E}(X - \mathbf{E}(X))^2 = \int (x - \mathbf{E}(X))^2 dF(x) .$$

3. The **value of DF at a given  $x \in \mathbb{R}$**  of RV  $X \sim F$  is also a function of DF  $F$ :

$$T(F) = F(x) .$$

Other functionals of  $F$  that depend on the quantile function  $F^{[-1]}$  are:

1. The  $q^{\text{th}}$  **quantile** of RV  $X \sim F$ :

$$T(F) = F^{[-1]}(q) \quad \text{where } q \in [0, 1] .$$

2. The **first quartile** or the  $0.25^{\text{th}}$  **quantile** of the RV  $X \sim F$ :

$$T(F) = F^{[-1]}(0.25) .$$

3. The **median** or the **second quartile** or the  $0.50^{\text{th}}$  **quantile** of the RV  $X \sim F$ :

$$T(F) = F^{[-1]}(0.50) .$$

4. The **third quartile** or the  $0.75^{\text{th}}$  **quantile** of the RV  $X \sim F$ :

$$T(F) = F^{[-1]}(0.75) .$$

**Definition 52 (Empirical Distribution Function (EDF or ECDF))** Suppose we have  $n$  IID RVs,  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} F$ , where  $F$  is a DF from the set of all DFs over the real line. Then, the  $n$ -sample empirical distribution function (EDF or ECDF) is the discrete distribution function  $\hat{F}_n$  that puts a probability mass of  $1/n$  at each sample or data point  $x_i$ :

$$\hat{F}_n(x) = \frac{\sum_{i=1}^n \mathbf{1}(X_i \leq x)}{n}, \quad \text{where} \quad \mathbf{1}(X_i \leq x) := \begin{cases} 1 & \text{if } x_i \leq x \\ 0 & \text{if } x_i > x \end{cases} \quad (4.8)$$

**Labwork 77 (Plot of empirical CDF)** Let us plot the ECDF for the five samples drawn from the  $\text{Uniform}(0, 1)$  RV in Labwork 74 using the MATLAB function ECDF (given in Labwork ??). Let us super-impose the samples and the true DF as depicted in Figure 4.3 with the following script:

---

```
plotunifecdf.m
xs = -1:0.01:2; % vector xs from -1 to 2 with increment .05 for x values
% get the [0,1] uniform DF or cdf of xs in vector cdf
cdf=zeros(size(xs));% initialise cdf as zero
indices = find(xs>=1); cdf(indices) = 1; % set cdf as 1 when xs >= 1
indices = find(xs>=0 & xs<=1); cdf(indices)=xs(indices); % cdf=xs when 0 <= xs <= 1
plot(xs,cdf,'r') % plot the DF
hold on; title('Uniform [0,1] DF and ECDF'); xlabel('x'); axis([-0.2 1.2 -0.2 1.2])
x=[0.5164, 0.5707, 0.0285, 0.1715, 0.6853]; % five samples
plot(x,zeros(1,5),'r+','LineWidth',2,'MarkerSize',10)% plot the data as red + marks
hold on; grid on; % turn on grid
ECDF(x,1,.2,.6);% ECDF (type help ECDF) plot is extended to left and right by .2 and .4, respectively.
```

---

**Definition 53 ( $q^{\text{th}}$  Sample Quantile)** For some  $q \in [0, 1]$  and  $n$  IID RVs  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} F$ , we can obtain the ECDF  $\hat{F}_n$  using (4.8). The  $q^{\text{th}}$  **sample quantile** is defined as the statistic (statistical functional):

$$T(\hat{F}_n) = \hat{F}_n^{[-1]}(q) := \inf \{x : \hat{F}_n^{[-1]}(x) \geq q\} . \quad (4.9)$$

By replacing  $q$  in this definition of the  $q^{\text{th}}$  sample quantile by 0.25, 0.5 or 0.75, we obtain the first, second (**sample median**) or third **sample quartile**, respectively.


The following algorithm can be used to obtain the  $q^{\text{th}}$  sample quantile of  $n$  IID samples  $(x_1, x_2, \dots, x_n)$  on the basis of their order statistics  $(x_{(1)}, x_{(2)}, \dots, x_{(n)})$ .

The  $q^{\text{th}}$  sample quantile,  $\hat{F}_n^{[-1]}(q)$ , is found by interpolation from the order statistics  $(x_{(1)}, x_{(2)}, \dots, x_{(n)})$  of the  $n$  data points  $(x_1, x_2, \dots, x_n)$ , using the formula:

$$\hat{F}_n^{[-1]}(q) = (1 - \delta)x_{(i+1)} + \delta x_{(i+2)}, \quad \text{where,} \quad i = \lfloor (n-1)q \rfloor \quad \text{and} \quad \delta = (n-1)q - \lfloor (n-1)q \rfloor .$$

Thus, the **sample minimum** of the data points  $(x_1, x_2, \dots, x_n)$  is given by  $\hat{F}_n^{[-1]}(0)$ , the **sample maximum** is given by  $\hat{F}_n^{[-1]}(1)$  and the **sample median** is given by  $\hat{F}_n^{[-1]}(0.5)$ , etc.

Figure 4.3: Plot of the DF of Uniform(0, 1), five IID samples from it, and the ECDF  $\hat{F}_5$  for these five data points  $x = (x_1, x_2, x_3, x_4, x_5) = (0.5164, 0.5707, 0.0285, 0.1715, 0.6853)$  that jumps by  $1/5 = 0.20$  at each of the five samples.




---

**Algorithm 1**  $q^{\text{th}}$  Sample Quantile of Order Statistics

---

1: *input:*

1.  $q$  in the  $q^{\text{th}}$  sample quantile, i.e. the argument  $q$  of  $\hat{F}_n^{[-1]}(q)$ ,
  2. order statistic  $(x_{(1)}, x_{(2)}, \dots, x_{(n)})$ , i.e. the sorted  $(x_1, x_2, \dots, x_n)$ , where  $n > 0$ .
- 2: *output:*  $\hat{F}_n^{[-1]}(q)$ , the  $q^{\text{th}}$  sample quantile
- 3:  $i \leftarrow \lfloor (n-1)q \rfloor$
  - 4:  $\delta \leftarrow (n-1)q - i$
  - 5: **if**  $i = n-1$  **then**
  - 6:    $\hat{F}_n^{[-1]}(q) \leftarrow x_{(i+1)}$
  - 7: **else**
  - 8:    $\hat{F}_n^{[-1]}(q) \leftarrow (1-\delta)x_{(i+1)} + \delta x_{(i+2)}$
  - 9: **end if**
- 10: *return:*  $\hat{F}_n^{[-1]}(q)$
-

**Labwork 78 (The  $q^{\text{th}}$  sample quantile)** Use the implementation of Algorithm 1 in Labwork ?? as the MATLAB function `qthSampleQuantile` to find the  $q^{\text{th}}$  sample quantile of two simulated data arrays:

1. `SortedXsFromUni01Twstr101`, the order statistics that was constructed in Labwork 76 and
2. Another sorted array of 7 samples called `SortedXs`

```
>> disp(SortedXsFromUni01Twstr101)
    0.0285    0.1715    0.5164    0.5707    0.6853
>> rand('twister',420);
>> SortedXs=sort(rand(1,7));
>> disp(SortedXs)
    0.1089    0.2670    0.3156    0.3525    0.4530    0.6297    0.8682
>> for q=[0, 0.25, 0.5, 0.75, 1.0]
    disp([q, qthSampleQuantile(q,SortedXsFromUni01Twstr101) ...
           qthSampleQuantile(q,SortedXs)])
end
      0    0.0285    0.1089
    0.2500    0.1715    0.2913
    0.5000    0.5164    0.3525
    0.7500    0.5707    0.5414
    1.0000    0.6853    0.8682
```

## 4.2 Exploring Data and Statistics

### 4.2.1 Univariate Data

A **histogram** is a graphical representation of the frequency with which elements of a data array:

$$x = (x_1, x_2, \dots, x_n) ,$$

of real numbers fall within each of the  $m$  intervals or **bins** of some **interval partition**:

$$b := (b_1, b_2, \dots, b_m) := ([\underline{b}_1, \bar{b}_1], [\underline{b}_2, \bar{b}_2], \dots, [\underline{b}_m, \bar{b}_m])$$

of the **data range** of  $x$  given by the closed interval:

$$\mathcal{R}(x) := [\min\{x_1, x_2, \dots, x_n\}, \max\{x_1, x_2, \dots, x_n\}] .$$

Elements of this partition  $b$  are called bins, their mid-points are called **bin centres**:

$$c := (c_1, c_2, \dots, c_m) := ((\underline{b}_1 + \bar{b}_1)/2, (\underline{b}_2 + \bar{b}_2)/2, \dots, (\underline{b}_m + \bar{b}_m)/2)$$

and their overlapping boundaries, i.e.  $\bar{b}_i = \underline{b}_{i+1}$  for  $1 \leq i < m$ , are called **bin edges**:

$$d := (d_1, d_2, \dots, d_{m+1}) := (\underline{b}_1, \underline{b}_2, \dots, \underline{b}_{m-1}, \underline{b}_m, \bar{b}_m) .$$

For a given partition of the data range  $\mathcal{R}(x)$  or some superset of  $\mathcal{R}(x)$ , three types of histograms are possible: frequency histogram, relative frequency histogram and density histogram. Typically, the partition  $b$  is assumed to be composed of  $m$  overlapping intervals of the same width  $w = \bar{b}_i - \underline{b}_i$  for all  $i = 1, 2, \dots, m$ . Thus, a histogram can be obtained by a set of bins along with their corresponding **heights**:

$$h = (h_1, h_2, \dots, h_m) , \text{ where } h_k := g(\#\{x_i : x_i \in b_k\})$$

Thus,  $h_k$ , the height of the  $k$ -th bin, is some function  $g$  of the number of data points that fall in the bin  $b_k$ . Formally, a histogram is a sequence of ordered pairs:

$$((b_1, h_1), (b_2, h_2), \dots, (b_m, h_m)) .$$

Given a partition  $b$ , a **frequency histogram** is the histogram:

$$((b_1, h_1), (b_2, h_2), \dots, (b_m, h_m)) , \text{ where } h_k := \#\{x_i : x_i \in b_k\} ,$$


a **relative frequency histogram** is the histogram:

$$((b_1, h_1), (b_2, h_2), \dots, (b_m, h_m)) , \text{ where } h_k := n^{-1} \#\{x_i : x_i \in b_k\} ,$$

and a **density histogram** is the histogram:

$$((b_1, h_1), (b_2, h_2), \dots, (b_m, h_m)) , \text{ where } h_k := (w_k n)^{-1} \#\{x_i : x_i \in b_k\} , w_k := \bar{b}_k - \underline{b}_k .$$

Figure 4.4: Frequency, Relative Frequency and Density Histograms



**Labwork 79 (Histograms with specified number of bins for univariate data)** Let us use samples from the `rand('twister',5489)` as our data set  $x$  and plot various histograms. Let us use `hist` function (read `help hist`) to make a default histogram with ten bins. Then we can make three types of histograms as shown in Figure 4.4 as follows:


```
>> rand('twister',5489);
>> x=rand(1,100); % generate 100 PRNs
>> hist(x) % see what default hist does in Figure Window
>> % Now let us look deeper into the last hist call
>> [Fs, Cs] = hist(x) % Cs is the bin centers and Fs is the frequencies of data set x
Fs =
    9     11     10      8      7     10      8     11     10     16
Cs =
    0.0598    0.1557    0.2516    0.3474    0.4433    0.5392    0.6351    0.7309    0.8268    0.9227
>> % produce a histogram plot the last argument 1 is the width value for immediately adjacent bars -- help bar
>> bar(Cs,Fs,1) % create a frequency histogram
>> bar(Cs,Fs/100,1) % create a relative frequency histogram
>> bar(Cs,Fs/(0.1*100),1) % create a density histogram (area of bars sum to 1)
>> sum(Fs/(0.1*100) .* ones(1,10)*0.1) % checking if area does sum to 1
>> ans = 1
```

Try making a density histogram with 1000 samples from `rand` with 15 bins. You can specify the number of bins by adding an extra argument to `hist`, for e.g. `[Fs, Cs] = hist(x,15)` will produce 15 bins of equal width over the data range  $\mathcal{R}(x)$ .

**Labwork 80 (Stem plots and ECDF plots for univariate data)** We can also visualise the 100 data points in the array  $x$  using stem plot and ECDF plot as shown in Figure 4.5 as follows:

```
>> rand('twister',5489);
>> x=rand(1,100); % produce 100 samples with rand
>> stem(x,'.') % make a stem plot of the 100 data points in x (the option '.' gives solid circles for x)
>>% ECDF (type help ECDF) plot is extended to left and right by .2 and .6, respectively
>>% (second parameter 6 makes the dots in the plot smaller).
>> ECDF(x,6,.2,.6);
```

Figure 4.5: Frequency, Relative Frequency and Density Histograms



We can also visually summarise univariate data using the **box plot** or **box-whisker plot** available in the Stats Toolbox of MATLAB. These family of plots display a set of sample quantiles, typically they are include, the median, the first and third quartiles and the minimum and maximum values of our data array  $x$ .

#### 4.2.2 Bivariate Data


By bivariate data array  $x$  we mean a  $2 \times n$  matrix of real numbers or equivalently  $n$  ordered pairs of points  $(x_{1,i}, x_{2,i})$  as  $i = 1, 2, \dots, n$ . The most elementary visualisation of these  $n$  ordered pairs is in orthogonal Cartesian co-ordinates. Such plots are termed **2D scatter plots** in statistics.

**Labwork 81 (Visualising bivariate data)** Let us generate a  $2 \times 5$  array representing samples of 5 ordered pairs sampled uniformly at random over the unit square  $[0, 1] \times [0, 1]$ . We can make 2D scatter plot as shown in Figure 4.6 as follows:

```
>> rand('twister',5489);
>> x=rand(2,5)% create a sequence of 5 ordered pairs uniformly from unit square [0,1]X[0,1]
x =
    0.8147    0.1270    0.6324    0.2785    0.9575
    0.9058    0.9134    0.0975    0.5469    0.9649
>> plot(x(1,:),x(2,:),'x') % a 2D scatter plot with marker cross or 'x'
>> plot(x(1,:),x(2,:),'x', 'MarkerSize',15) % a 2D scatter plot with marker cross or 'x' and larger Marker size
>> xlabel('x_1'); ylabel('x_2'); % label the axes
```

There are several other techniques for visualising bivariate data, including, 2D histograms, surface plots, heat plots, and we will encounter some of them in the sequel.

Figure 4.6: 2D Scatter Plot



### 4.2.3 Trivariate Data

Trivariate data is more difficult to visualise on paper but playing around with the rotate 3D feature in MATLAB's Figure window can help bring a lot more perspective.

**Labwork 82 (Visualising trivariate data)** We can make **3D scatter plots** as shown in Figure 4.7 as follows:

```
>> rand('twister',5489);
>> x=rand(3,5)% create a sequence of 5 ordered triples uniformly from unit cube [0,1]X[0,1]X[0,1]
x =
    0.8147    0.9134    0.2785    0.9649    0.9572
    0.9058    0.6324    0.5469    0.1576    0.4854
    0.1270    0.0975    0.9575    0.9706    0.8003
>> plot3(x(1,:),x(2,:),x(3,:),'x') % a simple 3D scatter plot with marker 'x'
>>% a more interesting one with options that control marker type, line-style,
>>% colour in [Red Green Blue] values and marker size - read help plot3 for more options
>> plot3(x(1,:),x(2,:),x(3,:),'Marker','*','LineStyle','none','Color',[1 0 1],'MarkerSize',15)
>> plot3(x(1,:),x(2,:),x(3,:),'m*','MarkerSize',15) % makes same figure as before but shorter to write
>> box on % turn on the box and see the effect on the Figure
>> grid on % turn on the grid and see the effect on the Figure
>> xlabel('x_1'); ylabel('x_2'); zlabel('x_3'); % assign labels to x,y and z axes
```


Repeat the visualisation below with a larger array, say  $x=\text{rand}(3,1000)$ , and use the rotate 3D feature in the Figure window to visually explore the samples in the unit cube. Do they seem to be uniformly distributed inside the unit cube?

There are several other techniques for visualising trivariate data, including, iso-surface plots, moving surface or heat plots, and you will encounter some of them in the future.


### 4.2.4 Multivariate Data

For high-dimensional data in  $d$ -dimensional space  $\mathbb{R}^d$  with  $d \geq 3$  you have to look at several lower dimensional projections of the data. We can simultaneously look at 2D scatter plots for every pair of


Figure 4.7: 3D Scatter Plot



co-ordinates  $\{(i, j) \in \{1, 2, \dots, d\}^2 : i \neq j\}$  and at histograms for every co-ordinate  $i \in \{1, 2, \dots, d\}$  of the  $n$  data points in  $\mathbb{R}^d$ . Such a set of low-dimensional projections can be conveniently represented in a  $d \times d$  matrix of plots called a **matrix plot**.

Figure 4.8: Plot Matrix of uniformly generated data in  $[0, 1]^5$ 

(a) First six samples



(b) All thousand samples

**Labwork 83** Let us make matrix plots from a uniformly generated sequence of 100 points in 5D unit cube  $[0, 1]^5$  as shown in Figure 4.8.

```
>> rand('twister',5489);
>> % generate a sequence of 1000 points uniformly distributed in 5D unit cube [0,1]X[0,1]X[0,1]X[0,1]X[0,1]
>> x=rand(1000,5);
>> x(1:6,:) % first six points in our 5D unit cube, i.e., the first six rows of x
ans =
```

```

0.8147    0.6312    0.7449    0.3796    0.4271
0.9058    0.3551    0.8923    0.3191    0.9554
0.1270    0.9970    0.2426    0.9861    0.7242
0.9134    0.2242    0.1296    0.7182    0.5809
0.6324    0.6525    0.2251    0.4132    0.5403
0.0975    0.6050    0.3500    0.0986    0.7054
>> plotmatrix(x(1:5,:),'r*') % make a plot matrix
>> plotmatrix(x) % make a plot matrix of all 1000 points

```

## 4.3 Loading and Exploring Real-world Data

All of the data we have played with so far were computer-generated. It is time to get our hands dirty with real-world data. The first step is to obtain the data. Often, publicly-funded institutions allow the public to access their databases. Such data can be fetched from appropriate URLs in one of the two following ways:

**Method A:** Manually download by filling the appropriate fields in an online request form.

**Method B:** Automagically download directly from your MATLAB session.

Then we want to inspect it for inconsistencies, missing values and replace them with `NaN` values in MATLAB that stand for not-any-number. Finally, we can visually explore, transform and interact with the data to discover interesting patterns that are hidden in the data. This process is called *exploratory data analysis* and is the foundational first step towards subsequent computational statistical experiments [John W. Tukey, *Exploratory Data Analysis*, Addison-Wesely, New York, 1977].

### 4.3.1 Geological Data

Let us focus on the data of earth quakes that heavily damaged Christchurch on February 22 2011. This data can be fetched from the URL <http://magma.geonet.org.nz/resources/quakesearch/> by Method A and loaded into MATLAB for exploratory data analysis as done in Labwork 84.

**Labwork 84** Let us go through the process one step at a time using Method A.

1. Download the data as a CSV or *comma separated variable* file in plain ASCII text (this has been done for this data already for you and saved as `NZ20110222earthquakes.csv` in the `CSEMatlabScripts` directory).
2. Open the file in a simple text editor such as `Note Pad` in Windows or one of the following editors in OS X, Unix, Solaris, Linux/GNU variants such as Ubuntu, SUSE, etc: `vi`, `vim`, `emacs`, `geany`, etc. The first three and last two lines of this file look as follows:

```

CUSP_ID,LAT,LONG,NZMGE,NZMGN,ORI_YEAR,ORI_MONTH,ORI_DAY,ORI_HOUR,ORI_MINUTE,ORI_SECOND,MAG,DEPTH
3481751,-43.55432,172.68898,2484890,5739375,2011,2,22,0,0,31.27814,3.79,5.8559,
3481760,-43.56579,172.70621,2486287,5738106,2011,2,22,0,0,43.70276,3.76,5.4045,
.
.
.
3469114,-43.58007,172.67126,2483470,5736509,2011,2,22,23,28,11.1014,3.117,3,
3469122,-43.55949,172.70396,2486103,5738805,2011,2,22,23,50,1.06171,3.136,12,

```

The thirteen columns correspond to fairly self-descriptive features of each measured earth quake given in the first line or row. They will become clear in the sequel. Note that the comma character (',') separates each unit or measurement or description in any CSV file.

3. The next set of commands show you how to load, manipulate and visually explore this data.

```
%>> %% Load the data from the comma delimited text file 'NZ20110222earthquakes.csv' with
%% the following column IDs
%% CUSP_ID,LAT,LONG,NZMGE,NZMGN,ORI_YEAR,ORI_MONTH,ORI_DAY,ORI_HOUR,ORI_MINUTE,ORI_SECOND,MAG,DEPTH
%% Using MATLAB's dlmread command we can assign the data as a matrix to EQ;
%% note that the option 1,0 to dlmread skips first row of column descriptors
%
% the variable EQall is about to be assigned the data as a matrix
EQall = dlmread('NZ20110222earthquakes.csv', ',', 1, 0);
size(EQall) % report the dimensions or size of the matrix EQall
ans =
    145      14
```

4. In order to understand the syntax in detail get help from MATLAB !

```
>> help dlmread
DLMREAD Read ASCII delimited file.
.
.
.
```

5. When there are units in the CSV file that can't be converted to floating-point numbers, it is customary to load them as a `NaN` or *Not-a-Number* value in MATLAB . So, let's check if there are any rows with `NaN` values and remove them from our analysis. Note that this is not the only way to deal with missing data! After that let's remove any locations outside Christchurch and its suburbs (we can find the latitude and longitude bounds from online resources easily) and finally view the 4-tuples of (latitude, longitude, magnitude, depth) for each measured earth quake in Christchurch on February 22 of 2011 as a scatter plot shown in Figure 4.9 (the axes labels were subsequently added from clicking <Edit> and <Figure Properties...> tabs of the output Figure Window).


```
>> EQall(any(isnan(EQall),2),:) = []; %Remove any rows containing NaNs from the matrix EQall
>> % report the size of EQall and see if it is different from before we removed and NaN containing rows
>> size(EQall)
ans = 145 14
>> % remove locations outside Chch and assign it to a new variable called EQ
>> EQ = EQall(-43.75<EQall(:,2) & EQall(:,2)<-43.45 ...
& 172.45<EQall(:,3) & EQall(:,3)<172.9 & EQall(:,12)>3, :);
>> % now report the size of the earthquakes in Christchurch in variable EQ
>> size(EQ)
ans = 124 14
>> % assign the four variables of interest
>> LatData=EQ(:,2); LonData=EQ(:,3); MagData=EQ(:,12); DepData=EQ(:,13);
>> % finally make a plot matrix of these 124 4-tuples as red points
>> plotmatrix([LatData,LonData,MagData,DepData], 'r.');
```

All of these commands have been put in a script M-file `NZEQChCch20110222.m` in Labwork ?? and you can simply call it from the command window to automatically load the data and assign it to the variables `EQAll` `EQ`, `LatData`, `LonData`, `MagData` and `DepData`, instead of retyping each command above every time you need these matrices in MATLAB , as follows:

```
>> NZEQChCch20110222
ans =    145    14
ans =    145    14
ans =    124    14
```

In fact, we will do exactly this to conduct more exploratory data analysis with these earth quake measurements in Labwork 85.

Figure 4.9: Matrix of Scatter Plots of the latitude, longitude, magnitude and depth of the 22-02-2011 earth quakes in Christchurch, New Zealand.



**Labwork 85** Try to understand how to manipulate time stamps of events in MATLAB and the Figures being output by following the comments in the script file `NZEQChCch20110222EDA.m`.

```
>> NZEQChCch20110222
ans =    145    14
ans =    145    14
ans =    124    14
ans =    145    14
ans =    145    14
ans =    124    14
ans = 22-Feb-2011 00:00:31
ans = 22-Feb-2011 23:50:01
```

---

```
----- NZEQChCch20110222EDA.m -----
%% Load the data from the comma delimited text file 'NZ20110222earthquakes.csv'
% using the script M-file NZEQChCch20110222.m
NZEQChCch20110222
%% working with time stamps is tricky
%% time is encoded by columns 6 through 11
%% as origin of earthquake in year, month, day, hour, minute, sec:
```

```

%% ORI_YEAR,ORI_MONTH,ORI_DAY,ORI_HOUR,ORI_MINUTE,ORI_SECOND
%% datenum is Matlab's date encoding function see help datenum
TimeData=datenum(EQ(:,6:11)); % assign origin times of earth quakes in datenum coordinates
MaxD=max(TimeData); % get the latest time of observation in the data
MinD=min(TimeData); % % get the earliest time of observation in the data
datestr(MinD) % a nice way to conver to calendar time!
datestr(MaxD) % ditto

% recall that there four variables were assigned in NZEQChCch20110222.m
% LatData=EQ(:,2); LonData=EQ(:,3); MagData=EQ(:,12); DepData=EQ(:,13);

%clear any existing Figure windows
clf
plot(TimeData,MagData,'o-') % plot origin time against magnitude of each earth quake

figure % tell matlab you are about to make another figure
plotmatrix([LatData,LonData,MagData,DepData],'r.');

figure % tell matlab you are about to make another figure
scatter(LonData,LatData,'.') % plot the LONGitude Vs. LATtitude

figure % tell matlab you are about to make another figure
% relative frequency histogram of magnitudes from 0 to 12 on Richter Scale with 15 bins
hist(MagData,15)

%max(MagData)

figure % tell matlab you are about to make another figure
semilogx(DepData,MagData,'.') % see the depth in log scale


%%%%%
% more advanced topic - uncomment and read help if bored
%tri = delaunay(LatData,LonData);
%triplot(tri,LatData,LonData,DepData);

```

---

## Geostatistical exploratory data analysis with Google Earth

Figure 4.10: Google Earth Visualisation of the earth quakes



A global search at <http://neic.usgs.gov/cgi-bin/epic/epic.cgi> with the following parame-

ters:

Date Range: 2011 2 22 to 2011 2 22

Catalog: USGS/NEIC (PDE-Q)

produced 43 earth quakes world-wide, including those in Christchurch as shown in Figure 4.10. One can do a lot more than a mere visualisation with the USGS/NEIC database of earth-quakes worldwide, the freely available Google earth software bundle <http://www.google.com/earth/index.html> and the freely available MATLAB package googleearth from [http://www.mathworks.com/matlabcentral/fx\\_files/12954/4/content/googleearth/html/html\\_product\\_page.html](http://www.mathworks.com/matlabcentral/fx_files/12954/4/content/googleearth/html/html_product_page.html).

### 4.3.2 Metereological Data

New Zealand's meteorological service NIWA provides weather data under its TERMS AND CONDITIONS FOR ACCESS TO DATA (See [http://cliflo.niwa.co.nz/doc/terms\\_print.html](http://cliflo.niwa.co.nz/doc/terms_print.html)). We will explore some data of rainfall and temperatures from NIWA.

#### Daily Rainfalls in Christchurch

Automagic downloading of the data by Method B can be done if the data provider allows automated queries. It can be accomplished by `urlread` for instance.

Paul Brouwers has a basic CliFlo datafeed on <http://www.math.canterbury.ac.nz/php/lib/cliflo/rainfall.php>. This returns the date and rainfall in milli meters as measured from the CHCH aeroclub station. It is assumed that days without readings would not be listed. The data doesn't go back much before 1944.

**Labwork 86** Understand how Figure 4.11 is obtained by the script file `RainFallsInChch.m` by typing and following the comments:

```
>> RainFallsInChch
RainFallsChch = [24312x1 int32] [24312x1 double]
ans = 24312 2
FirstDayOfData = 19430802
LastDayOfData = 20100721
```

---


```
RainFallsInChch.m
%% How to download data from an URL directly without having to manually
%% fill out forms
% first make a string of the data using urlread (read help urlread if you want details)
StringData = urlread('http://www.math.canterbury.ac.nz/php/lib/cliflo/rainfall.php');
RainFallsChch = textscan(StringData, '%d %f', 'delimiter', ',')
RC = [RainFallsChch{1} RainFallsChch{2}]; % assign Matlab cells as a matrix
size(RC) % find the size of the matrix

FirstDayOfData = min(RC(:,1))
LastDayOfData = max(RC(:,1))

plot(RC(:,2),'.')
xlabel('Days in Christchurch, NZ since August 2nd of 1943','FontSize',20);
ylabel('Rainfall in millimeters','FontSize',20)
```

---

Figure 4.11: Daily rainfalls in Christchurch since March 27 2010



### Daily Temperatures in Christchurch

**Labwork 87** Understand how Figure 4.12 is being generated by following the comments in the script file ChchTempsLoad.m by typing:

```
>> ChchTempsLoad
```

---

```
ChchTempsLoad.m
%% Load the data from the comma delimited text file 'NIWACliFloChchAeroClubStationTemps.txt'
%% with the following column IDs
%% Max_min: Daily Temperature in Christchurch New Zealand
%% Stationate(NZST),Tmax(C),Period(Hrs),Tmin(C),Period(Hrs),Tgmin(C),Period(Hrs),Tmean(C),RHmean(%),Period(Hrs)

% the matrix T is about to be assigned the data as a matrix; the option [27,1,20904,5] to
% specify the upper-left and lower-right corners of an imaginary rectangle
% over the text file 'NIWACliFloChchAeroClubStationTemps.txt'.
% here we start from line number 27 and end at the last line number 20904
% and we read only columns NZST,Tmax(C),Period(Hrs),Tmin(C),Period(Hrs)

T = dlmread('NIWACliFloChchAeroClubStationTemps.txt','','',[27,1,20904,5]);
% just keep column 1,2 and 4 named NZST,Tmax(C),Period(Hrs),Tmin(C),
% i.e. date in YYYYMMDD foramt, maximum temperature, minimum temperature
T = T(:,[1,2,4]); % just pull the time
% print size before removing missig data rows are removed
size(T) % report the dimensions or size of the matrix T

% This file has a lot of missing data points and they were replaced with
% NaN values - see the file for various manipulations that were done to the
% raw text file from NIWA (Copyright NIWA 2011 Subject to NIWA's Terms and
% Conditions. See: http://cliflo.niwa.co.nz/pls/niwp/doc/terms.html)
T(any(isnan(T),2),:) = [];% Remove any rows containing NaNs from a matrix

size(T) % if the matrix has a different size now then the data-less days now!
```

```


clf % clears all current figures

% Daily max and min temperature in the 100 days with good data
% before last date in this data, i.e., March 27 2011 in Christchurch NZ
H365Days = T(end-365:end,2);
L365Days = T(end-365:end,3);
F365Days = H365Days-L365Days; % assign the maximal fluctuation, i.e. max-min
plot(H365Days,'r*') % plot daily high or maximum temperature = Tmax
hold on; % hold the Figure so that we can overlay more plots on it
plot(L365Days,'b*') % plot daily low or minimum temperature = Tmin
plot(F365Days, 'g*') % plot daily Fluctuation = Tmax - Tmin
% filter for running means
windowSize = 7;
WeeklyHighs = filter(ones(1,windowSize)/windowSize,1,H365Days);
plot(WeeklyHighs,'r.-')
WeeklyLows = filter(ones(1,windowSize)/windowSize,1,L365Days);
plot(WeeklyLows,'b.-')
WeeklyFlucs = filter(ones(1,windowSize)/windowSize,1,F365Days);
plot(WeeklyFlucs,'g.-')
xlabel('Number of days since March 27 2010 in Christchurch, NZ','FontSize',20);
ylabel('Temperature in Celsius','FontSize',20)
MyLeg = legend('Daily High','Daily Low','Daily Fluc.', 'Weekly High','Weekly Low',...
    'Weekly Fluc.', 'Location','NorthEast')
% Create legend
% legend1 = legend(axes1,'show');
set(MyLeg,'FontSize',20);
xlim([0 365]); % set the limits or boundary on the x-axis of the plots
hold off % turn off holding so we stop overlaying new plots on this Figure

```

---

Figure 4.12: Daily temperatures in Christchurch for one year since March 27 2010



### 4.3.3 Textual Data

Processing and analysing textual data to make a decision is another important computational statistical experiment. An obvious example is machine translation and a less obvious one is exploratory data analysis of the textual content of


- a large document
  - twitter messages within an online social network of interest
  - etc.

An interesting document with a current affairs projection is the Joint Operating Environment 2010 Report by the US Department of Defense. This document was downloaded from [http://www.jfcom.mil/newslink/storyarchive/2010/JOE\\_2010\\_o.pdf](http://www.jfcom.mil/newslink/storyarchive/2010/JOE_2010_o.pdf). The first paragraph of this 74 page document (JOE 2010 Report) reads:

**ABOUT THIS STUDY** The Joint Operating Environment is intended to inform joint concept development and experimentation throughout the Department of Defense. It provides a perspective on future trends, shocks, contexts, and implications for future joint force commanders and other leaders and professionals in the national security field. This document is speculative in nature and does not suppose to predict what will happen in the next twenty-five years. Rather, it is intended to serve as a starting point for discussions about the future security environment at the operational level of war. Inquiries about the Joint Operating Environment should be directed to USJFCOM Public Affairs, 1562 Mitscher Avenue, Suite 200, Norfolk, VA 23551-2488, (757) 836-6555.

Distribution Statement A: Approved for Public Release

Figure 4.13: Wordle of JOE 2010



We can try to produce a statistic of this document by recording the frequency of words in its textual content. Then we can produce a “word histogram” or “word cloud” to explore the document visually at one of the coarsest possible resolutions of the textual content in the JOE 2010 Report. The “word cloud” shown in Figure 4.13 was produced by Phillip Wilson using *wordle* from <http://www.wordle.net/>. A description from the wordle URL says:

Wordle is a toy for generating word clouds from text that you provide. The clouds give greater prominence to words that appear more frequently in the source text. You can tweak your clouds with different fonts, layouts, and color schemes. The images you create with Wordle are yours to use however you like. You can print them out, or save them to the Wordle gallery to share with your friends.

**Labwork 88 (favourite word cloud)** This is just for fun. Produce a “word cloud” of your honours thesis or summer project or any other document that fancies your interest by using *wordle* from <http://www.wordle.net/>. Play with the aesthetic features to change colour, shapes, etc.

#### 4.3.4 Machine Sensor Data

Instrumentation of modern machines, such as planes, rockets and cars allow the sensors in the machines to collect live data and dynamically take *decisions* and subsequent *actions* by executing algorithms to drive their devices in response to the data that is streaming into their sensors. For example, a rocket may have to adjust its boosters to compensate for the prevailing directional changes in wind in order to keep going up and launch a satellite. These types of decisions and actions, theorised by *controlled Markov processes*, typically arise in various fields of engineering such as, aerospace, civil, electrical, mechanical, robotics, etc.

In an observational setting, without an associated control problem, one can use machine sensor data to get information about some state of the system or phenomenon, i.e., what is it doing? or where is it?, etc. Sometimes sensors are attached to a sample of individuals from a wild population, say Emperor Penguins in Antarctica where the phenomenon of interest may be the diving habits of this species after the eggs hatch. As an other example we can attach sensors to a double pendulum and find what it is doing when we give it a spin.

Based on such observational data the experimenter typically tries to learn about the behaviour of the system from the sensor data to estimate parameters, test hypotheses, etc. Such types of experiments are typically performed by scientists in various fields of science, such as, astronomy, biology, chemistry, geology, physics, etc.

#### Chaotic Time Series of a Double Pendulum




Figure 4.14: Double Pendulum

Sensors called *optical encoders* have been attached to the top end of each arm of a chaotic double pendulum in order to obtain the angular position of each arm through time as shown in Figure 4.14. Time series of the angular position of each arm for two trajectories that were initialized very similarly, say the angles of each arm of the double pendulum are almost the same at the initial time of release. Note how quickly the two trajectories diverge! System with such a sensitivity to initial conditions are said to be *chaotic*.

**Labwork 89 (A Challenging Task)** Try this if you are interested. Read any of the needed details about the design and fabrication of the double pendulum at <http://www.math.canterbury.ac.nz/~r.sainudiin/lmse/double-pendulum/>. Then use MATLAB to generate a plot similar to Figure 4.14(C) using time series data of trajectory 1 and trajectory 2 linked from the bottom of the above URL.

# Chapter 5

## Simulation

“Anyone who considers arithmetical methods of producing random digits is, of course, in a state of sin.” — John von Neumann (1951)

### 5.1 Physical Random Number Generators

Physical devices such as the BINGO machine demonstrated in class can be used to produce an integer uniformly at random from a finite set of possibilities. Such “ball bouncing machines” used in the British national lottery as well as the New Zealand LOTTO are complex nonlinear systems that are extremely sensitive to initial conditions (“chaotic” systems) and are physical approximations of the probability model called a “well-stirred urn” or an equi-probable de Moivre( $1/k, \dots, 1/k$ ) random variable.

Let us look at the New Zealand LOTTO draws at <http://lotto.nzpages.co.nz/statistics.html> and convince ourselves that all fourty numbers  $\{1, 2, \dots, 39, 40\}$  seem to be drawn uniformly at random. The British lottery animation at <http://understandinguncertainty.org/node/39> shows how often each of the 49 numbers came up in the first 1240 draws. Are these draws really random? We will answer these questions in the sequel (see <http://understandinguncertainty.org/node/40> if you can’t wait).

### 5.2 Pseudo-Random Number Generators

Our probability model and the elementary continuous Uniform(0, 1) RV are built from the abstract concept of a random variable over a probability triple. A direct implementation of these ideas on a computing machine is not possible. In practice, random variables are typically simulated by **deterministic** methods or algorithms. Such algorithms generate sequences of numbers whose behavior is virtually indistinguishable from that of truly random sequences. In computational statistics, simulating realisations from a given RV is usually done in two distinct steps. First, sequences of numbers that imitate independent and identically distributed (IID) Uniform(0, 1) RVs are generated. Second, appropriate transformations are made to these imitations of IID Uniform(0, 1) random variates in order to imitate IID random variates from other random variables or other random structures. These two steps are essentially independent and are studied by two non-overlapping groups of researchers. The formal term **pseudo-random number generator** (PRNG) or simply **random number generator** (RNG) usually refers to some deterministic algorithm used in the first step to produce pseudo-random numbers (PRNs) that imitate IID Uniform(0, 1) random variates.

In the following chapters, we focus on transforming IID Uniform(0, 1) variates to other non-uniform variates. In this chapter, we focus on the art of imitating IID Uniform(0, 1) variates using simple deterministic rules.

### 5.2.1 Linear Congruential Generators

The following procedure introduced by D. H. Lehmer in 1949 [*Proc. 2nd Symp. on Large-Scale Digital Calculating Machinery, Harvard Univ. Press, Cambridge, Mass., 1951, 141–146*] gives the simplest popular PRNG that can be useful in many statistical situations if used wisely.

---

**Algorithm 2** Linear Congruential Generator (LCG)

---

1: *input:* five suitable integers:

1.  $m$ , the modulus;  $0 < m$
2.  $a$ , the multiplier;  $0 \leq a < m$
3.  $c$ , the increment;  $0 \leq c < m$
4.  $x_0$ , the seed;  $0 \leq x_0 < m$
5.  $n$ , the number of desired pseudo-random numbers

2: *output:*  $(x_0, x_1, \dots, x_{n-1})$ , the linear congruential sequence of length  $n$

3: **for**  $i = 1$  to  $n - 1$  **do**

4:    $x_i \leftarrow (ax_{i-1} + c) \bmod m$

5: **end for**

6: *return:*  $(x_1, x_2, \dots, x_n)$

---

In order to implement LCGs we need to be able to do high precision exact integer arithmetic in MATLAB. We employ the Module `vpi` to implement variable precision integer arithmetic. You need to download this module for the next Labwork.

**Labwork 90 (Generic Linear Congruential Sequence)** Let us implement Algorithm 2 in MATLAB as follows.

---

```
function x = LinConGen(m,a,c,x0,n)
% Returns the linear congruential sequence
% Needs variable precision integer arithmetic in MATLAB!!!
% Usage: x = LinConGen(m,a,c,x0,n)
% Tested:
% Knuth3.3.4Table1.Line1: LinConGen(100000001,23,0,01234,10)
% Knuth3.3.4Table1.Line5: LinConGen(256,137,0,01234,10)
% Knuth3.3.4Table1.Line20: LinConGen(2147483647,48271,0,0123456,10)
% Knuth3.3.4Table1.Line21: LinConGen(2147483399,40692,0,0123456,10)

x=zeros(1,n); % initialize an array of zeros
X=vpi(x0); % X is a variable precision integer seed
x(1) = double(X); % convert to double
A=vpi(a); M=vpi(m); C=vpi(c); % A,M,C as variable precision integers
for i = 2:n % loop to generate the Linear congruential sequence
    % the linear congruential operation in variable precision integer
    % arithmetic
    % comment out the next ';' to get integer output
    X=mod(A * X + C, M);
```

---

```
x(i) = double(X); % convert to double
end
```

We can call it for some arbitrary input arguments as follows:

```
>> LinConGen(13,12,11,10,12)
ans =
    10     1     10     1     10     1     10     1     10     1     10     1
>> LinConGen(13,10,9,8,12)
ans =
     8     11     2     3     0     9     8     11     2     3     0     9
```

and observe that the generated sequences are not “random” for input values of  $(m, a, c, x_0, n)$  equalling  $(13, 12, 11, 10, 12)$  or  $(13, 10, 9, 8, 12)$ . Thus, we need to do some work to determine the *suitable* input integers  $(m, a, c, x_0, n)$ .


**Labwork 91 (LCG with period length of 32)** Consider the linear congruential sequence with  $(m, a, c, x_0, n) = (256, 137, 0, 123, 257)$  with period length of only  $32 < m = 256$ . We can visualise the sequence as plots in Figure 5.1 after calling the following M-file.

---

```
LinConGenKnuth334T1L5Plots.m
LCGSeq=LinConGen(256,137,0,123,257)
subplot(1,3,1)
plot(LCGSeq,'+')
axis([0 256 0 256]); axis square
LCGSeqIn01=LCGSeq ./ 256
subplot(1,3,2)
plot(LCGSeqIn01,'+')
axis([0 256 0 1]); axis square
subplot(1,3,3)
hist(LCGSeqIn01,15)
axis square
```

---

Figure 5.1: The linear congruential sequence of  $\text{LinConGen}(256, 137, 0, 123, 257)$  with non-maximal period length of 32 as a line plot over  $\{0, 1, \dots, 256\}$ , scaled over  $[0, 1]$  by a division by 256 and a histogram of the 256 points in  $[0, 1]$  with 15 bins.



### Choosing the *suitable* magic input $(m, a, c, x_0, n)$

The linear congruential generator is a special case of a *discrete dynamical system*:

$$x_i = f(x_{i-1}), \quad f : \{0, 1, 2, \dots, m-1\} \rightarrow \{0, 1, 2, \dots, m-1\} \text{ and } f(x_{i-1}) = (ax_{i-1} + c) \pmod{m}.$$

Since  $f$  maps a the finite set  $\{1, 2, \dots, m-1\}$  into itself, such systems are bound to have a repeating cycle of numbers called the **period**. In Labwork 90, the generator `LinConGen(13,12,11,10,12)` has period  $(10, 1)$  of length 2, the generator `LinConGen(13,10,9,8,12)` has period  $(8, 11, 2, 3, 0, 9)$  of length 6 and the generator `LinConGen(256,137,0,123,257)` has a period of length 32. All these generators have a non-maximal period length less than their modulus  $m$ . A good generator should have a maximal period of  $m$ . Let us try to implement a generator with a maximal period of  $m = 256$ .

The period of a general LCG is at most  $m$ , and for some choices of  $a$  the period can be much less than  $m$  as shown in the examples considered earlier. The LCG will have a full period if and only if:

1.  $c$  and  $m$  are relatively prime,
2.  $a - 1$  is divisible by all prime factors of  $m$ ,
3.  $a - 1$  is a multiple of 4 if  $m$  is a multiple of 4

**Labwork 92 (LCG with maximal period length of 256)** Consider the linear congruential sequence with  $(m, a, c, x_0, n) = (256, 137, 123, 13, 256)$ . First check that these parameters do indeed satisfy the three condition above and therefore can produce the maximal period length of only  $m = 256$ . Modify the input parameter to `LinConGen` and repeat Labwork 91 in order to first produce a sequence of length 257. Do you see that the period is of maximal length of 256 as opposed to the generator of Labwork 91? Next produce a Figure to visualise the sequence as done in Figure 5.1.

A useful sequence should clearly have a relatively long period, say at least  $2^{30}$ . Therefore, the **modulus  $m$  has to be rather large** because the **period** cannot have more than  $m$  elements. Moreover, the quality of pseudo-random numbers of a LCG is extremely sensitive to the choice of  $m$ ,  $a$  and  $c$  even if the maximal period is attained. The next example illustrates this point.

**Labwork 93 (The infamous RANDU)** RANDU is an infamous LCG, which has been used since the 1960s. It is widely considered to be one of the most ill-conceived random number generators designed. Notably, it fails the **spectral test** badly for dimensions greater than 2. The following commands help visualise the sequence of first 5001/3 triplets  $(x_i, x_{i+1}, x_{i+2})$  seeded from  $x_0 = 1$  (Figure 5.2). Read `help reshape` and `help plot3`.


```
>> x=reshape( (LinConGen(2147483648,65539,0,1,5001)./ 2147483648) ,3,[]);
>> plot3(x(1,:),x(2,:),x(3,:),'.'
```

**Labwork 94 (Fishman20 and Lecuyer21 LCGs)** The following two LCGs are recommended in Knuth's Art of Computer Programming, vol. 2, for generating pseudo-random numbers for simple simulation tasks.

```
>> LinConGen(2147483647,48271,0,08787458,10) ./ 2147483647
ans =    0.0041    0.5239    0.0755    0.7624    0.6496    0.0769    0.9030    0.4259    0.9948    0.8868

>> LinConGen(2147483399,40692,0,01234567,10) ./ 2147483399
ans =    0.0006    0.3934    0.4117    0.7893    0.3913    0.6942    0.6790    0.3337    0.2192    0.1883
```

Figure 5.2: The LCG called RANDU with  $(m, a, c) = (2147483648, 65539, 0)$  has strong correlation between three consecutive points as:  $x_{i+2} = 6x_{k+1} - 9x_k$ . The two plots are showing  $(x_i, x_{i+1}, x_{i+2})$  from two different view points. .



The number of random numbers  $n$  should at most be about  $m/1000$  in order to avoid the future sequence from behaving like the past. Thus, if  $m = 2^{32}$  then a new generator, with a new suitable set of  $(m, a, c, x_0, n)$  should be adopted after the consumption of every few million pseudo-random numbers.

The LCGs are the least sophisticated type of PRNGs. They are easier to understand but are not recommended for intensive simulation purposes. The next section briefly introduces a more sophisticated PRNG we will be using in this course. Moreover our implementation of LCGs using the variable precision integer package is extremely slow in MATLAB and is only of pedagogical interest.

### 5.2.2 Generalized Feedback Shift Register and the “Mersenne Twister” PRNG

The following generator termed `twister` in MATLAB is recommended for use in simulation. It has extremely long periods, low correlation and passes most statistical tests (the DIEHARD statistical tests). The `twister` random number generator of Makoto Matsumoto and Takuji Nishimura is a variant of the twisted generalized feedback shift-register algorithm, and is known as the “Mersenne Twister” generator [Makoto Matsumoto and Takuji Nishimura, *Mersenne Twister: A 623-dimensionally equidistributed uniform pseudorandom number generator*, ACM Transactions on Modeling and Computer Simulation, Vol. 8, No. 1 (Jan. 1998), Pages 3–30]. It has a Mersenne prime period of  $2^{19937} - 1$  (about  $10^{6000}$ ) and is **equi-distributed** in 623 dimensions. It uses 624 words of state per generator and is comparable in speed to the other generators. The recommended default seed is 5489. See <http://www.math.sci.hiroshima-u.ac.jp/~m-mat/MT/emt.html> and [http://en.wikipedia.org/wiki/Mersenne\\_twister](http://en.wikipedia.org/wiki/Mersenne_twister) for details.

Let us learn to implement the MATLAB function that generates PRNs. In MATLAB the function `rand` produces a deterministic PRN sequence. First, read `help rand`. We can generate PRNs as follows.

**Labwork 95 (Calling PRNG in MATLAB)** In MATLAB `rand` is basic PRNG command.

```
>> rand(1,10) % generate a 1 X 10 array of PRNs
ans =
    0.8147    0.9058    0.1270    0.9134    0.6324    0.0975    0.2785    0.5469    0.9575    0.9649
>> rand(1,10) % generate another 1 X 10 array of PRNs
ans =
    0.1576    0.9706    0.9572    0.4854    0.8003    0.1419    0.4218    0.9157    0.7922    0.9595
>> rand('twister',5489) % reset the PRNG to default state Mersenne Twister with seed=5489
>> rand(1,10) % reproduce the first array
ans =
    0.8147    0.9058    0.1270    0.9134    0.6324    0.0975    0.2785    0.5469    0.9575    0.9649
>> rand(1,10) % reproduce the second array
ans =
    0.1576    0.9706    0.9572    0.4854    0.8003    0.1419    0.4218    0.9157    0.7922    0.9595
```

In general, you can use any seed value to initiate your PRNG. You may use the `clock` command to set the seed:

```
>> SeedFromClock=sum(100*clock); % save the seed from clock
>> rand('twister',SeedFromClock) % initialize the PRNG
>> rand(1,10)
ans =
    0.3696    0.3974    0.6428    0.6651    0.6961    0.7311    0.8982    0.6656    0.6991    0.8606
>> rand(2,10)
ans =
    0.3432    0.9511    0.3477    0.1007    0.8880    0.0853    0.6067    0.6976    0.4756    0.1523
    0.5827    0.5685    0.0125    0.1555    0.5551    0.8994    0.2502    0.5955    0.5960    0.5700
>> rand('twister',SeedFromClock) % initialize the PRNG to same SeedFromClock
>> rand(1,10)
ans =
    0.3696    0.3974    0.6428    0.6651    0.6961    0.7311    0.8982    0.6656    0.6991    0.8606
```

**Labwork 96 (3D plots of triplets generated by the “Mersenne Twister”)** Try to find any correlation between triplets generated by the “Mersenne Twister” by rotating the 3D plot generated by the following code:


```
>> rand('twister',1234)
>> x=rand(3,2000); % store PRNs in a 3X2000 matrix named x
>> plot3(x(1,:),x(2,:),x(3,:),'.'
```

Compare this with the 3D plot of triplets from RANDU of Labwork 93. Which of these two PRNGs do you think is “more random” looking? and why?

Change the seed value to the recommended default by the authors and look at the point cloud (in red) relative to the previous point cloud (in blue). Rotate the plots to visualise from multiple angles. Are they still random looking?

```
>> rand('twister',1234)% same seed as before
>> x=rand(3,2000); % store PRNs in a 3X2000 matrix named x
>> rand('twister',5489)% the recommended default seed
>> y=rand(3,2000);% store PRNs seeded by 5489 in a 3X2000 matrix named y
>> plot3(x(1,:),x(2,:),x(3,:),'b.') % plot triplets as blue dots
>> hold on;
>> plot3(y(1,:),y(2,:),y(3,:),'r.') % plot triplets as red dots
```

Figure 5.3: Triplet point clouds from the “Mersenne Twister” with two different seeds (see Lab-work 96). .



### 5.3 Simulation of non-Uniform(0, 1) Random Variables

The Uniform(0, 1) RV of Model 2 forms the foundation for random variate generation and simulation. This is appropriately called the fundamental model or experiment, since every other experiment can be obtained from this one.

Next, we simulate or generate samples from other RVs by making the following two assumptions:

1. independent samples from the Uniform(0, 1) RV can be generated, and
2. real arithmetic can be performed exactly in a computer.

Both these assumptions are, in fact, not true and require a more careful treatment of the subject. We may return to these careful treatments later on.

#### 5.3.1 Inversion Sampler for Continuous Random Variables

**Proposition 54 (Inversion sampler)** Let  $F(x) := \int_{-\infty}^x f(y) dy : \mathbb{R} \rightarrow [0, 1]$  be a continuous DF with density  $f$ , and let its inverse  $F^{[-1]} : [0, 1] \rightarrow \mathbb{R}$  be:

$$F^{[-1]}(u) := \inf\{x : F(x) = u\} .$$

Then,  $F^{[-1]}(U)$  has the distribution function  $F$ , provided  $U$  is a Uniform(0, 1) RV. Recall  $\inf(A)$  or infimum of a set  $A$  of real numbers is the greatest lower bound of every element of  $A$ .

**Proof:** The “one-line proof” of the proposition is due to the following equalities:

$$\mathbf{P}(F^{[-1]}(U) \leq x) = \mathbf{P}(\inf\{y : F(y) = U\} \leq x) = \mathbf{P}(U \leq F(x)) = F(x), \quad \text{for all } x \in \mathbb{R}.$$

This yields the inversion sampler or the inverse (C)DF sampler, where we (i) generate  $u \sim \text{Uniform}(0, 1)$  and (ii) return  $x = F^{[-1]}(u)$ , as formalised by the following algorithm.

---

**Algorithm 3** Inversion Sampler or Inverse (C)DF Sampler


---

- 1: *input*: (1)  $F^{[-1]}(x)$ , inverse of the DF of the target RV  $X$ , (2) the fundamental sampler
  - 2: *initialise*: set the seed, if any, for the fundamental sampler
  - 3: *output*: a sample from  $X$  distributed according to  $F$
  - 4: *draw*  $u \sim \text{Uniform}(0, 1)$
  - 5: *return*:  $x = F^{[-1]}(u)$
- 

This algorithm emphasises the fundamental sampler’s availability in an *input* step, and its set-up needs in an *initialise* step. In the following sections, we will not mention these universal steps; they will be taken for granted. The direct applicability of Algorithm 3 is limited to univariate densities for which the inverse of the cumulative distribution function is explicitly known. The next section will consider some examples.

Recall the  $\text{Uniform}(\theta_1, \theta_2)$  RV of Model 4 with the following PDF, DF and inverse DF. Let us simulate from it using the inversion sampler.

Figure 5.4: A plot of the PDF, DF or CDF and inverse DF of the  $\text{Uniform}(-1, 1)$  RV  $X$ .



**Simulation 97** ( $\text{Uniform}(\theta_1, \theta_2)$ ) To simulate from  $\text{Uniform}(\theta_1, \theta_2)$  RV  $X$  using the Inversion Sampler, we first need to find  $F^{[-1]}(u)$  by solving for  $x$  in terms of  $u = F(x; \theta_1, \theta_2)$ :

$$u = \frac{x - \theta_1}{\theta_2 - \theta_1} \iff x = (\theta_2 - \theta_1)u + \theta_1 \iff F^{[-1]}(u; \theta_1, \theta_2) = \theta_1 + (\theta_2 - \theta_1)u$$

Here is a simple implementation of the Inversion Sampler for the  $\text{Uniform}(\theta_1, \theta_2)$  RV in MATLAB :

```
>> rand('twister', 786); % initialise the fundamental sampler for Uniform(0,1)
>> theta1=-1; theta2=1; % declare values for parameters theta1 and theta2
>> u=rand; % rand is the Fundamental Sampler and u is a sample from it
>> x=theta1+(theta2 - theta1)*u; % sample from Uniform(-1,1) RV
>> disp(x); % display the sample from Uniform[-1,,1] RV
0.5134
```

It is just as easy to draw  $n$  IID samples from  $\text{Uniform}(\theta_1, \theta_2)$  RV  $X$  by transforming  $n$  IID samples from the  $\text{Uniform}(0, 1)$  RV as follows:


```
>> rand('twister',786543); % initialise the fundamental sampler
>> theta1=-83; theta2=1004; % declare values for parameters a and b
>> u=rand(1,5); % now u is an array of 5 samples from Uniform(0,1)
>> x=theta1+(theta2 - theta1)*u; % x is an array of 5 samples from Uniform(-83,1004]) RV
>> disp(x); % display the 5 samples just drawn from Uniform(-83,1004) RV
465.3065 111.4994 14.3535 724.8881 254.0168
```

**Labwork 98 (Inversion Sampler Demo –  $\text{Uniform}(-5, 5)$ )** Let us comprehend the inversion sampler by calling the interactive visual cognitive tool built by Jennifer Harlow under a grant from University of Canterbury's Centre for Teaching and Learning (UCTL):

```
>> guiInversionSampler
```


The M-file `guiInversionSampler.m` will bring a graphical user interface (GUI) as shown in Figure 5.5. The default target distribution is  $\text{Uniform}(-5, 5)$ . Now repeatedly push the “Draw one sample” button several times and comprehend the simulation process. You can press “Draw 100 samples” to really comprehend the inversion sampler in action after 100 samples are drawn and depicted in the density histogram of the accumulating samples. Next try changing the numbers in the “Lower bound” and “Upper bound” boxes in order to alter the parameters  $\theta_1$  and  $\theta_2$  of  $\text{Uniform}(\theta_1, \theta_2)$  RV.

Figure 5.5: Visual Cognitive Tool GUI: Inversion Sampling from  $X \sim \text{Uniform}(-5, 5)$ .



Recall the  $\text{Exponential}(\lambda)$  RV of Model 3. Let us simulate from it using the inversion sampler.

Let us consider the problem of simulating from an  $\text{Exponential}(\lambda)$  RV with realisations in  $\mathbb{R}_+ := [0, \infty) := \{x : x \geq 0, x \in \mathbb{R}\}$  to model the waiting time for a bus at a bus stop.

Figure 5.6: The PDF  $f$ , DF  $F$ , and inverse DF  $F^{[-1]}$  of the Exponential( $\lambda = 1.0$ ) RV.

**Simulation 99** (Exponential( $\lambda$ )) For a given  $\lambda > 0$ , an Exponential( $\lambda$ ) RV has the following PDF  $f$ , DF  $F$  and inverse DF  $F^{[-1]}$ :

$$f(x; \lambda) = \lambda e^{-\lambda x} \quad F(x; \lambda) = 1 - e^{-\lambda x} \quad F^{[-1]}(u; \lambda) = \frac{-1}{\lambda} \log_e(1 - u) \quad (5.1)$$

We write the natural logarithm  $\log_e$  as  $\log$  for notational simplicity. An implementation of the Inversion Sampler for Exponential( $\lambda$ ) as a function in the M-file:

---

```
function x = ExpInvCDF(u,lambda);
% Return the Inverse CDF of Exponential(lambda) RV X
% Call Syntax: x = ExpInvCDF(u,lambda);
%           ExpInvCDF(u,lambda);
% Input      : lambda = rate parameter,
%             u = array of numbers in [0,1]
% Output     : x
x=-(1/lambda) * log(1-u);
```

---

We can simply call the function to draw a sample from, say the Exponential( $\lambda = 1.0$ ) RV by:

```
lambda=1.0; % some value for lambda
u=rand; % rand is the Fundamental Sampler
ExpInvCDF(u,lambda) % sample from Exponential(1) RV via function in ExpInvCDF.m
```

Because of the following (recall Example 41):

$$U \sim \text{Uniform}(0, 1) \implies -U \sim \text{Uniform}(-1, 0) \implies 1 - U \sim \text{Uniform}(0, 1),$$

we could save a subtraction operation in the above algorithm by replacing  $-(1/\lambda) * \log(1-u)$  by  $-(1/\lambda) * \log(u)$ . Recall that the transformation of  $U \sim \text{Uniform}(0, 1)$  by  $X = -(1/\lambda) \log(U)$  is exactly how we defined  $X$  as the Exponential( $\lambda$ ) RV in Model 3. This is implemented as the following function.

---

```
function x = ExpInvSam(u,lambda);
% Return the Inverse CDF based Sample from Exponential(lambda) RV X
% Call Syntax: x = ExpInvSam(u,lambda);
%           or ExpInvSam(u,lambda);
% Input      : lambda = rate parameter,
%             u = array of numbers in [0,1] from Uniform[0,1] RV
% Output     : x
x=-(1/lambda) * log(u);
```

---


```
>> rand('twister',46678); % initialise the fundamental sampler
>> Lambda=1.0; % declare Lambda=1.0
>> x=ExpInvSam(rand(1,5),Lambda); % pass an array of 5 Uniform(0,1) samples from rand
>> disp(x); % display the Exponential(1.0) distributed samples
0.5945    2.5956    0.9441    1.9015    1.3973
```

**Labwork 100 (Inversion Sampler Demo – Exponential(0.5))** Let us understand the inversion sampler by calling the interactive visual cognitive tool:

```
>> guiInversionSampler
```

The M-file `guiInversionSampler.m` will bring a graphical user interface (GUI) as shown in Figure 5.7. First change the target distribution from the default Uniform( $-5, 5$ ) to Exponential( $0.5$ ) from the drop-down menu. Now push the “Draw 10 samples” button and comprehend the simulation process. Next try changing the “Rate Parameter” from  $0.5$  to  $10.0$  for example and generate several inversion samples and see the density histogram of the accumulating samples. You can press “Draw one sample” to really comprehend the inversion sampler in action one step at a time.

Figure 5.7: Visual Cognitive Tool GUI: Inversion Sampling from  $X \sim \text{Exponential}(0.5)$ .



It is straightforward to do replicate experiments. Consider the experiment of drawing five independent samples from the  $\text{Exponential}(\lambda = 1.0)$  RV. Suppose we want to repeat or replicate this experiment seven times and find the sum of the five outcomes in each of these replicates. Then we may do the following:

```
>> rand('twister',1973); % initialise the fundamental sampler
>> % store 7 replications of 5 IID draws from Exponential(1.0) RV in array a
>> lambda=1.0; a= -1/lambda * log(rand(5,7)); disp(a);
0.7267    0.3226    1.2649    0.4786    0.3774    0.0394    1.8210
1.2698    0.4401    1.6745    1.4571    0.1786    0.4738    3.3690
```

```

0.4204    0.1219    2.2182    3.6692    0.9654    0.0093    1.7126
2.1427    0.1281    0.8500    1.4065    0.1160    0.1324    0.2635
0.6620    1.1729    0.6301    0.6375    0.3793    0.6525    0.8330
>> %sum up the outcomes of the sequence of 5 draws in each replicate
>> s=sum(a); disp(s);
5.2216    2.1856    6.6378    7.6490    2.0168    1.3073    7.9990

```

**Labwork 101 (Next seven buses at your bus-stop)** Consider the problem of modelling the arrival of buses at a bus stop. Suppose that the time between arrivals is an  $\text{Exponential}(\lambda = 0.1)$  RV  $X$  with a mean inter-arrival time of  $1/\lambda = 10$  minutes. Suppose you go to your bus stop and zero a stop-watch. Simulate the times of arrival for the next seven buses as indicated by your stop-watch. Seed the fundamental sampler by your Student ID (eg. if your ID is 11424620 then type `rand('twister', 11424620)`; just before the simulation). Hand in the code with the arrival times of the next seven buses at your ID-seeded bus stop.

The support of the  $\text{Exponential}(\lambda)$  RV is  $\mathbb{R}_+ := [0, \infty)$ . Let us consider a RV built by mirroring the  $\text{Exponential}(\lambda)$  RV about the origin with the entire real line as its support.

**Model 9 (Laplace( $\lambda$ ) or Double Exponential( $\lambda$ ) RV)** If a RV  $X$  is equally likely to be either positive or negative with an exponential density, then the Laplace( $\lambda$ ) or Double Exponential( $\lambda$ ) RV, with the rate parameter  $\lambda > 0, \lambda \in \mathbb{R}$ , may be used to model it. The density function for the Laplace( $\lambda$ ) RV given by  $f(x; \lambda)$  is

$$f(x; \lambda) = \frac{\lambda}{2} e^{-\lambda|x|} = \begin{cases} \frac{\lambda}{2} e^{\lambda x} & \text{if } x < 0 \\ \frac{\lambda}{2} e^{-\lambda x} & \text{if } x \geq 0 \end{cases}. \quad (5.2)$$

Let us define the sign of a real number  $x$  by

$$\text{sign}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}.$$

Then, the DF of the Laplace( $\lambda$ ) RV  $X$  is

$$F(x; \lambda) = \int_{-\infty}^x f(y; \lambda) dy = \frac{1}{2} \left( 1 + \text{sign}(x) \left( 1 - e^{-\lambda|x|} \right) \right), \quad (5.3)$$

and its inverse DF is

$$F^{[-1]}(u; \lambda) = -\frac{1}{\lambda} \text{sign} \left( u - \frac{1}{2} \right) \log \left( 1 - 2 \left| u - \frac{1}{2} \right| \right), \quad u \in [0, 1] \quad (5.4)$$

**Mean and Variance of Laplace( $\lambda$ ) RV  $X$ :** Show that the mean of a Laplace( $\lambda$ ) RV  $X$  is

$$\mathbf{E}(X) = \int_0^\infty x f(x; \lambda) dx = \int_0^\infty x \frac{\lambda}{2} e^{-\lambda|x|} dx = 0,$$

and the variance is

$$\mathbf{V}(X) = \left( \frac{1}{\lambda} \right)^2 + \left( \frac{1}{\lambda} \right)^2 = 2 \left( \frac{1}{\lambda} \right)^2.$$


Note that the mean is 0 due to the symmetry of the density about 0 and the variance is twice that of the  $\text{Exponential}(\lambda)$  RV.

**Labwork 102 (Rejection Sampler Demo – Laplace(5))** Let us comprehend the rejection sampler by calling the interactive visual cognitive tool:

```
>> guiInversionSampler
```

The M-file `guiInversionSampler.m` will bring a graphical user interface (GUI) as shown in Figure 5.8. Using the drop-down menu change from the default target distribution Uniform( $-5, 5$ ) to Laplace(5). Now repeatedly push the “Draw one sample” button several times and comprehend the simulation process. You can also press “Draw 1000 samples” and see the density histogram of the generated samples. Next try changing the numbers in the “Rate parameter” box from 5.00 to 1.00 in order to alter the parameter  $\lambda$  of Laplace( $\lambda$ ) RV. If you are more adventurous then try to alter the number in the “Location parameter” box from 0.00 to some thing else, say 10.00. Although our formulation of Laplace( $\lambda$ ) implicitly had a location parameter of 0.00, we can easily introduce a location parameter  $\mu$  into the PDF. With a pencil and paper try to rewrite the PDF in (5.2) with an additional location parameter  $\mu$ .

Figure 5.8: Visual Cognitive Tool GUI: Inversion Sampling from  $X \sim \text{Laplace}(5)$ .



**Simulation 103 (Laplace( $\lambda$ ))** Here is an implementation of an inversion sampler to draw IID samples from a Laplace( $\lambda$ ) RV  $X$  by transforming IID samples from the Uniform( $0, 1$ ) RV  $U$ :

---

```
function x = LaplaceInvCDF(u,lambda);
% Call Syntax: x = LaplaceInvCDF(u,lambda);
%               or LaplaceInvCDF(u,lambda);
%
% Input      : lambda = rate parameter > 0,
%               u = an 1 X n array of IID samples from Uniform[0,1] RV
% Output     : an 1Xn array x of IID samples from Laplace(lambda) RV
%               or Inverse CDF of Laplace(lambda) RV
x=-(1/lambda)*sign(u-0.5) .* log(1-2*abs(u-0.5));
```

---

We can simply call the function to draw a sample from, say the Laplace( $\lambda = 1.0$ ) RV by

```
>> lambda=1.0; % some value for lambda
>> rand('twister',6567); % initialize the fundamental sampler
>> u=rand(1,5); % draw 5 IID samples from Uniform(0,1) RV
>> disp(u);
0.6487    0.9003    0.3481    0.6524    0.8152

>> x=LaplaceInvCDF(u,lambda); % draw 5 samples from Laplace(1) RV using inverse CDF
>> disp(x);
0.3530    1.6127   -0.3621    0.3637    0.9953
```

Next, let us become familiar with an RV for which the expectation does not exist.

**Model 10 (Cauchy)** The density of the Cauchy RV  $Y$  is:

$$f(y) = \frac{1}{\pi(1+y^2)}, \quad -\infty < y < \infty , \quad (5.5)$$

and its DF is:

$$F(y) = \frac{1}{\pi} \tan^{-1}(y) + \frac{1}{2} . \quad (5.6)$$

Randomly spinning a LASER emitting improvisation of “Darth Maul’s double edged lightsaber” that is centered at  $(1, 0)$  in the plane  $\mathbb{R}^2$  and recording its intersection with the  $y$ -axis, in terms of the  $y$  coordinates of the point  $(0, y)$ , gives rise to the *Standard Cauchy* RV.

The Cauchy RV  $Y$  can be derived from a RV  $X \sim \text{Uniform}(-\pi/2, \pi/2)$  by the simple transformation  $Y = \tan(X)$  for the above construction. Since  $\tan(x)$  is one-to-one and monotone on the range of  $X$  given by  $(-\pi/2, \pi/2)$ , we can use the change of variable formula in Equation 3.10 to obtain the PDF  $f_Y(y)$  from the PDF  $f_X(x) = \frac{1}{\pi} \mathbb{1}_{(-\pi/2,\pi/2)}(x)$  as follows:

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| = f_X(\tan^{-1}(y)) \left| \frac{d}{dy} \tan^{-1}(y) \right| = \frac{1}{\pi} \left| \frac{1}{1+y^2} \right|$$

**Example 104 (Mean of Cauchy RV)** The expectation of the Cauchy RV  $X$ , obtained via integration by parts (set  $u = x$  and  $v = \tan^{-1}(x)$ ) does not exist, since:


$$\int |x| dF(x) = \frac{2}{\pi} \int_0^\infty \frac{x}{1+x^2} dx = (x \tan^{-1}(x)]_0^\infty - \int_0^\infty \tan^{-1}(x) dx = \infty . \quad (5.7)$$

Variance and higher moments cannot be defined when the expectation itself is undefined.

**Labwork 105 (Inversion Sampler Demo – Cauchy)** Let us comprehend the inversion sampler by calling the interactive visual cognitive tool:

```
>> guiInversionSampler
```

The M-file `guiInversionSampler.m` will bring a graphical user interface (GUI) as shown in Figure 5.9. Using the drop-down menu change from the default target distribution  $\text{Uniform}(-5, 5)$  to Cauchy. Now repeatedly push the “Draw one sample” button several times and comprehend the simulation process. You can also press “Draw 10 samples” several times and see the density histogram of the generated samples. Next try changing the numbers in the “Scale parameter” and “Location Parameter” boxes from the default values of 1.00 and 0.00, respectively. Although our

Figure 5.9: Visual Cognitive Tool GUI: Inversion Sampling from  $X \sim \text{Cauchy}$ .

formulation of Cauchy RV is also called *Standard Cauchy* as it implicitly had a location parameter of 0.00 and scale parameter of 1. With a pencil and paper (in conjunction with a wikipedia search if you have to) try to rewrite the PDF in (5.5) with an additional location parameter  $\mu$  and scale parameter  $\sigma$ .

**Simulation 106 (Cauchy)** We can draw  $n$  IID samples from the Cauchy RV  $X$  by transforming  $n$  IID samples from  $\text{Uniform}(0, 1)$  RV  $U$  using the inverse DF as follows:

```
>> rand('twister',2435567); % initialise the fundamental sampler
>> u=rand(1,5); % draw 5 IID samples from Uniform(0,1) RV
>> disp(u);
0.7176    0.6655    0.9405    0.9198    0.2598
>> x=tan(pi * u); % draw 5 samples from Standard cauchy RV using inverse CDF
>> disp(x); % display the samples in x
-1.2272   -1.7470   -0.1892   -0.2575   1.0634
```

Recall that the mean of the Cauchy RV  $X$  does not exist since  $\int |x| dF(x) = \infty$  (5.7). We will investigate this in Labwork 107.

**Labwork 107 (Running mean of the Standard Cauchy RV)** Let us see what happens when we plot the running sample mean for an increasing sequence of IID samples from the Standard Cauchy RV  $X$  by implementing the following script file:

---

```
% PlotStandardCauchyRunningMean.m
% script to plot the oscillating running mean of Std Cauchy samples
% relative to those for the Uniform(0,10) samples
rand('twister',25567); % initialize the fundamental sampler
for i=1:5
N = 10^5; % maximum sample size
u=rand(1,N); % draw N IID samples from Uniform(0,1)
```


```

x=tan(pi * u);      % draw N IID samples from Standard cauchy RV using inverse CDF
n=1:N;              % make a vector n of current sample size [1 2 3 ... N-1 N]
CSx=cumsum(x); % CSx is the cumulative sum of the array x (type 'help cumsum')
% Runnign Means <- vector division of cumulative sum of samples by n
RunningMeanStdCauchy = CSx ./ n; % Running Mean for Standard Cauchy samples
RunningMeanUnif010 = cumsum(u*10.0) ./ n; % Running Mean for Uniform(0,10) samples
semilogx(n, RunningMeanStdCauchy) %
hold on;
semilogx(n, RunningMeanUnif010, 'm')
end
xlabel('n = sample size');
ylabel('Running mean from n samples')

```

---

Figure 5.10: Unending fluctuations of the running means based on  $n$  IID samples from the Standard Cauchy RV  $X$  in each of five replicate simulations (blue lines). The running means, based on  $n$  IID samples from the Uniform(0, 10) RV, for each of five replicate simulations (magenta lines).



The resulting plot is shown in Figure 5.10. Notice that the running means or the sample mean of  $n$  samples as a function of  $n$ , for each of the five replicate simulations, never settles down to a particular value. This is because of the “thick tails” of the density function for this RV which produces extreme observations. Compare them with the running means, based on  $n$  IID samples from the Uniform(0, 10) RV, for each of five replicate simulations (magenta lines). The latter sample means have settled down stably to the mean value of 5 after about 700 samples.

### 5.3.2 Inversion Sampler for Discrete Random Variables

Next, consider the problem of **sampling from a random variable  $X$  with a discontinuous or discrete DF** using the inversion sampler. We need to define the inverse more carefully here.

**Proposition 55 (Inversion sampler with compact support)** Let the support of the RV  $X$  be over some real interval  $[a, b]$  and let its inverse DF be defined as follows:

$$F^{[-1]}(u) := \inf\{x \in [a, b] : F(x) \geq u, 0 \leq u \leq 1\} .$$

If  $U \sim \text{Uniform}(0, 1)$  then  $F^{[-1]}(U)$  has the DF  $F$ , i.e.  $F^{[-1]}(U) \sim F \sim X$ .

**Proof:** The proof is a consequence of the following equalities:

$$\mathbf{P}(F^{[-1]}(U) \leq x) = \mathbf{P}(U \leq F(x)) = F(x) := \mathbf{P}(X \leq x)$$

**Simulation 108** ( $\text{Bernoulli}(\theta)$ ) Consider the problem of simulating from a  $\text{Bernoulli}(\theta)$  RV based on an input from a  $\text{Uniform}(0, 1)$  RV. Recall that  $\lfloor x \rfloor$  (called the ‘floor of  $x$ ’) is the largest integer that is smaller than or equal to  $x$ , e.g.  $\lfloor 3.8 \rfloor = 3$ . Using the floor function, we can simulate a  $\text{Bernoulli}(\theta)$  RV  $X$  as follows:

```
>> theta = 0.3; % set theta = Prob(X=1)
% return x -- floor(y) is the largest integer less than or equal to y
>> x = floor(rand + theta) % rand is the Fundamental Sampler
>> disp(x) % display the outcome of the simulation
0
>> n=10; % set the number of IID Bernoulli(theta=0.3) trials you want to simulate
>> x = floor(rand(1,n)+theta); % vectorize the operation
>> disp(x) % display the outcomes of the simulation
0     0     1     0     0     0     0     0     1     1
```

Again, it is straightforward to do replicate experiments, e.g. to demonstrate the Central Limit Theorem for a sequence of  $n$  IID  $\text{Bernoulli}(\theta)$  trials.

```
>> % a demonstration of Central Limit Theorem --
>> % the sample mean of a sequence of n IID Bernoulli(theta) RVs is Gaussian(theta,theta(1-theta)/n)
>> theta=0.5; Reps=10000; n=10; hist(sum(floor(rand(n,Reps)+theta))/n)
>> theta=0.5; Reps=10000; n=100; hist(sum(floor(rand(n,Reps)+theta))/n,20)
>> theta=0.5; Reps=10000; n=1000; hist(sum(floor(rand(n,Reps)+theta))/n,30)
```

Recall the Point Mass( $\theta$ ) RV. Formally, we can simulate from it trivially as follows.

**Simulation 109** (Point Mass( $\theta$ )) Let us simulate a sample from the Point Mass( $\theta$ ) RV  $X$ . Since this RV produces the same realisation  $\theta$  we can implement it via the following M-file:

---

```
function x = Sim1PointMass(u,theta)
% Returns one sample from the Point Mass(theta) RV X
% Call Syntax: x = Sim1PointMass(u,theta);
% Input      : u = one uniform random number eg. rand()
%               theta = a real number (scalar)
% Output     : x = sample from X
x=theta;
```

---

Here is call to the function.

```
>> Sim1PointMass(rand(),2)
ans =
    2
>> % % we can use arrayfun to apply Sim1Pointmass to any array of Uniform(0,1) samples
>> arrayfun(@(u)(Sim1PointMass(u,17)),rand(2,10))
ans =
    17    17    17    17    17    17    17    17    17    17
    17    17    17    17    17    17    17    17    17    17
```

Note that it is not necessary to have input IID samples from  $\text{Uniform}(0, 1)$  RV via `rand` in order to draw samples from the Point Mass( $\theta$ ) RV. For instance, an input matrix of zeros can do the job:

```
>> arrayfun(@(u)(Sim1PointMass(u,17)),zeros(2,8))
ans =
    17    17    17    17    17    17    17    17
    17    17    17    17    17    17    17    17
```

Next let us consider a natural generalization of the Bernoulli( $\theta$ ) RV with more than two outcomes.

**Model 11** (de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ )) Given a specific point  $(\theta_1, \theta_2, \dots, \theta_k)$  in the  $k$ -Simplex:

$$\Delta_k := \{ (\theta_1, \theta_2, \dots, \theta_k) : \theta_1 \geq 0, \theta_2 \geq 0, \dots, \theta_k \geq 0, \sum_{i=1}^k \theta_i = 1 \} ,$$

we say that an RV  $X$  is de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) distributed if its PMF is:

$$f(x; \theta_1, \theta_2, \dots, \theta_k) = \begin{cases} 0 & \text{if } x \notin [k] := \{1, 2, \dots, k\}, \\ \theta_x & \text{if } x \in [k]. \end{cases}$$

The DF for de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) RV  $X$  is:

$$F(x; \theta_1, \theta_2, \dots, \theta_k) = \begin{cases} 0 & \text{if } -\infty < x < 1 \\ \theta_1 & \text{if } 1 \leq x < 2 \\ \theta_1 + \theta_2 & \text{if } 2 \leq x < 3 \\ \vdots & \\ \theta_1 + \theta_2 + \dots + \theta_{k-1} & \text{if } k-1 \leq x < k \\ \theta_1 + \theta_2 + \dots + \theta_{k-1} + \theta_k = 1 & \text{if } k \leq x < \infty \end{cases} \quad (5.8)$$

The de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) RV can be thought of as a probability model for “the outcome of rolling a polygonal cylindrical die with  $k$  rectangular faces that are marked with  $1, 2, \dots, k$ ”. The parameters  $\theta_1, \theta_2, \dots, \theta_k$  specify how the die is loaded and may be idealised as specifying the cylinder’s centre of mass with respect to the respective faces. Thus, when  $\theta_1 = \theta_2 = \dots = \theta_k = 1/k$ , we have a probability model for the outcomes of a fair die.

**Mean and variance of de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) RV:** The not too useful expressions for the first two moments of  $X \sim \text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$  are,

$$\mathbf{E}(X) = \sum_{x=1}^k x\theta(x) = \theta_1 + 2\theta_2 + \dots + k\theta_k , \text{ and}$$

$$\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = (\theta_1 + 2^2\theta_2 + \dots + k^2\theta_k) - (\theta_1 + 2\theta_2 + \dots + k\theta_k)^2 .$$

However, if  $X \sim \text{de Moivre}(1/k, 1/k, \dots, 1/k)$ , then the mean and variance for the fair  $k$ -faced die based on Faulhaber’s formula for  $\sum_{i=1}^k i^m$ , with  $m \in \{1, 2\}$ , are,

$$\mathbf{E}(X) = \frac{1}{k} (1 + 2 + \dots + k) = \frac{1}{k} \frac{k(k+1)}{2} = \frac{k+1}{2} ,$$

$$\mathbf{E}(X^2) = \frac{1}{k} (1^2 + 2^2 + \dots + k^2) = \frac{1}{k} \frac{k(k+1)(2k+1)}{6} = \frac{2k^2 + 3k + 1}{6} ,$$

$$\begin{aligned} \mathbf{V}(X) &= \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = \frac{2k^2 + 3k + 1}{6} - \left( \frac{k+1}{2} \right)^2 = \frac{2k^2 + 3k + 1}{6} - \left( \frac{k^2 + 2k + 1}{4} \right) \\ &= \frac{8k^2 + 12k + 4 - 6k^2 - 12k - 6}{24} = \frac{2k^2 - 2}{24} = \frac{k^2 - 1}{12} . \end{aligned}$$

Next we simulate from de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) RV  $X$  via its inverse DF


$$F^{[-1]} : [0, 1] \rightarrow [k] := \{1, 2, \dots, k\},$$

given by:

$$F^{[-1]}(u; \theta_1, \theta_2, \dots, \theta_k) = \begin{cases} 1 & \text{if } 0 \leq u < \theta_1 \\ 2 & \text{if } \theta_1 \leq u < \theta_1 + \theta_2 \\ 3 & \text{if } \theta_1 + \theta_2 \leq u < \theta_1 + \theta_2 + \theta_3 \\ \vdots & \vdots \\ k & \text{if } \theta_1 + \theta_2 + \dots + \theta_{k-1} \leq u < 1 \end{cases} \quad (5.9)$$

When  $k = 2$  in the de Moivre( $\theta_1, \theta_2$ ) model, we have an RV that is similar to the Bernoulli( $p = \theta_1$ ) RV. The DF  $F$  and its inverse  $F^{[-1]}$  for a specific  $\theta_1 = 0.3$  are depicted in Figure 5.11.

Figure 5.11: The DF  $F(x; 0.3, 0.7)$  of the de Moivre( $0.3, 0.7$ ) RV and its inverse  $F^{[-1]}(u; 0.3, 0.7)$ .



First we simulate from an equi-probable special case of the de Moivre( $\theta_1, \theta_2, \dots, \theta_k$ ) RV, with  $\theta_1 = \theta_2 = \dots = \theta_k = 1/k$ .

**Simulation 110** (de Moivre( $1/k, 1/k, \dots, 1/k$ )) The equi-probable *de Moivre*( $1/k, 1/k, \dots, 1/k$ ) RV  $X$  with a discrete uniform distribution over  $[k] = \{1, 2, \dots, k\}$  can be efficiently sampled using the ceiling function. Recall that  $\lceil y \rceil$  is the smallest integer larger than or equal to  $y$ , eg.  $\lceil 13.1 \rceil = 14$ . Algorithm 4 produces samples from the *de Moivre*( $1/k, 1/k, \dots, 1/k$ ) RV  $X$ .

---

**Algorithm 4** Inversion Sampler for de Moivre( $1/k, 1/k, \dots, 1/k$ ) RV

---

1: *input*:

1.  $k$  in de Moivre( $1/k, 1/k, \dots, 1/k$ ) RV  $X$
2.  $u \sim \text{Uniform}(0, 1)$

2: *output*: a sample from  $X$

3: *return*:  $x \leftarrow \lceil ku \rceil$

---

The M-file implementing Algorithm 4 is:

---

```
SimdeMoivreEqui.m
function x = SimdeMoivreEqui(u,k);
% return samples from de Moivre(1/k,1/k,...,1/k) RV X
% Call Syntax: x = SimdeMoivreEqui(u,k);
% Input      : u = array of uniform random numbers eg. rand
%               k = number of equi-probabble outcomes of X
% Output     : x = samples from X
x = ceil(k * u); % ceil(y) is the smallest integer larger than y
%x = floor(k * u); if outcomes are in {0,1,...,k-1}
```

---

Let us use the function `SimdeMoivreEqui` to draw five samples from a fair seven-faced cylindrical dice.

```
>> k=7; % number of faces of the fair dice
>> n=5; % number of trials
>> rand('twister',78657); % initialise the fundamental sampler
>> u=rand(1,n); % draw n samples from Uniform(0,1)
>> % inverse transform samples from Uniform(0,1) to samples
>> % from de Moivre(1/7,1/7,1/7,1/7,1/7,1/7,1/7)
>> outcomes=SimdeMoivreEqui(u,k); % save the outcomes in an array
>> disp(outcomes);
    6      5      5      5      2
```

Now, let us consider the more general problem of implementing a sampler for an arbitrary but specified  $\text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$  RV. That is, the values of  $\theta_i$  need not be equal to  $1/k$ .

**Simulation 111** ( $\text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$ ) We can generate samples from a  $\text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$  RV  $X$  when  $(\theta_1, \theta_2, \dots, \theta_k)$  are specifiable as an input vector via the following algorithm.

---

**Algorithm 5** Inversion Sampler for  $\text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$  RV  $X$

---

1: *input:*

1. parameter vector  $(\theta_1, \theta_2, \dots, \theta_k)$  of  $\text{de Moivre}(\theta_1, \theta_2, \dots, \theta_k)$  RV  $X$ .
2.  $u \sim \text{Uniform}(0, 1)$

2: *output:* a sample from  $X$

3: *initialise:*  $F \leftarrow \theta_1$ ,  $i \leftarrow 1$

4: **while**  $u > F$  **do**

5:    $i \leftarrow i + 1$

6:    $F \leftarrow F + \theta_i$

7: **end while**

8: *return:*  $x \leftarrow i$

---

The M-file implementing Algorithm 5 is:

---

```
SimdeMoivreOnce.m
function x = SimdeMoivreOnce(u,thetas)
% Returns a sample from the de Moivre(thetas=(theta_1,...,theta_k)) RV X
% Call Syntax: x = SimdeMoivreOnce(u,thetas);
%               deMoivreEqui(u,thetas);
% Input      : u = a uniform random number eg. rand
%               thetas = an array of probabilities thetas=[theta_1 ... theta_k]
% Output     : x = sample from X
x=1; % initial index is 1
cum_theta=thetas(x);
```

---

```

while u > cum_theta;
x=x+1;
cum_theta = cum_theta + thetas(x);
end

```

---

Let us use the function `deMoivreEqui` to draw five samples from a fair seven-faced dice.

```

>> k=7; % number of faces of the fair dice
>> n=5; % number of trials
>> rand('twister',78657); % initialise the fundamental sampler
>> Us=rand(1,n); % draw n samples from Uniform(0,1)
>> disp(Us);
    0.8330    0.6819    0.6468    0.6674    0.2577
>> % inverse transform samples from Uniform(0,1) to samples
>> % from de Moivre(1/7,1/7,1/7,1/7,1/7,1/7,1/7)
>> f=[1/7 1/7 1/7 1/7 1/7 1/7 1/7];
>> disp(f);
    0.1429    0.1429    0.1429    0.1429    0.1429    0.1429
>> % use funarray to apply function-handled SimdeMoivreOnce to
>> % each element of array Us and save it in array outcomes2
>> outcomes2=arrayfun(@(u)(SimdeMoivreOnce(u,f)),Us);
>> disp(outcomes2);
    6      5      5      5      2
>> disp(SimdeMoivreEqui(u,k)); % same result using the previous algorithm
    6      5      5      5      2

```

Clearly, Algorithm 5 may be used to sample from any  $\text{de Moivre}(\theta_1, \dots, \theta_k)$  RV  $X$ . We demonstrate this by producing five samples from a randomly generated PMF  $f_2$ .

```

>> rand('twister',1777); % initialise the fundamental sampler
>> f2=rand(1,10); % create an arbitrary array
>> f2=f2/sum(f2); % normalize to make a probability mass function
>> disp(f2); % display the weights of our 10-faced die
    0.0073    0.0188    0.1515    0.1311    0.1760    0.1121    ...
    0.1718    0.1213    0.0377    0.0723
>> disp(sum(f2)); % the weights sum to 1
    1.0000
>> disp(arrayfun(@(u)(SimdeMoivreOnce(u,f2)),rand(5,5))) % the samples from f2 are
    4      3      4      7      3
    6      7      4      5      3
    5      8      7     10      6
    2      3      5      7      7
    6      5      9      5      7

```

Note that the principal work here is the sequential search, in which the mean number of comparisons until success is:

$$1\theta_1 + 2\theta_2 + 3\theta_3 + \dots + k\theta_k = \sum_{i=1}^k i\theta_i$$

For the  $\text{de Moivre}(1/k, 1/k, \dots, 1/k)$  RV, the right-hand side of the above expression is:

$$\sum_{i=1}^k i \frac{1}{k} = \frac{1}{k} \sum_{i=1}^k i = \frac{1}{k} \frac{k(k+1)}{2} = \frac{k+1}{2},$$

indicating that the average-case efficiency is linear in  $k$ . This linear dependence on  $k$  is denoted by  $O(k)$ . In other words, as the number of faces  $k$  increases, one has to work linearly harder to get samples from  $\text{de Moivre}(1/k, 1/k, \dots, 1/k)$  RV using Algorithm 5. Using the simpler Algorithm 4, which exploits the fact that all values of  $\theta_i$  are equal, we generated samples in constant time, which is denoted by  $O(1)$ .

**Simulation 112** (Geometric( $\theta$ )) We can simulate a sample  $x$  from a Geometric( $\theta$ ) RV  $X$  using the following simple algorithm:

$$x \leftarrow \lfloor \log(u)/\log(1-\theta) \rfloor, \quad \text{where, } u \sim \text{Uniform}(0, 1).$$

To verify that the above procedure is valid, note that:

$$\begin{aligned} \lfloor \log(U)/\log(1-\theta) \rfloor = x &\iff x \leq \log(U)/\log(1-\theta) < x+1 \\ &\iff x \leq \log_{1-\theta}(U) < x+1 \\ &\iff (1-\theta)^x \geq U > (1-\theta)^{x+1} \end{aligned}$$

The inequalities are reversed since the base being exponentiated is  $1-\theta \leq 1$ . The uniform event  $(1-\theta)^x \geq U > (1-\theta)^{x+1}$  happens with the desired probability:

$$(1-\theta)^x - (1-\theta)^{x+1} = (1-\theta)^x(1-(1-\theta)) = \theta(1-\theta)^x =: f(x; \theta), \quad X \sim \text{Geometric}(\theta).$$

We implement the sampler to generate samples from Geometric( $\theta$ ) RV with  $\theta = 0.5$ , for instance:

```
>> theta=0.5; u=rand(); % choose some theta and uniform(0,1) variate
>> % Simulate from a Geomertic(theta) RV
>> floor(log (u) / log (1 - theta))
ans =
    0
>> floor(log ( rand(1,10) ) / log (1 - 0.5)) % theta=0.5, 10 samples
ans =
    0    0    1    0    2    1    0    0    0    0
```

### Labwork 113 (Compare PDF to the relative frequency histogram of simulated Geometric( $\theta$ ) RV)

It is a good idea to make a relative frequency histogram of a simulation algorithm and compare that to the PDF of the discrete RV we are simulating from. We use the following script to create Figure 3.8:

---

```
theta=0.5;
SampleSize=1000;
% simulate 1000 samples from Geometric(theta) RV
Samples=floor(log(rand(1,SampleSize))/ log (1-theta));
Xs = 0:10; % get some values for x
RelFreqs=hist(Samples,Xs)/SampleSize; % relative frequencies of Samples
stem(Xs,theta*((1-theta) .^ Xs),'*')% PDF of Geometric(theta) over Xs
hold on;
plot(Xs,RelFreqs,'o')% relative frequence histogram
RelFreqs100=hist(Samples(1:100),Xs)/100; % Relative Frequencies of first 100 samples
plot(Xs,RelFreqs100,'x')
legend('PDF of Geometric(0.5)', 'Relative freq. hist. (1000 samples)', ...
'Relative freq. hist. (100 samples)')
```

---

Let us simulate from the Binomial( $n, \theta$ ) RV of Model 7.

**Labwork 114 (Binomial coefficient)** We may implement the MATLAB function `BinomialCoefficient` to compute:

$$\binom{n}{x} = \frac{n!}{x!(n-x)!} = \frac{n(n-1)(n-2)\dots(n-x+1)}{x(x-1)(x-2)\dots(2)(1)} = \frac{\prod_{i=(n-x+1)}^n i}{\prod_{i=2}^x i},$$

with the following M-file:

---

```
function BC = BinomialCoefficient(n,x)
% returns the binomial coefficient of n choose x
% i.e. the combination of n objects taken x at a time
% x and n are scalar integers and 0 <= x <= n
NminusX = n-x;
NumeratorPostCancel = prod(n:-1:(max([NminusX,x])+1)) ;
DenominatorPostCancel = prod(2:min([NminusX, x]));
BC = NumeratorPostCancel/DenominatorPostCancel;
```

---

and call `BinomialCoefficient` in the function `BinomialPdf` to compute the PDF  $f(x; n, \theta)$  of the  $\text{Binomial}(n, \theta)$  RV  $X$  as follows:

---

```
function fx = BinomialPdf(x,n,theta)
% Binomial probability mass function. Needs BinomialCoefficient(n,x)
% f = BinomialPdf(x,n,theta)
% f is the prob mass function for the Binomial(x;n,theta) RV
% and x can be array of samples.
% Values of x are integers in [0,n] and theta is a number in [0,1]
fx = zeros(size(x));
fx = arrayfun(@(xi)(BinomialCoefficient(n,xi)),x);
fx = fx .* (theta .^ x) .* (1-theta) .^ (n-x);
```

---

For example, we can compute the desired PDF for an array of samples  $x$  from  $\text{Binomial}(8, 0.5)$  RV  $X$ , as follows:

```
>> x=0:1:8
x = 0 1 2 3 4 5 6 7 8
>> BinomialPdf(x,8,0.5)
ans = 0.0039 0.0312 0.1094 0.2188 0.2734 0.2188 0.1094 0.0312 0.0039
```

**Simulation 115** ( $\text{Binomial}(n, \theta)$  as  $\sum_{i=1}^n \text{Bernoulli}(\theta)$ ) Since the  $\text{Binomial}(n, \theta)$  RV  $X$  is the sum of  $n$  IID  $\text{Bernoulli}(\theta)$  RVs we can also simulate from  $X$  by first simulating  $n$  IID  $\text{Bernoulli}(\theta)$  RVs and then adding them up as follows:

```
>> rand('twister',17678);
>> theta=0.5; % give some desired theta value, say 0.5
>> n=5; % give the parameter n for Binomial(n,theta) RV X, say n=5
>> xis=floor(rand(1,n)+theta) % produce n IID samples from Bernoulli(theta=0.5) RVs X1,X2,...Xn
xis =
1 1 0 0 0
>> x=sum(xis) % sum up the xis to get a sample from Binomial(n=5,theta=0.5) RV X
x = 2
```

It is straightforward to produce more than one sample from  $X$  by exploiting the column-wise summing property of MATLAB's `sum` function when applied to a two-dimensional array:

```
>> rand('twister',17);
>> theta=0.25; % give some desired theta value, say 0.25 this time
>> n=3; % give the parameter n for Binomial(n,theta) RV X, say n=3 this time
>> xis10 = floor(rand(n,10)+theta) % produce an n by 10 array of IID samples from Bernoulli(theta=0.25) RVs
xis10 =
0 0 0 0 1 0 0 0 0 0
0 1 0 1 1 0 0 0 0 0
0 0 0 0 0 0 1 0 0 0
>> x=sum(xis10) % sum up the array column-wise to get 10 samples from Binomial(n=3,theta=0.25) RV X
x = 0 1 0 1 2 0 0 1 0 0
```

In Simulation 115, the number of IID  $\text{Bernoulli}(\theta)$  RVs needed to simulate one sample from the  $\text{Binomial}(n, \theta)$  RV is exactly  $n$ . Thus, as  $n$  increases, the amount of time needed to simulate from  $\text{Binomial}(n, \theta)$  is  $O(n)$ , i.e. linear in  $n$ . We can simulate more efficiently by exploiting a simple relationship between the  $\text{Geometric}(\theta)$  RV and the  $\text{Binomial}(n, \theta)$  RV.

The  $\text{Binomial}(n, \theta)$  RV  $X$  is related to the IID  $\text{Geometric}(\theta)$  RV  $Y_1, Y_2, \dots$ :  $X$  is the number of successful  $\text{Bernoulli}(\theta)$  outcomes (outcome is 1) that occur in a total of  $n$   $\text{Bernoulli}(\theta)$  trials, with the number of trials between consecutive successes distributed according to IID  $\text{Geometric}(\theta)$  RV.

**Simulation 116 (Binomial( $\theta$ ) from IID Geometric( $\theta$ ) RVs)** By this principle, we can simulate from the  $\text{Binomial}(\theta)$   $X$  by Step 1: generating IID  $\text{Geometric}(\theta)$  RVs  $Y_1, Y_2, \dots$ , Step 2: stopping as soon as  $\sum_{i=1}^k (Y_i + 1) > n$  and Step 3: setting  $x \leftarrow k - 1$ .

We implement the above algorithm via the following M-file:

---

```
Sim1BinomByGeoms.m
function x = Sim1BinomByGeoms(n,theta)
% Simulate one sample from Binomial(n,theta) via Geometric(theta) RVs
YSum=0; k=0; % initialise
while (YSum <= n),
    TrialsToSuccess=floor(log(rand)/log (1-theta)) + 1; % sample from Geometric(theta)+1
    YSum = YSum + TrialsToSuccess; % total number of trials
    k=k+1; % number of Bernoulli successes
end
x=k-1; % return x
```

---

Here is a call to simulate 12 samples from  $\text{Binomial}(n = 10, \theta = 0.5)$  RV:

```
>> theta=0.5; % declare theta
>> n=10; % say n=10
>> SampleSize=12;% say you want to simulate 12 samples
>> rand('twister',10001) % seed the fundamental sampler
>> Samples=arrayfun(@(T)Sim1BinomByGeoms(n,T),theta*ones(1,SampleSize))
Samples = 7 5 8 8 4 1 4 8 2 4 6 5
```

Figure 3.10 depicts a comparison of the PDF of  $\text{Binomial}(n = 10, \theta = 0.5)$  RV and a relative frequency histogram based on 100,000 simulations from it.

Let us simulate from the  $\text{Poisson}(\lambda)$  RV of Model 8 as shown in Figure 3.12.

**Simulation 117 (Poisson( $\lambda$ ) from IID Exponential( $\lambda$ ) RVs)** By this principle, we can simulate from the  $\text{Poisson}(\lambda)$   $X$  by Step 1: generating IID  $\text{Exponential}(\lambda)$  RVs  $Y_1, Y_2, \dots$ , Step 2: stopping as soon as  $\sum_{i=1}^k Y_i \geq 1$  and Step 3: setting  $x \leftarrow k - 1$ .

We implement the above algorithm via the following M-file:

---

```
Sim1Poisson.m
function x = Sim1Poisson(lambda)
% Simulate one sample from Poisson(lambda) via Exponentials
YSum=0; k=0; % initialise
while (YSum < 1),
    YSum = YSum + -(1/lambda) * log(rand);
    k=k+1;
end
x=k-1; % return x
```

---

Here is a call to simulate 10 samples from  $\text{Poisson}(\lambda = 10.0)$  and  $\text{Poisson}(\lambda = 0.1)$  RVs:

```
>> arrayfun(@(lambda)Sim1Poisson(lambda),10.0*ones(1,10)) % lambda=10.0
ans =
    14     7    10    13    11     3     6     5     8     5
>> arrayfun(@(lambda)Sim1Poisson(lambda),0.1*ones(1,10)) % lambda=0.1
ans =
     2     0     0     0     0     0     0     0     0     0
```

Figure 3.12 depicts a comparison of the PDF of  $\text{Poisson}(\lambda = 10)$  RV and a relative frequency histogram based on 1000 simulations from it.

Simulating from a  $\text{Poisson}(\lambda)$  RV is also a special case of simulating from the following more general RV.

**Model 12 ( $GD(\theta_0, \theta_1, \dots)$ )** We say  $X$  is a General Discrete( $\theta_0, \theta_1, \dots$ ) or  $GD(\theta_0, \theta_1, \dots)$  RV over the countable discrete state space  $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$  with parameters  $(\theta_0, \theta_1, \dots)$  if the PMF of  $X$  is defined as follows:

$$f(X = x; \theta_0, \theta_1, \dots) = \begin{cases} 0, & \text{if } x \notin \{0, 1, 2, \dots\} \\ \theta_0, & \text{if } x = 0 \\ \theta_1, & \text{if } x = 1 \\ \vdots \end{cases}$$

Algorithm 6 allows us to simulate from any member of the class of non-negative discrete RVs as specified by the probabilities  $(\theta_0, \theta_1, \dots)$ . When an RV  $X$  takes values in another countable set  $\mathbb{X} \neq \mathbb{Z}_+$ , then we can still use the above algorithm provided we have a one-to-one and onto mapping  $D$  from  $\mathbb{Z}_+$  to  $\mathbb{X}$  that allows us to think of  $\{0, 1, 2, \dots\}$  as indices of an array  $D$ .

---

**Algorithm 6** Inversion Sampler for  $GD(\theta_0, \theta_1, \dots)$  RV  $X$ 


---

1: *input:*

1.  $\theta_0$  and  $\{C(i) = \theta_i / \theta_{i-1}\}$  for any  $i \in \{1, 2, 3, \dots\}$ .
2.  $u \sim \text{Uniform}(0, 1)$

2: *output:* a sample from  $X$

3: *initialise:*  $p \leftarrow \theta_0$ ,  $q \leftarrow \theta_0$ ,  $i \leftarrow 0$

4: **while**  $u > q$  **do**

5:    $i \leftarrow i + 1$ ,  $p \leftarrow p C(i)$ ,  $q \leftarrow q + p$

6: **end while**

7: *return:*  $x = i$

---

**Simulation 118 ( $\text{Binomial}(n, \theta)$ )** To simulate from a  $\text{Binomial}(n, \theta)$  RV  $X$ , we can use Algorithm 6 with:

$$\theta_0 = (1 - \theta)^n, \quad C(x + 1) = \frac{\theta(n - x)}{(1 - \theta)(x + 1)}, \quad \text{Mean Efficiency: } O(1 + n\theta).$$

Similarly, with the appropriate  $\theta_0$  and  $C(x + 1)$ , we can also simulate from the  $\text{Geometric}(\theta)$  and  $\text{Poisson}(\lambda)$  RVs.

**Labwork 119** This is a challenging exercise for the student who is finding the other Labworks too easy. So those who are novice to MATLAB may skip this Labwork.

1. Implement Algorithm 6 via a function named `MyGenDiscInvSampler` in MATLAB. Hand in the M-file named `MyGenDiscInvSampler.m` giving detailed comments explaining your understanding of each step of the code. [Hint:  $C(i)$  should be implemented as a function (use function handles via `@`) that can be passed as a parameter to the function `MyGenDiscInvSampler`].
2. Show that your code works for drawing samples from a  $\text{Binomial}(n, p)$  RV by doing the following:
  - (a) Seed the fundamental sampler by your Student ID (if your ID is 11424620 then type `rand('twister', 11424620);`)
  - (b) Draw 100 samples from the  $\text{Binomial}(n = 20, p = 0.5)$  RV and report the results in an  $2 \times 2$  table with column headings `x` and No. of observations. [Hint: the inputs  $\theta_0$  and  $C(i)$  for the  $\text{Binomial}(n, p)$  RV is given above].
3. Show that your code works for drawing samples from a  $\text{Geometric}(p)$  RV by doing the following:
  - (a) Seed the fundamental sampler by your Student ID.
  - (b) Set the variable `Mytheta=rand`.
  - (c) Draw 100 samples from the  $\text{Geometric}(\text{Mytheta})$  RV and report the sample mean. [Note: the inputs  $\theta_0$  and  $C(i)$  for the  $\text{Geometric}(\theta)$  RV should be derived and the workings shown].

### 5.3.3 von Neumann Rejection Sampler (RS)

Rejection sampling [John von Neumann, 1947, in *Stanislaw Ulam 1909-1984*, a special issue of Los Alamos Science, Los Alamos National Lab., 1987, p. 135-136] is a Monte Carlo method to draw independent samples from a target RV  $X$  with probability density  $f(x)$ , where  $x \in \mathbb{X} \subset \mathbb{R}^k$ . Typically, the target density  $f$  is only known up to a constant and therefore the (normalised) density  $f$  itself may be unknown and it is difficult to generate samples directly from  $X$ .

Suppose we have another density or mass function  $g$  for which the following are true:

- (a) we can generate random variables from  $g$ ;
- (b) the support of  $g$  contains the support of  $f$ , i.e.  $\mathbb{Y} \supset \mathbb{X}$ ;
- (c) a constant  $a > 1$  exists, such that:

$$f(x) \leq ag(x). \quad (5.10)$$

for any  $x \in \mathbb{X}$ , the support of  $X$ . Then  $x$  can be generated from Algorithm 7.

**Proposition 56 (Fundamental Theorem of Simulation)** The von Neumann rejection sampler of Algorithm 7 produces a sample  $x$  from the random variable  $X$  with density  $f(x)$ .

**Proof:** We shall prove the result for the continuous case. For any real number  $t$ :

**Algorithm 7** Rejection Sampler (RS) of von Neumann

- 1: *input:*
- (1) a target density  $f(x)$ ,
  - (2) a proposal density  $g(x)$  satisfying (a), (b) and (c) above.

2: *output:* a sample  $x$  from RV  $X$  with density  $f$

3: **repeat**

4:   Generate  $y \sim g$  and  $u \sim \text{Uniform}(0, 1)$

5: **until**  $u \leq \frac{f(y)}{ag(y)}$

6: *return:*  $x \leftarrow y$

---

$$\begin{aligned}
 F(t) &= \mathbf{P}(X \leq t) = \mathbf{P}\left(Y \leq t \mid U \leq \frac{f(Y)}{ag(Y)}\right) = \frac{\mathbf{P}\left(Y \leq t, U \leq \frac{f(Y)}{ag(Y)}\right)}{\mathbf{P}\left(U \leq \frac{f(Y)}{ag(Y)}\right)} \\
 &= \frac{\int_{-\infty}^t \left( \int_0^{f(y)/ag(y)} 1 du \right) g(y) dy}{\int_{-\infty}^{\infty} \left( \int_0^{f(y)/ag(y)} 1 du \right) g(y) dy} = \frac{\int_{-\infty}^t \left( \frac{f(y)}{ag(y)} \right) g(y) dy}{\int_{-\infty}^{\infty} \left( \frac{f(y)}{ag(y)} \right) g(y) dy} \\
 &= \int_{-\infty}^t f(y) dy
 \end{aligned}$$

**Labwork 120 (Rejection Sampler Demo)** Let us understand the rejection sampler by calling the interactive visual cognitive tool:

```
>> guiRejections
```

The M-file `guiRejections.m` will bring a graphical user interface (GUI) as shown in Figure 5.12. Try various buttons and see how the output changes with explanations. Try switching the “Target distribution” to “Mywavy4” and generate several rejection samples and see the density histogram of the accumulating samples.

**Simulation 121 (Rejection Sampling Normal(0, 1) with Laplace(1) proposals)** Suppose we wish to generate from  $X \sim \text{Normal}(0, 1)$ . Consider using the rejection sampler with proposals from  $Y \sim \text{Laplace}(1)$  (using inversion sampler of Simulation 103). The support of both RVs is  $(-\infty, \infty)$ . Next:

$$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \text{ and } g(x) = \frac{1}{2} \exp(-|x|)$$

and therefore:


$$\frac{f(x)}{g(x)} = \sqrt{\frac{2}{\pi}} \exp\left(|x| - \frac{x^2}{2}\right) \leq \sqrt{\frac{2}{\pi}} \exp\left(\frac{1}{2}\right) = a \approx 1.3155 .$$

Hence, we can use the rejection method with:

$$\frac{f(y)}{ag(y)} = \frac{f(y)}{g(y)} \frac{1}{a} = \sqrt{\frac{2}{\pi}} \exp\left(|y| - \frac{y^2}{2}\right) \frac{1}{\sqrt{\frac{2}{\pi}} \exp\left(\frac{1}{2}\right)} = \exp\left(|y| - \frac{y^2}{2} - \frac{1}{2}\right)$$

Let us implement a rejection sampler as a function in the M-file `RejectionNormalLaplace.m` by reusing the function in `LaplaceInvCDF.m`.

Figure 5.12: Visual Cognitive Tool GUI: Rejection Sampling from  $X \sim \text{Normal}(0, 1)$  with PDF  $f$  based on proposals from  $Y \sim \text{Laplace}(1)$  with PDF  $g$ .




---

```
function x = RejectionNormalLaplace()
Accept = 0; % a binary variable to indicate whether a proposed point is accepted
while ~Accept % ~ is the logical NOT operation
    y = LaplaceInvCDF(rand(),1); % sample Laplace(1) RV
    Bound = exp( abs(y) - (y*y+1)/2 );
    u = rand();
    if u <= Bound
        x = y;
        Accept = 1;
    end % if
end % while
```

---


We may obtain a large number of samples and plot them as a histogram using the following commands:

```
>> % use funarray to convert 1000 zeros into samples from the Normal(0,1)
>> y=arrayfun(@(x)(RejectionNormalLaplace()),zeros(1,1000));
>> hist(y,20) % histogram with 20 bins
```

**Classwork 122 (A note on the proposal's tail in rejection sampling)** The condition  $f(x) \leq ag(x)$  is equivalent to  $f(x)/g(x) \leq a$ , which says that  $f(x)/g(x)$  must be bounded; therefore,  $g$  must have higher tails than  $f$ . The rejection method cannot be used to generate from a Cauchy distribution using a normal distribution, because the latter has lower tails than the former.

The next result tells us how many iterations of the algorithm are needed, on average, to get a sample value from a RV with PDF  $f$ .

Figure 5.13: Rejection Sampling from  $X \sim \text{Normal}(0, 1)$  with PDF  $f$  based on 100 proposals from  $Y \sim \text{Laplace}(1)$  with PDF  $g$ .



**Proposition 57 (Acceptance Probability of RS)** The expected number of iterations of the rejection algorithm to get a sample  $x$  is the constant  $a$ .

**Proof:** For the continuous case:

$$\mathbf{P}(\text{'accept } y) = \mathbf{P}\left(u \leq \frac{f(y)}{ag(y)}\right) = \int_{-\infty}^{\infty} \left( \int_0^{f(y)/ag(y)} du \right) g(y) dy = \int_{-\infty}^{\infty} \frac{f(y)}{ag(y)} g(y) dy = \frac{1}{a}.$$

And the number of proposals before acceptance is the Geometric( $1/a$ ) RV with expectation  $\frac{1}{1/a} = a$ .

The closer  $ag(x)$  is to  $f(x)$ , especially in the tails, the closer  $a$  will be to 1, and hence the more efficient the rejection method will be.

The rejection method can still be used only if the un-normalised form of  $f$  or  $g$  (or both) is known. In other words, if we use:

$$f(x) = \frac{\tilde{f}(x)}{\int \tilde{f}(x) dx} \text{ and } g(x) = \frac{\tilde{g}(x)}{\int \tilde{g}(x) dx}$$

we know only  $\tilde{f}(x)$  and/or  $\tilde{g}(x)$  in closed-form. Suppose the following are satisfied:

- (a) we can generate random variables from  $g$ ;
- (b) the support of  $g$  contains the support of  $f$ , i.e.  $\mathbb{Y} \supset \mathbb{X}$ ;
- (c) a constant  $\tilde{a} > 0$  exists, such that:

$$\tilde{f}(x) \leq \tilde{a}\tilde{g}(x), \quad (5.11)$$

for any  $x \in \mathbb{X}$ , the support of  $X$ . Then  $x$  can be generated from Algorithm 8.

**Algorithm 8** Rejection Sampler (RS) of von Neumann – target shape

- 1: *input:*
- (1) shape of a target density  $\tilde{f}(x) = \left( \int \tilde{f}(x)dx \right) f(x)$ ,
  - (2) a proposal density  $g(x)$  satisfying (a), (b) and (c) above.
- 2: *output:* a sample  $x$  from RV  $X$  with density  $f$
- 3: **repeat**
- 4: Generate  $y \sim g$  and  $u \sim \text{Uniform}(0, 1)$
  - 5: **until**  $u \leq \frac{\tilde{f}(y)}{\tilde{a}g(y)}$
- 6: *return:*  $x \leftarrow y$

Now, the expected number of iterations to get an  $x$  is no longer  $\tilde{a}$  but rather the integral ratio:

$$\left( \frac{\int_{\mathbb{X}} \tilde{f}(x)dx}{\int_{\mathbb{Y}} \tilde{a}g(y)dy} \right)^{-1}.$$

The **Ziggurat Method** [G. Marsaglia and W. W. Tsang, SIAM Journal of Scientific and Statistical Programming, volume 5, 1984] is a rejection sampler that can efficiently draw samples from the  $Z \sim \text{Normal}(0, 1)$  RV. The MATLAB function `randn` uses this method to produce samples from  $Z$ . See [http://www.mathworks.com/company/newsletters/news\\_notes/clevescorner/spring01\\_cleve.html](http://www.mathworks.com/company/newsletters/news_notes/clevescorner/spring01_cleve.html) or [http://en.wikipedia.org/wiki/Ziggurat\\_algorithm](http://en.wikipedia.org/wiki/Ziggurat_algorithm) for more details.

**Labwork 123 (Gaussian Sampling with `randn`)** We can use MATLAB function `randn` that implements the Ziggurat method to draw samples from an RV  $Z \sim \text{Normal}(0, 1)$  as follows:

```
>> randn('state',67678); % initialise the seed at 67678 and method as Ziggurat -- TYPE help randn
>> randn % produce 1 sample from Normal(0,1) RV
ans =
    1.5587
>> randn(2,8) % produce an 2 X 8 array of samples from Normal(0,1) RV
ans =
    1.2558    0.7834    0.6612    0.3247    0.1407    1.0562    0.8034    1.2970
   -0.5317    0.0417   -0.3454    0.6182   -1.4162    0.4796   -1.5015    0.3718
```

If we want to produce samples from  $X \sim \text{Normal}(\mu, \sigma^2)$  with some user-specified  $\mu$  and  $\sigma$ , then we can use the following relationship between  $X$  and  $Z \sim \text{Normal}(0, 1)$ :

$$X \leftarrow \mu + \sigma Z, \quad Z \sim \text{Normal}(0, 1).$$

Suppose we want samples from  $X \sim \text{Normal}(\mu = \pi, \sigma^2 = 2)$ , then we can do the following:

```
>> randn('state',679); % initialise the seed at 679 and method as Ziggurat -- TYPE help randn
>> mu=pi % set the desired mean parameter mu
mu =
    3.1416
>> sigma=sqrt(2) % set the desired standard deviation parameter sigma
sigma =
    1.4142
>> mu + sigma * randn(2,8) % produces a 2 X 8 array of samples from Normal(3.1416,1.4.42)
ans =
    1.3955    1.7107    3.9572    3.2618    6.1652    2.6971    2.4940    4.5928
    0.8442    4.7617    3.5397    5.0282    1.6139    5.0977    2.0477    2.3286
```

**Labwork 124 (Sampling from truncated normal distributions)** [Christian P. Robert, Simulation of truncated normal variables, Statistics and Computing (1995) 5, 121-125] Let  $N_+(\mu, \tau, \sigma^2)$  denote the left-truncated normal distribution with truncation point  $\tau$  and density given by

$$f(x|\mu, \tau, \sigma^2) = \frac{\exp(-(x-\mu)^2/2\sigma^2)}{\sqrt{2\pi}\sigma[1 - \Phi((\tau-\mu)/\sigma)]} \mathbb{1}_{x \geq \tau}.$$

When  $\tau < \mu$ , the rejection sampler can readily be used to simulate from  $N_+(\mu, \tau, \sigma^2)$  by simulating from  $\text{Normal}(\mu, \sigma^2)$  until a number larger than  $\tau$  is obtained. When  $\tau > \mu$ , however, this can be inefficient and increasingly so as  $\tau$  gets further out into the right tail. In this case, a more efficient approach is to use the rejection sampler with the following translated exponential distribution as the proposal distribution:

$$g(y|\lambda, \tau) = \lambda \exp(-\lambda(y-\tau)) \mathbb{1}_{y \geq \tau}.$$

1. Show that for simulating from  $N_+(\mu = 0, \tau, \sigma^2 = 1)$  when  $\tau \geq 0$ , the best choice of  $\lambda$  that maximizes the expected acceptance probability for the rejection sampler is given by

$$\lambda = \frac{\tau + \sqrt{\tau^2 + 4}}{2}$$

2. Find the maximum expected acceptance probabilities for the following truncation points,  $\tau = 0, 0.5, 1, 1.5, 2, 2.5$  and  $3$ . What can you conclude about efficiency as  $\tau$  gets further out into the right tail?
3. Describe how samples from  $N_+(\mu, \tau, \sigma^2)$  can be obtained by simulating from  $N_+(\mu = 0, \tau, \sigma^2 = 1)$  and using location-scale transformation.
4. A related distribution, denoted by  $N_-(\mu, \tau, \sigma^2)$ , is the right-truncated normal distribution truncated on the right at  $\tau$ . Describe how samples from  $N_-(\mu, \tau, \sigma^2)$  can be obtained by simulating from an appropriate left-truncated normal distribution.
5. Write a MATLAB function that provides samples from a truncated normal distribution. The function should have the following inputs: number of samples required, left or right truncation,  $\mu, \sigma^2$  and  $\tau$ .

# Chapter 6

## Limit Laws of Statistics

**Proposition 58 (Weak Law of Large Numbers (WLLN))** If we are given a sequence of independent and identically distributed RVs,  $X_1, X_2, \dots \stackrel{\text{IID}}{\sim} X_1$  and if  $\mathbf{E}(X_1)$  exists, as per (3.27), then the sample mean  $\bar{X}_n$  converges in probability to the expectation of any one of the IID RVs, say  $\mathbf{E}(X_1)$  by convention. More formally, we write:

$$\text{If } X_1, X_2, \dots \stackrel{\text{IID}}{\sim} X_1 \text{ and if } \mathbf{E}(X_1) \text{ exists, then } \bar{X}_n \xrightarrow{P} \mathbf{E}(X_1) .$$

**Proof:** For simplicity, we will prove a slightly weaker result by assuming finite variance of  $X_1$ . Suppose  $\mathbf{V}(X_1) < \infty$ , then:

$$\begin{aligned} \mathbf{P}(|\bar{X}_n - \mathbf{E}(\bar{X}_n)| \geq \epsilon) &= \frac{\mathbf{V}(\bar{X}_n)}{\epsilon^2} && [\text{by applying Chebychev's inequality (3.42) to the RV } \bar{X}_n] \\ &= \frac{\frac{1}{n}\mathbf{V}(X_1)}{\epsilon^2} && [\text{by the IID assumption of } X_1, X_2, \dots \text{ we can apply (4.3)}] \end{aligned}$$

Therefore, for any given  $\epsilon > 0$ ,

$$\begin{aligned} \mathbf{P}(|\bar{X}_n - \mathbf{E}(\bar{X}_n)| \geq \epsilon) &= \mathbf{P}(|\bar{X}_n - \mathbf{E}(\bar{X}_n)| \geq \epsilon) && [\text{by the IID assumption of } X_1, X_2, \dots, \mathbf{E}(\bar{X}_n) = \mathbf{E}(X_1), \text{ as per (4.2)}] \\ &= \frac{\frac{1}{n}\mathbf{V}(X_1)}{\epsilon^2} \rightarrow 0, \quad \text{as } n \rightarrow \infty , \end{aligned}$$

or equivalently,  $\lim_{n \rightarrow \infty} \mathbf{P}(|\bar{X}_n - \mathbf{E}(\bar{X}_n)| \geq \epsilon) = 0$ . And the last statement is the definition of the claim made by the weak law of large numbers (WLLN), namely that  $\bar{X}_n \xrightarrow{P} \mathbf{E}(X_1)$ .

**Heuristic Interpretation of WLLN:** The distribution of the sample mean RV  $\bar{X}_n$  obtained from an independent and identically distributed sequence of RVs  $X_1, X_2, \dots$  [i.e. all the RVs  $X_i$ 's are independent of one another and have the same distribution function, and thereby the same expectation, variance and higher moments], concentrates around the expectation of any one of the RVs in the sequence, say that of the first one  $\mathbf{E}(X_1)$  [without loss of generality], as  $n$  approaches infinity.

**Example 125 (Bernoulli WLLN and Galton's Quincunx)** We can appreciate the WLLN for  $\bar{X}_n = n^{-1}S_n = \sum_{i=1}^n X_i$ , where  $X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} \text{Bernoulli}(p)$  using the paths of balls dropped into a device built by Galton called the Quincunx.

**Proposition 59 (Central Limit Theorem (CLT))** Let  $X_1, X_2, \dots \stackrel{\text{IID}}{\sim} X_1$  and suppose  $\mathbf{E}(X_1)$

and  $\mathbf{V}(X_1)$  exists, then

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i \rightsquigarrow X \sim \text{Normal} \left( \mathbf{E}(X_1), \frac{\mathbf{V}(X_1)}{n} \right) , \quad (6.1)$$

$$\bar{X}_n - \mathbf{E}(X_1) \rightsquigarrow X - \mathbf{E}(X_1) \sim \text{Normal} \left( 0, \frac{\mathbf{V}(X_1)}{n} \right) , \quad (6.2)$$

$$\sqrt{n} (\bar{X}_n - \mathbf{E}(X_1)) \rightsquigarrow \sqrt{n} (X - \mathbf{E}(X_1)) \sim \text{Normal}(0, \mathbf{V}(X_1)) , \quad (6.3)$$

$$Z_n := \frac{\bar{X}_n - \mathbf{E}(\bar{X}_n)}{\sqrt{\mathbf{V}(\bar{X}_n)}} = \frac{\sqrt{n} (\bar{X}_n - \mathbf{E}(X_1))}{\sqrt{\mathbf{V}(X_1)}} \rightsquigarrow Z \sim \text{Normal}(0, 1) , \quad (6.4)$$

$$\lim_{n \rightarrow \infty} P \left( \frac{\bar{X}_n - \mathbf{E}(\bar{X}_n)}{\sqrt{\mathbf{V}(\bar{X}_n)}} \leq z \right) = \lim_{n \rightarrow \infty} \mathbf{P}(Z_n \leq z) = \Phi(z) := \int_{-\infty}^z \left( \frac{1}{\sqrt{2\pi}} \exp \left( \frac{-x^2}{2} \right) \right) dx . \quad (6.5)$$

Thus, for sufficiently large  $n$  (say  $n > 30$ ) we can make the following approximation:

$$P \left( \frac{\bar{X}_n - \mathbf{E}(\bar{X}_n)}{\sqrt{\mathbf{V}(\bar{X}_n)}} \leq z \right) \approx \mathbf{P}(Z \leq z) = \Phi(z) := \int_{-\infty}^z \left( \frac{1}{\sqrt{2\pi}} \exp \left( \frac{-x^2}{2} \right) \right) dx . \quad (6.6)$$

**Proof:** See any intermediate to advanced undergraduate text in Probability. Start from the index looking for “Central Limit Theorem” to find the page number for the proof ....

**Heuristic Interpretation of CLT:** Probability statements about the sample mean RV  $\bar{X}_n$  can be approximated using a Normal distribution.

Here is a simulation showing CLT in action.

```
>> % a demonstration of Central Limit Theorem --
>> % the sample mean of a sequence of n IID Exponential(lambda) RVs
>> % itself a Gaussian(1/lambda,lambda/n) RV
>> lambda=0.1; Reps=10000; n=10; hist(sum(-1/lambda * log(rand(n,Reps))/n)
>> lambda=0.1; Reps=10000; n=100; hist(sum(-1/lambda * log(rand(n,Reps))/n,20)
>> lambda=0.1; Reps=10000; n=1000; hist(sum(-1/lambda * log(rand(n,Reps))/n,20)
```

Let us look at an example that makes use of the CLT next.

**Example 126 (Errors in computer code (Wasserman03, p. 78))** Suppose the collection of RVs  $X_1, X_2, \dots, X_n$  model the number of errors in  $n$  computer programs named  $1, 2, \dots, n$ , respectively. Suppose that the RV  $X_i$  modeling the number of errors in the  $i$ -th program is the  $\text{Poisson}(\lambda = 5)$  for any  $i = 1, 2, \dots, n$ . Further suppose that they are independently distributed. Succinctly, we suppose that

$$X_1, X_2, \dots, X_n \stackrel{\text{IID}}{\sim} \text{Poisson}(\lambda = 5) .$$

Suppose we have  $n = 125$  programs and want to make a probability statement about  $\bar{X}_n$  which is the average error per program out of these 125 programs. Since  $\mathbf{E}(X_i) = \lambda = 5$  and  $\mathbf{V}(X_i) = \lambda = 5$ , we may want to know how often our sample mean  $\bar{X}_{125}$  differs from the expectation of 5 errors per

program. Using the CLT we can approximate  $\mathbf{P}(\bar{X}_n < 5.5)$ , for instance, as follows:

$$\begin{aligned}
 \mathbf{P}(\bar{X}_n < 5.5) &= P\left(\frac{\sqrt{n}(\bar{X}_n - \mathbf{E}(X_1))}{\sqrt{\mathbf{V}(X_1)}} < \frac{\sqrt{n}(5.5 - \mathbf{E}(X_1))}{\sqrt{\mathbf{V}(X_1)}}\right) \\
 &\approx P\left(Z < \frac{\sqrt{n}(5.5 - \lambda)}{\sqrt{\lambda}}\right) \quad [\text{by (6.6), and } \mathbf{E}(X_1) = \mathbf{V}(X_1) = \lambda] \\
 &= P\left(Z < \frac{\sqrt{125}(5.5 - 5)}{\sqrt{5}}\right) \quad [\text{Since, } \lambda = 5 \text{ and } n = 125 \text{ in this Example}] \\
 &= \mathbf{P}(Z \leq 2.5) = \Phi(2.5) = \int_{-\infty}^{2.5} \left( \frac{1}{\sqrt{2\pi}} \exp\left(\frac{-x^2}{2}\right) \right) dx \approx 0.993790334674224 .
 \end{aligned}$$

The last number above needed the following:

**Labwork 127 (Numerical approximation of  $\Phi(2.5)$ )** The numerical approximation of  $\Phi(2.5)$  was obtained via the following call to our erf-based `NormalCdf` function from ??.

```

>> format long
>> disp(NormalCdf(2.5,0,1))
0.993790334674224

```

The CLT says that if  $X_1, X_2, \dots \stackrel{\text{IID}}{\sim} X_1$ , then  $Z_n := \sqrt{n}(\bar{X}_n - \mathbf{E}(X_1))/\sqrt{\mathbf{V}(X_1)}$  is approximately distributed as  $\text{Normal}(0, 1)$ . In Example 126, we knew  $\sqrt{\mathbf{V}(X_1)}$ . However, in general, we may not know  $\sqrt{\mathbf{V}(X_1)}$ . The next proposition says that we may estimate  $\sqrt{\mathbf{V}(X_1)}$  using the sample standard deviation  $S_n$  of  $X_1, X_2, \dots, X_n$ , according to (4.5), and still make probability statements about the sample mean  $\bar{X}_n$  using a Normal distribution.

**Proposition 60 (CLT based on Sample Variance)** Let  $X_1, X_2, \dots \stackrel{\text{IID}}{\sim} X_1$  and suppose  $\mathbf{E}(X_1)$  and  $\mathbf{V}(X_1)$  exists, then

$$\frac{\sqrt{n}(\bar{X}_n - \mathbf{E}(X_1))}{S_n} \rightsquigarrow \text{Normal}(0, 1) . \quad (6.7)$$

We will use (6.7) for statistical estimation in the sequel.


# Chapter 7

## Answers to Selected Exercises


**Answer (Ex. 1.1)** — By operating with  $\Omega$ ,  $T$ ,  $L$  and  $S$  we can obtain the answers as follows:

- |   |  |
|---|--|
| (a) $T \cap L = \{L_3\}$                          | (f) $S \cap L = \emptyset$                             |
| (b) $T \cap S = \emptyset$                        | (g) $S^c \cap L = \{L_1, L_2, L_3\} = L$               |
| (c) $T \cup L = \{T_1, T_2, T_3, L_3, L_1, L_2\}$ | (h) $T^c = \{L_1, L_2, S_1, S_2, S_3, \dots, S_{50}\}$ |
| (d) $T \cup L \cup S = \Omega$                    | (i) $T^c \cap L = \{L_1, L_2\}$                        |
| (e) $S^c = \{T_1, T_2, T_3, L_3, L_1, L_2\}$      | (j) $T^c \cap T = \emptyset$                           |

**Answer (Ex. 1.3)** — We can check  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$  from the following sketch:



**Answer (Ex. 1.3)** — We can check  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$  from the following sketch:  
We can check  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$  from the following sketch:




**Answer (Ex. 1.4)** — To illustrate the idea that  $A \subseteq B$  if and only if  $A \cup B = B$ , we need to illustrate two implications:

1.if  $A \subseteq B$  then  $A \cup B = B$  and

2.if  $A \cup B = B$  then  $A \subseteq B$ .

The following Venn diagram illustrates the two implications clearly.



**Answer (Ex. 2.1) —** (a)  $\mathbf{P}(\{Z\}) = 0.1\% = \frac{0.1}{100} = 0.001$

(b)  $\mathbf{P}(\text{'picking any letter'}) = \mathbf{P}(\Omega) = 1$

(c)  $\mathbf{P}(\{E, Z\}) = \mathbf{P}(\{E\} \cup \{Z\}) = \mathbf{P}(\{E\}) + \mathbf{P}(\{Z\}) = 0.13 + 0.001 = 0.131$ , by Axiom (3)

(d)  $\mathbf{P}(\text{'picking a vowel'}) = \mathbf{P}(\{A, E, I, O, U\}) = (7.3\% + 13.0\% + 7.4\% + 7.4\% + 2.7\%) = 37.8\%$ , by the addition rule for mutually exclusive events, rule (2).

(e)  $\mathbf{P}(\text{'picking any letter in the word WAZZZUP'}) = \mathbf{P}(\{W, A, Z, U, P\}) = 14.4\%$ , by the addition rule for mutually exclusive events, rule (2).

(f)  $\mathbf{P}(\text{'picking any letter in the word WAZZZUP or a vowel'}) = \mathbf{P}(\{W, A, Z, U, P\}) + \mathbf{P}(\{A, E, I, O, U\}) - \mathbf{P}(\{A, U\}) = 14.4\% + 37.8\% - 10\% = 42.2\%$ , by the addition rule for two arbitrary events, rule (3).

**Answer (Ex. 2.2) —** 1. {BB, BW, WB, WW}

2. {RRRR, RRRL, RRRL, RLRR, LRRR, RLRL, RRLL, LLRR, LRLR, LRRL, RLLL, LLLL, LLLR, LLRL, LRLL, RLLL}

3. {6, 16, 26, 36, 46, 56, 116, 126, 136, 146, 156, 216, 226, 236, 246, 256, ...}

**Answer (Ex. 2.3) —** 1. The sample space  $\Omega = \{W, A, I, M, K, R\}$ .

2. Since there are eleven letters in WAIMAKARIRI the probabilities are:

$$\mathbf{P}(\{W\}) = \frac{1}{11}, \mathbf{P}(\{A\}) = \frac{3}{11}, \mathbf{P}(\{I\}) = \frac{3}{11}, \mathbf{P}(\{M\}) = \frac{1}{11}, \mathbf{P}(\{K\}) = \frac{1}{11}, \mathbf{P}(\{R\}) = \frac{2}{11}.$$

3. By the complementation rule, the probability of not choosing the letter R is:

$$1 - \mathbf{P}(\text{choosing the letter R}) = 1 - \frac{2}{11} = \frac{9}{11}.$$

**Answer (Ex. 2.4) —** 1. First, the sample space is:  $\Omega = \{B, I, N, G, O\}$  .

2. The probabilities of simple events are:

$$\mathbf{P}(B) = \mathbf{P}(I) = \mathbf{P}(N) = \mathbf{P}(G) = \mathbf{P}(O) = \frac{15}{75} = \frac{1}{5}.$$

3. Using the addition rule for mutually exclusive events,

$$\begin{aligned}
 \mathbf{P}(\Omega) &= \mathbf{P}(\{\mathbf{B}, \mathbf{I}, \mathbf{N}, \mathbf{G}, \mathbf{O}\}) \\
 &= \mathbf{P}(\{\mathbf{B}\} \cup \{\mathbf{I}\} \cup \{\mathbf{N}\} \cup \{\mathbf{G}\} \cup \{\mathbf{O}\}) \\
 &= \mathbf{P}(\mathbf{B}) + \mathbf{P}(\mathbf{I}) + \mathbf{P}(\mathbf{N}) + \mathbf{P}(\mathbf{G}) + \mathbf{P}(\mathbf{O}) \quad \text{simplifying notation} \\
 &= \frac{1}{5} + \frac{1}{5} + \frac{1}{5} + \frac{1}{5} + \frac{1}{5} \\
 &= 1
 \end{aligned}$$

4. Since the events  $\{\mathbf{B}\}$  and  $\{\mathbf{I}\}$  are disjoint,

$$\mathbf{P}(\{\mathbf{B}\} \cup \{\mathbf{I}\}) = \mathbf{P}(\mathbf{B}) + \mathbf{P}(\mathbf{I}) = \frac{1}{5} + \frac{1}{5} = \frac{2}{5}.$$

5. Using the addition rule for two arbitrary events we get,

$$\begin{aligned}
 \mathbf{P}(C \cup D) &= \mathbf{P}(C) + \mathbf{P}(D) - \mathbf{P}(C \cap D) \\
 &= \mathbf{P}(\{\mathbf{B}, \mathbf{I}, \mathbf{G}\}) + \mathbf{P}(\{\mathbf{G}, \mathbf{I}, \mathbf{N}\}) - \mathbf{P}(\{\mathbf{G}, \mathbf{I}\}) \\
 &= \frac{3}{5} + \frac{3}{5} - \frac{2}{5} \\
 &= \frac{4}{5}.
 \end{aligned}$$

**Answer (Ex. 2.5)** — We can assume that the first shot is independent of the second shot so we can multiply the probabilities here.

For case A, there is only one shot so the probability of hitting at least once is  $\frac{1}{2}$ .

For case B, the probability of missing both shots is  $\frac{2}{3} \cdot \frac{2}{3} = \frac{4}{9}$ , so the probability hitting some target at least once is

$$1 - \mathbf{P}(\text{missing the target both times}) = 1 - \frac{4}{9} = \frac{5}{9}$$

Therefore, case B has the greater probability of hitting the target at least once.

**Answer (Ex. 2.6)** — 1. The sample space is

$$\begin{aligned}
 &\{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\
 &(3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\
 &(5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)\}
 \end{aligned}$$

Note: Order matters here. For example, the outcome “16” refers to a “1” on the first die and a “6” on the second, whereas the outcome “61” refers to a “6” on the first die and a “1” on the second.

2. First tabulate all possible sums as follows:

+	1	2	3	4	5	6
1	2	3	4	<b>5</b>	<b>6</b>	7
2	3	4	<b>5</b>	<b>6</b>	7	8
3	4	<b>5</b>	<b>6</b>	7	8	9
4	<b>5</b>	<b>6</b>	7	8	9	10
5	<b>6</b>	7	8	9	10	11
6	7	8	9	10	11	12


Let  $A$  be the event *the sum is 5* and  $B$  be the event *the sum is 6*, then  $A$  and  $B$  are mutually exclusive events with probabilities

$$\mathbf{P}(A) = \frac{4}{36} \quad \text{and} \quad \mathbf{P}(B) = \frac{5}{36}.$$

Therefore,

$$\mathbf{P}(4 < \text{sum} < 7) = \mathbf{P}(A \cup B) = \mathbf{P}(A) + \mathbf{P}(B) = \frac{4}{36} + \frac{5}{36} = \frac{1}{4}$$

**Answer (Ex. 2.7)** — First draw a tree with the first split based on the outcome of the midterm test and the second on the outcome of the final exam. Note that the probabilities involved in this second branch are *conditional* probabilities that depend on the outcome of the midterm test. Let  $A$  be the event that the student passes the final exam and let  $B$  be the event that the student passes the midterm test.



Then the probability of passing the final exam is:

$$\mathbf{P}(A) = 0.56 + 0.12 = 0.68.$$

To do this with formulae, partitioning according to the midterm test result and using the multiplication rule, we get:

$$\begin{aligned}\mathbf{P}(A) &= \mathbf{P}(A \cap B) + \mathbf{P}(A \cap B^c) \\ &= \mathbf{P}(A|B)\mathbf{P}(B) + \mathbf{P}(A|B^c)\mathbf{P}(B^c) \\ &= (0.8)(0.7) + (0.4)(0.3) = 0.68\end{aligned}$$

**Answer (Ex. 2.8)** — Let  $A$  be the event that bottles are produced by machine 1; and  $A^c$  is the event that bottles are produced by machine 2.  $R$  denotes the event that the bottles are rejected; and  $R^c$  denotes the event that the bottles are accepted. We know the following probabilities:

$$\mathbf{P}(A) = 0.75 \quad \text{and} \quad \mathbf{P}(A^c) = 0.25$$

$$\mathbf{P}(R|A) = \frac{1}{20} \quad \text{and} \quad \mathbf{P}(R^c|A) = \frac{19}{20}$$

$$\mathbf{P}(R|A^c) = \frac{1}{30} \quad \text{and} \quad \mathbf{P}(R^c|A^c) = \frac{29}{30}$$

We want  $\mathbf{P}(A|R^c)$  which is given by

$$\mathbf{P}(A|R^c) = \frac{\mathbf{P}(R^c \cap A)}{\mathbf{P}(R^c)} = \frac{\mathbf{P}(R^c \cap A)}{\mathbf{P}(R^c \cap A) + \mathbf{P}(R^c \cap A^c)}$$

where,

$$\mathbf{P}(R^c \cap A) = \mathbf{P}(R^c|A)\mathbf{P}(A) = \frac{19}{20} \times 0.75$$


and,

$$\mathbf{P}(R^c \cap A^c) = \mathbf{P}(R^c|A^c)\mathbf{P}(A^c) = \frac{29}{30} \times 0.25$$

Therefore,

$$\mathbf{P}(A|R^c) = \frac{\frac{19}{20} \times 0.75}{\frac{19}{20} \times 0.75 + \frac{29}{30} \times 0.25} \approx 0.747$$

The tree diagram for this problem is:



So the required probability is

$$\mathbf{P}(A|R^c) = \frac{\mathbf{P}(R^c \cap A)}{\mathbf{P}(R^c)} = \frac{\frac{19}{20} \times 0.75}{\frac{19}{20} \times 0.75 + \frac{29}{30} \times 0.25} \approx 0.747$$

**Answer (Ex. 2.9)** — Let the event that a micro-chip is defective be  $D$ , and the event that the test is correct be  $C$ . So the probability that the micro-chip is defective is  $P(D) = 0.05$ , and the probability that it is effective is  $P(D^c) = 0.95$ .

The probability that the test correctly detects a defective micro-chip is the conditional probability  $P(C|D) = 0.8$ , and the probability that if a good micro-chip is tested but the test declares it is defective is the conditional probability  $P(C^c|D^c) = 0.1$ . Therefore, we also have the probabilities  $P(C^c|D) = 0.2$ , and  $P(C|D^c) = 0.9$ .

Moreover, the probability that a micro-chip is defective, and has been declared as defective is

$$P(C \cap D) = P(C|D)P(D) = 0.8 \times 0.05 = 0.04.$$

The probability that a micro-chip is effective, and has been declared as effective is

$$P(C \cap D^c) = P(C|D^c)P(D^c) = 0.9 \times 0.95 = 0.855.$$

The probability that a micro-chip is defective, and has been declared as effective is


$$P(C^c \cap D) = P(C^c|D)P(D) = 0.2 \times 0.05 = 0.01.$$

The probability that a micro-chip is effective, and has been declared as defective is

$$P(C^c \cap D^c) = P(C^c|D^c)P(D^c) = 0.1 \times 0.95 = 0.095.$$

The tree diagram for these events and probabilities is:

Defective      Tested Correctly



- (a) If a micro-chip is tested to be good, it could be defective but tested incorrectly, or it could be effective and tested correctly. Therefore, the probability that the micro-chip is tested good, but it is actually defective is

$$\frac{P(C^c \cap D)}{P(C^c \cap D) + P(C \cap D^c)} = \frac{0.01}{0.01 + 0.855} \approx 0.012$$

- (b) Similarly, the probability that a micro-chip is tested to be defective, but it was good is

$$\frac{P(C^c \cap D^c)}{P(C \cap D) + P(C^c \cap D^c)} = \frac{0.095}{0.095 + 0.04} \approx 0.704$$

- (c) The probability that both the micro-chips are effective, and have been tested and determined to be good, is

$$\left( \frac{P(C \cap D^c)}{P(C^c \cap D) + P(C \cap D^c)} \right)^2$$

and so the probability that at least one is defective is:

$$1 - \left( \frac{P(C \cap D^c)}{P(C^c \cap D) + P(C \cap D^c)} \right)^2 = 1 - \left( \frac{0.855}{0.01 + 0.855} \right)^2 \approx 0.023$$

**Answer (Ex. 2.10)** — (a) Let  $F_1$  be the event a gale of force 1 occurs, let  $F_2$  be the event a gale of force 2 occurs and  $F_3$  be the event a gale of force 3 occurs. Now we know that

$$P(F_1) = \frac{2}{3}, \quad P(F_2) = \frac{1}{4}, \quad P(F_3) = \frac{1}{12}.$$

If  $D$  is the event that a gale causes damage, then we also know the following conditional probabilities:

$$P(D|F_1) = \frac{1}{4}, \quad P(D|F_2) = \frac{2}{3}, \quad P(D|F_3) = \frac{5}{6}.$$

The probability that a reported gale causes damage is

$$P(D) = P(D \cap F_1) + P(D \cap F_2) + P(D \cap F_3)$$

where

$$P(D \cap F1) = P(D|F1)P(F1) = \frac{1}{4} \times \frac{2}{3} = \frac{1}{6},$$

$$P(D \cap F2) = P(D|F2)P(F2) = \frac{2}{3} \times \frac{1}{4} = \frac{1}{6},$$

and

$$P(D \cap F3) = P(D|F3)P(F3) = \frac{5}{6} \times \frac{1}{12} = \frac{5}{72}.$$

Hence

$$P(D) = \frac{1}{6} + \frac{1}{6} + \frac{5}{72} = \frac{29}{72}$$

(b) Knowing that the gale did cause damage we can calculate the probabilities that it was of the various forces using the probabilities in (a) as follows (Note:  $P(D \cap F1) = P(F1 \cap D)$  etc.):

$$P(F1|D) = \frac{P(F1 \cap D)}{P(D)} = \frac{1/6}{29/72} = \frac{12}{29}$$

$$P(F2|D) = \frac{P(F2 \cap D)}{P(D)} = \frac{1/6}{29/72} = \frac{12}{29}$$

$$P(F3|D) = \frac{P(F3 \cap D)}{P(D)} = \frac{5/72}{29/72} = \frac{5}{29}$$

(c) First note that the probability that a reported gale does NOT cause damage is:

$$P(D^c) = 1 - P(D) = 1 - \frac{29}{72} = \frac{43}{72}.$$

Now we need to find probabilities like  $P(F1 \cap D^c)$ . The best way to do this is to use the partitioning idea of the “Total Probability Theorem”, and write:

$$P(F1) = P(F1 \cap D^c) + P(F1 \cap D),$$

Rearranging this gives

$$P(F1 \cap D^c) = P(F1) - P(F1 \cap D)$$

and so

$$P(F1|D^c) = \frac{P(F1 \cap D^c)}{P(D^c)} = \frac{P(F1) - P(F1 \cap D)}{P(D^c)} = \frac{2/3 - 1/6}{43/72} = \frac{36}{43}.$$

Similarly,

$$P(F2|D^c) = \frac{P(F2 \cap D^c)}{P(D^c)} = \frac{P(F2) - P(F2 \cap D)}{P(D^c)} = \frac{1/4 - 1/6}{43/72} = \frac{6}{43},$$

and

$$P(F3|D^c) = \frac{P(F3 \cap D^c)}{P(D^c)} = \frac{P(F3) - P(F3 \cap D)}{P(D^c)} = \frac{1/12 - 5/72}{43/72} = \frac{1}{43}.$$

**Answer (Ex. 3.1)** —  $\mathbf{P}(X = 3)$  does not satisfy the condition that  $0 \leq \mathbf{P}(A) \leq 1$  for any event  $A$ . If  $\Omega$  is the sample space, then  $\mathbf{P}(\Omega) = 1$  and so the correct probability is

$$\mathbf{P}(X = 3) = 1 - 0.07 - 0.10 - 0.32 - 0.40 = 0.11 .$$

**Answer (Ex. 3.2)** — 1. Tabulate the values for the probability mass function as follows:

$x$	1	2	3	4	5
$\mathbf{P}(X = x)$	0.1	0.2	0.2	0.2	0.3

so the distribution function is:

$$F(x) = \mathbf{P}(X \leq x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 \\ 0.1 & \text{if } 1 \leq x < 2 \\ 0.3 & \text{if } 2 \leq x < 3 \\ 0.5 & \text{if } 3 \leq x < 4 \\ 0.7 & \text{if } 4 \leq x < 5 \\ 1 & \text{if } x \geq 5 \end{cases}$$

2. The probability that the machine needs to be replaced during the first 3 years is:

$$\mathbf{P}(X \leq 3) = \mathbf{P}(X = 1) + \mathbf{P}(X = 2) + \mathbf{P}(X = 3) = 0.1 + 0.2 + 0.2 = 0.5 .$$

(This answer is easily seen from the distribution function of  $X$ .)

3. The probability that the machine needs no replacement during the first three years is

$$\mathbf{P}(X > 3) = 1 - \mathbf{P}(X \leq 3) = 0.5 .$$

**Answer (Ex. 3.3)** — Assuming that the probability model is being built from the observed relative frequencies, the probability mass function is:

$$f(x) = \begin{cases} \frac{176}{200} & x = 1 \\ \frac{22}{200} & x = 2 \\ \frac{2}{200} & x = 3 \end{cases}$$

**Answer (Ex. 3.4)** — (a)

$x$	3	4	5	6	7	8	9	10	11	12	13
$F(x) = \mathbf{P}(X \leq x)$	0.07	0.08	0.17	0.18	0.34	0.59	0.79	0.82	0.84	0.95	1.00

$$(b) \quad (i) \mathbf{P}(X \leq 5) = F(5) = 0.17$$

$$(ii) \mathbf{P}(X < 12) = \mathbf{P}(X \leq 11) = F(11) = 0.84$$

$$(iii) \mathbf{P}(X > 9) = 1 - \mathbf{P}(X \leq 9) = 1 - F(9) = 1 - 0.79 = 0.21$$

$$(iv) \mathbf{P}(X \geq 9) = 1 - \mathbf{P}(X < 9) = 1 - \mathbf{P}(X \leq 8) = 1 - 0.59 = 0.41$$

$$(v) \mathbf{P}(4 < X \leq 9) = F(9) - F(4) = 0.79 - 0.08 = 0.71$$

$$(vi) \mathbf{P}(4 < X < 11) = \mathbf{P}(4 < X \leq 10) = F(10) - F(4) = 0.82 - 0.08 = 0.74$$

**Answer (Ex. 3.5)** — Since we are sampling without replacement,

$$\begin{aligned} \mathbf{P}(X = 0) &= \frac{4}{10} \cdot \frac{3}{9} = \frac{2}{15} \quad (\text{one way of drawing two right screws}), \\ \mathbf{P}(X = 1) &= \frac{6}{10} \cdot \frac{4}{9} + \frac{4}{10} \cdot \frac{6}{9} = \frac{8}{15} \quad (\text{two ways of drawing one left and one right screw}), \\ \mathbf{P}(X = 2) &= \frac{6}{10} \cdot \frac{5}{9} = \frac{1}{3} \quad (\text{one way of drawing two left screws}). \end{aligned}$$

So the probability mass function of  $X$  is:

$$f(x) = \mathbf{P}(X = x) = \begin{cases} \frac{2}{15} & \text{if } x = 0 \\ \frac{8}{15} & \text{if } x = 1 \\ \frac{1}{3} & \text{if } x = 2 \end{cases}$$

The required probabilities are:

1.

$$\mathbf{P}(X \leq 1) = \mathbf{P}(X = 0) + \mathbf{P}(X = 1) = \frac{2}{15} + \frac{8}{15} = \frac{2}{3}$$

2.

$$\mathbf{P}(X \geq 1) = \mathbf{P}(X = 1) + \mathbf{P}(X = 2) = \frac{8}{15} + \frac{1}{3} = \frac{13}{15}$$

3.

$$\mathbf{P}(X > 1) = \mathbf{P}(X = 2) = \frac{1}{3}$$

**Answer (Ex. 3.6)** — Using Equation (3.9), we can tabulate as follows:

$y$	0	1	4	9
$f_Y(y)$	$f_X(3) = \frac{1}{6}$	$f_X(2) + f_X(4) = \frac{2}{6}$	$f_X(1) + f_X(5) = \frac{2}{6}$	$f_X(6) = \frac{1}{6}$

**Answer (Ex. 3.7)** — The probability mass function  $f_Y(y; n)$  for  $Y = |X|$ , the absolute value of  $X$ , comes from applying the formula:

$$f_Y(y; n) = \sum_{x \in \{x: g(x)=y\}} f_X(x; n) ,$$

as follows:

$$f_Y(y) = \begin{cases} \sum_{x \in \{x: |x|=0\}} f_X(x; n) = f_X(0; n) = \frac{1}{2n+1} & \text{if } y = 0 \\ \sum_{x \in \{x: |x|=y\}} f_X(x; n) = (f_X(y; n) + f_X(-y; n)) = \frac{2}{2n+1} & \text{if } y \in \{1, 2, \dots, n\} \\ 0 & \text{otherwise} \end{cases}$$

**Answer (Ex. 3.8) —** 1. Since  $f$  is a probability mass function,

$$\sum_{x=0}^{\infty} \frac{k}{2^x} = 1, \quad \text{that is,} \quad k \sum_{x=0}^{\infty} \frac{1}{2^x} = 1.$$

Now  $\sum_{x=0}^{\infty} \frac{1}{2^x}$  is a geometric series with common ratio  $r = \frac{1}{2}$  and first term  $a = 1$ , and so has sum

$$S = \frac{a}{1-r} = \frac{1}{1-\frac{1}{2}} = 2$$

Therefore,

$$2k = 1, \quad \text{that is,} \quad k = \frac{1}{2}.$$

2. From (a), the probability mass function of  $f$  is

$$f(x) = \frac{\frac{1}{2}}{2^x} = \frac{1}{2^{x+1}}. \quad (x = 0, 1, 2, \dots)$$

Now

$$\mathbf{P}(X \geq 4) = 1 - \mathbf{P}(X < 4) = 1 - \mathbf{P}(X \leq 3)$$

where

$$\begin{aligned} \mathbf{P}(X \leq 3) &= \sum_{x=0}^3 \frac{1}{2^{x+1}} \\ &= \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} \\ &= \frac{8}{16} + \frac{4}{16} + \frac{2}{16} + \frac{1}{16} \\ &= \frac{15}{16}. \end{aligned}$$

That is,  $\mathbf{P}(X \geq 4) = \frac{1}{16}$ .

**Answer (Ex. 3.9) —** We are given that  $Y = 2^{-X}$ . Define the function

$$g : \{1, 2, 3, \dots\} \rightarrow \{2^{-1}, 2^{-2}, 2^{-3}, \dots\}$$

by  $y = g(x) = 2^{-x}$ . Then  $g$  is one-to-one and onto and so by Equation (3.9),

$$f_Y(y) = \sum_{x \in g^{-1}(y)} f_X(x) = \sum_{x \in g^{-1}(y)} f_X(x) = \sum_{x \in \{-\log_2(y)\}} f_X(x) = f_X(-\log_2(y)).$$

Note that the second equality above is emphasizing that the inverse image  $g[-1](y)$  is indeed the inverse function  $g^{-1}(y)$  for this  $g : \{1, 2, 3, \dots\} \rightarrow \{2^{-1}, 2^{-2}, 2^{-3}, \dots\}$ . Therefore,

$$f_Y(y) = \begin{cases} f_X(-\log_2(y)) = \theta(1-\theta)^{-\log_2(y)-1} & \text{if } y \in \{2^{-1}, 2^{-2}, 2^{-3}, \dots\} \\ 0 & \text{otherwise.} \end{cases}$$

**Answer (Ex. 3.10)** — Since  $y = g(x) = e^x$  is a monotone increasing function for  $x \geq 0$ , we can apply the change of variable formula.

Now  $x = g^{-1}(y) = \log_e(y)$  is a monotone increasing function for  $y$  in  $[1, \infty)$

$$\left| \frac{d}{dy} (g^{-1}(y)) \right| = \left| \frac{d}{dy} (\log_e(y)) \right| = \frac{1}{y}.$$

Therefore

$$f_Y(y) = f_X(\log_e(y)) \times \left| \frac{1}{y} \right| = \log_e(y) e^{-\log_e(y)} \times \frac{1}{y} = \log_e(y) \frac{1}{y^2}$$

since  $e^{-\log_e(y)} = e^{\log_e(y^{-1})} = y^{-1}$ .

So the probability density function of  $Y$  is given by

$$f_Y(y) = \begin{cases} \frac{\log_e(y)}{y^2} & \text{if } y \geq 1 \\ 0 & \text{otherwise} \end{cases}.$$

**Answer (Ex. 3.11)** — Note that  $\theta = \frac{1}{2}$  here.

1.  $X$  has probability mass function

$$f(x) = \begin{cases} \binom{4}{0} \frac{1}{2}^0 \frac{1}{2}^4 = \frac{1}{16} & x = 0 \\ \binom{4}{1} \frac{1}{2}^1 \frac{1}{2}^3 = \frac{4}{16} & x = 1 \\ \binom{4}{2} \frac{1}{2}^2 \frac{1}{2}^2 = \frac{6}{16} & x = 2 \\ \binom{4}{3} \frac{1}{2}^3 \frac{1}{2}^1 = \frac{4}{16} & x = 3 \\ \binom{4}{4} \frac{1}{2}^4 \frac{1}{2}^0 = \frac{1}{16} & x = 4 \end{cases}$$

2. The required probabilities are:

$$\mathbf{P}(X = 0) = f(0) = \frac{1}{16}$$

$$\mathbf{P}(X = 1) = f(1) = \frac{4}{16}$$

$$\mathbf{P}(X \geq 1) = 1 - \mathbf{P}(X = 0) = 1 - f(0) = \frac{15}{16}$$

$$\mathbf{P}(X \leq 3) = f(0) + f(1) + f(2) + f(3) = \frac{15}{16}$$

**Answer (Ex. 3.12)** — 1. If the random variable  $X$  denotes the number of type  $AB$  blood donors in the sample of 15, then  $X$  has a binomial distribution with  $n = 15$  and  $\theta = 0.05$ . Therefore

$$\mathbf{P}(X = 1) = \binom{15}{1} (0.05)^1 (0.95)^{14} = 0.366 \quad (3 \text{ sig. fig.}).$$

2.If the random variable  $X$  denotes the number of type  $B$  blood donors in the sample of 15, then  $X$  has a binomial distribution with  $n = 15$  and  $\theta = 0.10$ . Therefore

$$\begin{aligned}\mathbf{P}(X \geq 3) &= 1 - \mathbf{P}(X = 0) - \mathbf{P}(X = 1) - \mathbf{P}(X = 2) \\&= 1 - \binom{15}{0}(0.1)^0(0.9)^{15} - \binom{15}{1}(0.1)^1(0.9)^{14} - \binom{15}{2}(0.1)^2(0.9)^{13} \\&= 1 - 0.2059 - 0.3432 - 0.2669 \\&= 0.184 \quad (\text{to 3 sig. fig.)}\end{aligned}$$

3.If the random variable  $X$  denotes the number of type  $O$  or type  $A$  blood donors in the sample of 15, then  $X$  has a binomial distribution with  $n = 15$  and  $\theta = 0.85$ . Therefore

$$\begin{aligned}\mathbf{P}(X > 10) &= \mathbf{P}(X = 11) + \mathbf{P}(X = 12) + \mathbf{P}(X = 13) + \mathbf{P}(X = 14) + \mathbf{P}(X = 15) \\&= \binom{15}{11}(0.85)^{11}(0.15)^4 + \binom{15}{12}(0.85)^{12}(0.15)^3 \\&\quad + \binom{15}{13}(0.85)^{13}(0.15)^2 + \binom{15}{14}(0.85)^{14}(0.15)^1 + \binom{15}{15}(0.85)^{15}(0.15)^0 \\&= 0.1156 + 0.2184 + 0.2856 + 0.2312 + 0.0874 \\&= 0.938 \quad (\text{to 3 sig. fig.)}\end{aligned}$$

4.If the random variable  $X$  denotes the number of blood donors that are *not* of type  $A$  blood donors in the sample of 15, then  $X$  has a binomial distribution with  $n = 15$  and  $\theta = 0.6$ . Therefore

$$\begin{aligned}\mathbf{P}(X < 5) &= \mathbf{P}(X = 0) + \mathbf{P}(X = 1) + \mathbf{P}(X = 2) + \mathbf{P}(X = 3) + \mathbf{P}(X = 4) \\&= \binom{15}{0}(0.6)^0(0.4)^{15} + \binom{15}{1}(0.6)^1(0.4)^{14} + \binom{15}{2}(0.6)^2(0.4)^{13} \\&\quad + \binom{15}{3}(0.6)^3(0.4)^{12} + \binom{15}{4}(0.6)^4(0.4)^{11} \\&= 0.0000 + 0.0000 + 0.0003 + 0.0016 + 0.0074 \\&= 0.009 \quad (\text{to 3 DP.)}\end{aligned}$$

**Answer (Ex. 3.13)** — This is a Binomial experiment with parameters  $\theta = 0.1$  and  $n = 10$ , and so

$$\mathbf{P}(X \geq 1) = 1 - \mathbf{P}(X < 1) = 1 - \mathbf{P}(X = 0) ,$$

where

$$\mathbf{P}(X = 0) = \binom{10}{0}0.1^00.9^{10} \approx 0.3487 .$$

Therefore, the probability that the target will be hit at least once is

$$1 - 0.3487 \approx 0.6513 .$$

**Answer (Ex. 3.14) —** 1. Since  $f(x)$  is a (continuous) probability density function which integrates to one,

$$\int_{-4}^4 kdx = 1.$$

That is,

$$\begin{aligned} kx \Big|_{-4}^4 &= 1 \\ k(4 - (-4)) &= 1 \\ 8k &= 1 \\ k &= \frac{1}{8} \end{aligned}$$

2. First note that if  $x < -4$ , then

$$F(x) = \int_{-\infty}^x 0 dv = 0.$$

If  $-4 \leq x \leq 4$ , then

$$\begin{aligned} F(x) &= \int_{-\infty}^{-4} 0 dv + \int_{-4}^x \frac{1}{8} dv \\ &= 0 + \left[ \frac{1}{8} v \right]_{-4}^x \\ &= \frac{1}{8}(x + 4) \end{aligned}$$

If  $x \geq 4$ , then

$$\begin{aligned} F(x) &= \int_{-\infty}^{-4} 0 dv + \int_{-4}^4 \frac{1}{8} dv + \int_4^x 0 dv \\ &= 0 + \left[ \frac{1}{8} v \right]_{-4}^4 + 0 \\ &= 1 \end{aligned}$$

Hence

$$F(x) = \begin{cases} 0 & x < -4 \\ \frac{1}{8}(x + 4) & -4 \leq x \leq 4 \\ 1 & x \geq 4 \end{cases}$$

3. The graphs of  $f(x)$  and  $F(x)$  for random variable  $X$  are as follows:

**Answer (Ex. 3.15) —** 1. Since the distribution function is  $F(t; \lambda) = 1 - \exp(-\lambda t)$ ,

$$\mathbf{P}(t > \tau) = 1 - \mathbf{P}(t < \tau) = 1 - F(\tau; \lambda = 0.01) = 1 - (1 - e^{-0.01\tau}) = e^{-0.01\tau}.$$

2. Set

$$\mathbf{P}(t > \tau) = e^{-0.01\tau} = \frac{1}{2}$$

and solve for  $\tau$  to get then  $\tau = -100 \times \log(0.5) = 69.3$  (3 sig. fig.).

**Answer (Ex. 3.16)** — We are given that 537 flying bombs hit an area  $A$  of south London made up of  $24 \times 24 = 576$  small equal-sized areas, say  $A_1, A_2, \dots, A_{576}$ . Assuming the hits were purely random over  $A$  the probability that a particular bomb will hit a given small area, say  $A_i$ , is  $\frac{1}{576}$ . Let  $X$  denote the number of hits that a small area  $A_i$  receives in this German raid. Since 537 bombs fell over  $A$ , we can model  $X$  as  $\text{Binomial}(n = 537, \theta = \frac{1}{576})$  that is counting the number of ‘successes’ (for German bombers) with probability  $\theta$  in a sequence of  $n = 537$  independent  $\text{Bernoulli}(\theta)$  trials. Finally, we can approximate this  $\text{Binomial}(n = 537, \theta = \frac{1}{576})$  random variable by  $\text{Poisson}(\lambda)$  random variable with  $\lambda = n\theta = \frac{537}{576} \approx 0.933$ . Using the probability mass function formula for  $\text{Poisson}(\lambda = 0.933)$  random variable  $X$  we can obtain the probabilities and compare them with the relative frequencies from the data as follows:

$x$	observed frequency	observed relative frequency	Prob of $x$ hits
0	229	$229/576 = 0.398$	$f(0; 0.933) = 0.394$
1	211	$211/576 = 0.366$	$f(1; 0.933) = 0.367$
2	93	$93/576 = 0.161$	$f(2; 0.933) = 0.171$
3	35	$35/576 = 0.0608$	$f(3; 0.933) = 0.0532$
4	7	$7/576 = 0.0122$	$f(4; 0.933) = 0.0124$
$\geq 5$	1	$1/576 = 0.00174$	$1 - \sum_{x=0}^4 f(x; 0.933) = 0.00275$

**Answer (Ex. 3.17)** — Since 2 defects exist on every 100 meters, we would expect 6 defects on a 300 meter tape. If  $X$  is the number of defects on a 300 meter tape, then  $X$  is Poisson with  $\lambda = 6$  and so the probability of zero defects is

$$\mathbf{P}(X = 0; 6) = \frac{6^0}{0!} e^{-6} = 0.0025 .$$

**Answer (Ex. 3.18)** — Since  $X$  is  $\text{Poisson}(\lambda)$  random variable with  $\lambda = 0.5$ ,  $\mathbf{P}(X \geq 2)$  is the probability of observing two or more particles during any given second.

$$\mathbf{P}(X \geq 2) = 1 - \mathbf{P}(X < 2) = 1 - \mathbf{P}(X = 1) - \mathbf{P}(X = 0) ,$$

where  $\mathbf{P}(X = 1)$  and  $\mathbf{P}(X = 0)$  can be carried out by the Poisson probability mass function

$$\mathbf{P}(X = x) = f(x) = \frac{\lambda^x}{x!} e^{-\lambda} .$$

Now

$$\mathbf{P}(X = 0) = \frac{0.5^0}{0!} \times e^{-0.5} = 0.6065$$

and

$$\mathbf{P}(X = 1) = \frac{0.5^1}{1!} \times e^{-0.5} = 0.3033$$

and so

$$\mathbf{P}(X \geq 2) = 1 - 0.9098 = 0.0902 .$$

**Answer (Ex. 3.19)** — 1. The Probability mass function for  $\text{Poisson}(\lambda)$  random variable  $X$  is

$$\mathbf{P}(X = x) = f(x; \lambda) = \frac{e^{-\lambda} \lambda^x}{x!}$$

where  $\lambda$  is the mean number of lacunae per specimen and  $X$  is the random variable “number of lacunae on a specimen”.

2.If  $x = 0$  then  $x! = 0! = 1$  and  $\lambda^x = \lambda^0 = 1$ , and the formula becomes  $\mathbf{P}(X = 0) = e^{-\lambda}$ .

3.Since  $\mathbf{P}(X \geq 1) = 0.1$ ,

$$\mathbf{P}(X = 0) = 1 - \mathbf{P}(X \geq 1) = 0.9.$$

Using (b) and solving for  $\lambda$  gives:

$$e^{-\lambda} = 0.9 \quad \text{that is, } \lambda = -\ln(0.9) = 0.1 \text{ (approximately.)}$$

Hence

$$\mathbf{P}(X = 2) = \frac{e^{-0.1}(0.1)^2}{2!} = 0.45\% \text{ (approximately.)}$$

4.Occurrence of lacunae may not always be independent. For example, a machine malfunction may cause them to be clumped.

**Answer (Ex. 3.20)** — Since  $X$  is a Poisson( $\lambda$ ) random variable (by suppressing the ‘;  $\lambda$ ’ in the argument to  $f_X(\cdot)$  for notational ease), we get

$$f_X(x) = \mathbf{P}(X = x) = \begin{cases} \frac{\lambda^x e^{-\lambda}}{x!} & \text{for } x = 0, 1, 2, \dots \\ 0 & \text{otherwise.} \end{cases}$$

If  $Y = (X + 1)^{-2}$  then

$$Y \in \left\{ 1, \frac{1}{4}, \frac{1}{9}, \dots \right\}$$

and since  $y = g(x) = (x + 1)^{-2}$  has  $g^{-1}(y) = y^{-1/2} - 1$  as its inverse function, we get:

$$f_Y(y) = \mathbf{P}(Y = y) = \sum_{\{x:g(x)=y\}} f_X(x) = f_X\left(\frac{1}{\sqrt{y}} - 1\right) = \frac{\lambda^{(\frac{1}{\sqrt{y}}-1)} e^{-\lambda}}{(\frac{1}{\sqrt{y}}-1)!}$$

for  $y = 1, \frac{1}{4}, \frac{1}{9}, \dots$ , and 0 otherwise.

**Answer (Ex. 3.21)** — The probability that *one* light bulb doesn’t need to be replaced in 1200 hours is:

$$\begin{aligned} \mathbf{P}(X > 1.2) &= 1 - \mathbf{P}(X < 1.2) \\ &= 1 - \int_1^{1.2} 6(0.25 - (x - 1.5)^2) dx \\ &= 1 - \int_1^{1.2} 6(0.25 - x^2 + 3x - 2.25) dx \\ &= 1 - \int_1^{1.2} (-6x^2 + 18x - 12) dx \\ &= 1 - [-2x^3 + 9x^2 - 12x]_1^{1.2} \\ &= 1 - 0.1040 \\ &= 0.8960 \end{aligned}$$

Assuming that the three light bulbs function independently of each other, the probability that none of them need to be replaced in the first 1200 hours is

$$\mathbf{P}(\{X_1 > 1.2\} \cap \{X_2 > 1.2\} \cap \{X_3 > 1.2\}) = 0.8960^3 = 0.7193$$

where  $X_i$  is the length of time that bulb  $i$  lasts.

**Answer (Ex. 3.22) —** 1.

$$\begin{aligned} \int_0^2 k e^{-x} dx &= 1 \\ [-k e^{-x}]_0^2 &= 1 \\ k(-e^{-2} + 1) &= 1 \\ k = \frac{1}{1 - e^{-2}} &\quad (\approx 1.1565) \end{aligned}$$

2.

$$\begin{aligned} \mathbf{P}(X \geq 1) &= 1 - \mathbf{P}(X < 1) \\ &= 1 - \int_0^1 k e^{-x} dx \\ &= 1 + k(e^{-x}]_0^1 \\ &= 1 + \frac{e^{-1} - 1}{1 - e^{-2}} \\ &\approx 0.2689 \end{aligned}$$

**Answer (Ex. 3.23) —** Since  $y = g(x) = \sqrt{x}$  is a monotone increasing function for  $x \geq 0$ , we can apply the change of variable formula.

Now  $x = g^{-1}(y) = y^2$  is a monotone increasing function for  $y \geq 0$  so on this interval

$$\left| \frac{d}{dy} (g^{-1}(y)) \right| = \left| \frac{d}{dy} (y^2) \right| = 2y.$$

Therefore

$$f_Y(y) = f_X(y^2) \times |2y| = \lambda e^{-\lambda y^2} \times 2y = 2\lambda y e^{-\lambda y^2}.$$

So the probability density function of  $Y$  is given by

$$f_Y(y) = \begin{cases} 2\lambda y e^{-\lambda y^2} & y \geq 0 \\ 0 & y < 0 \end{cases}.$$

**Answer (Ex. 3.24) —** First note that  $y = g(x) = \log_e(x)$  is a monotone increasing function over  $a \leq x \leq b$ , so we can apply the change of variable formula.

$x = g^{-1}(y) = e^y$  is a monotone increasing function over  $\log_e(a) \leq \log_e(x) \leq \log_e(b)$ , that is, over  $\log_e(a) \leq y \leq \log_e(b)$ .

For  $\log_e(a) \leq y \leq \log_e(b)$ ,

$$\left| \frac{d}{dy} (g^{-1}(y)) \right| = \left| \frac{d}{dy} (e^y) \right| = e^y.$$

Therefore

$$f_Y(y) = f_X(g^{-1}(y)) \times \left| \frac{d}{dy} (g^{-1}(y)) \right| = \frac{1}{b-a} \times e^y.$$

So the probability density function of  $Y$  is given by

$$f_Y(y) = \begin{cases} \frac{e^y}{b-a} & \log_e(a) \leq y \leq \log_e(b) \\ 0 & \text{otherwise} \end{cases}.$$

**Answer (Ex. 3.25)** — 1. The expected number of conditioners that the store sells daily is

$$\begin{aligned} \mathbf{E}(X) &= \sum_{i=1}^n x_i p_{x_i} \\ &= (10 \times 0.1 + 11 \times 0.3 + 12 \times 0.4 + 13 \times 0.2) \\ &= 1 + 3.3 + 4.8 + 2.6 \\ &= 11.7. \end{aligned}$$

2. The profit per conditioner is \$55, and so the expected daily profit given by

$$E(55X) = 55E(X) = 55 \times 11.7 = 643.50,$$

is \$643.50.

**Answer (Ex. 3.26)** — The expected value  $\mathbf{E}(X)$  is

$$\begin{aligned} \mathbf{E}(X) &= \int_0^1 6x(1-x)x \, dx \\ &= \int_0^1 (6x^2 - 6x^3) \, dx \\ &= 2x^3 - \frac{6}{4}x^4 \Big|_0^1 \\ &= 2(1^3 - 0) - \frac{6}{4}(1^4 - 0) \\ &= 0.5 \end{aligned}$$

$$\begin{aligned} \mathbf{E}(X^2) &= \int_0^1 6x(1-x)x^2 \, dx \\ &= \int_0^1 6x^3 - 6x^4 \, dx \\ &= \frac{6}{4}x^4 - \frac{6}{5}x^5 \Big|_0^1 \\ &= \frac{6}{4}(1^4 - 0) - \frac{6}{5}(1^5 - 0) \\ &= 0.3 \end{aligned}$$

the variance is  $\mathbf{V}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = 0.3 - 0.5^2 = 0.05$

**Answer (Ex. 3.27)** — This was already done in Section 3.6.2 on Properties of Expectation. Make sure you understand each step.

**Answer (Ex. 3.28)** — Using the definition of variance, expectations and by completing the square, we get:

$$\begin{aligned} V(aX + b) &= E((aX + b)^2) - (E(aX + b))^2 = E((aX)^2 + 2aXb + b^2) - (aE(X) + b)^2 \\ &= a^2 E(X^2) + 2abE(X) + b^2 - a^2(E(X))^2 - 2abE(X) - b^2 = a^2 (E(X^2) - (E(X))^2) = a^2 V(X). \end{aligned}$$

**Answer (Ex. 3.30)** — 1. The probability mass function of  $X$  is

$$f(x) = \begin{cases} \frac{1}{6} & x = 1 \\ \frac{1}{6} & x = 2 \\ \frac{1}{6} & x = 3 \\ \frac{1}{6} & x = 4 \\ \frac{1}{6} & x = 5 \\ \frac{1}{6} & x = 6 \end{cases}$$

and so

$$\mathbf{E}(X) = \sum_{i=1}^6 x_i f(x_i) = \frac{1}{6} + \frac{2}{6} + \frac{3}{6} + \frac{4}{6} + \frac{5}{6} + \frac{6}{6} = 3.5.$$

Now,

$$\mathbf{E}(X^2) = \sum_{i=1}^6 x_i^2 f(x_i) = \frac{1}{6} + \frac{4}{6} + \frac{9}{6} + \frac{16}{6} + \frac{25}{6} + \frac{36}{6} = \frac{91}{6} = 15.1667$$

and so the variance is

$$V(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2 = 15.1667 - 3.5^2 = 2.9167.$$

2. The density function of the uniform distribution,  $X$ , on  $[0, 8]$  is

$$f(x) = \begin{cases} \frac{1}{8} & 0 < x < 8 \\ 0 & \text{otherwise} \end{cases}$$

Therefore,

$$\mathbf{E}(X) = \frac{8+0}{2} = 4,$$

and

$$\mathbf{V}(X) = \frac{(8-0)^2}{12} = \frac{16}{3}$$

3. Since  $f(x)$  is the density function of an Exponential( $\lambda$ ) random variable with parameter  $\lambda = 2$ , we can use earlier results to get

$$\mathbf{E}(X) = \frac{1}{2} \quad \text{and} \quad \mathbf{V}(X) = \frac{1}{4}.$$

(You should also be able to do this by integration!)

**Answer (Ex. 3.31) —**

1. To find  $a$  we simply set  $1 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx dy$  and solve for  $a$  as follows:

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx dy = \int_0^1 \int_0^1 a(x^2 + y) dx dy \\ &= a \int_0^1 \int_0^1 (x^2 + y) dx dy = a \int_0^1 \left[ \frac{1}{3}x^3 + yx \right]_{x=0}^1 dy \\ &= a \int_0^1 \left( \frac{1}{3} + y - 0 \right) dy = a \left[ \frac{y}{3} + \frac{1}{2}y^2 \right]_{y=0}^1 \\ &= a \left( 0 - \left( \frac{1}{3} + \frac{1}{2}1^2 \right) \right) = a \left( \frac{1}{3} + \frac{1}{2} \right) \\ &= a \left( \frac{5}{6} \right) \end{aligned}$$

Therefore  $a = 6/5$  and the joint PDF is

$$f_{X,Y}(x,y) = \begin{cases} \frac{6}{5}(x^2 + y) & \text{if } 0 < x < 1 \text{ and } 0 < y < 1 \\ 0 & \text{otherwise} . \end{cases}$$

2. First compute the marginal PDF  $f_X(x)$  for any  $x \in (0, 1)$  by integrating over  $y$

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy = \int_0^1 \frac{6}{5}(x^2 + y) dy = \left[ \frac{6}{5}(yx^2 + y^2/2) \right]_{y=0}^1 \\ &= \frac{6}{5} ((1 \times x^2 + 1^2/2) - 0) = \frac{6}{5} \left( x^2 + \frac{1}{2} \right) \end{aligned}$$

Finally, the marginal PDF of the RV  $X$  in the first component of the  $\vec{V}(X, Y)$  is

$$f_X(x) = \begin{cases} \frac{6}{5} \left( x^2 + \frac{1}{2} \right) & \text{if } 0 < x < 1 \\ 0 & \text{otherwise} . \end{cases}$$

3. Similarly, the marginal PDF  $f_Y(y)$  for any  $y \in (0, 1)$  by integrating over  $x$  is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx = \int_0^1 \frac{6}{5}(x^2 + y) dx = \frac{6}{5} \left[ x^3/3 + yx \right]_{x=0}^1 = \frac{6}{5}(y + 1/3).$$

Finally, the marginal PDF of the RV  $Y$  in the second component of the  $\vec{V}(X, Y)$  is

$$f_Y(y) = \begin{cases} \frac{6}{5} \left( y + \frac{1}{3} \right) & \text{if } 0 < y < 1 \\ 0 & \text{otherwise} . \end{cases}$$

4. The product of marginal PDFs of  $X$  and  $Y$  does not equal the joint PDF of  $(X, Y)$  for values of  $(x, y) \in (0, 1)^2$

$$f_X(x)f_Y(y) = \frac{6}{5} \frac{6}{5} \left( y + \frac{1}{3} \right) \left( x^2 + \frac{1}{2} \right) = \frac{6}{25} (6x^2y + 2x^2 + 3y + 1) \neq \frac{6}{5}(x^2 + y) = f_{X,Y}(x,y)$$

Therefore  $X$  and  $Y$  are not independent random variables (they are dependent!).

5. The joint distribution function  $F_{X,Y}(x,y)$  for any  $(x,y) \in (0,1)^2$  is

$$\begin{aligned} F_{X,Y}(x,y) &= \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u,v)dudv = \int_0^y \int_0^x \frac{6}{5}(u^2 + v)dudv = \frac{6}{5} \int_0^y \left[ \frac{u^3}{3} + vu \right]_{u=0}^x dv \\ &= \frac{6}{5} \int_0^y \left( \frac{x^3}{3} + vx - 0 \right) dv = \frac{6}{5} \left[ \frac{x^3 v}{3} + \frac{v^2 x}{2} \right]_{v=0}^y = \frac{6}{5} \left( \frac{x^3 y}{3} + \frac{y^2 x}{2} - 0 \right) \\ &= \frac{6}{5} \left( \frac{x^3 y}{3} + \frac{y^2 x}{2} - 0 \right) \end{aligned}$$

6.

$$\begin{aligned} \mathbf{P}(X > 0.5, Y < 0.6) &= \int_{-\infty}^{0.6} \int_{0.5}^{\infty} f_{X,Y}(x,y)dxdy = \int_0^{0.6} \int_{0.5}^1 \frac{6}{5}(x^2 + y)dxdy \\ &= \frac{6}{5} \int_0^{0.6} \left[ \frac{x^3}{3} + yx \right]_{x=0.5}^1 dy = \frac{6}{5} \int_0^{0.6} \left( \frac{7}{24} + \frac{y}{2} \right) dy \\ &= \frac{6}{5} \left[ \frac{7}{24}y + \frac{y^2}{2} \right]_{y=0}^{0.6} = \frac{6}{5} \left( \frac{7}{24} \times \frac{6}{10} + \frac{36}{400} \right) = 0.318 \end{aligned}$$

7.

$$\begin{aligned} E(X) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xf_{X,Y}(x,y)dxdy \\ &= \frac{6}{5} \int_0^1 \int_0^1 x(x^2 + y)dxdy = \frac{6}{5} \int_0^1 \int_0^1 (x^3 + xy)dxdy = \frac{6}{5} \int_0^1 \left[ \frac{x^4}{4} + \frac{1}{2}x^2 y \right]_{x=0}^1 dy \\ &= \frac{6}{5} \int_0^1 \left( \frac{1}{4} + \frac{y}{2} - 0 - 0 \right) dy = \frac{6}{5} \left[ \frac{y}{4} + \frac{y^2}{4} \right]_{y=0}^1 = \frac{6}{5} \left( \frac{1}{4} + \frac{1}{4} - 0 - 0 \right) = \frac{6}{5} \times \frac{1}{2} = \frac{3}{5} \end{aligned}$$

8.

$$\begin{aligned} E(Y) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} yf_{X,Y}(x,y)dxdy \\ &= \int_0^1 \int_0^1 y \frac{6}{5}(x^2 + y)dxdy = \frac{6}{5} \int_0^1 \int_0^1 (x^2 y + y^2)dxdy = \frac{6}{5} \int_0^1 \left[ \frac{x^3 y}{3} + y^2 x \right]_{x=0}^1 dy \\ &= \frac{6}{5} \int_0^1 \left( \frac{y}{3} + y^2 - 0 - 0 \right) dy = \frac{6}{5} \left[ \frac{y^2}{6} + \frac{y^3}{3} \right]_{y=0}^1 = \frac{6}{5} \left( \frac{1}{6} + \frac{1}{3} - 0 - 0 \right) = \frac{6}{5} \times \frac{3}{6} = \frac{3}{5} \end{aligned}$$

9.

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf_{X,Y}(x,y)dxdy \\ &= \frac{6}{5} \int_0^1 \int_0^1 xy(x^2 + y)dxdy = \frac{6}{5} \int_0^1 \int_0^1 x^3 y + xy^2 dxdy = \frac{6}{5} \int_0^1 \left[ \frac{x^4}{4} y + \frac{x^2 y^2}{2} \right]_{x=0}^1 dy \\ &= \frac{6}{5} \int_0^1 \left( \frac{y}{4} + \frac{y^2}{2} - 0 - 0 \right) dy = \frac{6}{5} \left[ \frac{y^2}{8} + \frac{y^3}{6} \right]_{y=0}^1 = \frac{6}{5} \left( \frac{1}{8} + \frac{1}{6} - 0 - 0 \right) \\ &= \frac{6}{5} \left( \frac{3}{24} + \frac{4}{24} \right) = \frac{6}{5} \times \frac{7}{24} = \frac{7}{20} \end{aligned}$$

10.

$$\mathbf{Cov}(X, Y) = E(XY) - E(X)E(Y) = \frac{7}{20} - \left( \frac{3}{5} \times \frac{3}{5} \right) = \frac{7}{20} - \frac{9}{25} = \frac{35}{100} - \frac{36}{100} = -\frac{1}{100}$$

**Answer (Ex. 3.32)** — Note that  $Y = (X - \mu)^2$  is not on-to-one so it is better to use the direct method by differentiating the distribution function of  $Y$ ,  $F_Y(y)$ , to obtain  $f_Y(y)$ .

If  $y \geq 0$ ,

$$\begin{aligned} F_Y(y) &= \mathbf{P}(Y \leq y) \\ &= \mathbf{P}((X - \mu)^2 \leq y) \\ &= \mathbf{P}(-\sqrt{y} \leq X - \mu \leq \sqrt{y}) \\ &= \mathbf{P}(\mu - \sqrt{y} \leq X \leq \mu + \sqrt{y}) \\ &= F_X(\mu + \sqrt{y}) - F_X(\mu - \sqrt{y}) \end{aligned}$$

Differentiating this expression gives

$$\begin{aligned} f_Y(y) &= \frac{d}{dy} (F_X(\mu + \sqrt{y}) - F_X(\mu - \sqrt{y})) \\ &= \frac{1}{2} y^{-\frac{1}{2}} f_X(\mu + \sqrt{y}) - \left( -\frac{1}{2} y^{-\frac{1}{2}} \right) f_X(\mu - \sqrt{y}) \\ &= \frac{1}{2\sqrt{y}} (f_X(\mu + \sqrt{y}) + f_X(\mu - \sqrt{y})) \end{aligned}$$

Note: If  $y < 0$  then  $f_Y(y) = 0$  since  $F_Y(y) = 0$  in this case.

**Answer (Ex. 3.33)** —

First derive the marginal PMF of  $X$  and  $Y$  and then check if the JPMF is the product of the marginal PMFs.

$$f_X(0) = \sum_{y \in \mathcal{S}_{X,Y}} f_{X,Y}(0, y) = f_{X,Y}(0, 0) + f_{X,Y}(0, 1) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

and

$$f_X(1) = \sum_{y \in \mathcal{S}_{X,Y}} f_{X,Y}(1, y) = f_{X,Y}(1, 0) + f_{X,Y}(1, 1) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

Thus,

$$f_X(x) = \begin{cases} \frac{1}{2} & \text{if } x = 0 \\ \frac{1}{2} & \text{if } x = 1 \\ 0 & \text{otherwise} . \end{cases}$$

Similarly,

$$f_Y(y) = \begin{cases} \sum_{x \in \mathcal{S}_{X,Y}} f_{X,Y}(x, 0) = f_{X,Y}(0, 0) + f_{X,Y}(1, 0) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2} & \text{if } y = 0 \\ \sum_{x \in \mathcal{S}_{X,Y}} f_{X,Y}(x, 1) = f_{X,Y}(0, 1) + f_{X,Y}(1, 1) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2} & \text{if } y = 1 \\ 0 & \text{otherwise} . \end{cases}$$

Finally, the product of  $f_X(x)$  and  $f_Y(y)$  is

$$f_X(x) \times f_Y(y) = \begin{cases} \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} & \text{if } (x, y) = (0, 0) \\ \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} & \text{if } (x, y) = (0, 1) \\ \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} & \text{if } (x, y) = (1, 0) \\ \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} & \text{if } (x, y) = (1, 1) \\ 0 & \text{otherwise} . \end{cases}$$

which in turn is equal to the JPMF  $f_{X,Y}(x,y)$  in the question. Therefore we have shown that the component RVs  $X$  and  $Y$  in the R $\vec{V}$  ( $X, Y$ ) are indeed independent.

**Answer (Ex. 3.34) —**

Let  $X_1, X_2, X_3$  be independent RVs that denote the thickness of the first, second and third layer, respectively. Let  $X$  denote the thickness of the final product. Then

$$X = X_1 + X_2 + X_3$$

By the property that  $V(\sum_{i=1}^n a_i X_i) = \sum_{i=1}^n a_i^2 V(X_i)$ , Variance of  $X$  is

$$V(X) = 1^2 V(X_1) + 1^2 V(X_2) + 1^2 V(X_3) = 25 + 40 + 30 = 95 \text{ nm}^2 .$$

This shows how the variance in each layer is propagated to the variance of the final product.

**Answer (Ex. 3.35) —**

Find  $E(XY)$ ,  $E(X)$  and  $E(Y)$  to get  $\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$  as follows:

$$\begin{aligned} E(XY) &= \sum_{(x,y) \in \mathcal{S}_{X,Y}} x \times y \times f_{X,Y}(x,y) \\ &= 0 \times 0 \times 0.2 + 1 \times 1 \times 0.1 + 1 \times 2 \times 0.1 + 2 \times 1 \times 0.1 + 2 \times 2 \times 0.1 + 3 \times 3 \times 0.4 = 4.5 \end{aligned}$$

$$\begin{aligned} E((X, Y)) &= \sum_{(x,y) \in \mathcal{S}_{X,Y}} (x, y) \times f_{X,Y}(x,y) \\ &= (0, 0) \times 0.2 + (1, 1) \times 0.1 + (1, 2) \times 0.1 + (2, 1) \times 0.1 + (2, 2) \times 0.1 + (3, 3) \times 0.4 \\ &= (0, 0) + (0.1, 0.1) + (0.1, 0.2) + (0.2, 0.1) + (0.2, 0.2) + (1.2, 1.2) \\ &= (1.8, 1.8) \end{aligned}$$

Since addition is component-wise  $E((X, Y)) = (E(X), E(Y))$  and therefore  $E(X) = E(Y) = 1.8$ . Alternatively, you can first find the marginal PMFs  $f_X$  and  $f_Y$  for  $X$  and  $Y$  and then take the expectations  $E(X) = \sum_x x \times f_X(x)$  and  $E(Y) = \sum_y y \times f_Y(y)$ .

Finally,

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = 4.5 - 1.8^2 = 1.26 .$$

**Answer (Ex. 3.36) —**

Since  $X$  and  $Y$  are independent,  $F_{X,Y}(x,y) = F_X(x)F_Y(y)$  for all  $(x,y) \in \mathbb{R}^2$ , and we get:

$$F_{X,Y}(x,y) = \begin{cases} 0 & \text{if } x < 1 \text{ or } y < 0 \\ \frac{1}{2} \left(1 - \frac{1}{2}e^{-y} - \frac{1}{2}e^{-2y}\right) & \text{if } 1 \leq x < 2 \text{ and } y \geq 0 \\ 1 - \frac{1}{2}e^{-y} - \frac{1}{2}e^{-2y} & \text{if } x \geq 2 \text{ and } y \geq 0 \end{cases}$$

You can arrive at the answer by partitioning  $x$ -axis into  $(-\infty, 1)$ ,  $[1, 2)$  and  $[2, \infty)$  where  $F_X(x)$  takes distinct values. Similarly, partition the  $y$ -axis into  $(-\infty, 0)$  and  $[0, \infty)$  where  $F_Y(y)$  takes distinct values. Now  $(x,y)$  can take values in one of these  $3 \times 2 = 6$  partitions of the  $x \times y$  plane as follows (make a picture!):

$$(-\infty, 1) \times (-\infty, 0), [1, 2) \times (-\infty, 0), [2, \infty) \times (-\infty, 0), (-\infty, 1) \times [0, \infty), [1, 2) \times [0, \infty), [2, \infty) \times [0, \infty) .$$

Now work out what  $F_{X,Y}(x,y) = F_X(x)F_Y(y)$  is for  $(x,y)$  in each of the above six partitions of the plane and you will get the expression for  $F_{X,Y}(x,y)$  given above.

**Answer (Ex. 3.37) —**

First obtain marginal PDF of  $Y$ . If  $y \in [2, 3]$  then

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)dx = \int_0^{\infty} e^{-x}dx = [-e^{-x}]_0^{\infty} = 0 - (-1) = 1 .$$

Therefore,

$$f_Y(y) = \begin{cases} 1 & \text{if } y \in [2, 3] \\ 0 & \text{otherwise} . \end{cases}$$

Now, obtain the marginal PDF of  $X$ . If  $x \in [0, \infty)$  then

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)dy = \int_2^3 e^{-x}dy = e^{-x} \int_2^3 1dy = e^{-x} [y]_2^3 = e^{-x}(3-2) = e^{-x} .$$

Therefore,

$$f_X(x) = \begin{cases} e^{-x} & \text{if } x \in [0, \infty) \\ 0 & \text{otherwise} . \end{cases}$$

Finally, verifying that  $f_{X,Y}(x,y) = f_X(x)f_Y(y)$  for any  $(x,y) \in \mathbb{R}^2$  is done case by case. Draw a picture on the plane to work out the cases from the distinct expressions taken by  $f_{X,Y}(x,y)$ . There are only two cases to consider (when  $f_{X,Y}(x,y)$  takes zero values and when  $f_{X,Y}(x,y)$  takes non-zero values):

1. If  $x \notin [0, \infty)$  or  $y \notin [2, 3]$  then  $f_X(x)f_Y(y) = 0 = f_{X,Y}(x,y)$
2. If  $x \in [0, \infty)$  and  $y \in [2, 3]$  then  $f_X(x)f_Y(y) = e^{-x} \times 1 = e^{-x} = f_{X,Y}(x,y)$ .

Thus,  $X$  and  $Y$  are independent.

**Answer (Ex. 3.38) —**

$$\begin{aligned} & \mathbf{P}(X_1 > 1000, X_2 > 1000, X_3 > 1000, X_4 > 1000) \\ &= \int_{1000}^{\infty} \int_{1000}^{\infty} \int_{1000}^{\infty} \int_{1000}^{\infty} 9 \times 10^{-12} e^{-0.001x_1 - 0.002x_2 - 0.0015x_3 - 0.003x_4} dx_1 dx_2 dx_3 dx_4 \\ &= 9 \times 10^{-12} \int_{1000}^{\infty} \int_{1000}^{\infty} \int_{1000}^{\infty} \int_{1000}^{\infty} e^{-0.001x_1} e^{-0.002x_2} e^{-0.0015x_3} e^{-0.003x_4} dx_1 dx_2 dx_3 dx_4 \\ &= 9 \times 10^{-12} \int_{1000}^{\infty} e^{-0.001x_1} \int_{1000}^{\infty} e^{-0.002x_2} \int_{1000}^{\infty} e^{-0.0015x_3} \int_{1000}^{\infty} e^{-0.003x_4} dx_4 dx_3 dx_2 dx_1 \end{aligned}$$

Since

$$\int_{1000}^{\infty} e^{-ax_i} dx_i = \left[ \frac{e^{-ax_i}}{-a} \right]_{1000}^{\infty} = 0 + \frac{e^{-1000 \times a}}{a} ,$$

the above quadruply iterated integral becomes

$$\begin{aligned} & 9 \times 10^{-12} \times \frac{e^{-1000 \times 0.001}}{0.001} \times \frac{e^{-1000 \times 0.002}}{0.002} \times \frac{e^{-1000 \times 0.0015}}{0.0015} \times \frac{e^{-1000 \times 0.003}}{0.003} \\ &= 9 \times 10^{-12} \times \frac{1000}{1} \times \frac{1000}{2} \times \frac{1000}{1.5} \times \frac{1000}{3} \times e^{-1} \times e^{-2} \times e^{-1.5} \times e^{-3} \\ &= 9 \times 10^{-12} \times \frac{1}{9} \times 10^{12} \times e^{-7.5} = e^{-7.5} \approx 0.00055 . \end{aligned}$$

**Answer (Ex. 3.39) —**

Due to independence of  $Y_1$ ,  $Y_2$  and  $Y_3$

$$\begin{aligned}\mathbf{P}(9500 < Y_1 < 10500, 950 < Y_2 < 1050, 75 < Y_3 < 85) \\ &= \mathbf{P}(9500 < Y_1 < 10500)\mathbf{P}(950 < Y_2 < 1050)\mathbf{P}(75 < Y_3 < 85)\end{aligned}$$

After standardizing each Normal RV (subtracting its mean and dividing by its standard deviation) we get

$$\begin{aligned}&\mathbf{P}(9500 < Y_1 < 10500)\mathbf{P}(950 < Y_2 < 1050)\mathbf{P}(75 < Y_3 < 85) \\ &= P\left(\frac{9500 - 10000}{250} < Z < \frac{10500 - 10000}{250}\right)P\left(\frac{950 - 1000}{20} < Z < \frac{1050 - 1000}{20}\right) \\ &\quad P\left(\frac{75 - 80}{4} < Z < \frac{85 - 80}{4}\right) \\ &= \mathbf{P}(-2.0 < Z < 2.0)\mathbf{P}(-2.5 < Z < 2.5)\mathbf{P}(-1.25 < Z < 1.25) \\ &= (\Phi(2.0) - (1 - \Phi(2.0))) \times (\Phi(2.5) - (1 - \Phi(2.5))) \times (\Phi(1.25) - (1 - \Phi(1.25))) \\ &= (2\Phi(2.0) - 1) \times (2\Phi(2.5) - 1) \times (2\Phi(1.25) - 1) \\ &= ((2 \times 0.9772) - 1) \times ((2 \times 0.9938) - 1) \times ((2 \times 0.8944) - 1) \quad \text{using Table for } \Phi(z) \\ &= 0.9544 \times 0.9876 \times 0.7888 = 0.7435\end{aligned}$$

The values for the distribution function  $\Phi(z)$  of the  $\text{Normal}(0, 1)$  RV  $Z$  are in the table on page 67. The thickness of the coating layer represented by  $Y_3$  has the least probability (0.7888) of meeting specifications. Consequently, a priority should be to reduce variability in this part of the process.

**Answer (Ex. 3.40) —**

Let  $X_1, X_2, \dots, X_{10}$  denote the fill volumes of 10 cans. The average fill volume is the sample mean

$$\bar{X}_{10} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

By property of Expectations and Variances for linear combinations

$$E(\bar{X}_{10}) = E\left(\frac{1}{10} \sum_{i=1}^{10} X_i\right) = \frac{1}{10} \sum_{i=1}^n E(X_i) = \frac{1}{10} \sum_{i=1}^n E(X_1) = \frac{1}{10} \times 10 \times E(X_1) = E(X_1) = 12.1$$

Or by directly using the “formula”  $E(\bar{X}_{10}) = E(X_1) = 12.1$  for these 10 identically distributed RVs. Similarly,

$$V(\bar{X}_{10}) = V\left(\frac{1}{10} \sum_{i=1}^{10} X_i\right) = 10 \times \frac{1}{10^2} V(X_1) = \frac{1}{10} \times 0.01 = 0.001$$

Or by directly using the “formula”  $V(\bar{X}_{10}) = V(X_1)/10$  for these 10 independently and identically distributed RVs.

By the special property of Normal RVs – a linear combination of independent normal RVs is also normal – we know that  $\bar{X}_{10}$  is a  $\text{Normal}(12.1, 0.001)$  RV. Consequently, the probability of interest is

$$\begin{aligned}\mathbf{P}(\bar{X}_{10} < 12.01) &= P\left(\frac{\bar{X}_{10} - E(\bar{X}_{10})}{\sqrt{0.001}} < \frac{12.01 - E(\bar{X}_{10})}{\sqrt{0.001}}\right) = P\left(Z < \frac{12.01 - 12.1}{0.0316}\right) \\ &\approx \mathbf{P}(Z < -2.85) = 1 - \mathbf{P}(Z < 2.85) = 1 - \Phi(2.85) = 1 - 0.9978 = 0.0022\end{aligned}$$

**Answer (Ex. 3.41) —**

Using the Multinomial( $n = 10, \theta_1 = 0.6, \theta_2 = 0.3, \theta_3 = 0.08, \theta_4 = 0.02$ ) RV as our model

$$\begin{aligned} P((X_1, X_2, X_3, X_4) = (6, 2, 2, 0); n = 10, \theta_1 = 0.6, \theta_2 = 0.3, \theta_3 = 0.08, \theta_4 = 0.02) \\ = \frac{10!}{6! \times 2! \times 2! \times 0!} \times 0.6^6 \times 0.3^2 \times 0.08^2 \times 0.02^0 \\ = \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1}{(6 \times 5 \times 4 \times 3 \times 2 \times 1) \times (2 \times 1) \times (2 \times 1) \times 1} \times 0.6^6 \times 0.3^2 \times 0.08^2 \times 1 \cong 0.03386 \end{aligned}$$

**Answer (Ex. 3.42) —**

1. The CF of the discrete RV  $X$  is

$$\begin{aligned} \varphi_X(t) = E(e^{itX}) &= \sum_{x \in \{0, 1, 2\}} e^{itx} f_X(x) = e^{it \times 0} \times \frac{1}{3} + e^{it \times 1} \times \frac{1}{3} + e^{it \times 2} \times \frac{1}{3} \\ &= \frac{1}{3} (1 + e^{it} + e^{i2t}) . \end{aligned}$$

2. To find  $V(X)$  using  $V(X) = E(X^2) - (E(X))^2$  we need the first two moments of  $X$ . First, differentiate  $\varphi_X(t)$  w.r.t.  $t$

$$\frac{d}{dt} \varphi_X(t) = \frac{d}{dt} \left( \frac{1}{3} (1 + e^{it} + e^{i2t}) \right) = \frac{1}{3} (0 + ie^{it} + 2ie^{i2t}) = \frac{i}{3} (e^{it} + 2e^{i2t})$$

We get the  $k$ -th moment  $E(X^k)$  by multiplying the  $k$ -th derivative of  $\varphi_X(t)$  evaluated at  $t = 0$  by  $\frac{1}{i^k}$  as follows:

$$E(X) = \frac{1}{i} \left[ \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{i} \left[ \frac{i}{3} (e^{it} + 2e^{i2t}) \right]_{t=0} = \frac{1}{i} \frac{i}{3} (e^0 + 2e^0) = \frac{1}{3}(1+2) = 1 .$$

$$\begin{aligned} E(X^2) &= \frac{1}{i^2} \left[ \frac{d^2}{dt^2} \varphi_X(t) \right]_{t=0} = \frac{1}{i^2} \left[ \frac{d}{dt} \frac{i}{3} (e^{it} + 2e^{i2t}) \right]_{t=0} = \frac{1}{i^2} \left[ \frac{i}{3} (ie^{it} + 4ie^{i2t}) \right]_{t=0} \\ &= \frac{1}{3} (e^0 + 4e^0) = \frac{5}{3} . \end{aligned}$$

Finally,

$$V(X) = E(X^2) - (E(X))^2 = \frac{5}{3} - 1^2 = \frac{5-3}{3} = \frac{2}{3} .$$

**Answer (Ex. 3.43) —**

- 1.

$$\varphi_X(t) = E(e^{itX}) = \sum_{x=0}^{\infty} e^{itx} \theta(1-\theta)^x = \sum_{x=0}^{\infty} \theta (e^{it}(1-\theta))^x = \frac{\theta}{1 - e^{it}(1-\theta)} .$$

The last equality is due to  $\sum_{x=0}^{\infty} ar^x = \frac{a}{1-r}$  with  $a = \theta$  and  $r = e^{it}(1-\theta)$ .

2.

$$\begin{aligned}
E(X) &= \frac{1}{i} \left[ \frac{d}{dt} \left( \frac{\theta}{1 - e^{it}(1 - \theta)} \right) \right]_{t=0} = \frac{1}{i} \theta \left[ \frac{d}{dt} \left( (1 - e^{it}(1 - \theta))^{-1} \right) \right]_{t=0} \\
&= \frac{1}{i} \theta \left[ - (1 - e^{it}(1 - \theta))^{-2} \frac{d}{dt} (1 - e^{it}(1 - \theta)) \right]_{t=0} \\
&= \frac{1}{i} \theta \left[ - (1 - e^{it}(1 - \theta))^{-2} (0 - ie^{it}(1 - \theta)) \right]_{t=0} \\
&= \frac{1}{i} \theta \left( - (1 - e^0(1 - \theta))^{-2} (-ie^0(1 - \theta)) \right) \\
&= \frac{1}{i} \theta \left( - (1 - 1 + \theta)^{-2} (-i(1 - \theta)) \right) \\
&= \frac{1 - \theta}{\theta} .
\end{aligned}$$

**Answer (Ex. 3.44)** —

$$\begin{aligned}
\varphi_X(t) &= E(e^{itX}) = \int_{-\infty}^{\infty} e^{itx} f_X(x; a, b) dx = \int_a^b e^{itx} \frac{1}{b-a} dx = \frac{1}{b-a} \int_a^b e^{itx} dx = \frac{1}{b-a} \left[ \frac{e^{itx}}{it} \right]_{x=a}^b \\
&= \frac{1}{(b-a)it} (e^{itb} - e^{ita}) .
\end{aligned}$$

**Answer (Ex. 3.45)** —

1.

$$\varphi_X(t) = E(e^{itX}) = \sum_{x=0}^{\infty} e^{itx} \frac{\lambda^x}{x!} e^{-\lambda} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{e^{itx} \lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{(\lambda e^{it})^x}{x!} = e^{-\lambda} e^{\lambda e^{it}} = e^{\lambda e^{it} - \lambda} .$$

The second-last equality above is using  $e^\alpha = \sum_{x=0}^{\infty} \frac{\alpha^x}{x!}$  with  $\alpha = \lambda e^{it}$ .

2. To find  $V(X)$  using  $\varphi_X(t)$  we need  $E(X)$  and  $E(X^2)$ .

$$\begin{aligned}
E(X) &= \frac{1}{i} \left[ \frac{d}{dt} \varphi_X(t) \right]_{t=0} = \frac{1}{i} \left[ \frac{d}{dt} e^{\lambda e^{it} - \lambda} \right]_{t=0} = \frac{1}{i} \left[ e^{\lambda e^{it} - \lambda} \frac{d}{dt} (\lambda e^{it} - \lambda) \right]_{t=0} \\
&= \frac{1}{i} \left[ e^{\lambda e^{it} - \lambda} \lambda i e^{it} \right]_{t=0} = \frac{1}{i} (e^{\lambda - \lambda} \lambda i) = \frac{1}{i} (\lambda i) = \lambda .
\end{aligned}$$

$$\begin{aligned}
E(X^2) &= \frac{1}{i^2} \left[ \frac{d^2}{dt^2} \varphi_X(t) \right]_{t=0} = \frac{1}{i^2} \left[ \frac{d}{dt} \left( e^{\lambda e^{it} - \lambda} \lambda i e^{it} \right) \right]_{t=0} = \frac{1}{i^2} \left[ \frac{d}{dt} \left( \lambda i e^{\lambda e^{it} - \lambda + it} \right) \right]_{t=0} \\
&= \frac{1}{i^2} \left[ \lambda i e^{\lambda e^{it} - \lambda + it} \frac{d}{dt} (\lambda e^{it} - \lambda + it) \right]_{t=0} = \frac{1}{i^2} \left[ \lambda i e^{\lambda e^{it} - \lambda + it} (\lambda i e^{it} - 0 + i) \right]_{t=0} \\
&= \frac{1}{i^2} (\lambda i e^{\lambda e^0 - \lambda + 0} (\lambda i e^0 + i)) = \frac{1}{i^2} (\lambda i (\lambda i + i)) = \frac{1}{i^2} (\lambda i^2 (\lambda + 1)) = \lambda^2 + \lambda .
\end{aligned}$$

Finally,

$$V(X) = E(X^2) - (E(X))^2 = \lambda^2 + \lambda - \lambda^2 = \lambda .$$

**Answer (Ex. 3.46) —**

$$\varphi_W(t) = \varphi_{X+Y}(t) = \varphi_X(t) \times \varphi_Y(t) = e^{\lambda e^{it} - \lambda} \times e^{\mu e^{it} - \mu} = e^{\lambda e^{it} - \lambda + \mu e^{it} - \mu} = e^{(\lambda + \mu)e^{it} - (\lambda + \mu)}.$$

So,  $W$  is a Poisson( $\lambda + \mu$ ) RV. Thus the sum of two independent Poisson RVs is also a Poisson RV with parameter given by the sum of the parameters of the two RVs being added. The same idea generalizes to the sum of more than two Poisson RVs.

**Answer (Ex. 3.47) —**

We can use the facts by noting  $Y = -Z = 0 + (-1) \times Z$ , with  $a = 0$  and  $b = -1$  in  $Y = a + bX$  and get

$$\varphi_Y(t) = e^{i \times 0 \times t} \varphi_Z(-1 \times t) = \varphi_Z(-t) = e^{-((-1) \times t)^2 / 2} = e^{-t^2 / 2} = \varphi_Z(t)$$

Thus,  $\varphi_{-Z}(t) = \varphi_Z(t)$  and therefore the distributions of  $Z$  and  $-Z$  are the same. This should make sense because by switching signs of a symmetric (about 0) RV you have not changed its distribution! Note: we are not saying  $Z = -Z$  but just that their distributions are the same, i.e.,  $F_Z(z) = F_{-Z}(z)$  for every  $z \in \mathbb{R}$ .