

# Modeling social network dynamics in the spotted hyena (*Crocuta crocuta*)

Conrad Beckmann<sup>1</sup>, Judy Bloom<sup>2</sup>, Jacob Lambert<sup>3</sup>, Rachael Ward<sup>4</sup>, Amiyaal Ilany<sup>5</sup>, and Jiang Jiang<sup>5</sup>

<sup>1</sup>University of Vermont, <sup>2</sup>Eastern Connecticut State University, <sup>3</sup>University of Tennessee, <sup>4</sup>Rhodes College

<sup>5</sup>National Institute for Mathematical and Biological Synthesis

## Introduction

- Spotted hyenas (*Crocuta crocuta*) maintain a complex social structure.
- Sub-Saharan African hyenas live in large clans consisting of 10 – 100 individuals.
- Females are dominant to all males and the social structure is matrilineal.
- Each hyena has a specific rank determined by its matriline, which affects the individual's access to resources.

- We focused on the Talek clan in Maasai Mara National Reserve, Kenya.
- Over the 22 year study, 80,000 observations of hyena interactions were collected, along with profiles of each hyena.
- We are developing a model which will show how matrilineal rank, fitness, and connectivity of the network changes over time.

## Methods

- All data were divided by year for females in the Talek clan.
- Dyadic connections were calculated using the Twice-Weight Association Index.

$$AI_{a,b} = \frac{obs_{a\_with\_b}}{obs_{a\_without\_b} + obs_{b\_without\_a} + obs_{a\_with\_b}}$$

**Equation 1.** Calculated from observations of two hyenas *a* and *b*

- Matrilineal rank was determined by maternal data.
- Ranks were determined from dyadic aggressive interactions.

$$CB = \frac{1 + B + \sum_{i=1}^B b_i}{1 + L + \sum_{i=1}^L l_i}$$

**Equation 2.** The Clutton-Brock Index uses dyadic aggressive behavior to calculate social rank.

- Fitness was calculated for all individuals *h* and matrilineal rank *m*.

$$fitness_h = \frac{births_h}{\sum_i births_i} \quad fitness_m = \frac{births_m}{\sum_i births_i}$$

**Equations 3 and 4.** Ratio of individual and matrilineal births to total births.




## Preliminary Results

- Connectivity and matrilineal rank appear to affect trends in rank over time.
- Matrilineal rank of higher ranks maintained their rank over time.
- Lower ranked matrilineal lines were less stable and more likely to disappear or increase in rank.
- Increases in rank appear to be affected by connections with higher ranked individuals.

## Future Work

- Create a mathematical model which will predict fitness over time based on rank and network connectivity.
- Compare the effects of internal connectivity within a matriline and external connectivity between matrilineal lines.
- Generalize the model to apply to other complex social network structures.



**Figure 1.** Social network between spotted hyenas for different years. Node color represents matriline. Node size is proportional to rank; highest rank is the largest. Edge-thickness is proportional to strength of the association.

## References

- Holekamp K.E., Smith J.E., Strelhoff C.C., van Horn R.C., and H.E. Watts. 2012. Society, demography, and genetic structure in the spotted hyena. *Molecular Ecology* 21:613-632.
- Lewis K., Gonzalez M., and J. Kaufman. 2011. Social selection and peer influence in an online social network. *Proceedings of the National Academy of Sciences* 109(1):68-72.
- Hobson E.A., Avery M.L., and T.F. Wright. 2013. An analytical framework for quantifying and testing patterns of temporal dynamics in social networks. *Animal Behaviour* 85:83-96.

**Acknowledgments:** This work was conducted with funding from the National Institute for Mathematical and Biological Synthesis, an Institute sponsored by the National Science Foundation, the U.S. Department of Homeland Security, and the U.S. Department of Agriculture through NSF Award #EF-0832858, with additional support from The University of Tennessee, Knoxville. We thank Kay Holekamp for allowing us to use the data collected from her lab.