

# The untyped lambda calculus

CIS 6700, Spring 2023

June 28, 2023

## 1 Untyped lambda calculus

This document describes the untyped lambda-calculus, with the following grammar. References are to definitions in Barendregt, “The Lambda Calculus: Its Syntax and Semantics”.

$tm, t, u, v, w$	$::=$	terms
		$x$ variables
		$\lambda x.t$ abstractions
		$t u$ function application
		$k$
		<b>add</b>
		<b>nil</b>
		$t, u$

In  $\beta$ -reduction, the argument of an application is substituted for the bound variable in an abstraction. We use the  $t[x \rightsquigarrow u]$  notation for substituting the term  $u$  for the variable  $x$  in the term  $t$ .

$t$   $\beta$ -reduces to  $u$  (Definition 3.1.3)

BETA-REDUCT

$$\frac{}{(\lambda x.t) u \beta\text{-reduces to } t[x \rightsquigarrow u]}$$

We can define a deterministic, small-step evaluation relation by reducing in the heads of applications. This is a *call-by-name* semantics and performs head-reduction.

$t \rightsquigarrow t'$  (Small-step evaluation)

S-APP1 $\frac{t \rightsquigarrow t'}{t u \rightsquigarrow t' t}$	S-BETA $\frac{}{(\lambda x.t) u \rightsquigarrow t[x \rightsquigarrow u]}$	S-CONS1 $\frac{t \rightsquigarrow t'}{t, u \rightsquigarrow t', t}$	S-CONS2 $\frac{t \rightsquigarrow t' \quad \mathbf{value} \ v}{v, t \rightsquigarrow v, t'}$
S-PRJ-ZERO $\frac{\mathbf{value} \ (v, w)}{(v, w) \ 0 \rightsquigarrow v}$	S-PRJ-SUC $\frac{\mathbf{value} \ (v, w)}{(v, w) \ (1 + k) \rightsquigarrow (v, w) \ k}$	S-ADD $\frac{}{\mathbf{add} \ (k_1, k_2) \rightsquigarrow k_1 + k_2}$	

We can also define a nondeterministic single-step full-reduction by performing  $\beta$ -reduction in any subterm. Iterating this reduction will convert a term into its  $\beta$ -normal form.

$$\boxed{t \longrightarrow_{\beta} u} \quad (\text{Full CBV } \beta\text{-reduction})$$

$\frac{\text{F-BETA} \quad \text{value } v}{(\lambda x.t) v \longrightarrow_{\beta} t[x \rightsquigarrow v]}$	$\frac{\text{F-ABS} \quad t \longrightarrow_{\beta} t'}{\lambda x.t \longrightarrow_{\beta} \lambda x.t'}$	$\frac{\text{F-APP1} \quad t \longrightarrow_{\beta} t'}{t u \longrightarrow_{\beta} t' u}$
$\frac{\text{F-APP2} \quad u \longrightarrow_{\beta} u'}{t u \longrightarrow_{\beta} t u'}$	$\frac{\text{F-CONS1} \quad t \longrightarrow_{\beta} t'}{t, u \longrightarrow_{\beta} t', u}$	$\frac{\text{F-CONS2} \quad u \longrightarrow_{\beta} u'}{t, u \longrightarrow_{\beta} t, u'}$
	$\frac{\text{F-PRJ-ZERO} \quad \text{value } v}{(v, w) 0 \longrightarrow_{\beta} v}$	
$\frac{\text{F-PRJ-SUC} \quad \text{value } v}{(v, w) (1 + k) \longrightarrow_{\beta} (v, w) k}$	$\frac{\text{F-ADD}}{\text{add } (k_1, k_2) \longrightarrow_{\beta} k_1 + k_2}$	

We can define when two terms are equivalent up to  $\beta$ .

$$\boxed{t \equiv_{\beta} u} \quad (\beta\text{-conversion or } \beta\text{-equality (Definition 2.1.4)})$$

$\frac{\text{EQ-BETA}}{(\lambda x.t) u \equiv_{\beta} t[x \rightsquigarrow u]}$	$\frac{\text{EQ-REFL}}{t \equiv_{\beta} t}$	$\frac{\text{EQ-SYM} \quad u \equiv_{\beta} t}{t \equiv_{\beta} u}$	$\frac{\text{EQ-TRANS} \quad t_1 \equiv_{\beta} t_2 \quad t_2 \equiv_{\beta} t_3}{t_1 \equiv_{\beta} t_3}$
$\frac{\text{EQ-APP1} \quad t \equiv_{\beta} t'}{t u \equiv_{\beta} t' u}$	$\frac{\text{EQ-APP2} \quad u \equiv_{\beta} u'}{t u \equiv_{\beta} t u'}$	$\frac{\text{EQ-ABS} \quad t \equiv_{\beta} t'}{\lambda x.t \equiv_{\beta} \lambda x.t'}$	$\frac{\text{EQ-CONS1} \quad t \equiv_{\beta} t'}{t, u \equiv_{\beta} t', u}$
	$\frac{\text{EQ-CONS2} \quad u \equiv_{\beta} u'}{t, u \equiv_{\beta} t, u'}$		

Finally, the Church-Rosser Theorem relies on the definition of parallel reduction. This is a version of reduction that is confluent.

$$\boxed{t \Longrightarrow_{\beta} u} \quad (\text{Parallel reduction (3.2.3)})$$

$\frac{\text{P-BETA} \quad t \Longrightarrow_{\beta} \lambda x.t' \quad u \Longrightarrow_{\beta} v \quad \text{value } v}{t u \Longrightarrow_{\beta} t'[x \rightsquigarrow v]}$	$\frac{\text{P-VAR}}{x \Longrightarrow_{\beta} x}$	$\frac{\text{P-ABS} \quad t \Longrightarrow_{\beta} t'}{\lambda x.t \Longrightarrow_{\beta} \lambda x.t'}$	$\frac{\text{P-APP} \quad t \Longrightarrow_{\beta} t' \quad u \Longrightarrow_{\beta} u'}{t u \Longrightarrow_{\beta} t' u'}$
---	--	--	--

$$\begin{array}{c}
\text{P-APP-0} \\
\frac{t \Rightarrow_{\beta} v, w \quad u \Rightarrow_{\beta} 0 \quad \text{value } v}{t \, u \Rightarrow_{\beta} v}
\end{array}
\qquad
\begin{array}{c}
\text{P-APP-N} \\
\frac{t \Rightarrow_{\beta} v, w \quad u \Rightarrow_{\beta} (1 + k) \quad w \, k \Rightarrow_{\beta} t' \quad \text{value } v}{t \, u \Rightarrow_{\beta} t'}
\end{array}
\qquad
\begin{array}{c}
\text{P-CONS} \\
\frac{t \Rightarrow_{\beta} t' \quad u \Rightarrow_{\beta} u'}{t, u \Rightarrow_{\beta} t', u'}
\end{array}$$

$$\begin{array}{c}
\text{P-NIL} \\
\frac{}{\text{nil} \Rightarrow_{\beta} \text{nil}}
\end{array}
\qquad
\begin{array}{c}
\text{P-ADD} \\
\frac{}{\text{add} \Rightarrow_{\beta} \text{add}}
\end{array}
\qquad
\begin{array}{c}
\text{P-NAT} \\
\frac{}{k \Rightarrow_{\beta} k}
\end{array}$$

## 2 Relation operations

Many of the operations above can be generated by applying the following closure operations to the  $\beta$  reduction relation. These operations are parameterized by an arbitrary relation  $R$ .

Note that  $t \rightarrow_R u$  with  $R$  equal to  $\beta$  is the same relation as  $t \rightarrow_{\beta} u$ . And, the compatible, reflexive, symmetric and transitive closure of  $\beta$  is the same relation as  $t \equiv_{\beta} u$ .

$\boxed{t \rightarrow_R u}$  (*Compatible closure of  $R$  (aka one-step reduction (3.1.5))*)

$$\begin{array}{c}
\text{CC-REL} \\
\frac{R \, t \, u}{t \rightarrow_R u}
\end{array}
\qquad
\begin{array}{c}
\text{CC-ABS} \\
\frac{t \rightarrow_R u}{\lambda x. t \rightarrow_R \lambda x. u}
\end{array}
\qquad
\begin{array}{c}
\text{CC-APP1} \\
\frac{t \rightarrow_R t'}{t \, u \rightarrow_R t' \, u}
\end{array}
\qquad
\begin{array}{c}
\text{CC-APP2} \\
\frac{u \rightarrow_R u'}{t \, u \rightarrow_R t \, u'}
\end{array}$$

$$\begin{array}{c}
\text{CC-CONS1} \\
\frac{t \rightarrow_R t'}{t, u \rightarrow_R t', u}
\end{array}
\qquad
\begin{array}{c}
\text{CC-CONS2} \\
\frac{u \rightarrow_R u'}{t, u \rightarrow_R t, u'}
\end{array}$$

$\boxed{t \rightarrow_{\overline{\overline{R}}} u}$  (*reflexive closure of  $R$  (3.1.4)*)

$$\begin{array}{c}
\text{R-REL} \\
\frac{R \, t \, u}{t \rightarrow_{\overline{\overline{R}}} u}
\end{array}
\qquad
\begin{array}{c}
\text{R-REFL} \\
\frac{}{t \rightarrow_{\overline{\overline{R}}} t}
\end{array}$$

$\boxed{t \rightarrow_R^+ u}$  (*transitive closure of  $R$  (3.1.4)*)

$$\begin{array}{c}
\text{T-REL} \\
\frac{R \, t \, u}{t \rightarrow_R^+ u}
\end{array}
\qquad
\begin{array}{c}
\text{T-TRANS} \\
\frac{t_1 \rightarrow_R^+ t_2 \quad t_2 \rightarrow_R^+ t_3}{t_1 \rightarrow_R^+ t_3}
\end{array}$$

$$\boxed{t \rightarrow_R^* u} \quad (\text{reflexive-transitive closure of } R)$$

$$\frac{\text{RT-REL} \quad R t u}{t \rightarrow_R^* u}$$

$$\frac{\text{RT-REFL}}{t \rightarrow_R^* t}$$

$$\frac{\text{RT-TRANS} \quad t_1 \rightarrow_R^* t_2 \quad t_2 \rightarrow_R^* t_3}{t_1 \rightarrow_R^* t_3}$$

$$\boxed{t \leftrightarrow_R u} \quad (\text{symmetric-transitive closure of } R)$$

$$\frac{\text{ST-REL} \quad R t u}{t \leftrightarrow_R u}$$

$$\frac{\text{ST-SYM} \quad u \leftrightarrow_R t}{t \leftrightarrow_R u}$$

$$\frac{\text{ST-TRANS} \quad t_1 \leftrightarrow_R t_2 \quad t_2 \leftrightarrow_R t_3}{t_1 \leftrightarrow_R t_3}$$

### 3 Eta-reduction

By changing the definition of the underlying primitive reduction, we can also reason about  $\eta$ -reduction and  $\beta\eta$ -equivalence. Note that the rule for  $\eta$ -reduction has been stated in a way that generates the appropriate output for the locally-nameless representation in Coq. In this output, the fact that  $x \notin \text{fv}t$  is implicit and does not need to be added as a precondition to the rule.

$$\boxed{t \text{ } \eta\text{-reduces to } u} \quad (\text{Definition 3.1.1 (i)})$$

$$\frac{\text{ETA-REDUCT} \quad t' = t x}{\lambda x. t' \text{ } \eta\text{-reduces to } t}$$

$$\boxed{t \text{ } \beta\eta\text{-reduces to } u} \quad (\text{Either } \beta \text{ or } \eta \text{ reduction (Definition 3.1.1 (ii))})$$

$$\frac{\text{ETA-BETA} \quad t \text{ } \beta\text{-reduces to } u}{t \text{ } \beta\eta\text{-reduces to } u}$$

$$\frac{\text{ETA-ETA} \quad t \text{ } \eta\text{-reduces to } u}{t \text{ } \beta\eta\text{-reduces to } u}$$