

Extremes of Stationary Time Series – Day 1.

Dr. Ioannis Papastathopoulos (with illustrations by Lambert de Monte)

February 27, 2023



Introduction

- Introduction and motivation
- Notation for stochastic processes
- Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

Introduction

- In basic data analysis we often assume that observations are independent or even independent and identically distributed,

$$X_1, \dots, X_n \stackrel{\text{ind}}{\sim} F_1, \dots, F_n, \quad X_1, \dots, X_n \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2),$$

- Time series is the study of observations that arise in some order (usually time) and so are dependent: in other words

one d*** thing after another!**

- There are many more ways to be dependent than to be independent, and almost all data are collected in time order, so time series arise in a vast range of disciplines: climatology; economics; finance; marketing; epidemiology; biomedicine; genomics; environmental science; computer science; electrical engineering; physics; ...

Introduction

Introduction and motivation

Notation for stochastic processes

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

Definition.

(a) A **stochastic process** $\{X_t\}_{t \in \mathcal{T}}$ with index set \mathcal{T} is a family of random variables defined on a probability space (Ω, \mathcal{F}, P) .

(b) A **realisation** of $\{X_t\}$ is the outcome $\{x_t\}_{t \in \mathcal{T}} = \{X_t(\omega)\}_{t \in \mathcal{T}}$ for some $\omega \in \Omega$.

About the index set

- In time series usually $\mathcal{T} = \mathbb{R}, \mathbb{R}_+$ or \mathbb{Z} ;
- owing to digitisation, \mathcal{T} cannot in practice contain a sub-interval of \mathbb{R} , but the time step Δt can be very small in some applications;
- (almost-)continuous time series can be thinned by subsampling at the points of a grid, or, in some cases, by integration over intervals of width h (e.g., rainfall data);
- for general discussion, we take $\mathcal{T} = \mathbb{Z}$, so that X_t is recorded at times $0, \pm 1, \pm 2, \dots$, and write a realisation of the process observed for finite period as x_1, \dots, x_n

Introduction

Introduction and motivation

Notation for stochastic processes

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- We will explore the impact of dependence between values in the series on the extreme values when observed at long-range and short-range.

- Generally, one considers how dependence affects:
 - $M_n = \max(X_1, \dots, X_n)$;

 - exceedances of a high threshold.

Introduction

Introduction and motivation

Notation for stochastic processes

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- We will explore the impact of dependence between values in the series on the extreme values when observed at long-range and short-range.

- Generally, one considers how dependence affects:
 - $M_n = \max(X_1, \dots, X_n)$;

 - exceedances of a high threshold.

Introduction

Introduction and motivation

Notation for stochastic processes

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- We will explore the impact of dependence between values in the series on the extreme values when observed at long-range and short-range.
- Generally, one considers how dependence affects:
 - $M_n = \max(X_1, \dots, X_n)$;
 - exceedances of a high threshold.

Introduction
Stationarity and stationary sequences
Definition of Stationarity
Long-range asymptotic independence condition
Long-range asymptotic independence condition
Long-range asymptotic independence condition
Asymptotic independence of Maxima (AIM)
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

Stationarity and stationary sequences

Introduction

Stationarity and stationary sequences

Definition of Stationarity

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Asymptotic independence of Maxima (AIM)

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- A process $\{X_t\}$ is said to be a stationary process if the joint distributions of

$$(X_{t_1}, \dots, X_{t_k}) \quad \text{and} \quad (X_{t_1+\tau}, \dots, X_{t_k+\tau})$$

are the same for any k, t_1, \dots, t_k , and τ .

- Throughout we will assume that the univariate marginal distribution function is F , *i.e.*,

$$F(x) = \mathbb{P}(X_t \leq x)$$

for all t .

Long-range asymptotic independence condition

Introduction

Stationarity and stationary sequences

Definition of Stationarity

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Asymptotic independence of Maxima (AIM)

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

We suppose that there exist normalising sequences $a_n > 0$ and b_n such that

$$\frac{M_n - b_n}{a_n}$$

has a non-degenerate limit distribution.

We want to characterise the limit behaviour of M_n .

Recall: $M_n = \max(X_1, \dots, X_n)$.

Long-range asymptotic independence condition

Introduction

Stationarity and stationary sequences

Definition of Stationarity

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Asymptotic independence of Maxima (AIM)

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

In the absence of any conditions that limit the amount of long-range dependence that can be present in the values of the series, any limit distribution can be obtained.

For example let $X_t = X_1$ for all t , then for all n ,

$$\mathbb{P}(M_n \leq x) = \mathbb{P}(X_1 \leq x) = F(x),$$

so the class of limit distributions covers all distributions.

Long-range asymptotic independence condition

Introduction

Stationarity and stationary sequences

Definition of Stationarity

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Asymptotic independence of Maxima (AIM)

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- In the absence of any conditions that limit the amount of long-range dependence that can be present in the values of the series, any limit distribution can be obtained.

- For example let $X_t = X_1$ for all t , then for all n ,

$$\mathbb{P}(M_n \leq x) = \mathbb{P}(X_1 \leq x) = F(x),$$

so the class of limit distributions covers all distributions.

Asymptotic independence of Maxima (AIM)

Introduction

Stationarity and stationary sequences

Definition of Stationarity

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Long-range asymptotic independence condition

Asymptotic independence of Maxima (AIM)

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- Let $M_{i,j} = \max(X_i, \dots, X_j)$ and $u_n = a_n x + b_n$ for a_n, b_n defined above and x any real number.
- **AIM(u_n) condition:** there exists a sequence q_n of positive integers with $q_n = o(n)$ such that for all i and j

$$|\mathbb{P}(M_{1,i} \leq u_n, M_{i+q_n, i+q_n+j} \leq u_n) - \mathbb{P}(M_{1,i} \leq u_n)\mathbb{P}(M_{1,j} \leq u_n)| \rightarrow 0,$$

as $n \rightarrow \infty$.

- The condition ensures that separated groups of extreme points become increasingly close to being independent as their separation and level both increase at appropriate rates.

See Leadbetter et al. [1983] and O'Brien [1987] for other similar conditions.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Unified Extremal Types Theorem
Outline strategy of proof
Notes on the Theorem
Extremes of Fréchet marginal variables
Extremes of Fréchet marginal variables
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of
stationary processes

Examples

References

Theorem Suppose that there exist normalising sequences a_n and b_n such that

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \rightarrow H(x), \quad n \rightarrow \infty,$$

where H is a non-degenerate distribution, and that the $\text{AIM}(a_n x + b_n)$ condition holds, then H is of the same type¹ as

$$\exp[-(1 + \xi x)_+^{-1/\xi}],$$

i.e. a GEV limit distribution.

¹Recall: We say that distribution functions F_1 and F_2 are of the same type if there exist constants $a > 0$ and b such that $F_2(ax + b) = F_1(x)$ for all x .

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
 - Unified Extremal Types Theorem
 - Outline strategy of proof
 - Notes on the Theorem
 - Extremes of Fréchet marginal variables
 - Extremes of Fréchet marginal variables
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
- References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of
stationary processes

Examples

References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
 - Unified Extremal Types Theorem
 - Outline strategy of proof**
 - Notes on the Theorem
 - Extremes of Fréchet marginal variables
 - Extremes of Fréchet marginal variables
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
- References

1. Partition time into a series of alternating long and short blocks, k of each.
2. Let the numbers in long and short blocks be r_n and q_n respectively with $r_n \rightarrow \infty$, $q_n \rightarrow \infty$ and $q_n/r_n \rightarrow 0$.
3. In a neighbouring pair of long and short blocks, the maximum over the two blocks is likely to be in the long block.
4. All maxima tend to fall into the long blocks.
5. Long block maxima are approximately independent as they are separated in time by the short blocks, so satisfy $\text{AIM}(u_n)$ condition.
6. Long block maxima satisfy the max-stability property, and hence are GEV.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of
stationary processes

Examples

References

- At a practical level this is all we need, the maximum of a stationary sequence that has some independence at long-range follows a GEV distribution.
- The result does not show how the dependence changes the behaviour of M_n .
- **We separate marginal and dependence features** for the remainder and subsequently fix the marginal distribution F of the $\{X_t\}$ to be Fréchet.

Extremes of Fréchet marginal variables

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Unified Extremal Types Theorem

Outline strategy of proof

Notes on the Theorem

Extremes of Fréchet marginal variables

Extremes of Fréchet marginal variables

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of
stationary processes

Examples

References

Let $\{X_t\}$ be a stationary process which has Fréchet marginal distributions, *i.e.*

$$F(x) = \exp(-1/x) \text{ for } x > 0.$$

Let \widehat{M}_n denote the maximum of n IID variables with marginal distribution F , then

$$\mathbb{P} \left(\frac{\widehat{M}_n}{n} \leq x \right) = \{F(nx)\}^n = \exp(-1/x) = G(x),$$

i.e., \widehat{M}_n/n has a Fréchet limit distribution for all n and we denote the limit distribution of the maximum of the IID variables by G .

Extremes of Fréchet marginal variables

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
 - Unified Extremal Types Theorem
 - Outline strategy of proof
 - Notes on the Theorem
 - Extremes of Fréchet marginal variables
 - Extremes of Fréchet marginal variables
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
- References

Suppose that $M_n = \max(X_1, \dots, X_n)$, that $\{X_t\}$ satisfies the AIM(nx) condition and that

$$\mathbb{P}\left(\frac{M_n}{n} \leq x\right) \rightarrow H(x),$$

then we can assess the effect of dependence on M_n by looking at the difference between H and G .


Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Moving Maxima
Marginal properties
Moving maxima dependence properties
Derivation of H
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

Motivating example

Moving Maxima

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Moving Maxima
Marginal properties
Moving maxima dependence properties
Derivation of H
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

Let $\{Y_t\}$ be IID with $F_Y(y) = \exp[-1/(2y)]$, *i.e.*, Fréchet type marginals. Define, for all t , $X_t = \max(Y_t, Y_{t-1})$.



[Introduction](#)

[Stationarity and stationary sequences](#)

[Unified Extremal Types Theorem](#)

[Motivating example](#)[Moving Maxima](#)[Marginal properties](#)[Moving maxima dependence properties](#)[Derivation of \$H\$](#)

[Extremal index](#)

[Extreme value distributions](#)

[Statistical inference for extremes of stationary processes](#)

[Examples](#)

[References](#)

The marginal distribution of the $\{X_t\}$ process is:

$$\begin{aligned}\mathbb{P}(X_t \leq x) &= \mathbb{P}(Y_t \leq x, Y_{t-1} \leq x) \\ &= \{F_Y(x)\}^2 \\ &= \exp[-2/(2x)] \\ &= \exp(-1/x),\end{aligned}$$

i.e., Fréchet distributed.

Moving maxima dependence properties

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Moving Maxima

Marginal properties

Moving maxima dependence properties

Derivation of H

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- For $\tau > 1$, X_t is independent of $X_{t+\tau}$ as the associated Y variables are all different for each of the X 's and hence the X 's are independent.
- For $\tau = 1$, X_t is dependent with $X_{t+\tau}$ as both X_t and X_{t+1} are functions of Y_t .
- It is hence clear that long-range independence holds.
- Extreme values occur as groups or clusters of independent values, with two values per cluster for all the largest values.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Moving Maxima
Marginal properties
Moving maxima dependence properties
Derivation of H
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

$$\begin{aligned}\mathbb{P}\left(\frac{M_n}{n} \leq x\right) &= \mathbb{P}(X_1 \leq nx, \dots, X_n \leq nx) = \mathbb{P}(Y_0 \leq nx, Y_1 \leq nx, \dots, Y_n \leq nx) \\ &= [F_Y(nx)]^{n+1} = [\exp\{-1/(2nx)\}]^{n+1} \\ &= \exp\{-(n+1)/(2nx)\} \rightarrow \exp\{-1/(2x)\} \quad \text{as } n \rightarrow \infty \\ &= \{G(x)\}^{1/2} = H(x).\end{aligned}$$

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extremal index
Features of the extremal index
Extremal index for moving maxima example
Implications of short-range extremal dependence
Implications of short-range extremal dependence
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

Extremal index

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
 - Extremal index
 - Features of the extremal index
 - Extremal index for moving maxima example
 - Implications of short-range extremal dependence
 - Implications of short-range extremal dependence
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
- References

- The previous example shows that short-range dependence in the extreme values of the process affects the limiting distribution of M_n .
- First we present a measure of short-range extremal dependence, termed the extremal index.
- Define the extremal index θ , for variables with Fréchet marginals, by

$$\begin{aligned}\theta &= \lim_{n \rightarrow \infty} \mathbb{P}(M_{2,p_n} \leq n | X_1 > n) \\ &= \lim_{n \rightarrow \infty} \{1 - \mathbb{P}(M_{2,p_n} > n | X_1 > n)\}\end{aligned}$$

where $p_n = o(n)$.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extremal index

Features of the extremal index

Extremal index for moving maxima example

Implications of short-range extremal dependence

Implications of short-range extremal dependence

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

- $0 \leq \theta \leq 1$.
- Larger values of θ correspond to weaker short-range extremal dependence.
- For an IID process $\theta = 1$ as

$$\begin{aligned}\theta &= \lim_{n \rightarrow \infty} \{F(n)\}^{p_n} \\ &= \lim_{n \rightarrow \infty} \exp(-p_n/n) \\ &= 1,\end{aligned}$$

since $p_n/n \rightarrow 0$.

- If a process is independent for all lags $\tau \geq m$, for some finite m then $1/m \leq \theta \leq 1$.

Extremal index for moving maxima example

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
 - Extremal index
 - Features of the extremal index
 - Extremal index for moving maxima example
 - Implications of short-range extremal dependence
 - Implications of short-range extremal dependence
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
- References

■ From the realisation of the process it is clear that $\theta = 1/2$.

■ A derivation is

$$\begin{aligned}\mathbb{P}(M_{2,p_n} \leq n | X_1 > n) &= \mathbb{P}(Y_1 \leq n, \dots, Y_{p_n} \leq n | \max(Y_0, Y_1) > n) \\ &\approx \mathbb{P}(Y_1 \leq n, \dots, Y_{p_n} \leq n | Y_0 > n) \frac{1}{2} + \mathbb{P}(Y_1 \leq n, \dots, Y_{p_n} \leq n | Y_1 > n) \frac{1}{2} \\ &\rightarrow 1 \times \frac{1}{2} + 0 \times \frac{1}{2}, \quad n \rightarrow \infty \\ &= \frac{1}{2} = \theta.\end{aligned}$$

Implications of short-range extremal dependence

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extremal index

Features of the extremal index

Extremal index for moving maxima example

Implications of short-range extremal dependence

Implications of short-range extremal dependence

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

Combining the above results and examples, the following result should appear to be quite natural.

For variables with Fréchet marginal distributions, provided that

$$\mathbb{P} \left(\frac{M_n}{n} \leq x \right) \rightarrow H(x), \quad n \rightarrow \infty$$

to a non-degenerate limit H and that

■ AIM(nx) condition holds;

■ θ exists,

then

$$H(x) = \{G(x)\}^\theta.$$

Implications of short-range extremal dependence

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extremal index

Features of the extremal index

Extremal index for moving maxima example

Implications of short-range extremal dependence

Implications of short-range extremal dependence

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

References

The implications of this result are that, except for a really weak long-range dependence condition:

1. for stationary processes the effect of dependence is through θ only;
2. θ can be absorbed into the normalising constants, so dependence does not change the limiting type;
3. the limit suggests

$$H(x) = \lim_{n \rightarrow \infty} [\{F(nx)\}^n]^\theta = \lim_{n \rightarrow \infty} F(nx)^{n\theta},$$

so $n\theta$ can be thought of as an effective number of independent variables.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Distribution of the maximum
Threshold exceedances
Extremal index
Theorem
Comments on Theorem
Implications of the limit process: clusters
Implications of the limit process: exceedances
Statistical inference for extremes of stationary processes
Examples
References

Extreme value distributions

[Introduction](#)

[Stationarity and stationary sequences](#)

[Unified Extremal Types Theorem](#)

[Motivating example](#)

[Extremal index](#)

[Extreme value distributions](#)

[Distribution of the maximum](#)

[Threshold exceedances](#)

[Extremal index](#)

[Theorem](#)

[Comments on Theorem](#)

[Implications of the limit process: clusters](#)

[Implications of the limit process:
exceedances](#)

[Statistical inference for extremes of
stationary processes](#)

[Examples](#)

[References](#)

- Consider random variables X_1, \dots, X_n with arbitrary distribution function F and which satisfy the $\text{AIM}(u_n)$ condition.

- As $n \rightarrow \infty$,

$$\mathbb{P} \left(\frac{M_n - b_n}{a_n} \leq y \right) \rightarrow \exp \left[-\theta \left\{ 1 + \xi \left(\frac{y - \mu}{\sigma} \right) \right\}_+^{-1/\xi} \right],$$

where if the variables are IID then $\theta = 1$.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Distribution of the maximum
Threshold exceedances
Extremal index
Theorem
Comments on Theorem
Implications of the limit process: clusters
Implications of the limit process: exceedances
Statistical inference for extremes of stationary processes
Examples
References

We assume that

- $\{X_t\}$ is a stationary process with arbitrary distribution function F ;
- the required norming constants for an IID process with marginal F are a_n and b_n with limit distribution G , a $\text{GEV}(0, 1, \xi)$ distribution;
- a long-range asymptotic independence condition (similar to the $\text{AIM}(a_n x + b_n)$ condition) holds.

Consider the point processes

$$P_n = \left\{ \left(\frac{i}{n+1}, \frac{X_i - b_n}{a_n} \right) : i = 1, \dots, n \right\}$$


on $[0, 1] \times \mathbb{R}$.

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
 - Distribution of the maximum
 - Threshold exceedances
- Extremal index
 - Theorem
 - Comments on Theorem
 - Implications of the limit process: clusters
 - Implications of the limit process: exceedances
- Statistical inference for extremes of stationary processes
- Examples
- References

Point process P_n for $n = 10, 100, 1000, 10000$ respectively with

$$X_i \sim \text{Moving Max}(\alpha_0, \alpha_1, \alpha_2) = \max(\alpha_0 Y_i, \alpha_1 Y_{i-1}, \alpha_2 Y_{i-2}),$$

with $\alpha_0 = 1/3, \alpha_1 = 1/2, \alpha_2 = 1/6$.



Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Distribution of the maximum
Threshold exceedances
Extremal index
Theorem
Comments on Theorem
Implications of the limit process: clusters
Implications of the limit process: exceedances
Statistical inference for extremes of stationary processes
Examples
References

- Under the above conditions on P_n , on the set $[0, 1] \times (u, \infty)$, where $u > b_l = \lim_{n \rightarrow \infty} (x_F - b_n)/a_n$, then [Hsing et al., 1988, Rootzén, 1988]

$$P_n \xrightarrow{d} P \quad \text{as } n \rightarrow \infty,$$

where P is a clustered non-homogeneous Poisson process with cluster distribution π .

- The cluster distribution π is concentrated on $\{1, 2, \dots\}$ and has probability mass function

$$\pi(j) = \lim_{n \rightarrow \infty} \pi_n(j) = \lim_{n \rightarrow \infty} \mathbb{P} \left[\sum_{i=1}^{p_n} I \left(\frac{X_i - b_n}{a_n} > x \right) = j \mid \sum_{i=1}^{p_n} I \left(\frac{X_i - b_n}{a_n} > x \right) > 0 \right],$$

where $p_n = o(n)$ and $x > b_l$.

- The Laplace functional of the limiting Poisson process P is

$$-\log \mathbb{E} e^{-Pf} = \theta(1 + \xi y)^{-1/\xi} \int_0^1 [1 - \phi(f(t))] dt, \quad f \geq 0, \text{ Borel}$$

where $\phi(s) = \sum_{j=1}^{\infty} e^{-sj} \pi(j)$ is the Laplace transform of the cluster distribution π .

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
 - Distribution of the maximum
 - Threshold exceedances
 - Extremal index
 - Theorem
- Comments on Theorem
- Implications of the limit process: clusters
- Implications of the limit process: exceedances
- Statistical inference for extremes of stationary processes
- Examples
- References

- The extremal index satisfies $\theta^{-1} = \sum_{j=1}^{\infty} j\pi(j)$.
- The set of cluster maxima (the largest value in each vertical string of points in the point process) form a non-homogeneous Poisson process with intensity

$$\lambda(t, y) = \theta(1 + \xi y)_+^{-1-1/\xi}.$$

- For data over a high threshold u we absorb the location b_n and scale a_n normalisation into the parameters to give the intensity of the non-homogeneous Poisson Process for the cluster maxima as

$$\lambda(t, y) = \theta\sigma^{-1} \left[1 + \xi \left(\frac{y - \mu}{\sigma} \right) \right]_+^{-1-1/\xi}.$$

Implications of the limit process: clusters

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
 - Distribution of the maximum
 - Threshold exceedances
 - Extremal index
 - Theorem
 - Comments on Theorem
- Implications of the limit process: clusters
- Implications of the limit process: exceedances
- Statistical inference for extremes of stationary processes
- Examples
- References

Fixing a threshold level $u > b_l$ we have that:

1. clusters are those set of points exceeding u which occur at the same normalised time. In the original series these are large points which occur within a short time period of one another.
2. the expected number of exceedances of u per cluster (for which the cluster maxima exceeds u) is θ^{-1} , irrespective of the level u ;
3. relative to independent series, when $\theta < 1$ there are fewer clusters (by a factor θ) and more exceedances per cluster (by a factor θ^{-1});
4. values in one cluster are independent of values in another cluster;
5. values within a cluster are dependent.

Implications of the limit process: exceedances

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
 - Distribution of the maximum
 - Threshold exceedances
 - Extremal index
 - Theorem
 - Comments on Theorem
- Implications of the limit process: clusters
- Implications of the limit process: exceedances
- Statistical inference for extremes of stationary processes
- Examples
- References

Fixing a threshold level $u > b_l$ we have that:

- 1. the rate of cluster maxima exceeding the threshold u is

$$\theta[1 + \xi(u - \mu)/\sigma]_+^{-1/\xi};$$

- 2. the rate of arbitrary values exceeding the threshold u is

$$[1 + \xi(u - \mu)/\sigma]_+^{-1/\xi};$$

- 3. cluster maxima excesses over u are independent, and follow a $\text{GPD}(\sigma_u, \xi)$, with shape parameter ξ ; $\text{GPD}(\sigma_u, \xi)$ model for cluster maxima excesses over u , where

$$\sigma_u = \sigma + \xi(u - \mu);$$

- 4. arbitrary (in time) excesses over u also follow a $\text{GPD}(\sigma_u, \xi)$, *i.e.* with the same parameters as the cluster maxima.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Outline of inference procedure
Identifying independent clusters
The runs method: declustering
The runs method: extremal index estimation
Intervals estimator based on interexceedances
Intervals estimator based on interexceedances
Extremal index estimation based on interexceedance times
Extremal index: quantifying uncertainty
Examples
References

Statistical inference for extremes of stationary processes

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

We consider the additional practical difficulties encountered due to dependence in analysing the extreme values of a stationary process.

In a practical data analysis, we must:

- select a high threshold;
- identify clusters of extreme values which are independent from one another;
- estimate the extremal index.

When clusters are defined we

- extract the cluster maxima and fit the GPD distribution to the cluster maxima over the threshold;
- estimate characteristics of the cluster;
- assess sensitivity of cluster characteristics to threshold selection.

Identifying independent clusters

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

There are a number of ways of identifying independent clusters of extreme values.
Here we will focus on:

- the runs method, proposed by Smith and Weissman [1994];
- the blocks and intervals methods, see Ferro and Segers [2003];
- the identification of conditions that can be tested for independence between clusters, see Ledford and Tawn [2003].

The runs method: declustering

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

For a selected high threshold u we conclude that if consecutive exceedances of the threshold u are separated by a set of r consecutive observations below the threshold u then the exceedances belong to separate clusters.

Similarly, exceedances separated by less than r consecutive non-exceedances are deemed to be in the same cluster.

The choice of u and r is critical, r needs to be the time lag when the process is “independent in the extremes”.

The runs method: extremal index estimation

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

These motivate the runs estimator of the extremal index θ :

- Use the empirical form the extremal index,

$$\hat{\theta}(u, r) = \hat{\mathbb{P}}(M_{2,r+1} \leq u \mid X_1 > u),$$

using the empirical probability.

- $\hat{\theta}(u, r)$ is identical to $1/(\text{mean cluster size})$ with clusters defined by the runs method.
- $\hat{\theta}(u, r)$ can be obtained using the function `exi(data, u, r)` from the `evd` R package.

Intervals estimator based on interexceedances

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

- Another estimator can be constructed by exploiting the fact that the extremal index is related to the times between threshold exceedances.
- Let $\{X_t\}_{t \geq 1}$ be a process for which we consider the exceedances of a threshold u . Let F denote the marginal distribution of the X_i .
- Assume W.L.O.G. that $X_1 > u$, and let $T(u)$ denote a random waiting time until the next exceedance of u .
- Then, $T(u) = \min\{t \geq 1 : X_{t+1} > u\}$ and

$$\mathbb{P}(T(u) > n) = \mathbb{P}(\max\{X_2, \dots, X_{n+1}\} \leq u \mid X_1 > u).$$

- If the X_i are IID, then

$$\mathbb{P}(T(u) > n) = F(u)^n,$$

and it can be shown that $\{1 - F(u)\}T(u)$ is asymptotically unit exponential.

Intervals estimator based on interexceedances

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
 - Outline of inference procedure
 - Identifying independent clusters
 - The runs method: declustering
 - The runs method: extremal index estimation
 - Intervals estimator based on interexceedances
 - Intervals estimator based on interexceedances
 - Extremal index estimation based on interexceedance times
 - Extremal index: quantifying uncertainty
- Examples
- References

- When the X_i are not independent, *i.e.*, $\theta \in (0, 1)$, it can be shown that

$$\{1 - F(u)\}T(u) \xrightarrow{d} T_\theta,$$

where T_θ denotes a random variable distributed according to the mixture

$$(1 - \theta)\varepsilon_0 + \theta\mu_\theta$$

where ε_0 is the degenerate probability distribution at 0 and μ_θ is the exponential distribution with mean θ^{-1} .

- Note: θ is both the proportion of non-zero interexceedance times and the reciprocal of the mean of the non-zero interexceedance times.

Extremal index estimation based on interexceedance times

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

- Suppose that we observe n_u exceedances of u at times $1 \leq S_1 < \dots < S_{n_u} \leq n$.

- Compute the interexceedance times $T_i = S_{i+1} - S_i$ for $i = 1, \dots, n_u - 1$.

- Set

$$\hat{\theta}_n^\delta(u) = \frac{2 \left(\sum_{i=1}^{n_u-1} T_i \right)^2}{(n_u - 1) \sum_{i=1}^{n_u-1} T_i^2} \quad \text{and} \quad \hat{\theta}_n^\star(u) = \frac{2 \left(\sum_{i=1}^{n_u-1} T_i - 1 \right)^2}{(n_u - 1) \sum_{i=1}^{n_u-1} (T_i - 1)(T_i - 2)}.$$

- Then, the Ferro & Segers estimator $\hat{\theta}_{FS}$ is given by

$$\hat{\theta}_{FS} = \begin{cases} \min\{1, \hat{\theta}_n^\delta(u)\} & \text{if } \max\{T_i : 1 \leq i \leq n_u - 1\} \leq 2 \\ \min\{1, \hat{\theta}_n^\star(u)\} & \text{if } \max\{T_i : 1 \leq i \leq n_u - 1\} > 2 \end{cases}$$

- $\hat{\theta}_{FS}$ can be obtained using the function `exi(data, u, r=0)` from the `evd` R package.

Extremal index: quantifying uncertainty

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Outline of inference procedure

Identifying independent clusters

The runs method: declustering

The runs method: extremal index estimation

Intervals estimator based on interexceedances

Intervals estimator based on interexceedances

Extremal index estimation based on interexceedance times

Extremal index: quantifying uncertainty

Examples

References

- A convenient way to estimate uncertainty for an estimate $\hat{\theta}$ is to perform a bootstrap procedure.
- In the case of stationary sequences, a block bootstrap procedure is chosen to account for short range dependence structures.
 - Consider n realisations $\mathbf{x}_{1:n} = \{x_t\}_{t=1}^n$ from a time-series $\{X_t : t = 1, 2, \dots\}$.
 - Create a collection of moving blocks of size b , *e.g.*, $C = \{\mathbf{x}_{1:b}, \dots, \mathbf{x}_{i:(i+b-1)}, \dots, \mathbf{x}_{n:b}\}$.
 - Create a block bootstrap sample $\mathbf{X}_{1:n}$ by sampling n/b blocks from C with replacement.
- For each of m block bootstrap samples, estimate the extremal index, resulting in a collection $\hat{\boldsymbol{\theta}} = \{\hat{\theta}_1, \dots, \hat{\theta}_m\}$.
- Construct a quantile-based confidence interval from $\hat{\boldsymbol{\theta}}$.

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
Example: daily rainfall data
Extremal index for the rain data
Example: financial time series data
Example: financial time series data
Extremal index for the squared S&P500 returns
Squared standardised returns
Extremal index for the squared standardised returns
Modelling cluster maxima and all exceedances
Clusters and cluster maxima
Point process intensity estimation
Example: Squared S&P500 returns
Diagnostics of GPD fit
References

Examples

Example: daily rainfall data

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima


Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

We illustrate the use of these cluster identification and extremal index estimation using a time series of daily rainfall accumulations at a location in south-west England, recorded during 1914-1962.



The series is assumed to be stationary over this observation period.

Extremal index for the rain data

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances



Clusters and cluster maxima

Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit


References



- $\hat{\theta}(u, r)$ are threshold dependent;
- increasing r does not materially change our estimate $\hat{\theta}(u, r)$;
- $r = 1$ is appropriate for cluster identification;
- the value of $\hat{\theta}(u, r) \approx 1$ indicates weak short-range dependence with a limiting cluster size of 1 for higher thresholds;
- rainfall episodes tend to last a number of days, but this analysis tells us that extreme daily rainfall events tend to be isolated.

Example: financial time series data

We now turn to estimating clustering for the financial time series data.




- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
 - Example: daily rainfall data
 - Extremal index for the rain data
 - Example: financial time series data
 - Example: financial time series data
 - Extremal index for the squared S&P₅₀₀ returns
 - Squared standardised returns
 - Extremal index for the squared standardised returns
 - Modelling cluster maxima and all exceedances
 - Clusters and cluster maxima
 - Point process intensity estimation
 - Example: Squared S&P₅₀₀ returns
 - Diagnostics of GPD fit
- References

Example: financial time series data

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
 - Example: daily rainfall data
 - Extremal index for the rain data
 - Example: financial time series data
 - Example: financial time series data
 - Extremal index for the squared S&P500 returns
 - Squared standardised returns
 - Extremal index for the squared standardised returns
 - Modelling cluster maxima and all exceedances
 - Clusters and cluster maxima
 - Point process intensity estimation
 - Example: Squared S&P500 returns
 - Diagnostics of GPD fit
- References

We focus on the squared returns from the S&P500 share index.



Recall: The return R_t for day t is $R_t = (X_{t-1} - X_t)/X_{t-1}$.

Extremal index for the squared S&P500 returns

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima


Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

We compute the extremal index of the squared returns based on a runs estimator with $r = 10$.



- $\hat{\theta}(u, r)$ is stable for $u \geq 0.0003$ (dotted line) corresponding to the 0.92 quantile of the squared returns.
- The value of $\hat{\theta}(u \geq 0.0003, r) \approx 0.270$ (red line) suggests that the mean cluster size is around 3.70.
- $r = 10$ seems fine for cluster identification.

Squared standardised returns

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima


Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

We now investigate the source of serial dependence being the changing volatility of the process. We work with the squared standardised series:




We standardise the returns series by

- subtracting the local mean;
- dividing by the local standard deviation.

Local means and standard deviations were calculated using the $h = 21$ observations centred on the value to be standardised.

Extremal index for the squared standardised returns



- Estimates of $\hat{\theta}(u, r)$ are threshold dependent.
- Increasing r does not materially change our conclusions, $r = 1$ is fine for cluster identification.
- The limiting value of $\theta = 1$ indicates weak short-range independence with a limiting cluster size of 1.
- By undertaking the more sophisticated analysis and standardising the series we appear to have accounted for all the short-range dependence in the series.

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima

Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

Modelling cluster maxima and all exceedances

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima

Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

For a specific threshold u , we can

- model the cluster maxima:
 - identify the clusters for a given threshold u and run length r using the `clusters()` function from the `evd` package in R.
 - fit the $\text{GPD}(\sigma_u, \xi)$ model for cluster maxima excesses over u ;
 - standard errors can be obtained by usual methods as cluster maxima are independent.
- model all exceedances:
 - fit the $\text{GPD}(\sigma_u, \xi)$ model for the excess of all exceedances over u ;
 - the data are dependent so block bootstrap methods are needed for standard error evaluation.

Clusters and cluster maxima

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances


Clusters and cluster maxima

Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References



- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
 - Example: daily rainfall data
 - Extremal index for the rain data
 - Example: financial time series data
 - Example: financial time series data
 - Extremal index for the squared S&P500 returns
 - Squared standardised returns
 - Extremal index for the squared standardised returns
 - Modelling cluster maxima and all exceedances
 - Clusters and cluster maxima
 - Point process intensity estimation
 - Example: Squared S&P500 returns
 - Diagnostics of GPD fit
- References

To estimate μ and σ in the point process intensity, we need information in addition to that given by our estimate of σ_u and ξ .

This information is given by the expected number of cluster maxima exceeding u .

Equating the point process and sample values gives

$$\theta[1 + \xi(u - \mu)/\sigma]_+^{-1/\xi} = n_{\text{cl},u}.$$

It follows that

$$\widehat{\sigma} = \widehat{\sigma}_u(n_{\text{exc},u})^{\widehat{\xi}}$$

and

$$\widehat{\mu} = u + (\widehat{\sigma} - \widehat{\sigma}_u)/\widehat{\xi}.$$

θ is estimated by $\widehat{\theta}_u = n_{\text{cl},u}/n_{\text{exc},u}$, *i.e.*, the number of clusters above u divided by the number of exceedances of u .

Example: Squared S&P500 returns

Introduction

Stationarity and stationary sequences

Unified Extremal Types Theorem

Motivating example

Extremal index

Extreme value distributions

Statistical inference for extremes of stationary processes

Examples

Example: daily rainfall data

Extremal index for the rain data

Example: financial time series data

Example: financial time series data

Extremal index for the squared S&P500 returns

Squared standardised returns

Extremal index for the squared standardised returns

Modelling cluster maxima and all exceedances

Clusters and cluster maxima

Point process intensity estimation

Example: Squared S&P500 returns

Diagnostics of GPD fit

References

For the squared S&P500 returns:



- using a declustering parameter of $r = 10$ and a threshold of $u = 0.0004$ corresponding to the 0.95 quantile of the squared returns gives $n_{\text{cl},u} = 35$ independent clusters;
- there are $n_{\text{exc},u} = 140$ exceedances of u ;
- we estimate the extremal index as $\hat{\theta} = 0.25$;
- fitting a GPD to cluster maxima gives $\hat{\sigma}_u = 0.000215(0.00002)$, $\hat{\xi} = 0.905(0.271)$;
- combining the above estimates gives

$$\hat{\sigma} = \hat{\sigma}_u (n_{\text{exc},u})^{\hat{\xi}} = 0.0189$$

and

$$\hat{\mu} = u + (\hat{\sigma} - \hat{\sigma}_u) / \hat{\xi} = 0.0211.$$

- Introduction
- Stationarity and stationary sequences
- Unified Extremal Types Theorem
- Motivating example
- Extremal index
- Extreme value distributions
- Statistical inference for extremes of stationary processes
- Examples
 - Example: daily rainfall data
 - Extremal index for the rain data
 - Example: financial time series data
 - Example: financial time series data
 - Extremal index for the squared S&P500 returns
 - Squared standardised returns
 - Extremal index for the squared standardised returns
 - Modelling cluster maxima and all exceedances
 - Clusters and cluster maxima
 - Point process intensity estimation
 - Example: Squared S&P500 returns
 - Diagnostics of GPD fit
- References



Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

References

Introduction
Stationarity and stationary sequences
Unified Extremal Types Theorem
Motivating example
Extremal index
Extreme value distributions
Statistical inference for extremes of stationary processes
Examples
References

C. A. T. Ferro and J. Segers. Inference for clusters of extreme values. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 65(2):545–556, 2003. doi: <https://doi.org/10.1111/1467-9868.00401>. URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/1467-9868.00401>.

T. Hsing, J. Hüsler, and M. Leadbetter. On the exceedance point process for a stationary sequence. *Probab. Theory Reltd Flds*, 78:97–112, 1988.

M. R. Leadbetter, G. Lindgren, and H. Rootzen. *Extremes and Related Properties of Random Sequences and Processes*. Springer Series in Statistics. Springer Verlag, 1983. ISBN 0387907319. <https://doi.org/10.1007/978-1-4612-5449-2>.

A. W. Ledford and J. A. Tawn. Diagnostics for dependence within time series extremes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 65(2):521–543, 2003. doi: <https://doi.org/10.1111/1467-9868.00400>. URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/1467-9868.00400>.

G. L. O’Brien. Extreme Values for Stationary and Markov Sequences. *The Annals of Probability*, 15(1):281 – 291, 1987. doi: 10.1214/aop/1176992270. URL <https://doi.org/10.1214/aop/1176992270>.

H. Rootzén. Maxima and exceedances of stationary Markov chains. *Adv. in Appl. Probab.*, 20:371–390, 1988.

R. L. Smith and I. Weissman. Estimating the extremal index. *Journal of the Royal Statistical Society: Series B (Methodological)*, 56(3):515–528, 1994. doi: <https://doi.org/10.1111/j.2517-6161.1994.tb01997.x>. URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1994.tb01997.x>.