

Extremes of Stationary Time Series – Day 2

Dr. Ioannis Papastathopoulos (with illustrations by Lambert de Monte)

February 28, 2023



Introduction

- Motivation
- Asymptotic dependence
- Another view of the extremal dependence in stationary time series
- χ_1 for Moving Maxima process
- Lag- t χ for stationary Gaussian autoregressive process
- Properties of extremal dependence coefficients
- Invariance of extremal index
- Some limitations
- Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

Introduction

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian autoregressive process

Properties of extremal dependence coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- So far have identified^I that dependence in time series affects the behaviour of the extreme values, the latter typically occur in clusters.
- The extremal index θ can be seen as a function of the (unknown) limiting distribution of the cluster of renormalized exceedances [Hsing et al., 1988]

$$\pi_n(j) = \mathbb{P} \left[\sum_{i=1}^{p_n} I \left(\frac{X_i - b_n}{a_n} > x \right) = j \mid \sum_{i=1}^{p_n} I \left(\frac{X_i - b_n}{a_n} > x \right) > 0 \right],$$

where $p_n = o(n)$ and $x > b_l$.

- The extremal index satisfies

$$\theta^{-1} = \sum_{j=1}^{\infty} j \lim_{n \rightarrow \infty} \pi_n(j)$$

^IUnder suitable mixing conditions such as the AIM condition.

- Consider the measure of extremal dependence

$$\chi = \lim_{u \rightarrow \infty} \mathbb{P}(X_2 > u \mid X_1 > u) \in [0, 1]$$

- $\chi > 0$: random variables are called **asymptotically dependent**
- $\chi = 0$: random variables are called **asymptotically independent**

- Interpretation: as u approaches ∞ ,

$$\Lambda(u) = \frac{\mathbb{P}\{\max(X_1, X_2) > u\}}{\mathbb{P}(X_1 > u)} \rightarrow (2 - \chi).$$

Hence, in common margins, the probability of the maximum of two variables exceeding an extreme level is a scale factor of the probability of one of the variables exceeding this level.

- X_1 and X_2 independent implies $\chi = 0$
- X_1 and X_2 perfectly dependent implies $\chi = 1$
- Reverse implications not true in general.

Another view of the extremal dependence in stationary time series

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian autoregressive process

Properties of extremal dependence coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- We distinguish between two classes of extremal dependence

- The lag- t coefficient of asymptotic dependence is defined by

$$\chi_t = \lim_{u \rightarrow \infty} \mathbb{P}(X_t > u \mid X_0 > u) \quad (\text{I})$$

- If there exists a $t \neq 0$ such that $\chi_t > 0$ then the process is said to be *asymptotically dependent* and *asymptotically independent* otherwise.

- Asymptotically dependent time series have $\theta \in [0, 1)$

- Asymptotically independent time series have $\theta = 1$

Another view of the extremal dependence in stationary time series

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian autoregressive process

Properties of extremal dependence coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- We distinguish between two classes of extremal dependence

- The lag- t coefficient of asymptotic dependence is defined by

$$\chi_t = \lim_{u \rightarrow \infty} \mathbb{P}(X_t > u \mid X_0 > u) \quad (\text{I})$$

- If there exists a $t \neq 0$ such that $\chi_t > 0$ then the process is said to be *asymptotically dependent* and *asymptotically independent* otherwise.
- Asymptotically dependent time series have $\theta \in [0, 1)$
- Asymptotically independent time series have $\theta = 1$

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence
in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian
autoregressive process

Properties of extremal dependence
coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov
processes

Statistical inference with time series
conditional extremes

References

Let $X_t = \max(Y_{t-1}, Y_t)$ where Y_t IID with $F_Y(y) = \exp[-1/(2y)]$. Then

$$\begin{aligned}\mathbb{P}(X_2 > u \mid X_1 > u) &= \frac{\mathbb{P}(X_1 > u, X_2 > u)}{\mathbb{P}(X_1 > u)} \\&= \frac{1 - \mathbb{P}(X_1 \leq u) - \mathbb{P}(X_2 \leq u) + \mathbb{P}(X_1 \leq u, X_2 \leq u)}{\mathbb{P}(X_1 > u)} \\&= \frac{1 - 2e^{-1/u} + e^{-3/(2u)}}{1 - e^{-1/u}} \approx \frac{1 - 2(1 - \frac{1}{u}) + 1 - \frac{3}{2u}}{\frac{1}{u}} \quad \text{as } u \text{ approaches } \infty \\&\rightarrow \frac{1}{2}\end{aligned}$$

Recall $\mathbb{P}(A \cap B) = 1 - \mathbb{P}((A \cap B)^c) = 1 - \mathbb{P}(A^c) - \mathbb{P}(B^c) + \mathbb{P}(A^c \cap B^c)$

Lag- t χ for stationary Gaussian autoregressive process

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian autoregressive process

Properties of extremal dependence coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest


Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Let $X_t = \rho X_{t-1} + (1 - \rho^2)^{1/2} \varepsilon_t$ where $\varepsilon_t \sim \mathcal{N}(0, 1)$ with $\rho \in (-1, 1)$.
- Here $\chi_t = 0$ for all $t > 0$.
- Similarly, for this process we have $\theta = 1$, *i.e.*, extremes occur singly in the limit. Maximum is asymptotically the same as the maximum of IID variables with the same marginal distribution



Properties of extremal dependence coefficients

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian autoregressive process

Properties of extremal dependence coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Recall definition of the extremal index θ

$$\theta = \lim_{n \rightarrow \infty} \mathbb{P}(M_{2,p_n} \leq n | X_1 > n)$$

$$= \lim_{u \rightarrow \infty} \mathbb{P}(X_1 \leq u, \dots, X_{p_u} \leq u | X_1 > u)$$

where $p_n = o(n)$, for variables with Fréchet marginals.

- The extremal index and the coefficient of asymptotic dependence are invariant to changes in the choice of the marginal distribution.

Introduction

- Motivation
- Asymptotic dependence
- Another view of the extremal dependence in stationary time series
- χ_1 for Moving Maxima process
- Lag- t χ for stationary Gaussian autoregressive process
- Properties of extremal dependence coefficients
- Invariance of extremal index
- Some limitations
- Examples of functionals of interest

Conditional extreme value theory

- Conditional extremes of stationary Markov processes

- Statistical inference with time series conditional extremes

References

- Suppose $\{X_t\}$ is a stationary process with unit Fréchet margins satisfying the AIM(n) condition.

- Let $G(x) = 1 - \exp(-x)$, $x > 0$. Then the extremal index of the time series $\{X_t\}$ is

$$\theta = \lim_{n \rightarrow \infty} \mathbb{P} [X_i \leq u_n : i = 2, \dots, p_n \mid X_1 > u_n]$$

$$= \lim_{n \rightarrow \infty} \mathbb{P} [G^{\leftarrow} \circ F(X_i) \leq G^{\leftarrow} \circ F(u_n) : i = 2, \dots, p_n \mid G^{\leftarrow} \circ F(X_1) > G^{\leftarrow} \circ F(u_n)]$$

$$= \lim_{n \rightarrow \infty} \mathbb{P} [Y_i \leq u_n^G : i = 2, \dots, p_n \mid Y_1 > u_n^G] = \theta^G \quad (u_n^G \sim \log n).$$

i.e., equal to the extremal index of the process $\{Y_t\}$, where Y_t follows a unit-exponential for all t .

- More generally, a monotone increasing marginal transformation does not change the value of the extremal index;

- Introduction
- Motivation
- Asymptotic dependence
- Another view of the extremal dependence in stationary time series
- χ_1 for Moving Maxima process
- Lag- t χ for stationary Gaussian autoregressive process
- Properties of extremal dependence coefficients
- Invariance of extremal index
- Some limitations
- Examples of functionals of interest
- Conditional extreme value theory
- Conditional extremes of stationary Markov processes
- Statistical inference with time series conditional extremes
- References

- Limited information when $\theta = 1$ or when $\chi_t = 0$ for all $t > 0$.
- Mean cluster-size is informative, but in applications we may be interested in other characteristics of the cluster distribution, *e.g.*,

$$T(u, d) = \mathbb{E}(g(\mathbf{Y}_{1:d}) \mid Y_1 > u),$$

or more generally, in the cluster-size distribution itself.

Examples of functionals of interest

Introduction

Motivation

Asymptotic dependence

Another view of the extremal dependence
in stationary time series

χ_1 for Moving Maxima process

Lag- t χ for stationary Gaussian
autoregressive process

Properties of extremal dependence
coefficients

Invariance of extremal index

Some limitations

Examples of functionals of interest

Conditional extreme value theory

Conditional extremes of stationary Markov
processes

Statistical inference with time series
conditional extremes

References

- what is the expected size of the largest element in a short period of time after seeing an exceedance:

$$T_1(u, d) = \mathbb{E}(\max \mathbf{Y}_{1:d} \mid Y_1 > u),$$

- what is the expected energy content in a short period of time after seeing an exceedance:

$$T_2(u, d) = \mathbb{E} \left[\left(\frac{1}{d} \sum_{i=1}^d Y_i^2 \right)^{1/2} \mid Y_1 > u \right],$$

- what is the subasymptotic cluster-size distribution:

$$T_3(u, d) = \mathbb{P} \left(\sum_{i=1}^d \mathbf{1}[Y_i > u] = r \mid Y_1 > u \right),$$

- and finally, how about standard measures of extremal dependence, e.g.,

$$\theta(u, d) = \mathbb{P}(Y_2 \leq u, \dots, Y_d \leq u \mid Y_1 > u) \quad \text{and} \quad \chi(u, d) = \mathbb{P}(Y_{t+1} > u \mid Y_1 > u)?$$

Introduction
Conditional extreme value theory
Univariate exceedances revisited
EVT and extrapolation
Extremes in 2-dimensions
Reduction to common margins
Conditioned limit laws
An application of L'Hôpital's theorem
Example: max-stable distribution with Laplace margins
Example: bivariate normal distribution with Laplace margins
Example: inverted max-stable distribution with Laplace margins
Statistical model
Illustration
Sampling importance-resampling
Theoretical examples
Conditional extremes of stationary Markov processes
Statistical inference with time series conditional extremes
References

Conditional extreme value theory

- Under mild conditions on the survival function of a random variable X , there exists $\sigma(u) > 0$

$$\mathbb{P} \left\{ \frac{X - u}{\sigma(u)} > x \mid X > u \right\} \longrightarrow \{1 + \xi x\}^{-1/\xi}, \quad u \rightarrow x^*,$$

i.e., the scaled excess random variable converges in distribution to the generalised Pareto distribution [Pickands, 1975, Davison and Smith, 1990]

- Interest often is in estimating either v_p given p , or p given v_p , where p and v_p satisfy

$$\mathbb{P}(X > v_p) = p,$$

- Typically, in practice v_p would correspond to a quantile of the distribution above which data have not been observed.

- **BEWARE:** Extrapolation is dangerous, but EVT provides a principled approach based on asymptotic theory for the behaviour of extrema.
- Fundamental premise in all statistical extreme value modelling is that we can approximate the distribution of extreme values by the limiting theoretical forms. For $v_p > u$

$$\begin{aligned}\mathbb{P}(X > v_p) &= \mathbb{P}(X > v_p \mid X > u) \mathbb{P}(X > u) \\ &= \mathbb{P}\left(\frac{X - u}{\sigma(u)} > \frac{v_p - u}{\sigma(u)} \mid X > u\right) \mathbb{P}(X > u)\end{aligned}$$

Inverting leads to a closed-form expression for the $(1/p)$ -return level

$$v_p = u + \frac{\sigma(u)}{\xi} \left[\left(\frac{\pi_u}{p} \right)^\xi - 1 \right],$$

where $\pi_u = \mathbb{P}(X > u)$.

Extremes in 2-dimensions

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling



Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- In 2 dimensions, the notion of ordering is not the same as that in \mathbb{R} due to the lack of total order.
- This brings complications in terms of defining what extreme events are.
- For example, an observation in \mathbb{R}^2 may be regarded as extreme if either:
 - both components exceed high thresholds;
 - at least one component exceeds a high threshold.



Reduction to common margins

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling


Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Standard practical approach is to bring marginal distributions to a common scale



- We shall adopt this approach assuming that such standardizations are possible in practice

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

Random vector (X_1, X_2) in \mathbb{R}^2 with standard Laplace margins.

■ $a_{2|1} : \mathbb{R} \rightarrow \mathbb{R}$

■ $b_{2|1} : \mathbb{R} \rightarrow \mathbb{R}_+$

■ a non-degenerate distribution $G_{2|1}$ on \mathbb{R}

such that, as u approaches ∞ ,

$$\mathbb{P} \left(X_1 - u > x, \frac{X_2 - a_{2|1}(X_1)}{b_{2|1}(X_1)} < z \mid X_1 > u \right) \rightarrow \exp(-x) G_{2|1}(z)$$

at continuity points $(x, z) \in \mathbb{R}_+ \times \mathbb{R}$ of the limit distribution.

An application of L'Hôpital's theorem

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

When the joint density exists, limit formulation

$$\lim_{u \rightarrow \infty} \mathbb{P} \left(X_1 - u > x, \frac{X_2 - a_{2|1}(X_1)}{b_{2|1}(X_1)} < z \mid X_1 > u \right) = \exp(-x) G_{2|1}(z)$$

is equivalent to

$$\begin{aligned} \lim_{u \rightarrow \infty} \frac{\frac{\partial}{\partial u} \mathbb{P} \left(X_1 > u, \frac{X_2 - a_{2|1}(X_1)}{b_{2|1}(X_1)} < z \right)}{\frac{\partial}{\partial u} \Pr(X_1 > u)} &= \lim_{u \rightarrow \infty} \boxed{\mathbb{P} \left(\frac{X_2 - a_{2|1}(X_1)}{b_{2|1}(X_1)} < z \mid X_1 = u \right)} \\ &= G_{2|1}(z) \end{aligned}$$

In what follows, we shall use either $\{X_1 > u\}$ or $\{X_1 = u\}$, assuming

Example: max-stable distribution with Laplace margins

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Suppose (X, Y) follows a bivariate logistic max-stable distribution with standard Laplace margins and dependence parameter $\kappa \in (0, 1]$, that is,

$$\mathbb{P}(X \leq x, Y \leq y) = \exp\{-V(T(x), T(y))\},$$

where $T(x) = -1/\log(F_L(x))$,

$$V(x, y) = \left(x^{-1/\kappa} + y^{-1/\kappa}\right)^\kappa \quad \text{and} \quad V_1(x, y) = \frac{\partial}{\partial x} V(x, y).$$

- The conditional distribution for this example is

$$\mathbb{P}(Y \leq y \mid X = x) = -T(x)^2 e^{1/T(x)} V_1\{T(x), T(y)\} \exp[-V\{T(x), T(y)\}].$$

- Conditioned limit law: for $z \in \mathbb{R}$,

$$\mathbb{P}(Y - X \leq z \mid X = u) \rightarrow [1 + \exp(-z/\kappa)]^{\kappa-1}, \quad u \rightarrow \infty.$$

Example: bivariate normal distribution with Laplace margins

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Suppose (X, Y) follows a bivariate normal distribution with standard Laplace margins and correlation parameter $\rho \in (-1, 1)$, that is,

$$\mathbb{P}(X \leq x, Y \leq y) = \int_{-\infty}^{\Phi^{-1}(F_L(x))} \int_{-\infty}^{\Phi^{-1}(F_L(y))} \frac{du dv}{2\pi(1 - \rho^2)^{1/2}} \exp \left(-\frac{1}{2(1 - \rho^2)} u^2 + v^2 - 2\rho uv \right)$$

where Φ denotes the cdf of the standard normal distribution and Φ^{-1} is its inverse.

- The conditional distribution for this example is

$$\mathbb{P}(Y \leq y \mid X = x) = \Phi \left(\frac{\Phi^{-1}(F_L(y)) - \rho \Phi^{-1}(F_L(x))}{(1 - \rho^2)^{1/2}} \right).$$

- Conditioned limit law: for $z \in \mathbb{R}$,

$$\mathbb{P} \left(\frac{Y - \text{sign}(\rho)\rho^2 X}{X^{1/2}} \leq z \mid X = u \right) \rightarrow \Phi \left(\frac{z}{\sqrt{2\rho^2(1 - \rho^2)}} \right), \quad u \rightarrow \infty.$$

Example: inverted max-stable distribution with Laplace margins

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

- Suppose (X, Y) follows a bivariate inverted logistic max-stable distribution with unit exponential margins and dependence parameter $\kappa \in (0, 1]$, that is,

$$\mathbb{P}(X > x, Y > y) = \exp\{-V(1/x, 1/y)\},$$

where

$$V(x, y) = \left(x^{-1/\kappa} + y^{-1/\kappa}\right)^\kappa$$

- The conditional distribution for this example is

$$\mathbb{P}(Y \leq y \mid X = x) = 1 + V_1(1, x/y) \exp(x - xV(1, x/y))$$

- Conditioned limit law: for $z \in \mathbb{R}_+$,

$$\mathbb{P}\left(\frac{Y}{X^{1-\kappa}} \leq z \mid X = u\right) \rightarrow 1 - \exp(-\kappa z^{1/\kappa}) \quad u \rightarrow \infty.$$

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration

Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

The distribution of extreme values is approximated by their limiting theoretical form, for $X_1 > u$,

$$X_2 = a_{2|1}(X_1) + b_{2|1}(X_1)Z_{2|1}$$

where $Z_{2|1} \sim G_{2|1}$.

- Under mild assumptions $a_{2|1} \in \text{RV}_1$ and $b_{2|1} \in \text{RV}_{\beta_{2|1}}$, where $\beta_{2|1} \in (0, 1)$.
- Parametric models for location-scale, simplify a and b by
 - $a_{2|1}(x) = \alpha_{2|1}x$, where $\alpha_{2|1} \in [-1, 1]$;
 - $b_{2|1}(x) = x^{\beta_{2|1}}$.
- $G_{2|1}$ does not admit a simple parametric form. False working assumption: $Z_{2|1} \sim \mathcal{N}(\mu_{2|1}, \sigma_{2|1}^2)$.

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is regularly varying at ∞ with exponent ρ if $\lim_{u \rightarrow \infty} f(ux)/f(u) = x^\rho$ for all $x \geq 0$. Notation: $f \in \text{RV}_\rho$.

[Introduction](#)

[Conditional extreme value theory](#)

[Univariate exceedances revisited](#)

[EVT and extrapolation](#)

[Extremes in 2-dimensions](#)

[Reduction to common margins](#)

[Conditioned limit laws](#)

[An application of L'Hôpital's theorem](#)

[Example: max-stable distribution with Laplace margins](#)

[Example: bivariate normal distribution with Laplace margins](#)

[Example: inverted max-stable distribution with Laplace margins](#)

[Statistical model](#)

[Illustration](#)



[Sampling importance-resampling](#)

[Theoretical examples](#)

[Conditional extremes of stationary Markov processes](#)

[Statistical inference with time series conditional extremes](#)

[References](#)



Sampling importance-resampling

Introduction

Conditional extreme value theory

Univariate exceedances revisited

EVT and extrapolation

Extremes in 2-dimensions

Reduction to common margins

Conditioned limit laws

An application of L'Hôpital's theorem

Example: max-stable distribution with Laplace margins

Example: bivariate normal distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Statistical model

Illustration




Sampling importance-resampling

Theoretical examples

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References



- Introduction
- Conditional extreme value theory
 - Univariate exceedances revisited
 - EVT and extrapolation
 - Extremes in 2-dimensions
 - Reduction to common margins
 - Conditioned limit laws
 - An application of L'Hôpital's theorem
 - Example: max-stable distribution with Laplace margins
 - Example: bivariate normal distribution with Laplace margins
 - Example: inverted max-stable distribution with Laplace margins
 - Statistical model
 - Illustration
 - Sampling importance-resampling
 - Theoretical examples
- Conditional extremes of stationary Markov processes
- Statistical inference with time series conditional extremes
- References

Asymptotic dependence: $\alpha_{2|1} = 1$ or $\alpha_{2|1} = -1$, and $\beta_{2|1} = 0$;

Asymptotic independence: $\alpha_{2|1} \in (-1, 1)$ and $\beta_{2|1} \in (0, 1)$

Introduction
Conditional extreme value theory
Conditional extremes of stationary Markov processes
Notation
Conditional extremes of a Markov process
Recurrence relations
Tail-chains and hidden tail-chains
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Convergence of the transition probability distribution
Theorem
Example: max-stable distribution with Laplace margins
Example: inverted max-stable distribution with Laplace margins
Variety of behaviour of tail- and hidden-tail chains
Properties of normalizing functionals and structure of tail processes
Properties of normalizing functions and structure of tail processes
Comments on Theorem
Back-and-forth hidden tail-chain
Statistical inference with time series conditional extremes
References

Conditional extremes of stationary Markov processes

- Introduction
- Conditional extreme value theory
- Conditional extremes of stationary Markov processes
- Notation
- Conditional extremes of a Markov process
- Recurrence relations
- Tail-chains and hidden tail-chains
- Higher-order Markov processes
- Higher-order Markov processes
- Higher-order Markov processes
- Higher-order Markov processes
- Convergence of the transition probability distribution
- Theorem
- Example: max-stable distribution with Laplace margins
- Example: inverted max-stable distribution with Laplace margins
- Variety of behaviour of tail- and hidden-tail chains
- Properties of normalizing functionals and structure of tail processes
- Properties of normalizing functions and structure of tail processes
- Comments on Theorem
- Back-and-forth hidden tail-chain
- Statistical inference with time series conditional extremes
- References

■ Y_t : measurement process, assumed to be a stationary-time series with a continuous marginal distribution F_Y .

■ X_t : standardized time series $X_t = F_L^{-1}\{F_Y(Y_t)\}$ where

$$F_L(x) = \begin{cases} 1 - \exp(y)/2 & y \geq 0 \\ \exp(y)/2 & y < 0 \end{cases}$$

is the standard Laplace distribution function.

■ Standardization gives that $X_t \sim \text{Laplace}(0, 1)$ for all t .

Conditional extremes of a Markov process

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Suppose that F is a bivariate distribution with common standard Laplace margins admitting regular conditional probabilities.
- There exists a stationary Markov process $\{X_t : t \in \mathbb{N}\}$ such that $(X_t, X_{t+1}) \sim F$ for all t .
- Suppose further a conditioned limit law exists. For simplicity, suppose there exist $\alpha \in (0, 1]$, $\beta \in (0, 1)$ and a distribution G supported on \mathbb{R} such that

$$\left(X_0 - u, \frac{X_1 - \alpha X_0}{X_0^\beta} \right) \mid \{X_0 = u\} \xrightarrow{d} (E, Z_1)$$

where $E \sim \exp(1)$, $Z \sim G$, and E is independent of Z .

Recall: a process X_t is Markov if the distribution of $X_t \mid X_{t-1}, X_{t-2}, \dots$ is equal to the distribution of $X_t \mid X_{t-1}$ for all t .

When saying that a distribution is supported on a subset A of \mathbb{R}^k , we do not allow the distribution to place mass at the boundary ∂A of A .

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Conditionally on $X_0 = u$ and for large u

$$X_1 \approx \alpha X_0 + X_0^\beta Z_1, \quad Z_1 \sim G.$$

- For sufficiently large u , X_2 will be large too. From stationarity

$$X_2 \approx \alpha X_1 + X_1^\beta \varepsilon_2, \quad \varepsilon_2 \sim G$$

$$= \alpha(\alpha X_0 + X_0^\beta Z_1) + (\alpha X_0 + X_0^\beta Z_1)^\beta \varepsilon_2$$

$$\approx \alpha^2 X_0 + \alpha X_0^\beta Z_1 + (\alpha X_0)^\beta \varepsilon_2 + \alpha^\beta \beta X_0^{2\beta-1} Z_1 \varepsilon_2$$

- After rearrangement, we see that

$$\frac{X_2 - \alpha^2 X_0}{X_0^\beta} \Big| \{X_0 = u\} \xrightarrow{d} (Z_1, Z_2)$$

where $Z_2 = \alpha Z_1 + \alpha^\beta \varepsilon_2$, where ε_2 is independent of Z_1 .

- The heuristic derivations in the previous slide can be formalized under mild conditions that are realistic for practical applications.
- When appropriately renormalized, the Markov chain converges to a non-degenerate process in the sense that for all $t \geq 1$,

$$\left(X_0 - u, \frac{X_1 - \alpha_1 X_0}{X_0^{\beta_1}}, \dots, \frac{X_t - \alpha_t X_0}{X_0^{\beta_t}} \right) \mid \{X_0 > u\} \xrightarrow{d} (E, Z_1, \dots, Z_t)$$

where $\alpha_t = \alpha^t$, $\beta_t = \beta$, $E \sim \exp(1)$ is independent of (Z_1, \dots, Z_t) and

$$Z_t = \alpha Z_{t-1} + \alpha^\beta \varepsilon_t$$

where $Z_0 = 0$ a.s. and $\{\varepsilon_t\}_{t=1}^\infty$ is a sequence of IID random variables from G .

- Classification of tail processes:
 - If $|\alpha_t| = 1$ and $\beta_t = 0$ for all t , then the tail process $\{Z_t\}$ is termed the *tail-chain* of $\{X_t\}$.
 - If $|\alpha_t| \neq 1$ for all t , then the process is termed the *hidden tail-chain* of $\{X_t\}$.

Higher-order Markov processes

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- For higher-order Markov processes where $k > 1$ we need additional initial conditions.
- Think about analogy with ordinary differential equations: the general solution to a second-order ODE requires two initial conditions, one for the value of the function at some time, say $t = 0$, and the other for the derivative of the function at $t = 0$.
- Initial conditions for stochastic recurrence relation: we can find $\mathbf{a}_{1:k-1} : \mathbb{R}^{k-1} \rightarrow \mathbb{R}$ and $\mathbf{b}_{1:k-1} : \mathbb{R}^{k-1} \rightarrow \mathbb{R}_+$ such that

$$\frac{\mathbf{X}_{1:k-1} - \mathbf{a}_{1:k-1}(X_0)}{\mathbf{b}_{1:k-1}(X_0)} \Big| \{X_0 > u\} \xrightarrow{d} \mathbf{Z}_{1:k-1} \quad \text{as } u \rightarrow \infty,$$

where each random variable Z_i has a non-degenerate distribution supported on \mathbb{R} .

Vector algebra is interpreted element-wise

- Consider how a complete characterization may be given for higher-order Markov processes with $k > 1$ using induction on \mathbb{N} . Fix a $t \geq k > 1$ and assume there exist sequences of norming functions a_i and b_i , $i = 1, \dots, t-1$, such that,

$$\frac{\mathbf{X}_{1:t-1} - \mathbf{a}_{1:t-1}(X_0)}{b_{1:t-1}(X_0)} \Big| \{X_0 > u\} \xrightarrow{d} \mathbf{Z}_{1:t-1} \quad \text{as } u \rightarrow \infty,$$

where each Z_i is a random variable with a non-degenerate distribution on \mathbb{R} .

- Suffices to consider marginal convergence, that is,

$$\mathbb{P} \left(\frac{X_t - a_t(X_0)}{b_t(X_0)} \leq x_t \mid X_0 > u \right) = \frac{1}{\overline{F}_0(u)} \int_u^\infty \int_{\mathbb{R}^{t-1}} \mathbb{P}(\mathbf{X}_{0:t-1} \in d\mathbf{x}_{0:t-1}) \left[\int_{-\infty}^{x_t} \mathbb{P} \left(\frac{X_t - a_t(X_0)}{b_t(X_0)} \in dz_t \mid \mathbf{X}_{0:t-1} = \mathbf{x}_{0:t-1} \right) \right],$$

- Replace $a_t(X_0)$ by $a_t(x_0)$ by conditioning on the exact value of X_0 being equal to x_0 .

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Use the Markov property so that the conditioning on all previous states is reduced to conditioning on the previous k states.

- Change variables to $z_0 = x_0 - u$ and $z_i = \{x_i - a_i(x_0)\}/b_i(x_0)$ for $i = 1, \dots, t-1$, for any $t \geq k$, previous expression is equal to

$$\int_0^\infty \frac{F_0\{v_u(dz_0)\}}{\overline{F}_0(u)} \times \left[\int_{\mathbb{R}^{t-1}} \mathbb{P} \left(\frac{\mathbf{X}_{1:t-1} - \mathbf{a}_{1:t-1}\{v_u(z_0)\}}{\mathbf{b}_{1:t-1}\{v_u(z_0)\}} \in d\mathbf{z}_{1:t-1} \mid X_0 = v_u(z_0) \right) \times \right. \\ \left. \times \left[\int_{-\infty}^{x_t} \mathbb{P} \left(\frac{X_t - a_t\{v_u(z_0)\}}{b_t\{v_u(z_0)\}} \in dz_t \mid \frac{\mathbf{X}_{t-k:t-1} - \mathbf{a}_{t-k:t-1}\{v_u(z_0)\}}{\mathbf{b}_{t-k:t-1}\{v_u(z_0)\}} = \mathbf{z}_{t-k:t-1} \right) \right] \right],$$

where $a_0(x) = x$, $b_0(x) = 1$ for all $x \in \mathbb{R}$ and $v_u(z_0) = u + z_0$.

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- To ensure marginal convergence under the induction hypothesis, we require convergence of

$$\mathbb{P} \left(\frac{X_t - a_t(u)}{b_t(u)} \leq z_t \mid \frac{\mathbf{X}_{t-k:t-1} - \mathbf{a}_{t-k:t-1}(u)}{\mathbf{b}_{t-k:t-1}(u)} = \mathbf{z}_{t-k:t-1} \right)$$

- Let $\mathbf{A}_t(u, \mathbf{z}) = \mathbf{a}_{t-k:t-1}(u) + \mathbf{b}_{t-k:t-1}(u) \mathbf{z}$ for $\mathbf{z} \in \mathbb{R}^k$, and rewrite probability above as

$$\mathbb{P} \left[\frac{X_t - a(\mathbf{X}_{t-k:t-1})}{b(\mathbf{X}_{t-k:t-1})} \leq \frac{dz_t}{\psi_{t,u}^b(\mathbf{X}_{t-k:t-1})} - \psi_{t,u}^a(\mathbf{X}_{t-k:t-1}) \mid \mathbf{X}_{t-k:t-1} = \mathbf{A}_t(u, \mathbf{z}_{t-k:t-1}) \right].$$

- $a : \mathbb{R}^k \rightarrow \mathbb{R}$ and $b : \mathbb{R}^k \rightarrow \mathbb{R}_+$ are location and scale functionals and $\psi_{t,u}^a$ and $\psi_{t,u}^b$ satisfy

$$\psi_{t,u}^a(\mathbf{z}) = \frac{a\{\mathbf{A}_t(u, \mathbf{z})\} - a_t(u)}{b_t(u)} \quad \text{and} \quad \psi_{t,u}^b(\mathbf{z}) = \frac{b\{\mathbf{A}_t(u, \mathbf{z})\}}{b_t(u)}$$

Convergence of the transition probability distribution

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

In what follows, we assume that there exist:

- for $t = k, k + 1, \dots$,
 - location and scale functions $a_t : \mathbb{R} \rightarrow \mathbb{R}$ and $b_t : \mathbb{R} \rightarrow \mathbb{R}_+$,
 - continuous update functions $\psi_t^a : \mathbb{R}^k \rightarrow \mathbb{R}$ and $\psi_t^b : \mathbb{R}^k \rightarrow \mathbb{R}_+$,
 - and location and scale functionals $a : \mathbb{R}^k \rightarrow \mathbb{R}$ and $b : \mathbb{R}^k \rightarrow \mathbb{R}_+$,
- such that, for all $\mathbf{z} \in \mathbb{R}^k$,

$$\lim_{u \rightarrow \infty} \psi_{t,u}^a(\mathbf{z}_u) \rightarrow \psi_t^a(\mathbf{z}) \quad \text{and} \quad \lim_{u \rightarrow \infty} \psi_{t,u}^b(\mathbf{z}_u) \rightarrow \psi_t^b(\mathbf{z})$$

whenever $\mathbf{z}_u \rightarrow \mathbf{z} \in \mathbb{R}^k$.

- a non-degenerate distribution K supported on \mathbb{R} , such that for all $\mathbf{z} \in \mathbb{R}^k$,

$$\mathbb{P} \left[\frac{X_k - a(\mathbf{X}_{0:k-1})}{b(\mathbf{X}_{0:k-1})} \leq \varepsilon \mid \mathbf{X}_{0:k-1} = \mathbf{A}_t(u, \mathbf{z}_{0:k-1}) \right] \rightarrow K(\varepsilon)$$

Introduction
Conditional extreme value theory
Conditional extremes of stationary Markov processes
Notation
Conditional extremes of a Markov process
Recurrence relations
Tail-chains and hidden tail-chains
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Convergence of the transition probability distribution
Theorem
Example: max-stable distribution with Laplace margins
Example: inverted max-stable distribution with Laplace margins
Variety of behaviour of tail- and hidden-tail chains
Properties of normalizing functionals and structure of tail processes
Properties of normalizing functions and structure of tail processes
Comments on Theorem
Back-and-forth hidden tail-chain
Statistical inference with time series conditional extremes
References

If $\{X_t : t = 0, 1, \dots\}$ is a stationary Markov chain satisfying the aforementioned assumptions, then

$$\left(\frac{X_0 - u}{\sigma(u)}, \frac{X_1 - a_1(X_0)}{b_1(X_0)}, \dots, \frac{X_t - a_t(X_0)}{b_t(X_0)}\right) \mid \{X_0 > u\} \xrightarrow{d} (E_0, Z_1, \dots, Z_t), \quad t \geq k,$$

where

(i) $E_0 \sim H_0$ and $(Z_1, Z_2 \dots, Z_t)$ are independent,

(ii) $Z_0 = 0$ a.s., $(Z_1, \dots, Z_{k-1}) \sim G$ and

$$Z_s = \psi_s^a(\mathbf{Z}_{s-k:s-1}) + \psi_s^b(\mathbf{Z}_{s-k:s-1}) \varepsilon_s, \quad s = k, k + 1, \dots$$

for a sequence of IID random variables $\varepsilon_s \sim K$.

Example: max-stable distribution with Laplace margins

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Suppose $\mathbf{X}_{0:k}$ follows a $(k+1)$ -variate logistic max-stable distribution with standard Laplace margins and dependence parameter $\kappa \in (0, 1]$, that is,

$$\mathbb{P}(\mathbf{X}_{0:k} \leq \mathbf{x}_{0:k-1}) = \exp\{-V(T(x_0), \dots, T(x_k))\},$$

where $T(x) = -1/\log(F_L(x))$, and

$$V(x_0, \dots, x_k) = \left(x_0^{-1/\kappa} + \dots + x_k^{-1/\kappa}\right)^\kappa$$

- Define

$$a(x_0, \dots, x_{k-1}) = -\kappa \log(e^{-x_0/\kappa} + \dots + e^{-x_{k-1}/\kappa}) \quad \text{and} \quad b(x_0, \dots, x_{k-1}) = 1.$$

- Convergence of transition probability distribution: for $\varepsilon \in \mathbb{R}$,

$$\mathbb{P}\left(\frac{X_k - a(\mathbf{X}_{0:k-1})}{b(\mathbf{X}_{0:k-1})} \leq \varepsilon \mid \mathbf{X}_{0:k-1} = u\mathbf{1} + \mathbf{z}_{0:k-1}\right) \rightarrow [1 + \exp(-\varepsilon/\kappa)]^{\kappa-k}, \quad u \rightarrow \infty.$$

for all $\mathbf{z}_{0:k-1} \in \mathbb{R}^k$

Example: inverted max-stable distribution with Laplace margins

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Suppose $\mathbf{X}_{0:k}$ follows a $(k+1)$ -variate inverted max-stable logistic distribution with dependence parameter $\kappa \in (0, 1]$ and standard Laplace margins, that is,

$$\mathbb{P}(\mathbf{X}_{0:k} > \mathbf{x}_{0:k-1}) = \exp\{-V(1/x_0, \dots, 1/x_k)\},$$

where

$$V(x_0, \dots, x_k) = \left(x_0^{-1/\kappa} + \dots + x_k^{-1/\kappa}\right)^\kappa$$

- Define

$$a(x_0, \dots, x_{k-1}) = 0 \quad \text{and} \quad b(x_0, \dots, x_{k-1}) = (x_0^{1/\kappa} + \dots + x_{k-1}^{1/\kappa})^\kappa (1-\kappa).$$

- Convergence of transition probability distribution: for $\varepsilon \in \mathbb{R}_+$,

$$\mathbb{P} \left\{ \frac{X_k - a(\mathbf{X}_{0:k-1})}{b(\mathbf{X}_{0:k-1})} \leq z \mid \mathbf{X}_{0:k-1} = (u z_0, u^{1-k} z_{0:k-1}) \right\} \rightarrow 1 - \exp(-\kappa \varepsilon^{1/\kappa}), \quad u \rightarrow \infty.$$

for all $\mathbf{z}_{0:k-1} \in \mathbb{R}_+^k$

Variety of behaviour of tail- and hidden-tail chains




Figure 1: Pointwise 2.5% and 97.5% quantiles of the sampling distribution (shaded region), mean of the sampling distribution (dashed line) and one realization from the (hidden) tail-chain (solid line). The parameters for all copulas are chosen such that the coefficient of residual tail dependence η [Ledford and Tawn, 1997] and the extremal coefficient [Beirlant et al., 2004] are equal. The distribution of $\mathbf{X}_0:k$ that was used in each example is: (a) standard multivariate Gaussian, (b) inverted logistic, (c) max-stable logistic and (d) max-stable Hüsler–Reiss

Properties of normalizing functionals and structure of tail processes

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

- Asymptotically dependent process (Multivariate Regular Variation):
 - $a(\boldsymbol{x}) = \log g(e^{\boldsymbol{x}})$ where g is 1-homogeneous.
 - $b(\boldsymbol{x}) = 0$
 - $Z_t = a(\boldsymbol{Z}_{t-k:t-1}) + \varepsilon_t$
- Asymptotically independent process (Hidden Regular Variation) with a not zero:
 - a is 1-homogeneous with $a(\mathbf{1}) < 1$
 - b is β -homogeneous with $\beta \in (0, 1)$
 - $Z_t = \nabla a(\boldsymbol{\alpha}_{t-k:t-1})^\top \boldsymbol{Z}_{t-k:t-1} + b(\boldsymbol{\alpha}_{t-k:t-1}) \varepsilon_t$
- Asymptotically independent process (Hidden Regular Variation) with a zero:
 - b is β -homogeneous with $\beta \in (0, 1)$.
 - Structure of tail process is complicated.

Properties of normalizing functions and structure of tail processes

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Notation

Conditional extremes of a Markov process

Recurrence relations

Tail-chains and hidden tail-chains

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Higher-order Markov processes

Convergence of the transition probability distribution

Theorem

Example: max-stable distribution with Laplace margins

Example: inverted max-stable distribution with Laplace margins

Variety of behaviour of tail- and hidden-tail chains

Properties of normalizing functionals and structure of tail processes

Properties of normalizing functions and structure of tail processes

Comments on Theorem

Back-and-forth hidden tail-chain

Statistical inference with time series conditional extremes

References

■ Asymptotically dependent process (Multivariate Regular Variation):

- $a_t(x) = x$
- $b_t(x) = 0$
- $\mathbb{E}(Z_t) < 0$ for all t

■ Asymptotically independent process (Hidden Regular Variation) with a not zero:

- $a_t(x) = \alpha_t x$ where $\alpha_t = a(\boldsymbol{\alpha}_{t-k:t-1})$ and $\alpha_t \rightarrow 0$ as $t \rightarrow \infty$
- $b_t(x) = x^\beta$ where $\beta \in (0, 1)$
- $Z_t \xrightarrow{p} 0$ as $t \rightarrow \infty$

■ Asymptotically independent process (Hidden Regular Variation) with a zero:

- $\log \beta_t = \log \beta + \log (\max_{i=1,\dots,k} \beta_{t-i})$
- Structure of tail process is complicated.

- Introduction
- Conditional extreme value theory
- Conditional extremes of stationary Markov processes
- Notation
- Conditional extremes of a Markov process
- Recurrence relations
- Tail-chains and hidden tail-chains
- Higher-order Markov processes
- Higher-order Markov processes
- Higher-order Markov processes
- Higher-order Markov processes
- Convergence of the transition probability distribution
- Theorem
- Example: max-stable distribution with Laplace margins
- Example: inverted max-stable distribution with Laplace margins
- Variety of behaviour of tail- and hidden-tail chains
- Properties of normalizing functionals and structure of tail processes
- Properties of normalizing functions and structure of tail processes
- Comments on Theorem
- Back-and-forth hidden tail-chain
- Statistical inference with time series conditional extremes
- References

- The entities we have referred to as tail-chains and hidden tail-chains are in fact forward tail-chains and hidden tail-chains.
- These describe the behaviour of the Markov chain only forward in time from a large observation.
- There is also the parallel interest in a backward tail/hidden tail chain, to give how the chain evolves into an extreme event, and the joint behaviour of the two, known as back-and-forth tail processes.

Introduction
Conditional extreme value theory
Conditional extremes of stationary Markov processes
Notation
Conditional extremes of a Markov process
Recurrence relations
Tail-chains and hidden tail-chains
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Higher-order Markov processes
Convergence of the transition probability distribution
Theorem
Example: max-stable distribution with Laplace margins
Example: inverted max-stable distribution with Laplace margins
Variety of behaviour of tail- and hidden-tail chains
Properties of normalizing functionals and structure of tail processes
Properties of normalizing functions and structure of tail processes
Comments on Theorem
Back-and-forth hidden tail-chain
Statistical inference with time series conditional extremes
References

Statistical inference with time series conditional extremes

- Assume that $X_t \sim F_L$ where F_L denotes the standard Laplace distribution.
- Let $T_u = \{t \in \mathbb{Z} : X_t > u\}$ denote the set of times where the process exceeds the level u .
- We assume that there exist location and scale functions $a_i : \mathbb{R} \rightarrow \mathbb{R}$ and $b_i : \mathbb{R} \rightarrow \mathbb{R}_+$, such that for any $A \subset \mathbb{Z}$ with $|A| < \infty$ and for all $t \in T_u$,

$$\left(X_t - u, \frac{\mathbf{X}_A - \mathbf{a}_{A-t}(X_t)}{\mathbf{b}_{A-t}(X_t)} \right) \mid X_t > u \xrightarrow{d} (E_t, \mathbf{Z}_A^t), \quad (2)$$

where

- $E_t \sim \exp(1)$ is independent of $\mathbf{Z}_A^t \sim G_{A \setminus t}$;
- where $G_{A \setminus t}$ has non-degenerate margins;
- $a_0(x) = x$ and $b_0(x) = 1$. This gives $Z_t^t = 0$.

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

- The limit relation (2) is taken to hold exactly above a sufficiently large level u . That is, we assume that for $t \in T_u$

$$X_t = u + E_t$$

$$\mathbf{X}_{t-L:t+L} = \mathbf{a}_{-L:L}(X_t) + \mathbf{b}_{-L:L}(X_t) \mathbf{Z}_{t-L:t+L}^t,$$

$$\mathbf{Z}_{t-L:t+L}^t \sim G_{-L:L}$$

- $L \in \mathbb{N}$ chosen sufficiently large so that extreme episodes are contained within $t - L : t + L$.
- Context often dictates the choice of L .

Considerations for statistical modelling

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

- In addition to $G_{-L:L}$, there are $2 \times (2L - 1)$ functions to infer $(\mathbf{a}_{-L:L}$ and $\mathbf{b}_{-L:L}$).
- Structure of norming functions:
 - Asymptotic dependence gives that $a_{|i|}(x) = x$ and $b_{|i|}(x) = 0$.
 - Asymptotic independence with $a_{|i|}$ not zero gives $a_{|i|}(x) \rightarrow 0$ as $i \rightarrow \infty$. So for large $|i|$ we have that $Z_{t+i}^t \sim F_L$.
 -
- Multivariate distribution $G_{-L:L}$ admits no finite-dimensional parametric form. Require a flexible statistical model for $G_{-L:L}$ that facilitates inference and simulation (later).

Parametric forms for norming functions based on Markov chain theory

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

Recurrence based approach. For example, for a k th order Markov process, $\alpha_{1:k-1}$ are free parameters and subsequent values $\alpha_{k:L}$ obtained via $\alpha_t = a(\alpha_{t-k:t-1})$, for $t = k, \dots, L$.

■ Recurrence based on correlation model

□ Order-1 Markov chain

$$\alpha_t = \theta \alpha_{t-1}, \quad 1 \leq t \leq L, \quad \text{with } \alpha_0 = 1, \theta \in [-1, 1] \quad (3)$$

□ Order-2 Markov chain

$$\alpha_t = \theta_1 \alpha_{t-1} + \theta_2 \alpha_{t-2}, \quad 2 \leq t \leq k, \quad \text{with } \alpha_0 = 1, \alpha_1 = \theta_1 / (1 - \theta_2). \quad (4)$$

■ For asymptotically independent models, the parameters θ , θ_1 and θ_2 are subject to constraints to ensure $\alpha_t \rightarrow 0$ as $t \rightarrow \infty$. In particular, we have $\theta \in (-1, 1)$ and $r_1, r_2, r_3 \in (-1, 1)^3$ where

$$\theta_1 = r_1 - r_1 r_2 - r_2 r_3$$

$$\theta_2 = r_2 - r_1 r_3 + r_1 r_2 r_3$$

$$\theta_3 = r_3$$

Parametric forms for norming functions based on Markov chain theory

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

- Constraints for higher-order Markov processes may be obtained in a simpler way.

- Regarding the correlation model, we can rewrite this as

$$\alpha_t = \sum_{i=1}^k \theta_i \alpha_{t-i} = c \sum_i \gamma_i \alpha_{t-i}$$

where $c = \sum_i \theta_i$ and $\gamma_i = \theta_i / c$ so that $\sum_i \gamma_i = 1$.

- coefficients γ_i are free parameters on the simplex so can be reparametrized as

$$\gamma_i = \frac{\exp(\Gamma_i)}{\sum_{i=1}^k \exp(\Gamma_i)}, \quad 1 \leq i \leq L$$

where $\Gamma_i \in \mathbb{R}$, subject to a sum-to-zero identifiability constraint $\sum_{i=1}^k \Gamma_i = 0$.

- To ensure $\alpha_t \rightarrow 0$ as $t \rightarrow \infty$, we require $|c| < 1$

Marginal model for the residual

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

■ $\mathbf{Z}_{t-L:t+L}^t = (Z_{t-i:t+i}^t : i = -L, \dots, L)$ where $Z_{t+i}^t \sim G_i$ with density

$$g_i(z) = \frac{\delta_i}{2\sigma_i\Gamma(1/\delta_i)} \exp \left\{ - \left| \frac{z - \mu_i}{\sigma_i} \right|^{\delta_i} \right\}, \quad \mu_i \in \mathbb{R}, \quad \sigma_i > 0, \quad \delta_i > 0. \quad (5)$$

□ $(\mu_i, \sigma_i, \delta_i) = (0, 1, 2)$: standard Gaussian

□ $(\mu_i, \sigma_i, \delta_i) = (0, 1, 1)$: standard Laplace

Dependence model for the residual

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

- In general, the components of the residual vector $\mathbf{Z}_{t-L:t+L}^t$ will be dependent.
- Simplest approach is in style with copula modelling methods. Let Σ be a $(2L + 1) \times (2L + 1)$ covariance matrix and let \mathbf{Q} be Σ^{-1} without the row and column associated with Z_0^t .
- Model for residual vector $\mathbf{Z}_{t-L:t+L}^t \mid Z_0^t = 0$, based on conditional normal distribution. This leads to

$$\left(\Phi^{-1} \left\{ G_{t-i}^Z \left(Z_{t-i}^t \right) \right\} : i = -L, \dots, -1, 1, \dots, L \right) \sim \text{MVN} \left(\boldsymbol{\mu}, \mathbf{Q}^{-1} \right)$$

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

- Interested in estimating conditionals expectations of the form

$$\mathbb{E}(g(\mathbf{X}_{0:d}) \mid X_1 > v) \quad v > u, d \in \mathbb{N},$$

for some function g .

- Monte-Carlo based estimator by simulate forwards in time from extreme event

Forward sampling algorithm

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

Conditional extreme value limits

Statistical model

Considerations for statistical modelling

Parametric forms for norming functions based on Markov chain theory

Parametric forms for norming functions based on Markov chain theory

Marginal model for the residual

Dependence model for the residual

Forward sampling

Forward sampling algorithm

References

input : Threshold $v > u$, $d, N \in \mathbb{N}$ and constants $(\hat{\alpha}_{0:k}, \hat{\beta})$ from fitted conditional model.

output: An estimate of $\mathbb{E}(g(\mathbf{X}_{0:d}) \mid X_1 > v)$

1 for $i \leftarrow 1$ **to** N **do**

2 *simulate exceedance amount* $E \sim \text{Exp}(1)$;

3 *set* $X_1^i = v + E$;

4 *simulate residual* $\hat{\mathbf{Z}}_{2:d}^{(1)}$ *from fitted conditional model independently of* X_1^i ;

5 *set* $\mathbf{X}_{2:d}^i = \hat{\alpha}_{1:d-1} X_1^i + (X_1^i)^{\hat{\beta}} \hat{\mathbf{Z}}_{2:d}^{(1)}$;

6 *set* $\mathbf{X}^i = (X_1, \mathbf{X}_{2:d}^i)$;

7 end

8 return $\hat{\mathbb{E}}(g(\mathbf{X}_{1:d}) \mid X_1 > v) = N^{-1} \sum_{i=1}^N g(\mathbf{X}^i)$

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

References

Introduction

Conditional extreme value theory

Conditional extremes of stationary Markov processes

Statistical inference with time series conditional extremes

References

J. Beirlant, Y. Goegebeur, J. Segers, and J. Teugels. *Statistics of Extremes, Theory and Applications*. Wiley, 2004.

A. C. Davison and R. L. Smith. Models for exceedances over high thresholds. *J. Roy. Statist. Soc., B*, 52:393–442, 1990.

T. Hsing, J. Hüsler, and M. Leadbetter. On the exceedance point process for a stationary sequence. *Probab. Theory Reltd Flds*, 78:97–112, 1988.

A. W. Ledford and J. A. Tawn. Modelling dependence within joint tail regions. *J. Roy. Statist. Soc., B*, 59:475–499, 1997.

J. Pickands. Statistical inference using extreme order statistics. *Ann. Statist.*, 3:119–131, 1975.