

ABUNDANCE, HABITAT ASSOCIATIONS, AND
MIGRATORY CONNECTIVITY OF BIRD SPECIES
OF SPECIAL CONCERN ON JOINT BASE
ELMENDORF-RICHARDSON, ALASKA

2020 SUMMARY REPORT

Submitted to:

Cassandra Schoofs

Biological Scientist (Environmental)

USAF PACAF 673d CES/CEIEC

Joint Base Elmendorf-Richardson, AK

Prepared by:

Laura McDuffie

U.S. Fish and Wildlife Service, Migratory Bird Management

1011 East Tudor Road, MS 201,

Anchorage, Alaska 99503

laura_mcduffie@fws.gov

Contents

0.1	Acronyms and Abbreviations	4
0.2	Executive Summary	5
0.3	Introduction	6
0.4	Study Area	7
0.5	Methods	10
0.5.1	Wetland Area Surveys: Study Design	10
0.5.2	ERF Proposed Expansion Survey: Study Design	12
0.5.3	Automated Recording Units	15
0.5.4	Nest searching	15
0.5.5	Captures and GPS tag deployment	15
0.5.6	Data Management	20
0.6	Results	20
0.6.1	Wetland Area Surveys	20
0.6.2	Eagle River Flats Expansion: Point Counts	28
0.6.3	Nest Discoveries	35
0.6.4	Banding, Sampling, and Tracking	35
0.7	Interpretation	36
0.7.1	Species Richness Curves: Area Survey	36
0.7.2	Species Richness: ERF Point Counts	41
0.8	Products	49
0.8.1	Data and Reporting	49
0.9	Acknowledgements	49
0.10	References	49
0.11	Appendix A: Species List	50
0.12	Appendix B: Status, number of individuals, and relative occurrence of bird species detected during two breeding season survey periods on lowland wetlands on Joint Base Elmendorf-Richardson, 2020.	63
0.13	Appendix C: Lesser Yellowlegs Project Summary	84
0.13.1	Lesser Yellowlegs in Anchorage	85
0.13.2	Annual Survival Estimates	87
0.13.3	Migratory Connectivity	98

List of Figures

1	Joint Base Elmendorf-Richardson is comprised of variable habitat types from boreal wetlands and tidal influenced mudflats to alpine ridgelines. Eagle River Flats was the focus of Lesser Yellowlegs captures in late June 2020.	8
2	Alpine habitat comprises about 20% of Joint Base Elmendorf-Richardson. Snowhawk Valley is a critical nesting habitat for many alpine obligate species including American Pipit, Northern Wheatear, and Surfbird	9
3	Core boreal wetland survey units selected in 2020	11
4	Core boreal wetland survey units selected in 2020	12

5	Point count locations within the Eagle River Flats proposed expansion area .	14
6	ARU locations on Eagle River Flats in 2019 and 2020. Two units were placed in the same location on the eastern edge of Eagle Rier Flats in both years . .	16
7	Recently hatched Lesser Yellowlegs chicks in nest bowl at Ammo Country wetlands	17
8	Placemnet of TinyTag data logger probe in the center of a Lesser Yellowlegs nest	18
9	Leg bands: Metal, plastic and alphanumeric flags	19
10	Measuring the diagonal tarsus of a Lesser Yellowlegs	19
11	Blackpoll Warbler are a Neotropical migrant that breeding in the boreal forests of Alaska and Canada. This is a late arrive species that was only detected during the second survey period. The male’s striking black and white plumage is contrary to the females’ monocramatic olive-green plumage	22
12	Blackpoll Warbler and Rusty Blackbird nest within mixed forests along the edges of wetlands	28
13	Sandpiper species forage in the emergent vegetation along the edges of lakes and ponds.	29
14	A failed Lesser Yellowlegs nest found on the Ammo Country wetlands at Joint Base Elmendorf-Richardson, 2020. The 4mm diameter fissures and egg frangments suggest the nest was depredated by an avian predator.	35
15	Lesser Yellowlegs study sites in 2018 and 2019. PinPoint Argos GPS units were deployed and biological samples were collected at each study site. . . .	85
16	GPS locations of Lesser Yellowlegs on Joint Base Elmendorf-Richardson by year (top) and individual (bottom).	86
17	Lesser Yellowlegs banded with alphanumeric flag “TY” on Joint Base Elmendorf-Richardson, 2020.	88
18	A Lesser Yellowlegs carrying a PinPoint Argos GPS tag on Ile Niapiskau (Mingan Archipeligo), Quebec, 2020. Photo: Y. Rochepault.	99
19	Migratory track lines for Lesser Yellowlegs tagged with Lotek PinPoint GPS Argos Satellite tags, 2018-2020.	100
20	Fall migratory stopover locations for Lesser Yellowlegs fitted with PinPoint Argos GPS tags. Blue polygons represent U.S. military installations. These were the individuals closest to military installations in 2018-2020.	100

List of Tables

0.1 Acronyms and Abbreviations

ADF&G	Alaska Department of Fish and Game
ARU	 Automated Recording Unit
BBA	Breeding Bird Atlas
BBS	Breeding Bird Survey
BCR	Bird Conservation Region
BPIF	Boreal Partners in Flight
CEIEC	Natural Resources Element, Conservation and Compliance
CES	Civil Engineer Squadron
DoD	Department of Defense
DOI	Department of Interior
ERF	Eagle River Flats (Joint Base Elmendorf-Richardson)
GPS	Global Positioning System
JBER	Joint Base Elmendorf-Richardson
MBM	Migratory Bird Management (U.S. Fish and Wildlife Service)
MWH	MWH Americas, Inc.
NWI	National Wetlands Inventory
PIF	Partners in Flight
SOSC	Species of Special Concern
UAA	University of Alaska Anchorage
USDA	U.S. Department of Agriculture
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey

0.2 Executive Summary

Many of North America’s most rapidly declining avifauna breed in the boreal ecosystems of Alaska. Joint Base Elmendorf-Richardson (JBER) located in Anchorage, Alaska, provides critical nesting habitat for five avian Species of Special Concern (SOSC): Solitary Sandpiper *Tringa solitaria*, Lesser Yellowlegs *Tringa flavipes*, Olive-sided Flycatcher *Contopus cooperi*, Rusty Blackbird *Euphagus carolinus* and Blackpoll Warbler *Setophaga striata*. Joint Base Elmendorf-Richardson derived the Species of Special Concern list from the Birds of Conservation Concern (USFWS 2008) and the Department of Defense Mission-Sensitive Priority Bird Species (DoD PIF 2015) lists. Department of Defense Partners in Flight recognizes both Olive-sided Flycatcher and Rusty Blackbirds as species of concern, while Solitary Sandpiper, Lesser Yellowlegs and Blackpoll Warbler are listed as species of high conservation need. Additionally in Alaska, Short-billed Dowitcher *Limnodromus griseus* and Greater Yellowlegs *Tringa melanoleuca* breed almost exclusively in Southcentral Alaska and are known to nest in boreal wetland habitat within the boundaries of Joint Base Elmendorf-Richardson.

Beginning in 2007, the U.S. Fish and Wildlife Service and Department of Defense initiated a collaborative effort to fill knowledge gaps on Species of Special Concern and boreal wetland breeders on Joint Base Elmendorf-Richardson. Primary objectives include: (1) develop and implement a survey program to determine relative abundance, distribution, and habitat associations of Species of Special Concern and boreal breeders; (2) implement avian surveys to assist with Environmental Impact Statement (EIS) reports, (3) capture, individually mark, and monitor Species of Special Concern and boreal breeders to estimate annual adult survival and breeding site fidelity; and (4) deploy tracking devices on Species of Special Concern and boreal breeders to identify habitat selection on Joint Base Elmendorf-Richardson, annual migratory routes, and important stopover and wintering locations. Since 2016, four boreal wetland surveys, 1 montane survey, and two Eagle River Flats point count surveys have been completed. Additionally, we deployed tracking devices on Solitary Sandpiper, Lesser Yellowlegs, Greater Yellowlegs, Olive-sided Flycatcher, Rusty Blackbird, and Tree Swallow.

The success of these survey and tracking studies compliment multiple-state and continent-wide efforts to investigate factors contributing to long-term Species of Special Concern population declines. Species abundance estimates and information collected on important habitats and nesting sites is consistent with the military’s environmental stewardship goals as follows: (1) conduct inter-service and international readiness training while meeting the natural resource management goals of Joint Base Elmendorf-Richardson, (2) inform the Bird/Wildlife Aircraft Strike Hazard Program to improve the U.S. Bird Avoidance Model for the safety of both military and civilian aircraft, (3) inform the habitat and species recovery programs throughout the U.S. and Canada, (4) assist with the national and international efforts to understand the limiting factors driving avian population declines, and (5) model physical and genetic linkages between breeding and non-breeding locations to identify connectivity of breeding avian populations on DoD lands in Alaska to non-breeding populations internationally.

0.3 Introduction

The Sikes Act Section 101(a) (16 U.S. Code 670(a)) requires that Department of Defense installations prepare an Integrated Natural Resource Management Plan (INRMP) to conserve and restore natural resources on military installations, and ensure all military activities align with environmental stewardship, laws and regulations, and the mission of the Department of Defense (DoD). The United States Air Force (USAF) environmental program developed through the INRMP advocates for natural resource stewardship, including the conservation of avian communities and habitats. A plan recently outlined by the DoD, *Strategic Plan for Bird Conservation and Management on Department of Defense Lands* (DoD 2014) aims to manage and conserve resident and migratory bird species and their habitats on DoD lands, while maintaining mission readiness resources for military training purposes.

Avian species associated with boreal wetlands are among the continents most rapidly declining avifauna. Of particular concern are the precipitous range-wide declines since 1970 of Blackpoll Warblers (-4.5% / yr.), Rusty Blackbirds (-3.5% / yr.), Olive-sided Flycatchers (-3.0% / yr.), Lesser Yellowlegs (-2.2% / yr.), and Solitary Sandpipers (0.01% / yr.; Sauer et al. 2017). Between 199 - 2015 in the Northwestern Interior Forest Bird Conservation Region (BCR) of Alaska, the same species experienced population declines or minimal growth; Blackpoll Warblers (-2.3% / yr.), Rusty Blackbirds (0.4% / yr.), Olive-sided Flycatchers (-1.2% / yr.), Lesser Yellowlegs (-2.0% / yr.), and Solitary Sandpipers (-1.3% / yr.; Handel and Sauer 2017). These species, designated as Species of Special Concern (SOSC), have lost an estimated 50 - 90% of their populations and most are projected to lose an additional 50% within the next 15 - 25 years (Rosenberg et al. 2019, Rosenberg et al. 2016).

State and continental conservation planning efforts have emphasized the need to determine demographic deficits, key habitats, linkages between breeding and wintering populations, contaminant burdens, and the incidence of disease among SOSC populations (Alaska Shorebird Group 2008, DoD Partners in Flight [PIF] 2015, Alaska Department of Fish and Game 2015, U.S. Shorebird Conservation Plan Partnership 2016, Boreal Partners in Flight 2017). Such information is urgently required to identify the mechanisms driving declines, help direct conservation and management of important areas and habitats, and ultimately help reverse avian population declines before more costly recovery efforts are required.

Department of Defense has a long history of supporting bird studies in Alaska, with the overarching goal to maintain the health of migratory and resident bird populations and their supporting habitats to allow for flexibility in military operations (DoD PIF 2015). Efforts to characterize the Joint Base Elmendorf-Richardson (JBER) avifauna were first initiated by U.S. Fish and Wildlife Service (USFWS) in 1994 with the establishment of an intensive inventory and monitoring effort that consisted of North American Breeding Bird Surveys link, Breeding Bird Atlas link, and Monitoring Avian Productivity and Survivorship link programs (Browne and Andres 2002, Andres 2005).

In 2007, the Alaska Department of Fish and Game (ADF&G) and USFWS began a comprehensive research effort to describe breeding ecology and reproductive rates of Rusty Blackbirds on JBER. Several publications resulted from this effort, which highlighted the importance of DoD lands in Alaska for Rusty Blackbirds (Matsuoka et al. 2010a, Matsuoka et al. 2010b).

MWH Americas Inc. (MWH) was contracted by 673d Civil Engineering Squadron/Civil Engineering Installation Environmental Conservation (673d CES/CEIEC) to build upon these efforts by expanding the extent of surveys to include several additional SOSC primarily associated with boreal wetlands. With input from USFWS, MWH surveyed wetlands during 2014 - 2015 to further establish baseline information on the abundance and distribution of wetland-associated SOSC (MWH Americas, Inc. 2015).

Since 2015, a collaborative effort between ADF&G, DoD, and USFWS continues to fill important information gaps for five SOSC on JBER: Solitary Sandpiper, Lesser Yellowlegs, Olive-sided Flycatcher, Rusty Blackbird, and Blackpoll Warbler. Primary objectives include: (1) develop and implement a robust survey and monitoring program to determine relative abundance and critical nesting habitat for wetland-associated SOSC, (2) implement avian surveys to assist with Environmental Impact Statement (EIS) reports, (3) capture and individually band SOSC as a means to estimate apparent adult survival, and (4) attach small tracking devices to SOSC, as well as Greater Yellowlegs and Short-billed Dowitcher, to determine local habitat selection on JBER and migratory connectivity between breeding and non-breeding locations. In addition, we collect occupancy data on seven focal species with suspected or known population declines: Common Loon, Red-throated Loon, Red-necked Grebe, Lesser Scaup, Greater Yellowlegs, Short-billed Dowitcher, Long-billed Dowitcher, and Wilson's Snipe.

Results from monitoring and tracking efforts complement the Birds and Bogs program, a citizen science program organized by Audubon Alaska and the University of Alaska Anchorage (UAA) that aims to document the distribution and abundance of wetland-associated boreal bird species in Anchorage, JBER, and the Matanuska-Susitna Valley (Mat-Su). A current UAA graduate student will use the data collected by both studies to examine habitat use patterns among wetlands within urban (Anchorage and Mat-Su) and non-urban (JBER) areas. Information from the migration tracking are used to investigate range-wide patterns of migratory connectivity and habitat use, and will be published as part of a Masters thesis. All information collected during surveys and tracking studies are imported annually to the DoD avifaunal database (ADAML); a program used to organize, summarize, and display geographically referenced avian information for DoD installations in Alaska.

0.4 Study Area

Rising from sea level to an elevation over 1,500 m, JBER encompasses a variety of habitat types including estuarine and freshwater wetlands (Figure 1), shrublands, broadleaf and coniferous forests, and alpine tundra (Figure 2). Despite long-term training activities across the installation, JBER contains the largest continuous extent of natural, low-elevation habitats in Anchorage (Andres 2005). For a detailed description of habitats see Jorgenson et al. (1998), Browne and Andres (2002), Andres et al. (2005), Matsuoka et al. (2010), and MWH Americas Inc. (2015).




Figure 1: Joint Base Elmendorf-Richardson is comprised of variable habitat types from boreal wetlands and tidal influenced mudflats to alpine ridgelines. Eagle River Flats was the focus of Lesser Yellowlegs captures in late June 2020.




Figure 2: Alpine habitat comprises about 20% of Joint Base Elmendorf-Richardson. Snowhawk Valley is a critical nesting habitat for many alpine obligate species including American Pipit, Northern Wheatear, and Surfbird

0.5 Methods

0.5.1 Wetland Area Surveys: Study Design

Our sampling universe included freshwater wetlands ~150m in elevation within JBER boundaries. In the lowlands, we treated wetland bodies as sampling units because of their discreteness within a matrix of boreal forest habitat. We used a two-tiered approach to select wetland-sampling units: (1) non-random and (2) stratified random. Non-random units, also known as core units, were wetlands selected for their expected value to Rusty Blackbirds (see Matsuoka et al. 2010 for details). Typically, these wetlands were seasonally or permanently flooded and included shorelines of lakes, ponds, and rivers with emergent vegetation, seasonally flooded scrub-shrub, or forests. A 50-m buffer around wetlands added potential nesting sites for Rusty Blackbirds in adjacent non-flooded wetlands or upland forest. When buffers of adjacent units overlapped, they were combined into a single unit. This non-random selection process resulted in 27 lowland units that ranged in size from 1 - 139 ha (see Matsuoka et al. 2009; Figure 3). We only conducted breeding bird surveys on core units in 2020.

To select stratified random sampling units for the breeding bird surveys, we used a spatial layer created by the National Wetlands Inventory (NWI; Cowardin et al. 1979) that was modified by MWH Americas Inc (2015), and updated annually by JBER. NWI classes were combined into three general habitat types: (1) emergent vegetation (permanently flooded by wetlands), (2) scrub- shrub (willow [*Salix*] spp. and alder [*Alnus*] spp. shrublands occurring on saturated soils to areas seasonally flooded by freshwater), and (3) forested wetlands (black spruce [*Picea mariana*] dominated forest occurring on saturated soils to areas seasonally flooded by freshwater). We excluded two additional wetland habitat classes from the selection process: (1) estuarine (all wetlands influenced by marine waters year round), and (2) ponds, lakes, and rivers, because these open waterbodies do not provide potential nesting habitat for SOSC or focal species. The amount of each habitat type in each wetland was measured in hectares, and units were assigned to the habitat type that covered the largest area. We ranked the three general wetland classes based on their expected suitability to SOSC breeding requirements: emergent = good, scrub-shrub = medium, and forest = poor. Sample allocation was weighted so that emergent and scrub- shrub dominated wetlands were selected at a higher proportion relative to their availability compared to forested wetlands. We conducted surveys during two periods that corresponded with peak detectability of (1) migratory waterbirds, resident and Nearctic migratory passerines (15 - 17 May); and (2) Neotropical migratory passerines (29-31 May). We conducted surveys 6:00 - 12:00 during the first period and 5:00 - 11:00 during the second period. We postponed surveys during periods of fog, steady rain, and/or wind exceeding 20 km/h. Directional photographs were taken at the midpoint of each survey unit as a means to compare successional vegetation growth over time (Figure 4).

We used the area search method to survey breeding bird occupancy. Surveyors covered 10 ha/hr while attempting to pass within 100 m of any location inside a unit. Surveyors mapped observations of SOSC and focal species on aerial photographs, using codes for pairs, males, females, unknown sex, and groups. Surveyors noted behaviors to determine whether birds were territorial and breeding on the units or were non- breeding migrants. Surveyors differentiated between actual and estimated pairs. An actual pair was defined as two birds

together exhibiting breeding behavior. An estimated pair was defined as (1) a territorial individual (displaying shorebird, singing passerine, or alarm-calling individual of either sex), or (2) a single female (only applicable for Rusty Blackbirds and Blackpoll Warblers due to plumage sexual dimorphism). A bird was recorded on a unit if ~50% of its territory appeared to be within the unit boundary. Immediately after each survey, surveyors transcribed the number of individuals and pairs to a data sheet. In addition, surveyors recorded (but did not map) the total number of non-SOSC individuals detected within and outside the boundaries of each unit.




Figure 3: Core boreal wetland survey units selected in 2020



Figure 4: Core boreal wetland survey units selected in 2020

0.5.2 ERF Proposed Expansion Survey: Study Design

During 2020, the USFWS crew conducted avian point count surveys within a proposed expansion area on the north side of Eagle River flats. Point count surveys used the distance sampling methods (Buckland et al. 2001) following the Alaska Landbird Monitoring Survey (ALMS) protocol (Handel et al. 2019) for conducting off-road point counts in Alaska. The ALMS protocol was designed to monitor changes in bird populations over time, and has been used to provide population trend estimates of breeding birds throughout Alaska (Handel & Sauer 2017).

To select point count locations, we overlaid a 500m x 500m grid centered on the midpoint of the proposed expansion area. The size of the expansion area determined the number of survey points. We selected the 10-point count locations that fell within the expansion area boundary (Figure 5). Point selection was not influenced by accessibility or slope, and it was up to the observer's discretion whether or not a point was safe to access; however, all points were deemed safe during the initial survey period.

Each survey conducted was 10 minutes in length and completed twice in the season (23 May and 6 June). Surveys began at 5:30 and were completed prior to 8:00. We did not conduct surveys during weather conditions that influenced 1) a bird's behavior and 2) our ability to detect a bird. We chose to postpone any ongoing surveys during periods of fog, steady rain, and/or wind exceeding 15 mph. We recorded the temperature, wind speed, and sky conditions at the start and end of each survey period using a Kestrel Wind Meter and the Beaufort wind scale. Observers approached each survey point with minimal disturbance and began the survey immediately upon arrival. Each survey was conducted by a single observer. The observer recorded species, time interval (0-3 min, 3-5 min, 5-8 min, 8-10 min),

behavior (i.e. singing, calling, flight display, etc.), and distance interval for all individual birds or groups of birds associating together. Horizontal distance from the observer to the bird(s) were recorded in 10 m increments out to 100 m, 25 m increments from 101-150 m, and 50 m increments from 151 - 400 m. Any detections beyond 400 m were denoted as greater than 400 m. A range finder was used by all observers to accurately measure and estimate distances. Observers recorded all observations on a circular map during the survey and summarized observation in a table immediately following each survey. Directional photos (N, S, E, and W) were collected at each point as a habitat reference. Finally, each observer recorded the species of mammals and birds detected between point count locations.




Figure 5: Point count locations within the Eagle River Flats proposed expansion area

0.5.3 Automated Recording Units

In 2019 and 2020, the USFWS crew and 673d Civil Engineer Squadron biologist Chris Garner, deployed Automated Recording Units (ARU) within the boundary of Eagle River Flats. In 2019, four units were placed along Eagle River and 2 units were placed on the eastern edge of the flats. In 2020, seven units were deployed along the southern, eastern, and northern edges of the flats (Figure 6). All units were programmed to collect bird vocalizations 4 hours after sunrise, 12:00-13:00, 16:00-17:00, and 1 hour before sunset from May through September. The units collect daily bird species occupancy data and do not require personnel to be present.

0.5.4 Nest searching

In 2020, we spent the early portion of the breeding season (May 15- June 1) searching for Lesser Yellowlegs and Wilson’s Snipe nests. We searched for Lesser Yellowlegs as part of a continuing study aimed at understanding breeding success, adult survival and migratory connectivity (Figure 7). We searched for Wilson’s Snipe nests as part of a collaborative study with Max Planck Institute of Ornithology. The goal was to examine the incubation rates of uniparental shorebirds.

To find nests, we observed bird’s behavior from a distance; mapped their locations and movements; and followed them back to their nests as they returned to incubate eggs. We recorded the location of each nest in a Global Positioning System (GPS) and collected information included the number of eggs observed, dimensions of each egg, float angle (used to determine age of eggs and to estimate hatch), and habitat type. Once found, Lesser Yellowlegs nests were monitored by motion-activated game cameras placed near each nest and temperature data logger probes (TinyTag, Gemini Data Loggers Ltd, Chichester, West Sussex, UK) placed in the center of each nest (Figure 8). Cameras were used to help us determine the cause of a nest failure (e.g. predation), while the temperature loggers were used to determine how frequently birds incubate during the 23-day incubation period. To monitor Wilson’s Snipe nests, we deployed temperature loggers (MSR, MSR Electronics, Seuzach, Switzerland) that were programmed to record the duration of incubation during a 24-hour period. Nest cameras were not utilized for Wilson’s Snipe nest monitoring.

0.5.5 Captures and GPS tag deployment

In 2020, we captured and banded Lesser Yellowlegs, Greater Yellowlegs, Solitary Sandpiper, and Blackpoll Warbler. The majority of shorebird captures occurred 1-3 days post hatch when adults were most responsive to chick calls. During early brood rearing, adults are extremely responsive to recordings of conspecific chick alarm calls. We captured adults by laying a mist net flat over tall grass or perpendicular to the ground and playing a chick distress call near the center of the net. The adults either walk under or fly into the net. In rare instances, adults that will not land on the ground near the net are trapped using a mist net swung vertically as a bird flies over. In addition to on-brood captures, in 2020 we also attempted passive captures of foraging Lesser Yellowlegs on the edge of Eagle River Flats. We set-up six vertical mist nets in an area we observed to have a high concentration of foraging Lesser Yellowlegs. We placed a wireless speaker in the center of the net set-up




Figure 6: ARU locations on Eagle River Flats in 2019 and 2020. Two units were placed in the same location on the eastern edge of Eagle River Flats in both years



Figure 7: Recently hatched Lesser Yellowlegs chicks in nest bowl at Ammo Country wetlands

and played a foraging call as an attractant.

Following captures, we attached a single aluminum U.S. Geological Survey (USGS) band, green color band, and dark green flag engraved with a white two-character alpha code (following the Pan-American Shorebird Program 2016; Figure 9). We attached PinPoint Argos GPS tags via a leg loop harness backpack to birds weighing at or over 80 grams (Rappole 1991). Finally, we recorded morphometrics (Figure 10) and collected biological samples from all banded birds. From each bird, we collected three blood samples: (1) genetic analysis (genoscape), (2) mercury contamination testing, and (3) molecular sexing. Additionally, we collected secondary feather and claw samples for use in a future stable isotope analysis.



Figure 8: Placemnet of TinyTag data logger probe in the center of a Lesser Yellowlegs nest



Figure 9: Leg bands: Metal, plastic and alphanumeric flags



Figure 10: Measuring the diagonal tarsus of a Lesser Yellowlegs

0.5.6 Data Management

Quality assured and quality controlled survey and banding data, GIS projects and associated data layers, and digitized hardcopies of survey maps and data forms are archived at USFWS-Migratory Bird Management. The JBER breeding bird surveys, as well each SOSC species follow specific data management and metadata protocols implemented by USFWS Alaska Region. See Products section (## Products) for a full list of deliverables generated from this project.

0.6 Results

0.6.1 Wetland Area Surveys

During the two survey periods (15 - 17 May and 29 - 31 May) we surveyed 27 core wetland units. The first survey was completed between 6:00 and 12:00 am with a total survey effort of 22 hours and 54 minutes. The second survey was completed between 5:00 and 11:00 with a total survey effort of 20 hours and 55 minutes. Four USFWS personnel completed the surveys.

0.6.1.1 All Species detections

Inside survey unit boundaries, we detected a total of 2,674 individual birds comprised of 82 species. Taxonomic family groups included 15 waterfowl species, 1 crane species, 8 shorebird species, 2 loon and grebe species, 4 gull species, 5 hawk species, 4 woodpecker species, 1 falcon species, and 39 passerine species (Appendix A).

The number of species and their total abundances varied between survey periods. We detected 56 species comprising 1,555 individuals during the first survey. The most commonly detected species within unit boundaries were Lincoln's Sparrow (13% of all individuals, 100% of units), Yellow-rumped Warbler (11% of all individuals, 70% of units), and Wilson's Snipe (10% of all individuals, 70% of units). This closely corresponds with results from previous years, where Lincoln's Sparrow, Yellow-rumped Warbler, and Dark-eyed Junco were the most commonly detected species during the first survey period. Additionally, several species were detected only during the first survey, including: Bald Eagle, Barrow's Goldeneye, Black-capped Chickadee, Brown Creeper, Common Loon, Downy Woodpecker, Fox Sparrow, Gadwall, Greater Scaup, Northern Goshawk, Northern Pintail, Red-breasted Nuthatch, Sandhill Crane and Violet-green Swallow.

During the second survey period, we detected 43 species comprising 1,119 individuals. The most commonly detected individuals within unit boundaries were Lincoln's Sparrow (15% of all individuals, 81% of units), Wilson's Snipe (13% of all individuals, 78% of units), and Yellow-rumped Warbler (10% of all individuals, 67% of units). The results from the second survey follow a similar trend to previous survey years, and the earlier survey period in 2020, however, American Robin were detected more frequently in 2017 and 2019. Additionally, Blackpoll Warbler, Short-billed Dowitcher and Spotted Sandpiper were only detected during the second survey period.

The differences in species composition indicates that two survey periods are required in order to differentiate between resident and early-arriving species (Nearctic migrants, waterbirds, and shorebirds), and late-arriving species (Neotropical migrant passerines). Late-arriving species such as Olive-sided Flycatcher, Alder Flycatcher, Swainson's Thrush and Blackpoll Warbler, were either not detected during the first survey or were more numerous during the second survey.

```
spacc1.2020<- read.csv("specaccum1_2020.csv", header = TRUE)
spacc2.2020<- read.csv("specaccum2_2020.csv", header = TRUE)
```

```
s1 <-colSums(spacc1.2020, na.rm = FALSE, dims = 1)
s1[order(-s1)]
```

##	plot	lincolns.sparrow
##	11972	199
##	yellowrumped.warbler	wilsons.snipe
##	179	158
##	darkeyed.junco	rubycrowned.kinglet
##	140	128
##	american.robin	ringnecked.duck
##	102	75
##	orangecrowned.warbler	goldencrowned.kinglet
##	50	41
##	varried.thrush	rusty.blackbird
##	32	31
##	northern.waterthrush	boreal.chickadee
##	30	27
##	lesser.yellowlegs	common.redpoll
##	27	24
##	tree.swallow	canada.goose
##	24	22
##	savannah.sparrow	wilsons.warbler
##	21	20
##	greenwinged.teal	american.wigeon
##	18	17
##	bonapartes.gull	mallard
##	17	15
##	solitary.sandpiper	whitecrowned.sparrow
##	15	15
##	hermit.thrush	northern.shoveler
##	12	12
##	olivesided.flycatcher	canada.jay
##	10	9
##	greater.yellowlegs	blackcapped.chickadee
##	9	6



Figure 11: Blackpoll Warbler are a Neotropical migrant that breeding in the boreal forests of Alaska and Canada. This is a late arrive species that was only detected during the second survey period. The male's striking black and white plumage is contrary to the females' monochromatic olive-green plumage

##	gadwall	rednecked.grebe
##	6	6
##	mew.gull	greater.scaup
##	5	4
##	lesser.scaup	redbreasted.nuthatch
##	4	4
##	swainsons.thrush	westernwood.peewe
##	4	4
##	whitewinged.crossbill	common.loon
##	4	3
##	arctic.tern	american.threetoed.woodpecker
##	2	2
##	bald.eagle	barrows.goldeneye
##	2	2
##	brown.creeper	downy.woodpecker
##	2	2
##	fox.sparrow	hairy.woodpecker
##	2	2
##	northern.pintail	rednecked.phalarope
##	2	2
##	trumpeter.swan	violetgreen.swallow
##	2	2
##	northern.goshawk	sandhill.crane
##	1	1
##	alder.flycatcher	american.dipper
##	0	0
##	american.pipit	bank.swallow
##	0	0
##	blackbilled.magpie	belted.kingfisher
##	0	0
##	blackpoll.warbler	bluewinged.teal
##	0	0
##	bohemian.waxwing	canvasback
##	0	0
##	common.goldeneye	common.raven
##	0	0
##	goldencrowned.sparrow	herring.gull.hybrid
##	0	0
##	least.sandpiper	merlin
##	0	0
##	northern.flicker	northern.harrier
##	0	0
##	pine.siskin	redtailed.hawk
##	0	0
##	redthroated.loon	semipalmated.plover

```
##                                0                                0
##      shortbilled.dowitcher      spotted.sandpiper
##                                0                                0
##      sharpshinned.hawk          townsend.solitaire
##                                0                                0
##      townsend.warbler            yellow.warbler
##                                0                                0
```

```
s2 <-colSums(spacc2.2020, na.rm = FALSE, dims = 1)
s2[order(-s2)]
```

```
##                                plot                                lincolns.sparrow
##                                11972                                164
##      wilsons.snipe            yellowrumped.warbler
##                                142                                108
##      darkeyed.junco            rubycrowned.kinglet
##                                84                                76
##      american.robin            orangerowned.warbler
##                                59                                50
##      swainsons.thrush            northern.waterthrush
##                                46                                44
##      lesser.yellowlegs            bonapartes.gull
##                                36                                34
##      savannah.sparrow            canada.jay
##                                24                                23
##      tree.swallow            olivesided.flycatcher
##                                21                                18
##      rusty.blackbird            whitecrowned.sparrow
##                                16                                16
##      westernwood.peewe            varried.thrush
##                                16                                14
##      boreal.chickadee            goldencrowned.kinglet
##                                13                                11
##      mallard            wilsons.warbler
##                                11                                10
##      whitewinged.crossbill            canada.goose
##                                10                                9
##      greater.yellowlegs            arctic.tern
##                                9                                6
##      mew.gull            american.wigeon
##                                6                                5
##      hermit.thrush            common.redpoll
##                                5                                4
##      hairy.woodpecker            lesser.scaup
##                                4                                4
```


##	trumpeter.swan	greenwinged.teal
##	4	3
##	solitary.sandpiper	american.threetoed.woodpecker
##	3	2
##	blackpoll.warbler	ringnecked.duck
##	2	2
##	rednecked.grebe	rednecked.phalarope
##	2	1
##	shortbilled.dowitcher	spotted.sandpiper
##	1	1
##	alder.flycatcher	american.dipper
##	0	0
##	american.pipit	bald.eagle
##	0	0
##	barrows.goldeneye	bank.swallow
##	0	0
##	blackbilled.magpie	blackcapped.chickadee
##	0	0
##	belted.kingfisher	bluewinged.teal
##	0	0
##	bohemian.waxwing	brown.creeper
##	0	0
##	canvasback	common.goldeneye
##	0	0
##	common.loon	common.raven
##	0	0
##	downy.woodpecker	fox.sparrow
##	0	0
##	gadwall	goldencrowned.sparrow
##	0	0
##	greater.scaup	herring.gull.hybrid
##	0	0
##	least.sandpiper	merlin
##	0	0
##	northern.flicker	northern.goshawk
##	0	0
##	northern.harrier	northern.pintail
##	0	0
##	northern.shoveler	pine.siskin
##	0	0
##	redbreasted.nuthatch	redtailed.hawk
##	0	0
##	redthroated.loon	sandhill.crane
##	0	0
##	semipalmated.plover	sharpshinned.hawk

##	0	0
##	townsends.solitaire	townsends.warbler
##	0	0
##	violetgreen.swallow	yellow.warbler
##	0	0

0.6.1.2 Species of Special Concern Detections

We recorded all five SOSC during the two survey periods. Lesser Yellowlegs were the most numerous and widespread SOSC species across JBER wetlands (Table 1). Overall, the number of SOSC individuals was less than in previous years, which we attribute to fewer units surveyed and a lack of nesting and foraging habitat at Otter Lake due to flooding by beaver dams (Figure 11). Lesser Yellowlegs were the most frequently detected SOSC species (63 total individuals) during both surveys, while Solitary Sandpipers, Olive-sided Flycatchers and Rusty Blackbirds were less abundant (18, 28 and 47 individuals, respectively). All SOSC occurred in similar numbers during survey periods, except for the Blackpoll Warbler, a Neotropical migrant, which we only detected during the second survey (2 individuals within unit boundaries).

Table 1: Summary of Species of Special Concern detected on unit during two survey periods on Joint Base Elmendorf-Richardson, 2020.

Survey 1 (May 15-17)

Survey 2 (May 29-31)

Species

Individ.

Pairs

Prop*

Individ

Pairs

Prop

Solitary Sandpiper

15

7

18.5

3

1

7.4

Lesser Yellowlegs

27

13

37.0

36

18

29.6

Olive-sided Flycatcher

10

5

14.8

18

9

29.6

Rusty Blackbird

31

15

44.4

16

8

29.6

Blackpoll Warbler

0

0

0.0

2

1

3.7

* Proportion of survey units containing the indicated Species of Special Concern.

of Solitary Sandpiper, Lesser Yellowlegs, Olive-sided Flycatcher, Rusty Blackbird, Blackpoll Warbler) (15, 27, 10, 31, 01, 02, 13, 5, 15, 01, 18, 5, 37, 14, 8, 44, 4, 01, 02, 36, 18, 16, 21, 01, 18, 9, 8, 11, 07, 4, 29, 6, 29, 6, 29, 6, 3, 7)

Each SOSC's migration strategy and natural history characteristics clearly explain the patterns of detectability. Lesser Yellowlegs are both vocally and visually obvious throughout their breeding cycle, and arrive early. Solitary Sandpipers are much more reserved than Lesser Yellowlegs and are readily detectable only during displaying (advertising for mates) or defending broods. The Olive-sided Flycatcher is a long-distance migrant that winters in northern South America, and although highly detectable throughout their breeding cycle, they typically arrive in late May. Our data suggest that the first survey occurred before most individuals, likely females, had arrived to the nesting grounds. The Blackpoll Warbler is another long-distance migrant that winters in northern and central South America. Blackpoll Warblers are a late arriving species and are detectable when males are defending their territories in late-May and early-June. Rusty Blackbird, a relatively short-distance migrant that winters in the southern U.S., is an early-arriving species that is generally more obvious and vocal during courtship and nest building activities that overlapped most with the first survey. Again, survey results support the need for two survey periods to accurately detect this suite of SOSC.

Preliminary analysis of habitat association of the SOSC suggested that Solitary Sandpipers and Lesser Yellowlegs were associated with wetlands dominated by emergent vegetation, Olive-sided Flycatchers were associated with shrub-dominated wetlands, and Rusty Blackbirds and Blackpoll Warblers were associated with forested wetlands (Figure 12).



Figure 12: Blackpoll Warbler and Rusty Blackbird nest within mixed forests along the edges of wetlands

0.6.2 Eagle River Flats Expansion: Point Counts

In 2020, point count surveys were completed in the designated Eagle River Flats expansion area. The 10-point count locations were divided among four personnel and were completed



Figure 13: Sandpiper species forage in the emergent vegetation along the edges of lakes and ponds.

between 6:00 and 9:00 on May 23 and June 6. We recorded 309 individuals comprised of 28 species across both survey periods.

Detection rates were high for most species groups except for raptors (hawks, owls, and falcons) and loons and grebes, which were detected at 0% and 20% of points, respectively. Passerines were the most commonly detected species group and recorded at 100% of survey points. Swainson's Thrush were detected at all point count locations, followed by Yellow-rumped Warbler (Myrtle), American Robin and Ruby-crowned Kinglet (90% of points). Additionally, Sandhill Crane were detected at all point count locations, however this species was always recorded beyond 100 m. Wilson's Snipe, Red-necked Grebe, Common Loon, Green-winged Teal, American Wigeon, Western Wood-Pewee, Tree Swallow, Hairy Woodpecker, Golden-crowned Kinglet, Canada Jay, Boreal Chickadee, Black-capped Chickadee, and Alder Flycatcher were detected at only one point count location. These results only reflect species detected during point counts, and do not include species detected between points.

In addition to avian species, we collected information on mammals observed between point count locations. During the survey on 23 May, a pack of Grey Wolves was observed between survey points 1 and 3. On this same day, a moose with fresh wounds was observed standing in shallow water on the edge of Eagle River Flats ~155 m from survey point 1. When we returned to the location of the moose on 20 June, a moose carcass and wolf scat were present.

Table 2: Avian species detected on the proposed expansion area of Eagle River Flats during

two point counts periods, 2020.

Species

Point 1

Point 2

Point 3

Point 4

Point 5

Point 6

Point 7

Point 8

Point 9

Point 10

Canada Goose

X

X

X

X

X

American Wigeon

X

Green-winged Teal

X

Red-necked Grebe

X

X

X

X

Sandhill Crane

X

X

X

X

X

X

X

X

X

X

Short-billed Dowitcher

X

X

X

Wilson's Snipe

X

Lesser Yellowlegs

X

X

X

X

X

X

X

Mew Gull

X

X

Common Loon

X

American Three-toed Woodpecker

X

X

X

Hairy Woodpecker

X

Western Wood-Pewee

X

Alder Flycatcher

X

Canada Jay

X

Black-capped Chickadee

X

Boreal Chickadee

X

Tree Swallow

X

Golden-crowned Kinglet

X

Ruby-crowned Kinglet

X

X

X

X

X

X

X

X

X

Varied Thrush

X

X

X

X

X

Swainson's Thrush

X

X

X

X

X

X

X

X

X

X

Hermit Thrush

X

X

American Robin

X

X

X

X

X

X

X

X

X

Common Redpoll

X

X

X

X

X

X

X

X

Dark-eyed Junco

X

X

X

X

X

Orange-crowned Warbler

X

X

X

Yellow-rumped Warbler

X

X

X

X

X

X

X

X

X

Wilson's Warbler

X

X

X

X

X

©"Solitary Sandpiper", "Lesser Yellowlegs", "Olive-sided Flycatcher", "Rusty Blackbird", "Blackpoll Warbler") c(15, 27, 10, 31, 0) c(7, 13, 5, 15, 0) c(18, 5, 37, 14, 8, 44, 4, 0) c(3, 36, 14, 16, 2) c(1, 16, 9, 8, 11) c(7, 4, 29, 6, 29, 6, 3, 7)

0.6.3 Nest Discoveries

The USFWS crew successfully found 7 Lesser Yellowlegs, 1 Greater Yellowlegs, and 10 Wilson's Snipe nests on JBER, at various stages of development, ranging from initiation to hatch (Figure 11). Nest success for lesser yellowlegs was exceptionally high in 2020 (85.7%), compared to past years where hatch success was less than 30%. In 2020, nests were monitored with nest cameras and in-nest temperature loggers, which allowed personnel to limit the number of nest checks during the incubation period. The one nest that did fail was likely due to an avian predator, based on the shell fragments found near the nest cup (Figure 13). The single Greater Yellowlegs nest was abandoned due to unknown causes, and the Wilson's snipe nest success is currently being determined by our collaborator at Max Planck Institute.



Figure 14: A failed Lesser Yellowlegs nest found on the Ammo Country wetlands at Joint Base Elmendorf-Richardson, 2020. The 4mm diameter fissures and egg fragments suggest the nest was depredated by an avian predator.

0.6.4 Banding, Sampling, and Tracking

In 2020, we focused our banding efforts on Blackpoll Warbler, Lesser Yellowlegs, Greater Yellowlegs, Solitary Sandpiper, and Short-billed Dowitcher. We captured and banded 11

Blackpoll Warbler, 4 Greater Yellowlegs, 2 Solitary Sandpiper, and 49 Lesser Yellowlegs (29 adults, 20 chicks) on JBER (see Appendix C for full list of banded individuals). We did not capture any Short-billed Dowitcher within the JBER boundary.

Blackpoll Warbler were captured along Dena'ina Road on Elmendorf. All adults captured were territorial males and banded with a unique color band combination. We hope to resight individuals in future year to determine site fidelity and apparent adult survival.

Greater Yellowlegs were captured along Fish Lake road and the northern edge of Eagle River Flats. As with the Blackpoll Warbler, we banded all birds with a unique color band combination. Additionally, we deployed two PinPoint Argos GPS tags (Lotek Wireless, Inc., Newmarket, Ontario) on breeding adults. This was the first year tags were attached to the species and we look forward to sharing the results next season.

We captured and banded a breeding pair of Solitary Sandpiper at Otter Lake South during the brood rearing period. In 2019, the margins of Otter Lake were flooded due to a beaver dam, and this continued into the 2020 breeding season. Foraging and nesting habitat was reduced for many shorebird species, however, a pair of Solitary Sandpiper successfully nested in the surrounding black spruce forest. As with all banded birds, we hope to resight these individuals in future years to determine adult survival.

Lastly, the USFWS and ADF&G crews individually marked 103 Lesser Yellowlegs at sites on JBER, the Anchorage Coastal Wildlife Refuge and Far North Bicentennial Park. This was the fifth consecutive year of banding Lesser Yellowlegs on JBER, and the third consecutive year we deployed PinPoint Argos GPS on breeding adults. In 2016, we began deploying light-level geolocators on adult Lesser Yellowlegs as a tool to understand how Lesser Yellowlegs migrate, and the susceptibility of Anchorage breeders to unregulated shorebird harvest in South America and the Caribbean. Light-level geolocators are lightweight, >1g, and inexpensive, but exhibit high location error (<200 km latitude and <100 km longitude) and the bird must be recaptured to obtain the data. Lesser Yellowlegs prove difficult to recapture, therefore we transitioned to PinPoint Argos GPS in 2018, which are accurate to 10m and recapture is not required to retrieve data. In addition to tracking adult migration, we also banded 2/3 of adults with unique alpha character leg flags. Our goal is to determine breeding site fidelity, apparent adult survival, and nest success. A full summary of the Lesser Yellowlegs project, including preliminary results is described in appendix B.

0.7 Interpretation

0.7.1 Species Richness Curves: Area Survey

The understanding of species diversity across boreal wetlands on JBER is critical for managing and conserving bird species and their habitats, while sustaining mission readiness resources for military training purposes. Species Richness Curves, or Species Accumulation Curves are used to estimate the number of species in a particular area. During the JBER wetland surveys, common and widespread species with high detectability were encountered quickly, as reflected in the steep rise in number of species early in the survey effort (Figure 13). As the number of surveys completed increased, the rate at which new species were detected

slowed as only uncommon and less detectable species were available for detection. Among all survey units, species richness was lower during the second survey period than the first period (Figure 13). This is likely a function of breeding chronology. Male passerine and shorebird species that were singing and displaying during the first survey period were inconspicuous during the second period as pairs established nests and began incubation.

The trend in detection rate of species during the first and second survey periods was consistent across years for surveys completed on the 27 nonrandom units (Figure 14). Species richness increased quickly with survey effort during the first period and more slowly during the second period, based on the slope of each curve. Although the overall trend was consistent among years, 2017 experienced the lowest species richness during the first survey period, while 2020 experienced the lowest species richness during the second survey period. These curves do not take into account any differences in observer effort or bird identification skill. Each year of surveys, at least one observer was unique, meaning that they did not survey in any other years.

Figure XX: 2020 Survey




Figure XX: Survey Period 1






Figure XX: Survey Period 2



Species Richness Curves are valuable for examining actual detection trends, however, it is also useful to estimate how many species would be detected if our survey effort was increased. A Rarefaction Curve model estimates the total number of species detected representing any survey effort. For example, the rarefaction curves estimates that after 500 total individuals are observed, the predicted total number of species detected would be 45 for survey period 1 and 38 for survey period 2. Doubling the survey effort to 1,000 individuals predicts an expected total species number of 51 for period 1 and 39 for period 2. This a 13% increase for survey period 1 and a 3% increase for survey period 2. These low increase percentages suggests that additional species would likely not be breeders, and instead may represent early or late passage migrants, or wandering non-breeders, detected by chance while surveying breeding individuals. Increasing survey effort during migration periods in April and August will increase species totals among survey units. However, migratory birds use each unit differently depending on habitat type and proximity to Eagle River Flats, which supports migratory species (McDuffie et al. 2018).


Figure XX: Rerfaction Curves 2020



Avian Taxon Detection Trends: Area Survey With the completion of boreal breeding bird surveys in 2020, we have four years of data representing the same 27 nonrandom units. By comparing within year trends of raw individual observations, we determined that detections for all species groups except for loon and grebe, decreased from 2016 to 2017, but increased gradually from 2017 to 2020 (Figure XX). However, it is important to note that surveys were not completed in 2018.

Passerines were the most dominate species group across all survey years, followed by shorebirds and waterfowl. The habitat mosaics at each survey unit clearly determine species presence and absence. For example, large waterbodies within survey units were eliminated from all survey units because SOSC do not nest on the water. Many waterfowl species were on waterbodies (off unit) when detected and therefore, were not included in the final survey model.

Figure XX: Detection Trends



0.7.2 Species Richness: ERF Point Counts

Species richness and diversity varied across point count locations depending on habitat type and proximity to the edge of Eagle River Flats (Figure XX). For example, points located less than 500 m from the edge of the flats had the highest species richness, on average. This is likely because species vocalizations can carry long distances across the flats, and shorter distances through the hilly and forested terrain of the proposed expansion area. The further a point was located from the flats, the lower the species richness in most cases; however, habitat type clearly influenced species richness.

To examine the potential influence of habitat type on species richness, we subset the species data to only include individuals present 100m radius or less from the centroid of the survey point. Dominant vegetation type was determined using the directional photos taken at each point count location. Coniferous dominated habitats occurred where at least three photos showed thick spruce forest. Deciduous dominated habitats occurred where at least three photos showed open deciduous dominated forests with thin undercover. Finally, mixed deciduous dominated habitats occurred where at least three photos showed an equal mixture of spruce and deciduous trees. Although only 1 of the 10 points fell under the category of coniferous, this habitat classification supported the highest average species richness among the three habitat classifications (Figure XX).

point RICHNESS

```

## 1      1      16
## 2      2       7
## 3      3      11
## 4      4      12
## 5      5       9
## 6      6      12
## 7      7       8
## 8      8      19
## 9      9      11
## 10     10      15

## 'data.frame':  10 obs. of  10 variables:
## $ point      : chr  "Exp.1" "EXP.2" "Exp.3" "Exp.4" ...
## $ habitat    : chr  "Deciduous" "Mixed Deciduous" "Mixed Deciduous" "Mixed Deciduous"
## $ dist       : int   155 1155 200 700 1200 1700 160 660 1160 100
## $ indiv1     : int   23 9 12 15 19 12 23 17 16 36
## $ rich1      : int   13 5 7 9 7 7 13 14 10 14
## $ rich1001   : int    5 0 5 5 3 4 3 8 4 5
## $ indiv2     : int   17 9 10 14 9 13 18 23 14 23
## $ rich2      : int    9 4 5 6 4 7 10 9 8 11
## $ rich1002   : int    5 2 3 5 2 4 4 5 4 2
## $ diversity  : num   2.62 1.78 2.17 2.22 1.89 ...

##      point      habitat dist indiv1 rich1 rich1001 indiv2 rich2 rich1002
## 1  Exp.1      Deciduous  155     23    13         5     17     9         5
## 2  EXP.2 Mixed Deciduous 1155      9     5         0      9     4         2
## 3  Exp.3 Mixed Deciduous  200     12     7         5     10     5         3
## 4  Exp.4 Mixed Deciduous  700     15     9         5     14     6         5
## 5  Exp.5 Mixed Deciduous 1200     19     7         3      9     4         2
## 6  Exp.6      Deciduous 1700     12     7         4     13     7         4
## 7  Exp.7 Mixed Deciduous  160     23    13         3     18    10         4
## 8  Exp.8 Mixed Deciduous  660     17    14         8     23     9         5
## 9  Exp.9      Deciduous 1160     16    10         4     14     8         4
## 10 Exp.10     Coniferous  100     36    14         5     23    11         2

##      diversity
## 1      2.62
## 2      1.78
## 3      2.17
## 4      2.22
## 5      1.89
## 6      2.21
## 7      1.71
## 8      2.54
## 9      2.19
## 10     2.42

```

Table 3: Summary of species richness during point count surveys on the proposed Eagle River Flats expansion area .

Survey 1 (May 23)

Survey 2 (June 6)

Combined Surveys

Point

Habitat

Distance to ERF (m)

Indiv.

Sp. Richness

Sp. Richness 100m*

Indiv.

Sp. Richness

Sp. Richness 100m

Shannon Diversity Index

Exp.1

Deciduous

155

23

13

5

17

9

5

2.62

EXP.2

Mixed Deciduous

1155

9

5

0

9

4

2

1.78

Exp.3

Mixed Deciduous

200

12

7

5

10

5

3

2.17

Exp.4

Mixed Deciduous

700

15

9

5

14

6

5

2.22

Exp.5

Mixed Deciduous

1200

19

7

3

9

4

2

1.89

Exp.6

Deciduous

1700

12

7

4

13

7

4

2.21

Exp.7

Mixed Deciduous

160

23

13

3

18

10

4

1.71

Exp.8

Mixed Deciduous

660

17

14

8
 23
 9
 5
 2.54
 Exp.9
 Deciduous
 1160
 16
 10
 4
 14
 8
 4
 2.19
 Exp.10
 Coniferous
 100
 36
 14
 5
 23
 11
 2
 2.42

* Species richness within a 100m diameter of the survey point centroid.

c("Solitary Sandpiper", "Lesser Yellowlegs", "Olive-sided Flycatcher", "Rusty Blackbird", "Blackpoll Warbler") c(15, 27, 10, 31, 0) c(7, 13, 5, 15, 0) c(18, 5, 37, 14, 8, 44, 4, 0) c(3, 36, 18, 16, 2) c(1, 18, 8, 8, 1) c(7, 4, 29, 6, 29, 6, 3, 7)

##	point	habitat	dist	indiv1	rich1	rich1001	indiv2	rich2	rich1002
## 1	Exp.1	Deciduous	155	23	13	5	17	9	5
## 2	EXP.2 Mixed	Deciduous	1155	9	5	0	9	4	2
## 3	Exp.3 Mixed	Deciduous	200	12	7	5	10	5	3

## 4	Exp.4	Mixed Deciduous	700	15	9	5	14	6	5
## 5	Exp.5	Mixed Deciduous	1200	19	7	3	9	4	2
## 6	Exp.6	Deciduous	1700	12	7	4	13	7	4
## 7	Exp.7	Mixed Deciduous	160	23	13	3	18	10	4
## 8	Exp.8	Mixed Deciduous	660	17	14	8	23	9	5
## 9	Exp.9	Deciduous	1160	16	10	4	14	8	4
## 10	Exp.10	Coniferous	100	36	14	5	23	11	2
##	diversity average								
## 1		2.62	11.0						
## 2		1.78	4.5						
## 3		2.17	6.0						
## 4		2.22	7.5						
## 5		1.89	5.5						
## 6		2.21	7.0						
## 7		1.71	11.5						
## 8		2.54	11.5						
## 9		2.19	9.0						
## 10		2.42	12.5						

Figure XX: Point Count Species Richness as a funtion of distance to Eagle I





Figure XX: Point Count Species Richness as a function of habitat classification



Conclusions ###Proposed Next-Steps #####Surveys We propose to continue surveys of SOSC on lowland wetland units during 2021. Additional surveys will (1) support long-term DoD management objectives and (2) compliment the Birds and Bogs programs effort to determine differences in distribution, abundance, and habitat affinities of SOSC in Anchorage and on JBER. We recommend the scheduling of at least two survey periods during future years to accurately detect both early and late arriving species. Actual timing of surveys should be adjusted for spring phenology.

In 2021, we propose to begin monitoring efforts of breeding birds, including Surfbird, in Snowhawk Valley and surrounding alpine ridgelines. An update of the North American Breeding Bird Survey trend results (Sauer et al. 2017; Table 6.1.1.0) for the Northwestern Interior Forest BCR (Bird Conservation Region 4), which encompasses the Anchorage boreal and alpine region, underscores the need to expand monitoring efforts to include non-wetland habitats. For example, wetland-associated SOSC focused on during 2014 - 2020 continue to show long-term declines; however, species associated with montane habitats also show steep, long-term declines. #####Research Lesser Yellowlegs are one of the steepest declining shorebird species in North America (Sauer et al. 2017). Although the magnitude and number of threats during the annual cycle are likely to be numerous, one known risk is sport and subsistence hunting, which occurs on several Caribbean islands and in northeastern South America. Preliminary GPS tracking data indicate that Lesser Yellowlegs breeding in Anchorage do migrate through or stopover in areas practicing shorebird harvest.

In 2021, we will continue to coordinate biological sample collection across the species breeding

range (including JBER) to map genetically distinct populations. We will then compare samples from hunter-killed birds at non-breeding sites to determine whether breeding populations are disproportionately affected. Additionally, we will continue to expand our knowledge of Lesser Yellowlegs breeding behavior and productivity by monitoring chick growth and vital rates. We do not plan to deploy additional PinPoint Argos GPS on Lesser Yellowlegs at JBER in 2021, but will continue to analyze and organize results for publication.

0.8 Products

0.8.1 Data and Reporting

1. GIS project including (1) spatial layers used to design 2016, 2017, 2019, and 2020 surveys of SOSC on JBER, (2) capture locations of SOSC, (3) nest locations of SOSC, and (4) resighting locations of SOSC.
2. Electronic database containing 2016, 2017, 2018, 2019, and 2020 banding data and resighting data.
3. Electronic database containing 2016, 2017, 2019, and 2020 breeding bird survey “clean” data.
4. Field protocol documents for 2016-2020.
5. Access to Shiny App titled Avifaunal Database for Alaska Military Lands
###Follow-on Work
6. Analysis of apparent adult survival of Species of Special Concern.
7. Analysis of Lesser Yellowlegs productivity.
8. Analysis of Lesser Yellowlegs migratory connectivity among breeding and non-breeding sites.
9. Analysis of population-level genetics and contaminants of Lesser Yellowlegs.

0.9 Acknowledgements

We thank Zachary Pohlen, Callie Gesmundo, Hannah Vincelle (USFWS-MBM), Katie Christie, Danielle Eneix, Jeff Wagner, Shelby McCahon, Mitch Paisker, Rachel Gingras, and Courtney Brown (ADF&G), who provided assistance with Lesser Yellowlegs field work. Julie Hagelin (ADF&G), Audrey Taylor (UAA), and Lee Tibbitts (USGS-ASC) provided helpful advice on capturing SOSC. We are especially grateful to Cassandra Schoofs and her predecessor Kristy Craig, who have provided invaluable assistance and support that facilitated all aspects of this project. Funding for this project was provided by 673d CES/CEIEC, U.S. Department of Air Force (project numbers FXSB46058 and FXSB46058119); Alaska Department of Fish and Game; Threatened, Endangered, and Diversity Program; and U.S. Fish and Wildlife Service (WSFR-SWG Grants T-32-1, T-33-2020).

0.10 References

Andres, B. A. 2005. Breeding bird atlas of Fort Richardson, Alaska. Unpublished report, U. S. Fish and Wildlife Service. Migratory Bird Management, Anchorage, Alaska. 85 pp. Boreal Partners in Flight. 2017. In prep. Alaska Landbird Conservation Plan. Anchorage, AK.

Browne, B. T. and B. A. Andres. 2002. LCTA breeding bird survey Fort Richardson, Alaska, 1995-1997. Unpublished report, U.S. Fish and Wildlife Service, Anchorage, AK. 12 pp.

Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, Biological Services Program (FWS/OBS-79/31), Washington, DC.

Department of Defense Partners in Flight Program. 2015. Strategic Plan for Bird Conservation and Management on Department of Defense Lands. <https://www.denix.osd.mil/nr/priorities/birds/factsheets/dod-pif-mission-sensitive-priority-bird-species-october-2015-legacy-14-1717/>

Handel, C.M., J.R. Sauer. 2017. Combined analysis of roadside and off-road breeding bird survey data to assess population change in Alaska. *Condor* 119:557-575.

Hutt, M. B. 1991. Shooting of migrating shorebirds in Barbados. ICBP Technical Publication 12:77-91.

Jorgenson, M. T., J. E. Roth, M. K. Reynolds, M. D. Smith, W. Lentz, and A. L. Zusi-Cobb. 1998. An ecological land classification for Fort Wainwright, Alaska. Final report for U.S. Army CRELL, Hanover, NH.

Alaska Biological Research, Inc., Fairbanks, AK.

Matsuoka, S. M., Shaw, D., and Johnson, J. A. 2010a. Estimating the abundance of nesting Rusty Blackbirds in relation to wetland habitats in Alaska. *Condor* 112:825-833.

Matsuoka, S. M., D. Shaw, P. H. Sinclair, J. A. Johnson, R. M. Corcoran, N. C. Dau, P. M. Myers, and N. A. Rojek. 2010b. Nesting ecology of the Rusty Blackbird in Alaska and Canada. *Condor* 112:810-824.

MWH Americas, Inc. 2015. Species of special concern 2014 survey report. Prepared for Joint Base Elmendorf-Richardson, 673d CES/CEIEC. Unpublished report. 12 pp.

Rappole, J. H. and Tipton, A. R. 1991. New harness design for attachment of radio transmitters to small passerines. *J. Field Orn.* 62:335-337.

Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., Stanton, J. C., Panjabi, A., Helft, L., Parr, M., and Marra, P. P. 2019. Decline of North American avifauna. *Science* 366:120-124.

Rosenberg, K. V., Kennedy, J. A., Dettmers, R., Ford, R. P., Reynolds, D., Alexander, J. D., Beardmore, C. J., Blancher, P. J., Bogart, R. E., Butcher, G. S., Camfield, A. F., Couturier, A., Demarest, D. W., Easton, W. E., Giocomo, J. J., Keller, R. H., Mini, A. E., Panjabi, A. O., Pashley, D. N., Rich, T. D., Ruth, J. M., Stabins, H., Stanton, J., and Will, T. 2016. Partners in Flight Landbird Conservation Plan: 2016 Revision for Canada and Continental United States. Partners in Flight Science Committee. 119 pp.

Sauer, J. R., D. K. Niven, J. E. Hines, D. J. Ziolkowski, Jr, K. L. Pardieck, J. E. Fallon, and W. A. Link. 2017. The North American Breeding Bird Survey, Results and Analysis 1966 - 2015. Version 2.07.2017 USGS Patuxent Wildlife Research Center, Laurel, MD.

U.S. Fish and Wildlife Service. 2008. Birds of Conservation Concern 2008. United States Department of Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Arlington, Virginia. 85 pp. [Online version available at <http://www.fws.gov/migratorybirds/>]

U.S. Shorebird Conservation Plan Partnership. 2016. U.S. shorebirds of conservation concern 2016. <http://www.shorebirdplan.org/science/assessment-conservation-status-shorebirds/>. Assessed September 2017.

Watts, B. D., E. T. Reed, C. Turin. 2015. Estimating sustainable mortality limits for shorebirds using the Western Atlantic Flyway. *Wader Study* 122:37-53.

0.11 Appendix A: Species List

Appendix A. Common name, scientific name, and 4-letter species code for all birds present on Joint Base Elmendorf-Richardson

Taxonomic Group
Common Name
Scientific Name
Species Code
Waterfowl
Snow Goose
Anser caerulescens
SNGO
Greater White-fronted Goose
Anser albifrons
GWFG
Canada Goose
Branta canadensis
CANG
Trumpeter Swan
Cygnus buccinator
TRUS
Blue-winged Teal
Spatula discors
BWTE
Northern Shoveler
Spatula clypeata
NOSH
Gadwall
Mareca strepera
GADW
American Wigeon
Mareca americana
AMWI
Mallard

Anas platyrhynchos

MALL

Northern Pintail

Anas acuta

NOPI

Green-winged Teal

Anas crecca

GWTE

Canvasback

Aythya valisineria

CANV

Redhead

Aythya americana

REDH

Ring-necked Duck

Aythya collaris

RNDU

Greater Scaup

Aythya marila

GRSC

Lesser Scaup

Aythya affinis

LESC

Harlequin Duck

Histrionicus histrionicus

HADU

Surf Scoter

Melanitta perspicillata

SUSC

Bufflehead

Bucephala albeola
BUFF
Common Goldeneye
Bucephala clangula
COGO
Barrow's Goldeneye
Bucephala islandica
BAGO
Common Merganser
Mergus merganser
COME
Grouse & Ptarmigan
Spruce Grouse
Falci pennis canadensis
SPGR
Grebes
Horned Grebe
Podiceps auritus
HOGR
Red-necked Grebe
Podiceps grisegena
RNGR
Cranes
Sandhill Crane
Antigone canadensis
SACR
Plovers
Black-bellied Plover
Pluvialis squatarola
BBPL

American Golden-Plover
Pluvialis dominica
AMGP
Pacific Golden-Plover
Pluvialis fulva
PAGP
Semipalmated Plover
Charadrius semipalmatus
SEPL
Killdeer
Charadrius vociferus
KILL
Sandpipers
Whimbrel
Numenius phaeopus
WHIM
Hudsonian Godwit
Limosa haemastica
HUGO
Surfbird
Calidris virgata
SURF
Least Sandpiper
Calidris minutilla
LESA
Pectoral Sandpiper
Calidris melanotos
PESA
Short-billed Dowitcher
Limnodromus griseus

SBDO

Long-billed Dowitcher

Limnodromus scolopaceus

LBDO

Wilson's Snipe

Gallinago delicata

WISN

Red-necked Phalarope

Phalaropus lobatus

RNPH

Spotted Sandpiper

Actitis macularius

SPSA

Solitary Sandpiper

Tringa solitaria

SOSA

Wandering Tattler

Tringa incana

WATA

Greater Yellowlegs

Tringa melanoleuca

GRYE

Lesser Yellowlegs

Tringa flavipes

LEYE

Gulls & Terns

Bonaparte's Gull

Chroicocephalus philadelphia

BOGU

Mew Gull

Larus canus
MEGU
Herring/Glaucous-winged Gull Hybrid
Larus argentatus/Larus glaucescens
Hybrid
Arctic Tern
Sterna paradisaea
ARTE
Loons
Red-throated Loon
Gavia stellata
RTLO
Pacific Loon
Gavia pacifica
PALO
Common Loon
Gavia immer
COLO
Osprey
Osprey
Pandion haliaetus
OSPR
Hawks & Eagles
Golden Eagle
Aquila chrysaetos
GOEA
Northern Harrier
Circus hudsonius
NOHA
Sharp-shinned Hawk

Accipiter striatus
SSHA
Northern Goshawk
Accipiter gentilis
NOGO
Bald Eagle
Haliaeetus leucocephalus
BAEA
Red-tailed Hawk
Buteo jamaicensis
RTHA
Owls
Great Horned Owl
Bubo virginianus
GHOW
Great Gray Owl
Strix nebulosa
GGOW
Boreal Owl
Aegolius funereus
BOOW
Northern Saw-whet Owl
Aegolius acadicus
NSWO
Kingfishers
Belted Kingfisher
Megascops alcyon
BEKI
Woodpeckers
American Three-toed Woodpecker

Picoides dorsalis
ATTW
Black-backed Woodpecker
Picoides arcticus
BBWO
Downy Woodpecker
Dryobates pubescens
DOWO
Hairy Woodpecker
Dryobates villosus
HAWO
Northern Flicker
Colaptes auratus
NOFL
Falcons
Merlin
Falco columbarius
MERL
Passerines
Olive-sided Flycatcher
Contopus cooperi
OSFL
Western Wood-Pewee
Contopus sordidulus
WEWP
Alder Flycatcher
Empidonax alnorum
ALFL
Northern Shrike
Lanius borealis

NSHR

Canada Jay

Perisoreus canadensis

CAJA

Steller's Jay

Cyanocitta stelleri

STJA

Black-billed Magpie

Pica hudsonia

BBMA

Common Raven

Corvus corax

CORA

Black-capped Chickadee

Poecile atricapillus

BCCH

Boreal Chickadee

Poecile hudsonicus

BOCH

Horned Lark

Eremophila alpestris

HOLA

Tree Swallow

Tachycineta bicolor

TRES

Violet-green Swallow

Tachycineta thalassina

VGSW

Bank Swallow

Riparia riparia

BANS

Golden-crowned Kinglet

Regulus satrapa

GCKI

Ruby-crowned Kinglet

Regulus calendula

RCKI

Red-breasted Nuthatch

Sitta canadensis

RBNU

Brown Creeper

Certhia americana

BRCR

American Dipper

Cinclus mexicanus

AMDI

European Starling

Sturnus vulgaris

EUST

Townsend's Solitaire

Myadestes townsendi

TOSO

Varied Thrush

Ixoreus naevius

VATH

Gray-cheeked Thrush

Catharus minimus

GCTH

Swainson's Thrush

Catharus ustulatus

SWTH

Hermit Thrush

Catharus guttatus

HETH

American Robin

Turdus migratorius

AMRO

Bohemian Waxwing

Bombycilla garrulus

BOWA

American Pipit

Anthus rubescens

AMPI

Pine Grosbeak

Pinicola enucleator

PIGR

Common Redpoll

Acanthis flammea

CORE

White-winged Crossbill

Loxia leucoptera

WWCR

Pine Siskin

Spinus pinus

PISI

Fox Sparrow

Passerella iliaca

FOSP

Dark-eyed Junco

Junco hyemalis

DEJU

White-crowned Sparrow

Zonotrichia leucophrys

WCSP

Golden-crowned Sparrow

Zonotrichia atricapilla

GCSP

Savannah Sparrow

Passerculus sandwichensis

SAVS

Lincoln's Sparrow

Melospiza lincolnii

LISP

Rusty Blackbird

Euphagus carolinus

RUBL

Northern Waterthrush

Parkesia noveboracensis

NOWA

Orange-crowned Warbler

Leiothlypis celata

OCWA

Common Yellowthroat

Geothlypis trichas

COYE

Yellow Warbler

Setophaga petechia

YEWA

Blackpoll Warbler

Setophaga striata

BLPW

Yellow-rumped Warbler

Setophaga coronata

YRWA

Townsend's Warbler

Setophaga townsendi

TOWA

Wilson's Warbler

Cardellina pusilla

WIWA

0.12 Appendix B: Status, number of individuals, and relative occurrence of bird species detected during two breeding season survey periods on lowland wetlands on Joint Base Elmendorf-Richardson, 2020.

##	AMRO	AMWI	ARTE	ATTW	BAEA	BAGO	BCCH	BEKI
##	102	17	2	2	2	2	6	0
##	BOCH	BOGU	BRCR	CAJA	CANG	COGO	COLO	CORA
##	27	33	2	11	16	6	3	0
##	CORE	DEJU	DOWO	FOSP	GADW	GCKI	GCSP	GRSC
##	24	140	2	2	6	41	0	4
##	GRYE	GWGUxHERG	GWTE	HAWO	HETH	LESA	LESC	LEYE
##	9	0	18	2	12	0	4	27
##	LISP	MALL	MEGU	MERL	NOFL	NOGO	NOHA	NOPI
##	199	15	5	0	0	1	0	2
##	NOWA	NSHO	OCWA	OSFL	RBNU	RCKI	RNDU	RNGR
##	30	12	50	10	4	128	75	6
##	RNPH	RUBL	SACR	SAVS	SOSA	SSHA	SWTH	TRES
##	2	31	1	22	15	0	5	26
##	TRUS	VATH	VGSW	WCSP	WEWP	WISN	WIWA	WWCR
##	2	32	2	15	4	158	20	4
##	YRWA							
##	179							
##	AMRO	AMWI	ARTE	ATTW	BAEA	BAGO	BCCH	BEKI
##	43	7	1	1	0	1	3	0
##	BOCH	BOGU	BRCR	CAJA	CANG	COGO	COLO	CORA
##	9	12	1	3	3	3	1	0
##	CORE	DEJU	DOWO	FOSP	GADW	GCKI	GCSP	GRSC

##	2	66	0	1	3	19	0	2
##	GRYE	GWGUxHERG	GWTE	HAWO	HETH	LESA	LESC	LEYE
##	4	0	9	0	6	0	0	11
##	LISP	MALL	MEGU	MERL	NOFL	NOGO	NOHA	NOPI
##	92	5	1	0	0	0	0	0
##	NOWA	NSHO	OCWA	OSFL	RBNU	RCKI	RNDU	RNGR
##	15	6	25	5	2	63	37	3
##	RNPH	RUBL	SACR	SAVS	SOSA	SSHA	SWTH	TRES
##	1	14	0	11	7	0	2	6
##	TRUS	VATH	VGSW	WCSP	WEWP	WISN	WIWA	WWCR
##	1	16	1	7	2	76	10	0
##	YRWA							
##	88							

##	ALFL	AMRO	AMWI	ARTE	ATTW	BAEA	BAGO	BANS	BBMA	BCCH	BEKI
##	0	20	2	1	1	0	0	0	0	0	0
##	BLPW	BOCH	BOGU	CAJA	CANG	COGO	COLO	CORA	CORE	DEJU	FOSP
##	1	2	10	4	1	0	0	0	0	39	0
##	GADW	GCKI	GRYE	GULLsp	GWTE	HAWO	HETH	LESC	LEYE	LISP	MALL
##	0	4	4	0	1	2	2	0	18	79	4
##	MEGU	NOPI	NOWA	NSHO	OCWA	OSFL	PISI	RBNU	RCKI	RNDU	RNGR
##	1	0	22	1	28	9	0	0	39	1	1
##	RNPH	RUBL	SACR	SAVS	SBDO	SOSA	SPSA	SWTH	TOWA	TRES	TRUS
##	0	7	0	12	0	1	0	17	0	5	3
##	VATH	VGSW	WCSP	WEWP	WISN	WIWA	WWCR	YEWA	YRWA		
##	6	0	8	8	65	5	1	6	48		

##	ALFL	AMRO	AMWI	ARTE	ATTW	BAEA	BAGO	BANS	BBMA	BCCH	BEKI
##	0	59	5	6	2	0	0	0	0	0	0
##	BLPW	BOCH	BOGU	CAJA	CANG	COGO	COLO	CORA	CORE	DEJU	FOSP
##	2	13	34	23	9	0	0	0	4	84	0
##	GADW	GCKI	GRYE	GULLsp	GWTE	HAWO	HETH	LESC	LEYE	LISP	MALL
##	0	11	9	2	3	4	5	4	36	164	11
##	MEGU	NOPI	NOWA	NSHO	OCWA	OSFL	PISI	RBNU	RCKI	RNDU	RNGR
##	6	0	44	2	56	18	0	0	80	2	2
##	RNPH	RUBL	SACR	SAVS	SBDO	SOSA	SPSA	SWTH	TOWA	TRES	TRUS
##	1	16	0	24	1	3	1	46	0	21	6
##	VATH	VGSW	WCSP	WEWP	WISN	WIWA	WWCR	YEWA	YRWA		
##	14	0	16	16	142	10	10	12	96		

##	x freq	
## 1	AMRO	18
## 2	AMWI	8
## 3	ARTE	1
## 4	ATTW	1
## 5	BAEA	2

## 6	BAGO	1
## 7	BCCH	3
## 8	BOCH	11
## 9	BOGU	8
## 10	BRCR	1
## 11	CAJA	7
## 12	CANG	5
## 13	COGO	1
## 14	COLO	2
## 15	CORE	5
## 16	DEJU	14
## 17	DOWO	2
## 18	FOSP	1
## 19	GADW	3
## 20	GCKI	10
## 21	GRSC	1
## 22	GRYE	4
## 23	GWTE	4
## 24	HAWO	2
## 25	HETH	1
## 26	LESC	1
## 27	LEYE	10
## 28	LISP	27
## 29	MALL	6
## 30	MEGU	3
## 31	NOGO	1
## 32	NOPI	1
## 33	NOWA	8
## 34	NSHO	2
## 35	OCWA	10
## 36	OSFL	5
## 37	RBNU	2
## 38	RCKI	16
## 39	RNDU	5
## 40	RNGR	3
## 41	RNPH	1
## 42	RUBL	12
## 43	SACR	1
## 44	SAVS	5
## 45	SOSA	5
## 46	SWTH	2
## 47	TRES	10
## 48	TRUS	1
## 49	VATH	5
## 50	VGSW	1

## 51	WCSP	6
## 52	WEWP	2
## 53	WISN	19
## 54	WIWA	8
## 55	WWCR	2
## 56	YRWA	19
##	x freq	
## 1	AMRO	13
## 2	AMWI	2
## 3	ARTE	2
## 4	ATTW	1
## 5	BLPW	1
## 6	BOCH	9
## 7	BOGU	6
## 8	CAJA	9
## 9	CANG	3
## 10	CORE	2
## 11	DEJU	14
## 12	GCKI	4
## 13	GRYE	4
## 14	GULLsp	1
## 15	GWTE	2
## 16	HAWO	2
## 17	HETH	2
## 18	LESC	1
## 19	LEYE	8
## 20	LISP	22
## 21	MALL	4
## 22	MEGU	2
## 23	NOWA	8
## 24	NSHO	1
## 25	OCWA	12
## 26	OSFL	8
## 27	RCKI	14
## 28	RNDU	1
## 29	RNGR	1
## 30	RNPB	1
## 31	RUBL	8
## 32	SAVS	7
## 33	SBDO	1
## 34	SOSA	2
## 35	SPSA	1
## 36	SWTH	12
## 37	TRES	9

## 38	TRUS	3
## 39	VATH	4
## 40	WCSP	5
## 41	WEWP	4
## 42	WISN	21
## 43	WIWA	4
## 44	WWCR	2
## 45	YEWA	1
## 46	YRWA	17

Appendix B. Status, number of individuals, and relative occurrence of bird species detected during two breeding season survey periods on lowland wetlands on Joint Base Elmendorf-Richardson, 2020.

Number of Indiv. (On Unit)

Number of Pairs (On Unit)

Proportion of Units (%)

Species

Status

Survey 1

Survey 2

Survey 1

Survey 2

Survey 1

Survey 2

Canada Goose

Br.

16

9

3

1

18.5

11.1

Trumpeter Swan

Br.

2

6

1

3

3.7

11.1

Northern Shoveler

Br.

12

2

6

1

7.4

3.7

Gadwall

Br.

6

0

3

0

11.1

0.0

American Wigeon

Br.

17

5

7

2

29.6

7.4

Mallard

Br.

15

11

5

4

22.2

14.8

Northern Pintail

Br.

2

0

0

0

3.7

0.0

Green-winged Teal

Br.

18

3

9

1

14.8

7.4

Ring-necked Duck

Br.

75

2

37

1

18.5

3.7

Greater Scaup

Br.

4

0

2

0

3.7

0.0

Lesser Scaup

Br.

4

4

0

0

3.7

3.7

Barrow's Goldeneye

Br.

2

0

1

0

3.7

0.0

Common Goldeneye

Br.

6

0

3

0

3.7

0.0

Red-necked Grebe

Br.

6

2

3

1

11.1

3.7

Sandhill Crane

Br.

1

0

0

0

3.7

0.0

Least Sandpiper

Br.

0

4

0

0

0.0

3.7

Short-billed Dowitcher

Br.

0

1

0

0

0.0

3.7

Wilson's Snipe

Br.

158

142

76

65

70.4

77.8

Spotted Sandpiper

Br.

0

1

0

0

0.0

3.7

Solitary Sandpiper

Br.

15

3

7

1

18.5

7.4

Lesser Yellowlegs

Br.

27

36

11

18

37.0

29.6

Greater Yellowlegs

Br.

9

9

4

4

14.8

14.8

Red-necked Phalarope

Br.

2

1

1

0

3.7

3.7

Bonaparte's Gull

Br.

33

34

12

10

29.6

22.2

Mew Gull

Br.

5

6

1

1

11.1

7.4

Arctic Tern

Br.

2

6

1

1

3.7

7.4

Common Loon

Br.

3

0

1

0

7.4

0.0

Bald Eagle

Res.

2

0

0

0

7.4

0.0

Northern Goshawk

Br.

1

0

0

0

3.7

0.0

Downy Woodpecker

Res.

2

0

0

0

7.4

0.0

Hairy Woodpecker

Res.

2

4

0

2

7.4

7.4

American Three-toed Woodpecker

Res.

2

2

1

1

3.7

3.7

Olive-sided Flycatcher

Br.

10

18

5

9

18.5

29.6

Western Wood-Pewee

Br.

4

16

2

8

7.4

14.8

Alder Flycatcher

Br.

0

1

0

0

0.0

3.7

Canada Jay

Res.

11

23

3

4

25.9

33.3

Tree Swallow

Br.

26

21

6

5

37.0

33.3

Violet-green Swallow

Br.

2

0

1

0

3.7

0.0

Black-capped Chickadee

Res.

6

0

3

0

11.1

0.0

Boreal Chickadee

Res.

27

13

9

2

40.7

33.3

Red-breasted Nuthatch

Res.

4

0

2

0

7.4

0.0

Brown Creeper

Res.

2

0

1

0

3.7

0.0

Golden-crowned Kinglet

Br.

41

11

19

4

37.0

14.8

Ruby-crowned Kinglet

Br.

128

80

0

39

59.3

51.9

Swainson's Thrush

Br.

5

46

2

17

7.4

44.4

Hermit Thrush

Br.

12

5

6

2

3.7

7.4

American Robin

Br.

102

59

43

20

66.7

48.1

Varied Thrush

Br.

32

14

16

6

18.5

14.8

White-winged Crossbill

Res.

4

10

0

1

7.4

7.4

Common Redpoll

Res.

24

4

2

0

18.5

7.4

Savannah Sparrow

Br.

22

24

11

12

18.5

25.9

Fox Sparrow

Br.

2

0

1

0

3.7

0.0

Lincoln's Sparrow

Br.

199

164

92

79

100.0

81.5

White-crowned Sparrow

Br.

15

16

7

8

22.2

18.5

Dark-eyed Junco

Br.

140

84

66

39

51.9

51.9

Rusty Blackbird

Br.

31

16

14

7

44.4

29.6

Northern Waterthrush

Br.

30

44

15

22

29.6

29.6

Orange-crowned Warbler

Br.

50

56

25

28

37.0

44.4

Yellow Warbler

Br.

0

12

0

6

0.0

3.7

Blackpoll Warbler

Br.

0

2

0

1

0.0

3.7

Yellow-rumped Warbler

Br.

179

96

88

48

70.4

63.0

Wilson's Warbler

Br.

20

10

10

5

29.6

14.8

* Survey periods were 15-17 May and 29-31 May. Status is specified as resident (Res.) for taxa that occur year-round and nest on JBER, migrant (Mig.) for species that occur on JBER during migration but do not nest, and breeder (Br.) for species that occur on JBER only as breeders. Names of Species of Special Concern are in bold and focal species are italicized.

0.13 Appendix C: Lesser Yellowlegs Project Summary

The Lesser Yellowlegs is a Neotropical migrant shorebird that breeds across the boreal wetlands of North America from northwestern Alaska to central Quebec. In winter, Lesser Yellowlegs are found in a variety of wetland habitats throughout Central and South America, the Caribbean and the southern United States. Since 1966, the Lesser Yellowlegs population has declined by 5.3% per year across North America (Sauer et al. 2013). In Alaska, the population has declined 5.3% to 9.2% per year since 2003 (Handel and Sauer 2017). On their wintering grounds, Lesser Yellowlegs have incurred an 80% decline since the 1970s (Ottema and Ramcharan 2009).

The factors contributing to the decline are not well understood, however, wetland conversion, agricultural pollution, climate change, and most prominently, unregulated hunting, are all potential threats. As described by Clay et al. (2012), several Caribbean island shooting clubs have constructed wetlands, also known as, “shooting swamps”, to attract shorebirds during fall migration. Shorebird hunting is also prevalent along the northern coast of South America, where tens of thousands of shorebirds, including a high proportion of Lesser Yellowlegs are harvested annually, according to surveys of local hunters (Taylor 2016). The species is considered to be the most widely hunted shorebird species in the Atlantic Flyway with an estimated harvest of nearly 30,000 birds annually on Barbados and Guadeloupe alone (Hutt 1991, Watts et al. 2015, B. Andres, pers. comm.).

In 2018, USFWS Migratory Bird Management and ADF&G Threatened, Endangered and Diversity Program, collaborated with the goal of expanding the current JBER movement study to include populations across the geographical extent of the Lesser Yellowlegs breeding range. The project included sites in Anchorage, Alaska; Yellowknife, Northwest Territories; and James Bay, Ontario in 2018, and continued expansion in 2019, with the addition of Kanuti National Wildlife Refuge, Alaska; Churchill, Manitoba; and Mingan, Quebec (Figure XXX). The movement data collected by the GPS units will provide conservationists an understanding of the proportion and timing of Lesser Yellowlegs traveling through regions

practicing shorebird harvest, and enable the implementation of harvest regulations and site closures for the protection of Lesser Yellowlegs.

In addition to deploying tracking devices at each site, biological samples, including blood, feathers and claws, were collected from all captured individuals. Blood samples will be used to build a “genoscape” of the breeding population, examine potential mercury contamination in the body, and genetically sex each individual. A genoscape tests for genetic structure among geographically distinct breeding populations, that is, sequencing DNA to determine if the Alaska breeding populations are genetically distinct from the Canadian population. Additionally, feather and claw samples collected on the breeding grounds will be used to expand the Lesser Yellowlegs isotopic signature library and build upon a previous study focused on linking harvested birds in South America to their breeding origin.




Figure 15: Lesser Yellowlegs study sites in 2018 and 2019. PinPoint Argos GPS units were deployed and biological samples were collected at each study site.

0.13.1 Lesser Yellowlegs in Anchorage

In 2020, We captured brood-rearing and foraging adult Lesser Yellowlegs from 2 - 21 June. We tagged 12 adults with the sole focus of understanding local movements on JBER, and annual migratory movement patterns including primary stopover and nonbreeding locations. Results from the local movement analysis indicate that Eagle River Flats (ERF) is essential for adult and brood foraging. For example, the majority of adult Lesser Yellowlegs tagged on JBER in 2018, 2019, and 2020 either moved their broods to Eagle River Flats 1 day after hatch, or used the flats as a staging area before autumn migration (Figure XXX). Additionally, one adult banded in 2018 and one in 2019 on the Anchorage Coastal Wildlife Refuge, used ERF and Otter Lake as a staging area prior to migration. Staging adults are typically observed foraging on ERF in late June through early August.

Each PinPoint Argos GPS tag deployed was equipped with a user-defined data collection schedule. The project goals and objectives dictated how tags were programmed. Migration




Figure 16: GPS locations of Lesser Yellowlegs on Joint Base Elmendorf-Richardson by year (top) and individual (bottom).

tags were set to collect autumn and spring locations more frequently than winter locations, while local movement tags were programmed to frequently collect locations prior to autumn migration. We were successful in collecting data in 2018 and 2019 to meet the project objectives; therefore, in 2020, we programmed tags to collect locations less frequently during migration and more frequently during winter to help fill knowledge gaps. Additionally, in 2019 we recaptured two Lesser Yellowlegs carrying PinPoint Argos GPS tags from 2018. We removed and replaced the GPS tag on both individuals. Our goal is to identify potential variation in migration routes between years for a single individual.

0.13.2 Annual Survival Estimates

Since 2016, 182 adult Lesser Yellowlegs were banded in Anchorage, of which, 92 were banded on JBER (Figure XX). Each year crews conduct resighting surveys with the goal of estimating annual adult survival. In 2020, a crew of 12 biologists from USFWS and ADF&G resighted 25 previously banded birds between May and July for a total effort of 456 hours and 18 minutes (Table XX).

Although effort was similar to previous years, fewer birds were resighted at Otter Lake in 2020 and 2019 than any other year. This is likely due to suboptimal foraging and nesting conditions corresponding to increased beaver activity at the lake. Resighting records between years provide vital information on annual survival rates; however, if this normally site-faithful species breeds elsewhere due to a change in environmental conditions, survival rate estimates will be skewed. This was clearly the case for the Lesser Yellowlegs with flag code “AE”. This bird was banded at Otter Lake North in 2016 and was observed nesting at Otter Lake in 2017 and 2018. In 2019 and 2020, the bird was not resighted at Otter Lake; however, while deploying ARU units on the edge of Eagle River Flats in 2020, “AE” was observed foraging in the tidal ponds. This resighting confirmed that Lesser Yellowlegs can live longer than the expected 4 years. Further, a Lesser Yellowlegs banded as an adult on the Anchorage Coastal Wildlife Refuge in 2013, was observed in 2020.

Table 4: Summary of all Lesser Yellowlegs adults banded and resighted on Joint Base Elmendorf-Richardson, 2016-2020 .

Flag Code

Year Banded

Banding Location

Resight Year(s)

AU

2016

McVeigh Marsh

2017

AV



Figure 17: Lesser Yellowlegs banded with alphanumeric flag “TY” on Joint Base Elmendorf-Richardson, 2020.

2016

McVeigh Marsh

2017, 2018

AA

2016

Hillberg Lake

AK

2016

Otter Lake South

2017, 2018

AL

2016

Otter Lake South

2017

AC

2016

Otter Lake North

AE
2016
Otter Lake North
2017, 2018, 2020
AH
2016
Otter Lake North
2018
AJ
2016
Otter Lake North
2018
AX
2016
Otter Lake South
2017, 2018
AY
2016
Clunie South
2018
CA
2016
Clunie South
AM
2017
Ammo Country
AN
2017
Otter Lake North
2018

AP
2017
Otter Lake North
2018
AT
2017
Otter Lake North
2018
CC
2017
Otter Lake North
2018, 2020
EP
2017
Otter Lake South
CH
2017
Otter Lake South
2018, 2020
CJ
2017
Otter Lake South
2018, 2019
EC
2017
Otter Lake North
2018
EE
2017
Otter Lake North

EH
2017
Otter Lake North
2018
EJ
2017
Otter Lake North
2018
EK
2017
Lower Sixmile Lake
2018, 2020
EL
2017
Lower Sixmile Lake
EM
2017
Lower Sixmile Lake
2018, 2019, 2020
EN
2017
Otter Lake North
2018
ET
2017
Beebe Lake
EU
2017
Otter Lake South
2018, 2019

EV
2017
Otter Lake North
CU
2018
Web Pond
CV
2018
Ammo Country
CX
2018
Web Pond
CT
2018
Beebe Lake
HA
2018
Otter Lake North
HC
2018
Otter Lake North
HE
2018
Otter Lake North
HH
2018
Otter Lake North
HJ
2018
Beebe Lake

HL
2018
Otter Lake North
2019
HM
2018
Otter Lake South
HP
2018
Otter Lake North
2019
HT
2018
Otter Lake South
HU
2018
Otter Lake North
HV
2018
Otter Lake North
JJ
2018
Otter Lake North
JK
2018
Otter Lake North
JL
2018
Otter Lake North
JM

2018

Otter Lake South

JN

2018

Web Pond

JU

2018

Lower Sixmile Lake

MA

2019

Ammo Country

MC

2019

Route Bravo

ME

2019

Route Bravo

MJ

2019

Web Pond

ML

2019

Ammo Country

2020

MM

2019

Ammo Country

MP

2019

Ammo Country

2020
MT
2019
Ammo Country
2020
MU
2019
Ammo Country
MV
2019
Otter Lake North
UA
2020
Ammo Country
UC
2020
Dena'ina Road
UE
2020
Dena'ina Road
UH
2020
Ammo Country
UJ
2020
Ammo Country
UK
2020
Ammo Country
UL

2020
Route Bravo
UM
2020
Route Bravo
UP
2020
Web Pond
UN
2020
Otter Lake North
UT
2020
Otter Lake North
UU
2020
Ammo Country
UV
2020
Ammo Country
UX
2020
Half Moon
UY
2020
Half Moon
PX
2020
Web Pond
PY

2020
Fire Tower Road
TA
2020
Fire Tower Road
TC
2020
Eagle River Flats
TE
2020
Eagle River Flats
TH
2020
Eagle River Flats
TJ
2020
Eagle River Flats
TK
2020
Eagle River Flats
TP
2020
Eagle River Flats
TT
2020
Eagle River Flats
TU
2020
Eagle River Flats
TV

2020

Eagle River Flats

TX

2020

Eagle River Flats

TY

2020

Eagle River Flats

XC

2020

Eagle River Flats

0.13.3 Migratory Connectivity

The Lesser Yellowlegs migratory connectivity project consists of 110 adults individually marked with PinPoint Argos GPS (Figure XX). The GPS tags were distributed among the study sites in Alaska and Canada (Figure XX). During 2018 and 2019, several tags failed due to potential predation events, battery failure, or tag communication failure. A summary of Lesser Yellowlegs movements across years will be published as part of a Masters thesis and independent publications.

Preliminary results indicate that Lesser Yellowlegs exhibit moderate migratory connectivity between breeding sites and nonbreeding sites. Birds that breed at the same site spend the nonbreeding period in similar locations, but do not tightly cluster. During autumn migration in 2018 and 2019, birds from the same breeding cohort followed similar migration routes and spent the non-breeding period in similar locations (Figure XX). Additionally, birds from different breeding sites are shown to intermingle at stopover locations during autumn and spring migration, as well as at non-breeding sites. Preliminary results indicate that birds breeding in Alaska migrate using the Central Flyway during autumn migration, and overwinter further south than birds breeding across Canada. Central Canada breeding birds use the Central Flyway to reach stopover areas in Alberta, Saskatchewan, and North Dakota before switching to the Atlantic Flyway, while eastern Canada breeders use the Atlantic Flyway throughout migration. Not all birds breeding in the same location follow the same migratory paths. This is likely due to variation in gender, departure date, food resources, weather patterns, and individual preference.

Preliminary results of autumn migratory movements indicate that birds do not use U.S. military installations as stopover sites along their southbound trajectory (Figure XX). The majority of Lesser Yellowlegs use agricultural fields and ponds during migration, but these types of lands are not present on most active military installations. In 2013, U.S. Department of Agriculture (USDA), DoD and the Department of Interior (DOI) established the “Sentinel



Figure 18: A Lesser Yellowlegs carrying a PinPoint Argos GPS tag on Ile Niapiskau (Mingan Archipeligo), Quebec, 2020. Photo: Y. Rochepault.

Landscapes Partnerships” that promotes the preservation of farms, ranches, forests and open spaces, while allowing the continuation of vital training missions on military installations and surrounding landscapes. Although only seven landscapes in the U.S. have been designated thus far, there is potential for additional designations (sentinellandscapes.org). Protection of landscapes around military installations will certainly benefit Lesser Yellowlegs as this species is shown to stop near military lands across the United States.




Figure 19: Migratory track lines for Lesser Yellowlegs tagged with Lotek PinPoint GPS Argos Satellite tags, 2018-2020.




Figure 20: Fall migratory stopover locations for Lesser Yellowlegs fitted with PinPoint Argos GPS tags. Blue polygons represent U.S. military installations. These were the individuals closest to military installations in 2018-2020.