

Scuola di Scienze  
Dipartimento di Fisica e Astronomia  
Corso di Laurea in Fisica

# DEEP LEARNING ON ABSTRACT SIMPLICIAL COMPLEXES

**Relatore:**  
**Prof.ssa. Rita Fioresi**

**Presentata da:**  
**Tommaso Lamma**

Anno Accademico 2020/2021

# Contents

<b>1 Preliminaries on topology</b>	<b>1</b>
1.1 Simplicial Complexes . . . . .	1
1.2 Abstract Simplicial Complexes . . . . .	5
1.3 Simplicial Homology . . . . .	6
1.4 Simplicial Cohomology . . . . .	8
<b>2 Geometric Deep Learning</b>	<b>9</b>
2.1 Laplacian Operators . . . . .	9
2.2 Graphs . . . . .	11
<b>Conclusion</b>	<b>12</b>
<b>A Category Theory</b>	<b>13</b>
<b>Bibliography</b>	<b>14</b>

# Chapter 1

## Preliminaries on topology

The essential idea in algebraic topology is to convert problems about topological spaces and continuous functions into problems about algebraic objects and their homomorphisms, this way one hopes to end up with an easier problem to solve.

### 1.1 Simplicial Complexes

In this section we shall define structures called simplicial complexes and discuss some of their properties. In order to define these structures we need the definitions of convex hull and affine independence in  $\mathbb{R}^n$ . In this chapter we recall some notions of algebraic topology, such as simplicial complexes and homology. For more details we invite the reader to consult [3], a good reference also for the preliminary necessary notions of topology we are unable to treat here.

**Definition 1.1.** Let  $A \subset \mathbb{R}^n$ , we define  $A$  to be *convex* if

$$x, y \in A \Rightarrow tx + (1 - t)y \in A$$

for all  $t \in [0, 1]$ .

In Figure 1.1 can see in blue an example of a convex set, since we cannot find two points whose linking segment lies in part outside the set. Conversely, since the green set contains two points linked by a segment that partially lies outside the set, we call that set non-convex.




Fig. 1.1: Illustration of a convex (blue) and a non-convex (green) set.

**Definition 1.2.** Let  $\sigma := \{x_i\}_{i \in I}$  be a subset of  $\mathbb{R}^n$ , where  $I$  is a finite set of indexes, we define  $\sigma$  to be *affinely independent* if  $\{x_0 - x_i\}_{i \in I - \{0\}}$  is linearly independent.

We show now that the definition of affine independence of  $\sigma = \{x_i\}_{i \in I} \subset \mathbb{R}^n$  is independent of the choice of  $x_0$ .

**Proposition 1.3.** Let  $\sigma := \{x_i\}_{i \in I}$  be a finite subset of  $\mathbb{R}^n$ , let  $j \in I$  then, if  $\{x_j - x_i\}_{i \in I - \{j\}}$  is linearly independent, also  $\{x_0 - x_i\}_{i \in I - \{0\}}$  is.

*Proof.* If  $j = 0$  the statement is trivially true. Let  $j \neq 0$  and  $\lambda_i \in \mathbb{R}$  for all  $i \neq j$ , then

$$\sum_{i \in I - \{j\}} \lambda_i (x_j - x_i) = 0 \Rightarrow \lambda_i = 0 \quad \forall i \in I - \{j\}.$$

Let then  $\mu_i \in \mathbb{R}$  for all  $i \neq 0$ , and suppose

$$\sum_{i \in I - \{0\}} \mu_i (x_0 - x_i) = (x_0 - x_j) \sum_{i \in I - \{0\}} \mu_i + \sum_{i \in I - \{0\}} \mu_i (x_j - x_i) = 0.$$

If we define  $\mu_0 := -\sum_{i \in I - \{0\}} \mu_i$  we have that

$$0 = \sum_{i \in I} \mu_i (x_j - x_i) = \sum_{i \in I - \{j\}} \mu_i (x_j - x_i) \Rightarrow \mu_i = 0 \quad \forall i \in I - \{j\},$$

which proves our proposition. the definition of affine ind the definition of affine independence is well stated.  $\square$

**Definition 1.4.** Let  $\sigma := \{x_i\}_{i \in I}$  be a finite subset of  $\mathbb{R}^n$ , we define the *convex set generated by  $\sigma$*  to be the smallest convex set containing  $X$  according to the inclusion relation. We shall denote this set by  $[\sigma]$  and call it *convex hull* of  $\sigma$ .

Since the intersection of convex sets is convex, the convex set generated by  $\sigma$  can be equivalently defined as the intersection of all convex sets containing  $\sigma$ .

**Theorem 1.5.** Let  $\sigma := \{x_i\}_{i \in I}$  be a finite subset of  $\mathbb{R}^n$ , if  $\sigma$  is affinely independent then the convex set generated by  $\sigma$  is

$$[\sigma] = \left\{ \sum_{i \in I} \lambda_i x_i : \lambda_i \geq 0, \sum_{i \in I} \lambda_i = 1 \right\}.$$

Furthermore for any point  $x \in [\sigma]$  we have that

$$x = \sum_{i \in I} \lambda_i x_i = \sum_{i \in I} \mu_i x_i \Rightarrow \lambda_i = \mu_i \quad \forall i \in I,$$

where  $\lambda_i, \mu_i \geq 0$  and  $\sum_{i \in I} \lambda_i = \sum_{i \in I} \mu_i = 1$ .

*Proof.* Let  $C := \{\cap_{\alpha} C_{\alpha} : \sigma \subset C_{\alpha}, C_{\alpha} \text{ convex}\}$ , we divide the proof in three steps:

(i)  $C \subset [\sigma]$ .

This is true if  $[\sigma]$  is convex and contains  $\sigma$ . The proof that it contains  $\sigma$  is trivial. In fact for every vertex  $x_j = \sum_{i \in I} \delta_{ij} x_i$ , and  $\sum_{i \in I} \delta_{ij} = 1$ .

To prove that it is convex we chose two points  $a = \sum_{i \in I} a_i x_i, b = \sum_{i \in I} b_i x_i$  where  $a_i, b_i \geq 0 \quad \forall i \in I$  and  $\sum_{i \in I} a_i = \sum_{i \in I} b_i = 1$ . For  $t \in [0, 1]$

$$ta + (1-t)b = t \sum_{i \in I} a_i x_i + (1-t) \sum_{i \in I} b_i x_i = \sum_{i \in I} (ta_i + (1-t)b_i) x_i.$$

Since  $ta_i + (1-t)b_i \geq 0$  and  $\sum_{i \in I} (ta_i + (1-t)b_i) = t \sum_{i \in I} a_i + (1-t) \sum_{i \in I} b_i = 1$  for all  $i \in I$ , our statement is proven.

(ii)  $[\sigma] \subset C$ .

If all but one the  $\lambda_i$  are zero certainly  $\sum_{i \in I} \lambda_i x_i \in C$ , since  $C$  contains all the vertexes. The inductive hypothesis, by relabeling, is that if the first  $\lambda_0, \dots, \lambda_{n-1}$  are non-zero, hence

not even 1, then  $\sum_{i \in I} \lambda_i x_i \in C$ . We want to show that whenever  $\lambda_0, \dots, \lambda_n$  are non-zero then also  $\sum_{i \in I} \lambda_i x_i \in C$ , since  $\lambda_n \neq 1$  we have that

$$\sum_{i \in I} \lambda_i x_i = \sum_{i=0}^n \lambda_i x_i = \lambda_n x_n + \sum_{i=0}^{n-1} \lambda_i x_i = \lambda_n x_n + (1 - \lambda_n) \sum_{i=0}^{n-1} \frac{\lambda_i}{1 - \lambda_n} x_i.$$

Since  $\sum_{i=0}^{n-1} \frac{\lambda_i}{1 - \lambda_n} = 1$ , for the inductive hypothesis  $\sum_{i=0}^{n-1} \frac{\lambda_i}{1 - \lambda_n} x_i \in C$ . Also the vertex  $x_n$  is contained in  $C$  by definition, therefore, being  $C$  convex and  $\lambda_n \in [0, 1]$ , it follows that

$$\lambda_n x_n + (1 - \lambda_n) \sum_{i=0}^{n-1} \frac{\lambda_i}{1 - \lambda_n} x_i \in C.$$

Accordingly  $\sum_{i \in I} \lambda_i x_i \in C$ , by induction we conclude the proof.

(iii)  $\sum_{i \in I} \lambda_i x_i = \sum_{i \in I} \mu_i x_i \Rightarrow \lambda_i = \mu_i \forall i \in I$ .

Let  $\sum_{i \in I} \lambda_i x_i = \sum_{i \in I} \mu_i x_i$ , then also  $x_0 \sum_{i \in I} \lambda_i + \sum_{i \in I} \lambda_i (x_i - x_0) = x_0 \sum_{i \in I} \mu_i + \sum_{i \in I} \mu_i (x_i - x_0)$ , and since both  $\lambda_i$  and  $\mu_i$  are normalised we have that

$$\sum_{i \in I} (\lambda_i - \mu_i)(x_0 - x_i) = \sum_{i \in I - \{0\}} (\lambda_i - \mu_i)(x_0 - x_i) = 0 \Rightarrow \lambda_i = \mu_i \quad \forall i \in I - \{0\},$$

because of the affine independence. □

**Definition 1.6.** We define a  $p$ -simplex  $[\sigma]$  to be the convex hull of an affinely independent set  $\sigma := \{x_i\}_{i \in I} \subset \mathbb{R}^n$ , where  $p = |I| - 1$  is called dimension of the  $p$ -simplex.

Theorem 1.5 gives us the possibility to represent a point in a simplex  $[\sigma]$  via a finite set of real parameters defined in the range  $[0, 1]$  and satisfying the normalisation condition  $\sum_{i \in I} \lambda_i = 1$ . Such parameters are called *baricentric coordinates* of  $[\sigma]$ .

The points in  $\sigma$  are called *vertexes* of the simplex  $[\sigma]$ , accordingly we define the vertex set of a simplex  $[\sigma]$  to be  $Vert([\sigma]) = \sigma$ .

**Definition 1.7.** Let  $[\sigma]$  be a  $p$ -simplex and  $p, t \in \mathbb{N}$ , we say that another  $t$ -simplex  $[\tau]$  is a *face* of  $[\sigma]$  or equivalently that  $[\sigma]$  is a *coface* of  $[\tau]$ , and we write  $[\tau] \leq [\sigma]$ , if  $\tau \subset \sigma$ , where  $t \leq p$ .

Now we are ready for our main definitions.

**Definition 1.8.** We define a *simplicial complex*  $\mathcal{G}$  to be a collection of simplexes such that

- (i) if any simplex  $[\tau] \leq [\sigma]$  and  $[\sigma] \in \mathcal{G}$ , then  $[\tau] \in \mathcal{G}$ ,
- (ii) if  $[\sigma], [\tau] \in \mathcal{G}$ , then  $[\sigma] \cap [\tau] \in \mathcal{G}$ .

Figure 1.2 represents a simplicial complex, while Figure 1.3 represents a collection of simplexes which is not a simplicial complex.




Fig. 1.2: Example of simplicial complex.




Fig. 1.3: Set of simplexes which is not a simplicial complex.

In fact, we can see in Figure 1.3 that the intersection property of simplicial complexes is not satisfied.

## 1.2 Abstract Simplicial Complexes

In this first part we analyzed simplicial complexes as peculiar subsets of  $\mathbb{R}^n$ , which from now on we shall call *geometric simplicial complexes*. Although this approach provides simplicial complexes with the topology inherited from the metric space, it hides the power of simplicial complexes to describe networks and interactions which exist independently of that topology. To make this distinction clear we will treat simplicial complexes as a realization of more abstract objects called *abstract simplicial complexes*. A richer discussion of abstract simplicial complexes can be found in [2] at 3.1 or in [6] at 7.3.

**Definition 1.9.** Let  $\mathcal{V}$  be a finite set, we define an *abstract simplicial complex*  $\mathcal{A}$  to be a family of non empty subsets of  $\mathcal{V}$  such that:

- (i) if  $v \in \mathcal{V}$ , then  $\{v\} \in \mathcal{A}$ ,
- (ii) if  $\sigma \in \mathcal{A}$  and  $\tau \subset \sigma$ , then  $\tau \in \mathcal{A}$ .

We call the member of this family *abstract simplices*.

We call  $\mathcal{V}$  the *vertex set* of  $\mathcal{A}$  and denote it by  $\text{Vert}(\mathcal{A})$ ; since the vertex set is finite, every abstract simplex is finite, therefore we can use the notation  $\sigma = \{v_i\}_{i \in I}$ , to denote a simplex in  $\mathcal{A}$ .

**Definition 1.10.** Let  $\mathcal{A}$  be an abstract simplicial complex and  $\mathcal{G}$  a geometric simplicial complex, if for all  $\{x_i\}_{i \in I} \in \mathcal{A}$  also  $[x_i]_{i \in I} \in \mathcal{G}$  we say that  $\mathcal{G}$  is a *geometric realization* of  $\mathcal{A}$ .

While every geometric simplicial complex can be thought as a geometric realization of an abstract simplicial complex, the existence of a geometric realization for an arbitrary abstract simplicial complex is not trivial at all.

**Theorem 1.11.** *Let  $\mathcal{A}$  be an  $n$ -dimensional abstract simplicial complex, then it admits a geometric realization in  $\mathbb{R}^{2n+1}$ .*

A proof of this theorem can be found in [2] at 3.1.

Both for abstract and geometric simplicial complexes one can define maps called *simplicial maps*. We obtain a category whose equivalences are called isomorphisms. A short discussion of category theory can be found in Appendix A.

In the following sections we shall use abstract simplicial complexes, which can be always thought geometrically in the appropriate  $\mathbb{R}^{2d+1}$ .

## 1.3 Simplicial Homology

An important field in algebraic topology is homology theory. We shall discuss homology theory to the extent that allows us to define the laplacian operator on simplicial complexes, for supplementary readings see [7, 3]. First we want to equip our simplicial complexes with an orientation. So far we have considered the simplex  $\{x_i\}_{i \in I}$  to remain unchanged under arbitrary reorderings of the index set  $I$ , but in the most of the application this is not the case.

**Proposition 1.12.** *Let  $\{x_i\}_{i \in I}$  be a  $p$ -simplex and*

$$\{x_i\}_{i \in I} \sim \{x_i\}_{i \in \pi(I)} \iff \text{sgn}(\pi) = 1,$$

*where  $\pi : I \rightarrow I$  is a permutation of the indexes and  $\text{sgn}(\pi)$  its sign, then  $\sim$  is an equivalence relation.*

**Remark 1.13.** Let  $S_{p+1}$  be the group of permutations of a  $p$ -simplex, and  $\{-1, 1\}$  a multiplicative group, we recall the fact that  $\text{sgn} : S_{p+1} \rightarrow \{-1, 1\}$  is a group homomorphism, that is

$$\text{sgn}(\pi\eta) = \text{sgn}(\pi)\text{sgn}(\eta) \quad \forall \pi, \eta \in S_{p+1}.$$

*Proof.* We divide the proof in three steps:

- (i) Since  $\text{sgn}(\text{id}_I) = 1$  we have that  $\{x_i\}_{i \in I} \sim \{x_i\}_{i \in \text{id}_I(I)} = \{x_i\}_{i \in I}$ .
- (ii) Since  $\text{sgn}(\pi)\text{sgn}(\pi^{-1}) = \text{sgn}(\pi\pi^{-1}) = \text{sgn}(\text{id}_I) = 1$  we have that  $\text{sgn}(\pi) = \text{sgn}(\pi^{-1})$ , therefore  $\{x_i\}_{i \in I} \sim \{x_i\}_{i \in \pi(I)} \iff \{x_i\}_{i \in \pi(I)} \sim \{x_i\}_{i \in I}$ .
- (iii) The transitivity is a consequence of the fact that the product of two even permutations is also even.  $\square$

**Definition 1.14.** We define an *oriented simplex*  $|x_i\rangle_{i \in I}$  to be a simplex  $\{x_i\}_{i \in I}$  together with the choice of one out of the two equivalence classes with respect to  $\sim$ .

**Definition 1.15.** Let  $\mathcal{A}$  be a simplicial complex and let  $G$  denote an arbitrary additive abelian group. We define the *group of  $p$ -chains*

$$C_p(\mathcal{A}, G) := \frac{\{\sum_{i \in I} g_i |x_i\rangle : g_i \in G\}}{\{g_i |x_0, x_1, \dots, x_p\rangle + g_i |x_1, x_0, \dots, x_p\rangle : g_i \in G\}}.$$

Devo pensarci un po' a questo modo di definirlo, ma credo che intenda semplicemente di indicarle due orientazioni opposte mettendo come fattore l'inverso. Cercherò di approfondire i free abelian groups dal Lang ed aggiungere una descrizione più dettagliata.

For the most of the applications the groups  $\mathbb{Z}, \mathbb{R}, \mathbb{Z}_2$  are considered. To keep our notation light we shall write  $C_p$  instead of  $C_p(\mathcal{A}, G)$ .

**Proposition 1.16.**  $C_p(\mathcal{A}, G)$  is an abelian group.

A particularly relevant role in homology theory is played by the *boundary map*. First we define the boundary of an oriented simplex.

**Definition 1.17.** Let  $|\sigma\rangle = |x_0, \dots, x_{p+1}\rangle$  be an oriented  $(p+1)$ -simplex. The boundary  $\partial|\sigma\rangle$  of  $|\sigma\rangle$  is the  $p$ -chain defined by

$$\partial_{p+1}|\sigma\rangle := \sum_{i=0}^{p+1} (-1)^i |x_0, \dots, \hat{x}_i, \dots, x_{p+1}\rangle$$

where the  $\hat{\phantom{x}}$  over a symbol means that symbol is deleted.



*Remark 1.18.* Note that whenever we are able to construct a geometric realization for the oriented simplicial complex, the set  $\bigcup_{i=0}^{p+1} [x_0, \dots, \hat{x}_i, \dots, x_{p+1}]$  is the topological boundary of  $[\sigma]$ . Furthermore we are able to extend the boundary from simplexes to chains.

**Definition 1.19.** We define the *boundary map*  $\partial_{p+1} : C_{p+1} \rightarrow C_p$  to be the group homomorphism defined by

$$\partial_{p+1}(\sum_{i \in I} g_i |\sigma_i\rangle) := \sum_{i \in I} g_i \partial_{p+1} |\sigma_i\rangle.$$

An important property of boundary maps is the following, which we shall denote as *homology lemma*.

**Lemma 1.20.** *The boundary maps satisfy  $\partial_p \circ \partial_{p+1} = 0$ .*

*Proof.* Since the boundary maps are linear it is sufficient to check this on the generators. Let  $\partial_{p+1} |v_0, \dots, v_{p+1}\rangle = \sum_{i=0}^{p+1} (-1)^i |v_0, \dots, \hat{v}_i, \dots, v_{p+1}\rangle$  then

$$(\partial_p \circ \partial_{p+1}) |v_0, \dots, v_{p+1}\rangle = \sum_{j=0, j \neq i}^{p+1} \sum_{i=0}^{p+1} (-1)^{i+j} |v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_{p+1}\rangle = 0. \quad \square$$

The homology lemma is necessary to define the *homology group*.

**Definition 1.21.** We define the *p-homology group* to be

$$H_p := \frac{\ker \partial_p}{\operatorname{im} \partial_{p+1}},$$

where  $\operatorname{im} \partial_{p+1}$  is called the group of simplicial *p-cycles* and  $\ker \partial_p$  is called the group of simplicial *p-boundaries*.

The homology group is therefore, intuitively, the space of cycles that are not boundaries. Without the homology lemma the quotient would not well defined since  $\operatorname{im} \partial_{p+1} \subset \ker \partial_p$  would not be satisfied.

## 1.4 Simplicial Cohomology

In order to define data on simplicial complexes we are interested in studying the dual of the chains  $|\sigma\rangle$ , which we shall call cochains  $\langle\sigma|$ .

**Definition 1.22.** Let  $C_p$  be the group p-chains, we define the group of  $p$ -cochains to be  $C^p := \text{Hom}(C_p, G)$ , i.e. the dual space of  $C_p$ .

As well as with for chains, we have also for cochains a sequence of homeomorphisms called *coboundary maps*. The coboundary maps are defined to be the dual of the boundary maps, hence satisfying a dual version of the homology lemma called *cohomology lemma*.

**Definition 1.23.** The dual of the boundary maps which we shall call *coboundary maps*, is the group homomorphism defined by

$$d_{p+1} : C^p \rightarrow C^{p+1} \quad d_p(\langle\sigma|) := \langle\sigma|\partial_{p+1} \quad \forall \langle\sigma| \in C^p.$$

Therefore  $(d_{p+1}\langle\sigma|)|\tau\rangle = \langle\sigma|\partial_{p+1}|\tau\rangle \quad \forall |\tau\rangle \in C_p$ .

The proof of  $d^p + 1 \circ d_p$  follow directly from the homology lemma. The cohomology lemma allows us to define the *cohomology group*.

**Definition 1.24.** We define the  $p$ -cohomology group to be

$$H^p := \frac{\ker d_{p+1}}{\text{im } d_p},$$

where  $\text{im } d_{p-1}$  is the group of simplicial p-cocycles and  $\ker d_p$  is the group of simplicial p-coboundaries.

To have a specular intuition of what the coboundary operator actually does, we equip our simplicial chain groups with an inner product. This finite dimensional Hilbert space structure shall allow us to represent the cochain  $\langle\sigma|$  evaluated on the chain  $|\tau\rangle$  as the inner product  $\langle\sigma|\tau\rangle$ .

**Definition 1.25.** We define the scalar product  $\langle| \rangle : C_p \times C_p \rightarrow \mathbb{R}$  on the canonical basis of  $C_p$  to be

$$\langle i|j\rangle = \delta_{ij},$$

where  $i, j$  are any two p-simplexes in the canonical basis of  $C_p$  and  $\delta_{ij}$  is the Kronecker Delta.

The same Hilbert space structure allows us to represent the coboundary operator  $d$  acting on a cochain with its cochain representative.

**Definition 1.26.** Let  $d_{p+1} : C^p \rightarrow C^{p+1}$  be the coboundary operator, then for any  $\langle\sigma| \in C^p, \tau \in C_{p+1}$  we can define its dual representation  $\partial_{p+1}^\dagger$  by

$$(d_{p+1}\langle\sigma|)|\tau\rangle = \langle\sigma|\partial_{p+1}|\tau\rangle = \langle\tau|\partial_{p+1}^\dagger|\sigma\rangle \quad \forall |\tau\rangle \in C_{p+1}.$$

We can notice that our definitions lead to the restatement of the equivalent of the generalized Stokes' theorem on simplicial complexes according to the integration previously defined, i.e.  $(d\langle\sigma|)|\tau\rangle = \langle\sigma|\partial|\tau\rangle$ . One can therefore think of the coboundary operator as a discrete exterior derivative acting on cochains. Similarly we name the inner product *integration* due to the analogy previously stated, the choice of a measure of integration is equivalent to a choice of a basis for the chain group.

# Chapter 2

## Geometric Deep Learning

### 2.1 Laplacian Operators

An important role in the definition of a convolution on simplicial complexes is played by the Laplacian operator, especially by its eigenfunctions and spectrum. A more thorough discussion of this operator can be found in [4].

**Definition 2.1.** We define the  $p$ -Laplacian operator to be

$$\Delta_p := \partial_{p+1} \partial_{p+1}^\dagger + \partial_p^\dagger \partial_p =: \Delta_p^+ + \Delta_p^-.$$

The Laplacian operator is defined to be self-adjoint and positive definite.

**Proposition 2.2.** Let  $\Delta_p$  be a laplacian operator, then  $\Delta_p^\dagger = \Delta_p$ .

*Proof.* Let  $|\sigma\rangle, |\tau\rangle \in C_p$

$$\begin{aligned} \langle \sigma | \Delta_p | \tau \rangle &= \langle \sigma | (\partial_{p+1} \partial_{p+1}^\dagger + \partial_p^\dagger \partial_p) | \tau \rangle = \\ &= \langle \tau | (\partial_{p+1} \partial_{p+1}^\dagger + \partial_p^\dagger \partial_p)^\dagger | \sigma \rangle = \\ &= \langle \tau | (\partial_{p+1} \partial_{p+1}^\dagger + \partial_p^\dagger \partial_p) | \sigma \rangle = \langle \sigma | \Delta_p | \tau \rangle. \end{aligned}$$

□

According to the spectral theorem there exists a basis of eigenchains of the Laplacian, and since all  $\Delta_p, \Delta_p^+, \Delta_p^-$  are self-adjoint we can say that they all admit a basis of eigenchains.

**Proposition 2.3.** Let  $\Delta_p |\sigma\rangle = \lambda_\sigma |\sigma\rangle$  then  $\lambda_\sigma \geq 0$ .

*Proof.* Let  $\Delta_p^+ |\sigma\rangle = \lambda_\sigma^+ |\sigma\rangle$ , we see that  $\langle \sigma | \Delta_p^+ | \sigma \rangle = |\partial_{p+1} |\sigma\rangle|^2 \geq 0$ , and since  $\langle \sigma | \Delta_p^+ | \sigma \rangle = \lambda_\sigma^+ \langle \sigma | \sigma \rangle$  we have that  $\lambda_\sigma^+ \geq 0$ .

Let then  $\Delta_p^- |\sigma\rangle = \lambda_\sigma^- |\sigma\rangle$ , we see that  $\langle \sigma | \Delta_p^- | \sigma \rangle = |\partial_p |\sigma\rangle|^2 \geq 0$ , and since  $\langle \sigma | \Delta_p^- | \sigma \rangle = \lambda_\sigma^- \langle \sigma | \sigma \rangle$ , we also have that  $\lambda_\sigma^- \geq 0$ .

Furthermore, since  $\Delta_p^+ \Delta_p^- = \Delta_p^- \Delta_p^+ = 0$  we have that  $[\Delta_p^+, \Delta_p^-] = 0$ , thence  $[\Delta_p, \Delta_p^\pm] = 0$ , therefore  $\Delta_p, \Delta_p^+, \Delta_p^-$  share a basis of eigenchains. Let  $|\sigma\rangle$  be in that common basis then  $\Delta_p |\sigma\rangle = \lambda_\sigma |\sigma\rangle$ , where  $\lambda_\sigma = \lambda_\sigma^+ + \lambda_\sigma^- \geq 0$ . □

Another really interesting property that was first proven by Beno Eckmann in 1944, is that the kernel of the  $p$ -Laplacian is isomorphic to the  $p$ -homology group.

**Theorem 2.4.** Let  $\Delta_p$  be a laplacian operator, then  $\ker \Delta_p \simeq H_p$ .

*Proof.* We have

$$\Delta_p =: \partial_{p+1} \partial_{p+1}^\dagger + \partial_p^\dagger \partial_p =: \Delta_p^+ + \Delta_p^-.$$

Because of the homology lemma  $\Delta_p^+ \Delta_{-p} = \Delta_p^- \Delta_p^+ = 0$ , therefore  $\ker \Delta_p^\pm \subset \operatorname{im} \Delta_p^\mp$ .

It is trivial to see that, since  $\ker \Delta_p = \ker \Delta_p^+ \cap \ker \Delta_p^-$ , we have that  $\ker \partial_{p+1}^\dagger \cap \ker \partial_p \subset \ker \Delta_p$ .

Less trivial is the opposite inclusion

$$\partial_{p+1}^\dagger \sigma \in \operatorname{im} \partial_{p+1}^\dagger, \sigma \in \ker \Delta_p \implies \partial_{p+1}^\dagger \sigma \in \ker \partial_{p+1} = (\operatorname{im} \partial_{p+1}^\dagger)^\perp,$$

$$\partial_p \sigma \in \operatorname{im} \partial_p, \sigma \in \ker \Delta_p \implies \partial_p \sigma \in \ker \partial_p^\dagger = (\operatorname{im} \partial_p)^\perp,$$

$$\partial_p \sigma = \partial_{p+1}^\dagger \sigma \in \operatorname{im} \partial_{p+1}^\dagger \cap (\operatorname{im} \partial_{p+1}^\dagger)^\perp = \operatorname{im} \partial_p \cap (\operatorname{im} \partial_p)^\perp = 0,$$

therefore  $\ker \Delta_p \subset \ker \partial_{p+1}^\dagger \cap \ker \partial_p$ . Finally we notice that

$$\ker \Delta_p = \ker \partial_{p+1}^\dagger \cap \ker \partial_p = (\operatorname{im} \partial_{p+1})^\perp \cap \ker \partial_p \simeq H_p. \quad \square$$

## 2.2 Graphs

In questa sezione metterei la descrizione del grafo come fa bronstein con i due prodotti scalari e il laplaciano. I pesi alla fine sono solo una scelta della base del gruppo(in  $\mathbb{R}$  spazio) delle p-catene. Inoltre ne approfitterei per fare qui i disegni e gli esempi relativi a bordo e cobordo

# Conclusion

Most of the deep learning techniques used today are based on models which learn a partition of the set of smooth functions defined on euclidean domains into human friendly equivalence classes. Although this approach has been successful in modern machine learning, it only deals with a really small set of domains. The goal of geometric deep learning is to extend this method to data defined on manifolds and simplicial complexes.

Convolution on euclidean domains is itself based on the translation invariance of such domains. In fact the convolution of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , with some filter  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is

$$(f * g)(x) = \langle f, g \circ T^{-1} \rangle_{L^2},$$

where  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a translation represented by the vector  $x$ . Moreover, one could also consider such a convolution to be defined on the translation group itself, represented by some  $\mathbb{R}^n$ . In the same way one can define a convolution on the group  $(\mathbb{Z}, +)$  as  $(a * b)_n = \sum_{i \in \mathbb{Z}} a_i b_{n-i}$ . Similarly an interesting example is that of images, which are samples ad grids, any image can be thought as a function defined on the group  $(\mathbb{Z} \times \mathbb{Z}, +)$ . Such definitions of convolution operators are equivariant with respect to the action of the group they are defined upon. In image recognition translation equivariance is necessary, nevertheless the most common CNN's need to learn the rotations of the same filter as different filters in order to become rotation equivariant. Although manifolds and simplicial complexes are not in general groups, G-equivariant CNN's (see [1]) could be the key to reveal the secrets behind the success of such architectures.

# Appendix A

## Category Theory

**Definition A.1.** A category  $\mathbf{C}$  consists of three ingredients:

1. a class of *objects*  $Obj(\mathbf{C})$ ,
2. sets of *morphisms*  $Hom(A, B)$  for every ordered pair  $(A, B) \in Obj(\mathbf{C}) \times Obj(\mathbf{C})$ ,
3. a composition  $Hom(A, B) \times Hom(B, C) \rightarrow Hom(A, C)$ , denoted by  $(f, g) \mapsto f \circ g$  for every  $A, B, C \in Obj(\mathbf{C})$ , satisfying the following axioms:
  - (i) the family of  $Hom(A, B)$  is pairwise disjoint,
  - (ii) the composition, when defined, is associative,
  - (iii) for each  $A \in Obj(\mathbf{C})$  there exists an *identity*  $1_A \in Hom(A, A)$  such that for  $f \in Hom(A, B)$  and  $g \in Hom(C, A)$  we have that  $1_A \circ f = f$  and  $g \circ 1_A = g$ .

Instead of writing  $f \in Hom(A, B)$ , we usually write  $f : A \rightarrow B$ .

**Definition A.2.** Let  $\mathbf{A}$  and  $\mathbf{C}$  be categories, a *functor*  $T : \mathbf{A} \rightarrow \mathbf{C}$  is a function, that is,

- (i) for each  $A \in Obj(\mathbf{A})$  it assigns  $TA \in Obj(\mathbf{C})$ ,
- (ii) for each morphism  $f : A \rightarrow A'$  it assigns a morphism  $Tf : TA \rightarrow TA' \quad \forall A, A' \in Obj(\mathbf{A})$ ,
- (iii) if  $f, g$  are morphisms in  $\mathbf{A}$  for which  $g \circ f$  is defined, then  $T(g \circ f) = (Tg) \circ (Tf)$ ,
- (iv)  $T(1_A) = 1_{TA} \quad \forall A \in \mathbf{A}$ .

The property (iii) of the previous definition actually defines what we shall call *covariant functors*. If instead we require  $T(g \circ f) = (Tf) \circ (Tg)$ , we are defining a *contravariant functor*.

**Definition A.3.** An *equivalence* in a category  $\mathbf{C}$  is a morphism  $f : A \rightarrow B$  for all  $A, B \in Obj(\mathbf{C})$  for which there exists a morphism  $g : B \rightarrow A$  such that  $f \circ g = 1_B$  and  $g \circ f = 1_A$ .

**Theorem A.4.** If  $\mathbf{A}$  and  $\mathbf{C}$  are categories and  $T : \mathbf{A} \rightarrow \mathbf{C}$  is a functor of either variance, then whenever  $f$  is an equivalence on  $\mathbf{A}$  then  $Tf$  is an equivalence on  $\mathbf{C}$ .

*Proof.* We apply  $T$  to the equations  $f \circ g = 1_B$  and  $g \circ f = 1_A$ , that for a covariant functor leads to  $(Tf) \circ (Tg) = T(1_B) = 1_{TB}$  and  $(Tg) \circ (Tf) = T(1_A) = 1_{TA}$ .  $\square$

A category that will be used in the following section is the category of topological spaces and continuous functions.

**Proposition A.5.** Topological spaces and continuous functions are a category **Top**, whose equivalences are called homeomorphisms.

Other examples of categories can be found in [6] at 0.3 and in [5].

# Bibliography

- [1] Taco S. Cohen and Max Welling. *Group Equivariant Convolutional Networks*. 2016.
- [2] Herbert Edelsbrunner and John Harer. *Computational Topology - an Introduction*. American Mathematical Society, 2010.
- [3] Allen Hatcher. *Algebraic Topology*. Cambridge Univ. Press, 2000.
- [4] Danijela Horak and Juergen Jost. *Spectra of combinatorial Laplace operators on simplicial complexes*. 2011.
- [5] Saunders MacLane. *Categories for the Working Mathematician*. 1971.
- [6] Joseph J. Rotman. *An Introduction to Algebraic Topology*. Graduate Texts in Mathematics. Springer-Verlag New York, 1988.
- [7] I. M. Singer and J. A. Thorpe. *Lecture Notes on Elementary Topology and Geometry*. Springer, New York, NY, 1976.