# Inexact SARAH Algorithm for Stochastic Optimization

Lam M. Nguyen Katya

Katya Scheinberg

Martin Takáč

November 27, 2018

#### **Abstract**

We develop and analyze a variant of variance reducing stochastic gradient algorithm, known as SARAH [10], which does not require computation of the exact gradient. Thus this new method can be applied to general expectation minimization problems rather than only finite sum problems. While the original SARAH algorithm, as well as its predecessor, SVRG [2], require an exact gradient computation on each outer iteration, the inexact variant of SARAH (iSARAH), which we develop here, requires only stochastic gradient computed on a mini-batch of sufficient size. The proposed method combines variance reduction via sample size selection and iterative stochastic gradient updates. We analyze the convergence rate of the algorithms for strongly convex, convex, and nonconvex cases with appropriate mini-batch size selected for each case. We show that with an additional, reasonable, assumption iS-ARAH achieves the best known complexity among stochastic methods in the case of general convex case stochastic value functions.

## 1 Introduction

We consider the problem of stochastic optimization

$$\min_{w \in \mathbb{R}^d} \left\{ F(w) = \mathbb{E}[f(w; \xi)] \right\},\tag{1}$$

where  $\xi$  is a random variable. One of the most popular applications of this problem is expected risk minimization in supervised learning. In this case, random variable  $\xi$  represents a random data sample (x,y), or a set of such samples  $\{(x_i,y_i)\}_{i\in I}$ . We can consider a set of realization  $\{\xi_{[i]}\}_{i=1}^n$  of  $\xi$  corresponding to a set of random samples  $\{(x_i,y_i)\}_{i=1}^n$ , and define  $f_i(w):=f(w;\xi_{[i]})$ . Then the sample average approximation of F(w), known as empirical risk in supervised learning, is written as

$$\min_{w \in \mathbb{R}^d} \left\{ F(w) = \frac{1}{n} \sum_{i=1}^n f_i(w) \right\}. \tag{2}$$

Lam M. Nguyen, IBM Thomas J. Watson Research Center, NY, USA. Email: LamNguyen.MLTD@ibm.com

**Katya Scheinberg**, Industrial and Systems Engineering, Lehigh University, PA, USA. Email: katyas@lehigh.edu. The research of this author was partially supported by NSF Grants CCF 16-18717 and CCF 17-40796

Martin Takáč, Industrial and Systems Engineering, Lehigh University, PA, USA. Email: Takac.MT@gmail.com. The research of this author was partially supported by NSF Grants CCF 16-18717, CMMI 16-63256 and CCF 17-40796.

Throughout the paper, we assume the existence of unbiased gradient estimator, that is  $\mathbb{E}[\nabla f(w;\xi)] = \nabla F(w)$  for any fixed  $w \in \mathbb{R}^d$ . In addition we assume that there exists a lower bound of function F.

In recent years, a class of variance reduction methods [17, 5, 1, 2, 4, 10] has been proposed for problem (2) which have smaller computational complexity than both, the full gradient descent method and the stochastic gradient method. All these methods rely on the finite sum form of (2) and are, thus, not readily extendable to (1). In particular, SVRG [2] and SARAH [10] are two similar methods that consist of an outer loop, which includes one exact gradient computation at each outer iteration and an inner loop with multiple iterative stochastic gradient updates. The only difference between SVRG and SARAH is how the iterative updates are performed in the inner loop. The advantage of SARAH is that the inner loop itself results in a convergent stochastic gradient algorithm. Hence, it is possible to apply only one-loop of SARAH with sufficiently large number of steps to obtain an approximately optimal solution (in expectation). The convergence behavior of one-loop SARAH is similar to that of the standard stochastic gradient method [10]. The multiple-loop SARAH algorithm matches convergence rates of SVRG, however, due to its convergent inner loop, it has an additional practical advantage of being able to use an adaptive inner loop size (see [10] for details).

A version of SVRG algorithm, SCSG, has been recently proposed and analyzed in [6, 7]. While this method has been developed for (2) it can be directly applied to (1) because the exact gradient computation is replaced with a mini-batch stochastic gradient. The size of the inner loop of SCSG is then set to a geometrically distributed random variable with distribution dependent on the size of the mini-batch used in the outer iteration. In this paper, we propose and analyze an inexact version of SARAH (iSARAH) which can be applied to solve (1). Instead of exact gradient computation, a mini-batch gradient is computed using a sufficiently large sample size. We develop total sample complexity analysis for this method under various convexity assumptions on F(w). These complexity results are summarized in Tables 1-3 and are compared to the result for SCSG from [6, 7] when applied to (1). We also list the complexity bounds for SVRG, SARAH and SCSG when applied to finite sum problem (2).

As SVRG, SCSG and SARAH, iSARAH algorithm consists of the *outer loop*, which performs variance reduction by computing sufficiently accurate gradient estimate, and the *inner loop*, which performs the stochastic gradient updates. The main difference between SARAH and SVRG is that the inner loop of SARAH by itself is a convergent stochastic gradient algorithm, while the inner loop of SVRG is not. In other words, if only one outer iteration of SARAH is performed and then followed by sufficiently many inner iterations, we refer to this algorithm as one-loop SARAH. In [10] one-loop SARAH is analyzed and shown to match the complexity of stochastic gradient descent. Here along with multiple-loop iSARAH we analyze one-loop iSARAH, which is obtained from one-loop SARAH by replacing the first full gradient computation with a stochastic gradient based on a sufficiently large mini-batch.

We summarize our complexity results and compare them to those of other methods in Tables 1-3. All our complexity results present the bound on the number of iterations it takes to achieve  $\mathbb{E}[\|\nabla F(w)\|^2] \leq \epsilon$ . These complexity results are developed under the assumption that  $f(w,\xi)$  is L-smooth for every realization of the random variable  $\xi$ . Table 1 shows the complexity results in the case when F(w), but not necessarily every realization  $f(w,\xi)$ , is  $\mu$ -strongly convex, with  $\kappa = L/\mu$  denoting the condition number. Notice that results for one-loop iSARAH (and SARAH) are the same from strongly convex case as for simple convex case. The convergence rate of the multiple-loop iSARAH, on the other hand, is better, in terms of dependence on  $\kappa$ , than the rate achieved by SCSG, which is the only other variance reduction method of the type we consider here, that applies to (1). The general convex case is summarized in Table 2. In this case, the multiple-loop iSARAH achieves the best convergence rate with respect to  $\epsilon$  among the compared stochastic methods, but this rate is derived under an additional assumption (Assumption 4) which is discussed in detail

in Section 3. We note here that Assumption 4 is weaker than the, common in the analysis of stochastic algorithms, assumption that the iterates remain in a bounded set. In recent paper [14] a stochastic line search method is shown to have expected iteration complexity of  $\mathcal{O}\left(\frac{1}{\epsilon}\right)$  under the assumption that the iterates remain in a bounded set, however, no total sample complexity is derived in [14].

Finally, the results for nonconvex problems are presented in Table 3. In this case, SCSG achieves the best convergence rate under the bounded variance assumption, which requires that  $\mathbb{E}[\|\nabla f(w;\xi) - \nabla F(w)\|^2] \le C$ , for some C>0 and  $\forall w\in\mathbb{R}^d$ . While convergence rate of multiple-loop iSARAH for nonconvex function remains an open question (as it is for the original SARAH algorithm), we derive convergence rate for one-loop iSARAH without the bounded variance assumption. This convergence rate matches that of the general stochastic gradient algorithm, since the one-loop iSARAH method is not a variance reduction method. The one-loop iSARAH method can be viewed as a variant of a momentum SGD method.

Table 1: Comparison results (Strongly convex)

Method	Bound	Problem type
<b>SARAH (one-loop)</b> [10, 13]	$\mathcal{O}\left(n + \frac{1}{\epsilon^2}\right)$	Finite-sum
SARAH (multiple-loop) [10]	$\mathcal{O}\left((n+\kappa)\log\left(\frac{1}{\epsilon}\right)\right)$	Finite-sum
SVRG [2, 16]	$\mathcal{O}\left((n+\kappa)\log\left(\frac{1}{\epsilon}\right)\right)$	Finite-sum
SCSG [6, 7]	$\mathcal{O}\left(\left(\min\left\{\frac{\kappa}{\epsilon},n\right\}+\kappa\right)\log\left(\frac{1}{\epsilon}\right)\right)$	Finite-sum
SCSG	$\mathcal{O}\left(\left(\frac{\kappa}{\epsilon} + \kappa\right) \log\left(\frac{1}{\epsilon}\right)\right)$	Expectation
SGD [12]	$O\left(\frac{1}{\epsilon}\right)$	Expectation
iSARAH (one-loop)	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation
iSARAH (multiple-loop)	$\mathcal{O}\left(\max\left\{\frac{1}{\epsilon},\kappa\right\}\log\left(\frac{1}{\epsilon}\right)\right)$	Expectation

Table 2: Comparison results (General convex)

Method	Bound	Problem type	Additional assumption
<b>SARAH (one-loop)</b> [10, 13]	$\mathcal{O}\left(n + \frac{1}{\epsilon^2}\right)$	Finite-sum	None
SARAH (multiple-loop) [10]	$\mathcal{O}\left(\left(n+\frac{1}{\epsilon}\right)\log\left(\frac{1}{\epsilon}\right)\right)$	Finite-sum	Assumptions 4
SVRG [2, 16]	$\mathcal{O}\left(n + \frac{\sqrt{n}}{\epsilon}\right)$	Finite-sum	None
SCSG [6, 7]	$\mathcal{O}\left(\min\left\{\frac{1}{\epsilon^2}, \frac{n}{\epsilon}\right\}\right)$	Finite-sum	None
SCSG	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation	None
SGD	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation	Bounded variance
iSARAH (one-loop)	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation	None
iSARAH (multiple-loop)	$\mathcal{O}\left(\frac{1}{\epsilon}\log\left(\frac{1}{\epsilon}\right)\right)$	Expectation	Assumption 4

We summarize the key results we obtained in this paper as follows.

• We develop and analyze a stochastic variance reduction method for solving general stochastic optimization problems (1) which is an inexact version of SARAH algorithm [10].

Table 3: Comparison results (Nonconvex)

Method	Bound	Problem type	Additional assumption
<b>SARAH</b> (one-loop) [10, 13]	$\mathcal{O}\left(n + \frac{1}{\epsilon^2}\right)$	Finite-sum	None
SVRG [2, 16]	$\mathcal{O}\left(n + \frac{n^{2/3}}{\epsilon}\right)$	Finite-sum	None
SCSG [6, 7]	$\mathcal{O}\left(\min\left\{\frac{1}{\epsilon^{5/3}}, \frac{n^{2/3}}{\epsilon}\right\}\right)$	Finite-sum	Bounded variance
SCSG	$\mathcal{O}\left(\frac{1}{\epsilon^{5/3}}\right)$	Expectation	Bounded variance
SGD	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation	Bounded variance
iSARAH (one-loop)	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$	Expectation	None

- To the best of our knowledge, the proposed algorithm is the best in terms of sample complexity for convex problems than any other algorithm for (1), under an additional relatively mild assumption.
- The one-loop iSARAH presents a version of momentum SGD method. We analyze convergence rates of this algorithm in the general convex and nonconvex cases without assuming boundedness of the variance of the stochastic gradients.

#### 1.1 Paper Organization

The remainder of the paper is organized as follows. In Section 2, we describe Inexact SARAH (iSARAH) algorithm in detail. We provide the convergence analysis of iSARAH in Section 3 which includes the sample complexity bounds for the one-loop case when applied to strongly convex, convex, and nonconvex functions; and for the multiple-loop case when applied to strongly convex and convex functions. We conclude the paper and discuss future work in Section 4.

# 2 The Algorithm

Like SVRG and SARAH, iSARAH consists of the outer loop and the inner loop. The inner loop performs recursive stochastic gradient updates, while the outer loop of SVRG and SARAH compute the exact gradient. Specifically, given an iterate  $w_0$  at the beginning of each outer loop, SVRG and SARAH compute  $v_0 = \nabla F(w_0)$ . The only difference between SARAH and iSARAH is that the latter replaces the exact gradient computation by a gradient estimate based on a sample set of size b.

In other words, given a iterate  $w_0$  and a sample set size b,  $v_0$  is a random vector computed as

$$v_0 = \frac{1}{b} \sum_{i=1}^{b} \nabla f(w_0; \zeta_i),$$
 (3)

where  $\{\zeta_i\}_{i=1}^b$  are i.i.d.<sup>1</sup> and  $\mathbb{E}[\nabla f(w_0; \zeta_i)|w_0] = \nabla F(w_0)$ . We have  $\mathbb{E}[v_0|w_0] = \frac{1}{b}\sum_{i=1}^b \nabla F(w_0) = \nabla F(w_0)$ . The larger b is, the more accurately the gradient estimate  $v_0$  approximates  $\nabla F(w_0)$ . The key idea of the analysis of iSARAH is to establish bounds on b which ensure sufficient accuracy for recovering original SARAH convergence rate.

The key step of the algorithm is a recursive update of the stochastic gradient estimate (SARAH update)

$$v_t = \nabla f(w_t; \xi_t) - \nabla f(w_{t-1}; \xi_t) + v_{t-1}, \tag{4}$$

followed by the iterate update

$$w_{t+1} = w_t - \eta v_t. \tag{5}$$

Let  $\mathcal{F}_t = \sigma(w_0, w_1, \dots, w_t)$  be the  $\sigma$ -algebra generated by  $w_0, w_1, \dots, w_t$ . We note that  $\xi_t$  is independent of  $\mathcal{F}_t$  and that  $v_t$  is a *biased estimator* of the gradient  $\nabla F(w_t)$ .

$$\mathbb{E}[v_t|\mathcal{F}_t] = \nabla F(w_t) - \nabla F(w_{t-1}) + v_{t-1}.$$

<sup>&</sup>lt;sup>1</sup>Independent and identically distributed random variables. We note from probability theory that if  $\zeta_1, \ldots, \zeta_b$  are i.i.d. random variables then  $g(\zeta_1), \ldots, g(\zeta_b)$  are also i.i.d. random variables for any measurable function g.

In contrast, SVRG update is given by

$$v_t = \nabla f(w_t; \xi_t) - \nabla f(w_0; \xi_t) + v_0 \tag{6}$$

which implies that  $v_t$  is an unbiased estimator of  $\nabla F(w_t)$ .

The outer loop of iSARAH algorithm is summarized in Algorithm 1 and the inner loop is summarized in Algorithm 2.

#### Algorithm 1 Inexact SARAH (iSARAH)

```
Parameters: the learning rate \eta>0 and the inner loop size m, the sample set size b. Initialize: \tilde{w}_0. Iterate: for s=1,2,\ldots,\mathcal{T}, do \tilde{w}_s=\mathrm{iSARAH\text{-}IN}(\tilde{w}_{s-1},\eta,m,b). end for Output: \tilde{w}_{\mathcal{T}}.
```

# **Algorithm 2** iSARAH-IN $(w_0, \eta, m, b)$

```
Input: w_0(=\tilde{w}_{s-1}) the learning rate \eta>0, the inner loop size m, the sample set size b. Generate random variables \left\{\zeta_i\right\}_{i=1}^b i.i.d. Compute v_0=\frac{1}{b}\sum_{i=1}^b\nabla f(w_0;\zeta_i). w_1=w_0-\eta v_0. Iterate: for t=1,\ldots,m-1, do Generate a random variable \xi_t v_t=\nabla f(w_t;\xi_t)-\nabla f(w_{t-1};\xi_t)+v_{t-1}. w_{t+1}=w_t-\eta v_t. end for Set \tilde{w}=w_t with t chosen uniformly at random from \{0,1,\ldots,m\} Output: \tilde{w}
```

As a variant of SARAH, iSARAH inherits the special property that a one-loop iSARAH, which is the variant of Algorithm 1 with  $\mathcal{T}=1$  and  $m\to\infty$  is a convergent algorithm. In the next section we provide the analysis for both one-loop and multiple-loop versions of i-SARAH.

Convergence criteria. Our iteration complexity analysis aims to bound the total number of stochastic gradient evaluations needed to achieve a desired bound on the gradient norm. For that we will need to bound the number of outer iterations  $\mathcal{T}$  which is needed to guarantee that  $\|\nabla F(w_{\mathcal{T}})\|^2 \leq \epsilon$  and also to bound m and b. Since the algorithm is stochastic and  $w_{\mathcal{T}}$  is random the  $\epsilon$ -accurate solution is only achieved in expectation. i.e.,

$$\mathbb{E}[\|\nabla F(w_{\mathcal{T}})\|^2] \le \epsilon. \tag{7}$$

# 3 Convergence Analysis of iSARAH

## 3.1 Basic Assumptions

The analysis of the proposed algorithm will be performed under appropriate subset of the following key assumptions.

**Assumption 1** (*L*-smooth).  $f(w; \xi)$  is *L*-smooth for every realization of  $\xi$ , i.e., there exists a constant L > 0 such that

$$\|\nabla f(w;\xi) - \nabla f(w';\xi)\| \le L\|w - w'\|, \ \forall w, w' \in \mathbb{R}^d.$$
 (8)

Note that this assumption implies that  $F(w) = \mathbb{E}[f(w;\xi)]$  is also *L-smooth*. The following strong convexity assumption will be made for the appropriate parts of the analysis, otherwise, it would be dropped.

**Assumption 2** ( $\mu$ -strongly convex). The function  $F: \mathbb{R}^d \to \mathbb{R}$ , is  $\mu$ -strongly convex, i.e., there exists a constant  $\mu > 0$  such that  $\forall w, w' \in \mathbb{R}^d$ ,

$$F(w) \ge F(w') + \nabla F(w')^{\top} (w - w') + \frac{\mu}{2} ||w - w'||^2.$$

Under Assumption 2, let us define the (unique) optimal solution of (2) as  $w_*$ , Then strong convexity of F implies that

$$2\mu[F(w) - F(w_*)] \le \|\nabla F(w)\|^2, \ \forall w \in \mathbb{R}^d.$$
(9)

Under strong convexity assumption we will use  $\kappa$  to denote the condition number  $\kappa \stackrel{\text{def}}{=} L/\mu$ .

Finally, as a special case of the strong convexity with  $\mu=0$ , we state the general convexity assumption, which we will use for some of the convergence results.

**Assumption 3.**  $f(w; \xi)$  is convex for every realization of  $\xi$ , i.e.,  $\forall w, w' \in \mathbb{R}^d$ ,

$$f(w;\xi) \ge f(w';\xi) + \nabla f(w';\xi)^{\top} (w - w').$$

We note that Assumption 2 does not imply Assumption 3, because the latter applies to all realizations, while the former applied only to the expectation.

Hence in our analysis, depending on the result we aim at, we will require Assumption 3 to hold by itself, or Assumption 2 and Assumption 3 to hold together. We will always use Assumption 1.

## 3.2 Existing Results

We provide some well-known results from the existing literature that support our theoretical analysis as follows. First, we start introducing two standard lemmas in smooth convex optimization ([8]) for a general function f.

**Lemma 1** (Theorem 2.1.5 in [8]). Suppose that f is convex and L-smooth. Then, for any  $w, w' \in \mathbb{R}^d$ ,

$$f(w) \le f(w') + \nabla f(w')^T (w - w') + \frac{L}{2} ||w - w'||^2, \tag{10}$$

$$f(w) \ge f(w') + \nabla f(w')^{\top} (w - w') + \frac{1}{2L} \|\nabla f(w) - \nabla f(w')\|^2, \tag{11}$$

$$(\nabla f(w) - \nabla f(w'))^{\top}(w - w') \ge \frac{1}{L} \|\nabla f(w) - \nabla f(w')\|^2.$$
(12)

Note that (10) does not require the convexity of f.

**Lemma 2** (Theorem 2.1.11 in [8]). Suppose that f is  $\mu$ -strongly convex and L-smooth. Then, for any w,  $w' \in \mathbb{R}^d$ ,

$$(\nabla f(w) - \nabla f(w'))^{\top}(w - w') \ge \frac{\mu L}{\mu + L} \|w - w'\|^2 + \frac{1}{\mu + L} \|\nabla f(w) - \nabla f(w')\|^2. \tag{13}$$

The following existing results are more specific properties of component functions  $f(w; \xi)$ .

**Lemma 3** ([2]). Suppose that Assumptions 1 and 3 hold. Then,  $\forall w \in \mathbb{R}^d$ ,

$$\mathbb{E}[\|\nabla f(w;\xi) - \nabla f(w_*;\xi)\|^2] \le 2L[F(w) - F(w_*)],\tag{14}$$

where  $w_*$  is any optimal solution of F(w).

**Lemma 4** (Lemma 1 in [12]). Suppose that Assumptions 1 and 3 hold. Then, for  $\forall w \in \mathbb{R}^d$ ,

$$\mathbb{E}[\|\nabla f(w;\xi)\|^2] \le 4L[F(w) - F(w_*)] + 2\mathbb{E}[\|\nabla f(w_*;\xi)\|^2],\tag{15}$$

where  $w_*$  is any optimal solution of F(w).

**Lemma 5** (Lemma 1 in [11]). Let  $\xi$  and  $\{\xi_i\}_{i=1}^b$  be i.i.d. random variables with  $\mathbb{E}[\nabla f(w;\xi_i)] = \nabla F(w)$ ,  $i=1,\ldots,b$ , for all  $w\in\mathbb{R}^d$ . Then,

$$\mathbb{E}\left[\left\|\frac{1}{b}\sum_{i=1}^{b}\nabla f(w;\xi_{i}) - \nabla F(w)\right\|^{2}\right] = \frac{\mathbb{E}[\|\nabla f(w;\xi)\|^{2}] - \|\nabla F(w)\|^{2}}{b}.$$
 (16)

*Proof.* The proof of this Lemma is in [11]. We are going to use mathematical induction to prove the result. With b = 1, it is easy to see

$$\mathbb{E}\left[\|\nabla f(w;\xi_1) - \nabla F(w)\|^2\right] = \mathbb{E}[\|\nabla f(w;\xi_1)\|^2] - 2\|\nabla F(w)\|^2 + \|\nabla F(w)\|^2$$
$$= \mathbb{E}[\|\nabla f(w;\xi_1)\|^2] - \|\nabla F(w)\|^2.$$

Let assume that it is true with b = m - 1, we are going to show it is also true with b = m. We have

$$\mathbb{E}\left[\left\|\frac{1}{m}\sum_{i=1}^{m}\nabla f(w;\xi_{i})-\nabla F(w)\right\|^{2}\right]$$

$$\begin{split} &= \mathbb{E}\left[\left\|\frac{\sum_{i=1}^{m-1}\nabla f(w;\xi_{i}) - (m-1)\nabla F(w) + (\nabla f(w;\xi_{m}) - \nabla F(w))}{m}\right\|^{2}\right] \\ &= \frac{1}{m^{2}}\left(\mathbb{E}\left[\left\|\sum_{i=1}^{m-1}\nabla f(w;\xi_{i}) - (m-1)\nabla F(w)\right\|^{2}\right] + \mathbb{E}\left[\left\|\nabla f(w;\xi_{m}) - \nabla F(w)\right\|^{2}\right]\right) \\ &+ \frac{1}{m}\mathbb{E}\left[2\left(\sum_{i=1}^{m-1}\nabla f(w;\xi_{i}) - (m-1)\nabla F(w)\right)^{\top}\left(\nabla f(w;\xi_{m}) - \nabla F(w)\right)\right] \\ &= \frac{1}{m^{2}}\left(\mathbb{E}\left[\left\|\sum_{i=1}^{m-1}\nabla f(w;\xi_{i}) - (m-1)\nabla F(w)\right\|^{2}\right] + \mathbb{E}\left[\left\|\nabla f(w;\xi_{m}) - \nabla F(w)\right\|^{2}\right]\right) \\ &= \frac{1}{m^{2}}\left((m-1)\mathbb{E}[\left\|\nabla f(w;\xi_{1})\right\|^{2}] - (m-1)\left\|\nabla F(w)\right\|^{2} + \mathbb{E}[\left\|\nabla f(w;\xi_{m})\right\|^{2}] - \left\|\nabla F(w)\right\|^{2}\right) \\ &= \frac{1}{m}\left(\mathbb{E}[\left\|\nabla f(w;\xi_{1})\right\|^{2}] - \left\|\nabla F(w)\right\|^{2}\right). \end{split}$$

The third and the last equalities follow since  $\xi, \xi_1, \dots, \xi_b$  are i.i.d. with  $\mathbb{E}[\nabla f(w; \xi_i)] = \nabla F(w)$ . Therefore, the desired result is achieved.

Lemmas 4 and 5 clearly imply the following result.

**Corollary 1.** Suppose that Assumptions 1 and 3 hold. Let  $\xi$  and  $\{\xi_i\}_{i=1}^b$  be i.i.d. random variables with  $\mathbb{E}[\nabla f(w;\xi_i)] = \nabla F(w)$ ,  $i = 1, \ldots, b$ , for all  $w \in \mathbb{R}^d$ . Then,

$$\mathbb{E}\left[\left\|\frac{1}{b}\sum_{i=1}^{b}\nabla f(w;\xi_{i}) - \nabla F(w)\right\|^{2}\right] \leq \frac{4L[F(w) - F(w_{*})] + 2\mathbb{E}[\|\nabla f(w_{*};\xi)\|^{2}] - \|\nabla F(w)\|^{2}}{b}, \quad (17)$$

where  $w_*$  is any optimal solution of F(w).

Based on the above lemmas, we will show in detail how to achieve our main results in the following subsections.

#### 3.3 Special Property of SARAH Update

The most important property of the SVRG algorithm is the variance reduction of the steps. This property holds as the number of outer iteration grows, but it does not hold, if only the number of inner iterations increases. In other words, if we simply run the inner loop for many iterations (without executing additional outer loops), the variance of the steps does not reduce in the case of SVRG, while it goes to zero in the case of SARAH with large learning rate in the strongly convex case. We recall the SARAH update as follows.

$$v_t = \nabla f(w_t; \xi_t) - \nabla f(w_{t-1}; \xi_t) + v_{t-1}, \tag{18}$$

followed by the iterate update:

$$w_{t+1} = w_t - \eta v_t. (19)$$

We will now show that  $||v_t||^2$  is going to zero in expectation in the *strongly convex* case. These results substantiate our conclusion that SARAH uses more stable stochastic gradient estimates than SVRG.

**Proposition 1.** Suppose that Assumptions 1, 2 and 3 hold. Consider  $v_t$  defined by (18) with  $\eta < 2/L$  and any given  $v_0$ . Then, for any  $t \ge 1$ ,

$$\mathbb{E}[\|v_t\|^2] \le \left[1 - \left(\frac{2}{\eta L} - 1\right) \mu^2 \eta^2\right] \mathbb{E}[\|v_{t-1}\|^2]$$

$$\le \left[1 - \left(\frac{2}{\eta L} - 1\right) \mu^2 \eta^2\right]^t \|v_0\|^2.$$

The proof of this Proposition can be derived directly from Theorem 1b in [13]. This result implies that by choosing  $\eta = \mathcal{O}(1/L)$ , we obtain the linear convergence of  $||v_t||^2$  in expectation with the rate  $(1 - 1/\kappa^2)$ .

We will provide our convergence analysis in detail in next sub-section. We will divide our results into two parts: the *one-loop* results corresponding to iSARAH-IN (Algorithm 2) and the *multiple-loop* results corresponding to iSARAH (Algorithm 1).

#### 3.4 One-loop (iSARAH-IN) Results

We begin with providing two useful lemmas that do not require convexity assumption. Lemma 6 bounds the sum of expected values of  $\|\nabla F(w_t)\|^2$ ; and Lemma 7 expands the value of  $\mathbb{E}[\|\nabla F(w_t) - v_t\|^2]$ .

Lemma 6. Suppose that Assumption 1 holds. Consider iSARAH-IN (Algorithm 2). Then, we have

$$\sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t)\|^2] \le \frac{2}{\eta} \mathbb{E}[F(w_0) - F(w_*)] + \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_t\|^2], \quad (20)$$

where  $w_* = \arg\min_{w} F(w)$ .

*Proof.* By Assumption 1 and  $w_{t+1} = w_t - \eta v_t$ , we have

$$\mathbb{E}[F(w_{t+1})] \stackrel{(10)}{\leq} \mathbb{E}[F(w_t)] - \eta \mathbb{E}[\nabla F(w_t)^\top v_t] + \frac{L\eta^2}{2} \mathbb{E}[\|v_t\|^2]$$

$$= \mathbb{E}[F(w_t)] - \frac{\eta}{2} \mathbb{E}[\|\nabla F(w_t)\|^2] + \frac{\eta}{2} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] - \left(\frac{\eta}{2} - \frac{L\eta^2}{2}\right) \mathbb{E}[\|v_t\|^2],$$

where the last equality follows from the fact  $a^Tb=\frac{1}{2}\left[\|a\|^2+\|b\|^2-\|a-b\|^2\right]$  .

By summing over  $t = 0, \dots, m$ , we have

$$\mathbb{E}[F(w_{m+1})] \leq \mathbb{E}[F(w_0)] - \frac{\eta}{2} \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t)\|^2] + \frac{\eta}{2} \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] - \left(\frac{\eta}{2} - \frac{L\eta^2}{2}\right) \sum_{t=0}^{m} \mathbb{E}[\|v_t\|^2],$$

which is equivalent to  $(\eta > 0)$ :

$$\sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t)\|^2] \leq \frac{2}{\eta} \mathbb{E}[F(w_0) - F(w_m)] + \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_t\|^2] \\
\leq \frac{2}{\eta} \mathbb{E}[F(w_0) - F(w_*)] + \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_t\|^2],$$

where the second inequality follows since  $w_* = \arg\min_w F(w)$ .

**Lemma 7.** Suppose that Assumption 1 holds. Consider  $v_t$  defined by (4) in iSARAH-IN (Algorithm 2). Then for any  $t \ge 1$ ,

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] = \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] + \sum_{j=1}^t \mathbb{E}[\|v_j - v_{j-1}\|^2] - \sum_{j=1}^t \mathbb{E}[\|\nabla F(w_j) - \nabla F(w_{j-1})\|^2].$$

*Proof.* Let  $\mathcal{F}_j = \sigma(w_0, w_1, \dots, w_j)$  be the  $\sigma$ -algebra generated by  $w_0, w_1, \dots, w_j^2$ . We note that  $\xi_j$  is independent of  $\mathcal{F}_j$ . For  $j \geq 1$ , we have

$$\mathbb{E}[\|\nabla F(w_j) - v_j\|^2 | \mathcal{F}_j] = \mathbb{E}[\|[\nabla F(w_{j-1}) - v_{j-1}] + [\nabla F(w_j) - \nabla F(w_{j-1})] - [v_j - v_{j-1}]\|^2 | \mathcal{F}_j] \\
= \|\nabla F(w_{j-1}) - v_{j-1}\|^2 + \|\nabla F(w_j) - \nabla F(w_{j-1})\|^2 + \mathbb{E}[\|v_j - v_{j-1}\|^2 | \mathcal{F}_j] \\
+ 2(\nabla F(w_{j-1}) - v_{j-1})^{\top} (\nabla F(w_j) - \nabla F(w_{j-1})) \\
- 2(\nabla F(w_{j-1}) - v_{j-1})^{\top} \mathbb{E}[v_j - v_{j-1}| \mathcal{F}_j] \\
- 2(\nabla F(w_j) - \nabla F(w_{j-1}))^{\top} \mathbb{E}[v_j - v_{j-1}| \mathcal{F}_j] \\
= \|\nabla F(w_{j-1}) - v_{j-1}\|^2 - \|\nabla F(w_j) - \nabla F(w_{j-1})\|^2 + \mathbb{E}[\|v_j - v_{j-1}\|^2 | \mathcal{F}_j],$$

where the last equality follows from

$$\mathbb{E}[v_j - v_{j-1}|\mathcal{F}_j] \stackrel{\text{(4)}}{=} \mathbb{E}[\nabla f(w_j; \xi_j) - \nabla f(w_{j-1}; \xi_j)|\mathcal{F}_j] = \nabla F(w_j) - \nabla F(w_{j-1}).$$

By taking expectation for the above equation, we have

$$\mathbb{E}[\|\nabla F(w_i) - v_i\|^2] = \mathbb{E}[\|\nabla F(w_{i-1}) - v_{i-1}\|^2] - \mathbb{E}[\|\nabla F(w_i) - \nabla F(w_{i-1})\|^2] + \mathbb{E}[\|v_i - v_{i-1}\|^2].$$

By summing over  $j = 1, \dots, t \ (t \ge 1)$ , we have

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] = \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] + \sum_{j=1}^t \mathbb{E}[\|v_j - v_{j-1}\|^2] - \sum_{j=1}^t \mathbb{E}[\|\nabla F(w_j) - \nabla F(w_{j-1})\|^2].$$

#### 3.4.1 General Convex Case

In this subsection, we analyze one-loop results of Inexact SARAH (Algorithm 2) in the general convex case. We first derive the bound for  $\mathbb{E}[\|\nabla F(w_t) - v_t\|^2]$ .

**Lemma 8.** Suppose that Assumptions 1 and 3 hold. Consider  $v_t$  defined as (4) in SARAH (Algorithm 1) with  $\eta < 2/L$ . Then we have that for any  $t \ge 1$ ,

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] \le \frac{\eta L}{2 - \eta L} \Big[ \mathbb{E}[\|v_0\|^2] - \mathbb{E}[\|v_t\|^2] \Big] + \mathbb{E}[\|\nabla F(w_0) - v_0\|^2]. \tag{21}$$

 $<sup>{}^{2}\</sup>mathcal{F}_{j}$  contains all the information of  $w_{0},\ldots,w_{j}$  as well as  $v_{0},\ldots,v_{j-1}$ 

*Proof.* For  $j \geq 1$ , we have

$$\mathbb{E}[\|v_{j}\|^{2}|\mathcal{F}_{j}] \\
= \mathbb{E}[\|v_{j-1} - (\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j})\|^{2}|\mathcal{F}_{j}] \\
= \|v_{j-1}\|^{2} + \mathbb{E}\Big[\|\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j})\|^{2} - \frac{2}{\eta}(\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j}))^{\top}(w_{j-1} - w_{j})|\mathcal{F}_{j}\Big] \\
\stackrel{(12)}{\leq} \|v_{j-1}\|^{2} + \mathbb{E}\Big[\|\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j})\|^{2} - \frac{2}{L\eta}\|\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j})\|^{2}|\mathcal{F}_{j}\Big] \\
= \|v_{j-1}\|^{2} + \left(1 - \frac{2}{\eta L}\right)\mathbb{E}[\|\nabla f(w_{j-1};\xi_{j}) - \nabla f(w_{j};\xi_{j})\|^{2}|\mathcal{F}_{j}] \\
\stackrel{(4)}{=} \|v_{j-1}\|^{2} + \left(1 - \frac{2}{\eta L}\right)\mathbb{E}[\|v_{j} - v_{j-1}\|^{2}|\mathcal{F}_{j}],$$

which, if we take expectation, implies that

$$\mathbb{E}[\|v_j - v_{j-1}\|^2] \le \frac{\eta L}{2 - nL} \Big[ \mathbb{E}[\|v_{j-1}\|^2] - \mathbb{E}[\|v_j\|^2] \Big],$$

when  $\eta < 2/L$ .

By summing the above inequality over  $j = 1, \dots, t \ (t \ge 1)$ , we have

$$\sum_{j=1}^{t} \mathbb{E}[\|v_j - v_{j-1}\|^2] \le \frac{\eta L}{2 - \eta L} \Big[ \mathbb{E}[\|v_0\|^2] - \mathbb{E}[\|v_t\|^2] \Big]. \tag{22}$$

By Lemma 7, we have

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] \leq \sum_{j=1}^t \mathbb{E}[\|v_j - v_{j-1}\|^2] + \mathbb{E}[\|\nabla F(w_0) - v_0\|^2]$$

$$\stackrel{(22)}{\leq} \frac{\eta L}{2 - \eta L} \Big[ \mathbb{E}[\|v_0\|^2] - \mathbb{E}[\|v_t\|^2] \Big] + \mathbb{E}[\|\nabla F(w_0) - v_0\|^2].$$

**Lemma 9.** Suppose that Assumptions 1 and 3 hold. Consider  $v_0$  defined as (3) in SARAH (Algorithm 1). Then we have,

$$\frac{\eta L}{2 - \eta L} \mathbb{E}[\|v_0\|^2] + \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] 
\leq \frac{2}{2 - \eta L} \left( \frac{4L \mathbb{E}[F(w_0) - F(w_*)] + 2\mathbb{E}\left[\|\nabla f(w_*; \xi)\|^2\right] - \mathbb{E}[\|\nabla F(w_0)\|^2]}{b} \right) 
+ \frac{\eta L}{2 - \eta L} \mathbb{E}[\|\nabla F(w_0)\|^2].$$
(23)

Proof. By Corollary 1, we have

$$\frac{\eta L}{2 - \eta L} \mathbb{E}[\|v_0\|^2 | w_0] - \frac{\eta L}{2 - \eta L} \|\nabla F(w_0)\|^2 + \mathbb{E}[\|\nabla F(w_0) - v_0\|^2 | w_0] 
= \frac{2}{2 - \eta L} \Big[ \mathbb{E}[\|v_0\|^2 | w_0] - \|\nabla F(w_0)\|^2 \Big] 
= \frac{2}{2 - \eta L} \Big[ \mathbb{E}[\|v_0 - \nabla F(w_0)\|^2 | w_0 \Big] 
\stackrel{(17)}{\leq} \frac{2}{2 - \eta L} \left( \frac{4L[F(w_0) - F(w_*)] + 2\mathbb{E}\left[\|\nabla f(w_*; \xi)\|^2\right] - \|\nabla F(w_0)\|^2}{b} \right).$$

Taking the expectation and adding  $\frac{\eta L}{2-\eta L}\mathbb{E}[\|\nabla F(w_0)\|^2]$  for both sides, the desired result is achieved.  $\Box$ 

We then derive this basic result for the convex case by using Lemmas 8 and 9.

**Lemma 10.** Suppose that Assumptions 1 and 3 hold. Consider iSARAH-IN (Algorithm 2) with  $\eta \leq 1/L$ . Then, we have

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^{2}] \leq \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \mathbb{E}[\|\nabla F(w_{0})\|^{2}] + \frac{2}{2 - \eta L} \left(\frac{4L\mathbb{E}[F(w_{0}) - F(w_{*})] + 2\mathbb{E}\left[\|\nabla f(w_{*};\xi)\|^{2}\right] - \mathbb{E}[\|\nabla F(w_{0})\|^{2}]}{b}\right), (24)$$

where  $w_*$  is any optimal solution of F(w); and  $\xi$  is the random variable.

Proof. By Lemma 8, we have

$$\sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] \le \frac{m\eta L}{2 - \eta L} \mathbb{E}[\|v_0\|^2] + (m+1)\mathbb{E}[\|\nabla F(w_0) - v_0\|^2]. \tag{25}$$

Hence, by Lemma 6 with  $\eta < 1/L$ , we have

$$\sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t)\|^2] \leq \frac{2}{\eta} \mathbb{E}[F(w_0) - F(w_*)] + \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_t) - v_t\|^2] 
\leq \frac{2}{\eta} \mathbb{E}[F(w_0) - F(w_*)] + \frac{m\eta L}{2 - \eta L} \mathbb{E}[\|v_0\|^2] + (m+1)\mathbb{E}[\|\nabla F(w_0) - v_0\|^2].$$
(26)

Since  $\tilde{w} = w_t$ , where t is picked uniformly at random from  $\{0, 1, \dots, m\}$ . The following holds,

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^{2}] = \frac{1}{m+1} \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_{t})\|^{2}]$$

$$\stackrel{(26)}{\leq} \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \mathbb{E}[\|v_{0}\|^{2}] + \mathbb{E}[\|\nabla F(w_{0}) - v_{0}\|^{2}]$$

$$\stackrel{(23)}{\leq} \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \mathbb{E}[\|\nabla F(w_{0})\|^{2}]$$

$$+ \frac{2}{2 - \eta L} \left(\frac{4L\mathbb{E}[F(w_{0}) - F(w_{*})] + 2\mathbb{E}\left[\|\nabla f(w_{*}; \xi)\|^{2}\right] - \mathbb{E}[\|\nabla F(w_{0})\|^{2}]}{b}\right).$$

This expected bound for  $\|\nabla F(\tilde{w})\|^2$  will be used for deriving both one-loop and multiple-loop results in the convex case.

Lemma 10 can be used to get the following result for general convex.

**Theorem 1.** Suppose that Assumptions 1 and 3 hold. Consider iSARAH-IN (Algorithm 2) with  $\eta = \frac{1}{L\sqrt{m+1}} \le \frac{1}{L}$ ,  $b = 2\sqrt{m+1}$  and a given  $w_0$ . Then we have,

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^2] \le \frac{2}{\eta(m+1)} [F(w_0) - F(w_*)] + \frac{1}{\sqrt{m+1}} \left[ 4L[F(w_0) - F(w_*)] + 2\mathbb{E}[\|\nabla f(w_*; \xi)\|^2] \right], \tag{27}$$

where  $w_*$  is any optimal solution of F(w); and  $\xi$  is some random variable.

*Proof.* By Lemma 10, for any given  $w_0$ , we have

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^{2}] \leq \frac{2}{\eta(m+1)} [F(w_{0}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \|\nabla F(w_{0})\|^{2}$$

$$+ \frac{2}{2 - \eta L} \left( \frac{4L[F(w_{0}) - F(w_{*})] + 2\mathbb{E}\left[\|\nabla f(w_{*};\xi)\|^{2}\right] - \|\nabla F(w_{0})\|^{2}}{b} \right)$$

$$\leq \frac{2}{\eta(m+1)} [F(w_{0}) - F(w_{*})] + \frac{1}{2 - \eta L} \frac{4L}{\sqrt{m+1}} [F(w_{0}) - F(w_{*})]$$

$$+ \frac{2}{2 - \eta L} \frac{\mathbb{E}\left[\|\nabla f(w_{*};\xi)\|^{2}\right]}{\sqrt{m+1}}$$

$$\leq \frac{2}{\eta(m+1)} [F(w_{0}) - F(w_{*})] + \frac{1}{\sqrt{m+1}} \left[4L[F(w_{0}) - F(w_{*})] + 2\mathbb{E}[\|\nabla f(w_{*};\xi)\|^{2}]\right].$$

The last inequality follows since  $\eta \leq \frac{1}{L}$ , which implies  $\frac{1}{2-\eta L} \leq 1$ . The second last inequality follows since  $\eta \leq \frac{1}{L\sqrt{m+1}}$  and  $b = 2\sqrt{m+1}$ .

Based on Theorem 1, we are able to derive the following total complexity for iSARAH-IN in the general convex case.

**Corollary 2.** Suppose that Assumptions 1 and 3 hold. Consider iSARAH-IN (Algorithm 2) with the learning rate  $\eta = \mathcal{O}\left(\frac{1}{\sqrt{m+1}}\right)$  and the number of samples  $b = 2\sqrt{m+1}$ , where m is the total number of iterations, then  $\|\nabla F(\tilde{w})\|^2$  converges sublinearly in expectation with a rate of  $\mathcal{O}\left(\sqrt{\frac{1}{m+1}}\right)$ , and therefore, the total complexity to achieve an  $\epsilon$ -accurate solution is  $\mathcal{O}(1/\epsilon^2)$ .

*Proof.* It is easy to see that to achieve  $\mathbb{E}[\|\nabla F(\tilde{w})\|^2] \leq \epsilon$  we need  $m = \mathcal{O}(1/\epsilon^2)$  and hence the total work is  $2\sqrt{m} + 2m = \mathcal{O}\left(\frac{1}{\epsilon} + \frac{1}{\epsilon^2}\right) = \mathcal{O}\left(\frac{1}{\epsilon^2}\right)$ .

#### 3.4.2 Nonconvex Case

We now move to the nonconvex case. We begin by stating and proving a lemma similar to Lemma 8, bounding  $\mathbb{E}[\|\nabla F(w_t) - v_t\|^2]$ , but without Assumption 3.

**Lemma 11.** Suppose that Assumption 1 holds. Consider  $v_t$  defined as (4) in iSARAH-IN (Algorithm 2). Then for any  $t \ge 1$ ,

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] \le \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] + L^2 \eta^2 \sum_{j=1}^t \mathbb{E}[\|v_{j-1}\|^2]. \tag{28}$$

*Proof.* We have, for  $t \geq 1$ ,

$$\|v_t - v_{t-1}\|^2 \stackrel{\text{(4)}}{=} \|\nabla f(w_t; \xi_t) - \nabla f(w_{t-1}; \xi_t)\|^2 \stackrel{\text{(8)}}{\leq} L^2 \|w_t - w_{t-1}\|^2 = L^2 \eta^2 \|v_{t-1}\|^2. \tag{29}$$

Hence, by Lemma 7,

$$\mathbb{E}[\|\nabla F(w_t) - v_t\|^2] \leq \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] + \sum_{j=1}^t \mathbb{E}[\|v_j - v_{j-1}\|^2]$$

$$\stackrel{(29)}{\leq} \mathbb{E}[\|\nabla F(w_0) - v_0\|^2] + L^2 \eta^2 \sum_{j=1}^t \mathbb{E}[\|v_{j-1}\|^2].$$

**Lemma 12.** Suppose that Assumption 1 holds. Consider  $v_t$  defined as (4) in iSARAH-IN (Algorithm 2) with  $\eta \leq \frac{2}{L(\sqrt{1+4m+1})}$ . Then we have

$$L^{2}\eta^{2} \sum_{t=0}^{m} \sum_{j=1}^{t} \mathbb{E}[\|v_{j-1}\|^{2}] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_{t}\|^{2}] \le 0.$$
(30)

*Proof.* For  $\eta \leq \frac{2}{L(\sqrt{1+4m}+1)}$ , we have

$$L^{2}\eta^{2} \sum_{t=0}^{m} \sum_{j=1}^{t} \mathbb{E}[\|v_{j-1}\|^{2}] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_{t}\|^{2}]$$

$$= L^{2}\eta^{2} \Big[ m\mathbb{E}\|v_{0}\|^{2} + (m-1)\mathbb{E}\|v_{1}\|^{2} + \dots + \mathbb{E}\|v_{m-1}\|^{2} \Big]$$

$$- (1 - L\eta) \Big[ \mathbb{E}\|v_{0}\|^{2} + \mathbb{E}\|v_{1}\|^{2} + \dots + \mathbb{E}\|v_{m}\|^{2} \Big]$$

$$\leq [L^{2}\eta^{2}m - (1 - L\eta)] \sum_{t=1}^{m} \mathbb{E}[\|v_{t-1}\|^{2}] \leq 0,$$

since  $\eta = \frac{2}{L(\sqrt{1+4m+1})}$  is a root of the equation  $L^2\eta^2m - (1-L\eta) = 0$ .

With the help of the above lemmas, we are able to derive our result for nonconvex.

**Theorem 2.** Suppose that Assumption 1 holds. Consider iSARAH-IN (Algorithm 2) with  $\eta \leq \frac{2}{L(\sqrt{1+4m}+1)} \leq \frac{1}{L}$ ,  $b = \sqrt{m+1}$  and a given  $w_0$ . Then we have,

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^2] \le \frac{2}{\eta(m+1)} [F(w_0) - F^*] + \frac{1}{\sqrt{m+1}} \Big( \mathbb{E}[\|\nabla f(w_0; \xi)\|^2] \Big), \tag{31}$$

where  $F^*$  is any lower bound of F; and  $\xi$  is some random variable.

*Proof.* Let  $F^*$  be any lower bound of F. By Lemma 6 and since  $\tilde{w} = w_t$ , where t is picked uniformly at random from  $\{0, 1, \dots, m\}$ , we have

$$\mathbb{E}[\|\nabla F(\tilde{w})\|^{2}] = \frac{1}{m+1} \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_{t})\|^{2}]$$

$$\leq \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F^{*}] + \frac{1}{m+1} \left( \sum_{t=0}^{m} \mathbb{E}[\|\nabla F(w_{t}) - v_{t}\|^{2}] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_{t}\|^{2}] \right)$$

$$\stackrel{(28)}{\leq} \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F^{*}] + \mathbb{E}[\|\nabla F(w_{0}) - v_{0}\|^{2}]$$

$$+ \frac{1}{m+1} \left( L^{2}\eta^{2} \sum_{t=0}^{m} \sum_{j=1}^{t} \mathbb{E}[\|v_{j-1}\|^{2}] - (1 - L\eta) \sum_{t=0}^{m} \mathbb{E}[\|v_{t}\|^{2}] \right)$$

$$\stackrel{(30)}{\leq} \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F^{*}] + \mathbb{E}[\|\nabla F(w_{0}) - v_{0}\|^{2}]$$

$$\stackrel{(16)}{\leq} \frac{2}{\eta(m+1)} \mathbb{E}[F(w_{0}) - F^{*}] + \frac{1}{b} \mathbb{E}[\|\nabla f(w_{0}; \xi)\|^{2}].$$

For any given  $w_0$  and  $b = \sqrt{m+1}$ , we could achieve the desired result.

Based on Theorem 2, we are able to derive the following total complexity for iSARAH-IN in the nonconvex case.

**Corollary 3.** Suppose that Assumption 1 holds. Consider iSARAH-IN (Algorithm 2) with the learning rate  $\eta = \mathcal{O}\left(\frac{1}{\sqrt{m+1}}\right)$  and the number of samples  $b = \sqrt{m+1}$ , where m is the total number of iterations, then  $\|\nabla F(\tilde{w})\|^2$  converges sublinearly in expectation with a rate of  $\mathcal{O}\left(\sqrt{\frac{1}{m+1}}\right)$ , and therefore, the total complexity to achieve an  $\epsilon$ -accurate solution is  $\mathcal{O}(1/\epsilon^2)$ .

*Proof.* Same as general convex case, to achieve  $\mathbb{E}[\|\nabla F(\tilde{w})\|^2] \leq \epsilon$  we need  $m = \mathcal{O}(1/\epsilon^2)$  and hence the total work is  $\sqrt{m} + 2m = \mathcal{O}\left(\frac{1}{\epsilon} + \frac{1}{\epsilon^2}\right) = \mathcal{O}\left(\frac{1}{\epsilon^2}\right)$ .

## 3.5 Multiple-loop iSARAH Results

In this section, we analyze multiple-loop results of Inexact SARAH (Algorithm 1).

#### 3.5.1 Strongly Convex Case

We now turn to the discussion on the convergence of iSARAH under the strong convexity assumption on F.

**Theorem 3.** Suppose that Assumptions 1, 2 and 3 hold. Consider iSARAH (Algorithm 1) with the choice of  $\eta$ , m, and b such that

$$\alpha = \frac{1}{\mu \eta(m+1)} + \frac{\eta L}{2 - \eta L} + \frac{4\kappa - 2}{b(2 - \eta L)} < 1.$$

(Note that  $\kappa = L/\mu$ .) Then, we have

$$\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] - \Delta \le \alpha^s (\|\nabla F(\tilde{w}_0)\|^2 - \Delta), \tag{32}$$

where

$$\Delta = \frac{\delta}{1-\alpha} \text{ and } \delta = \frac{4}{b(2-\eta L)} \mathbb{E}\left[\|\nabla f(w_*;\xi)\|^2\right].$$

*Proof.* By Lemma 10, with  $\tilde{w} = \tilde{w}_s$  and  $w_0 = \tilde{w}_{s-1}$ , we have

$$\mathbb{E}[\|\nabla F(\tilde{w}_{s})\|^{2}] \leq \frac{2}{\eta(m+1)} \mathbb{E}[F(\tilde{w}_{s-1}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}]$$

$$+ \frac{2}{2 - \eta L} \left( \frac{4L\mathbb{E}[F(\tilde{w}_{s-1}) - F(w_{*})] + 2\mathbb{E}\left[\|\nabla f(w_{*}; \xi)\|^{2}\right] - \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}]}{b} \right)$$

$$\stackrel{(9)}{\leq} \left( \frac{1}{\mu \eta(m+1)} + \frac{\eta L}{2 - \eta L} + \frac{4\kappa - 2}{b(2 - \eta L)} \right) \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}]$$

$$+ \frac{4}{b(2 - \eta L)} \mathbb{E}\left[\|\nabla f(w_{*}; \xi)\|^{2}\right]$$

$$= \alpha \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}] + \delta$$

$$\leq \alpha^{s} \|\nabla F(\tilde{w}_{0})\|^{2} + \alpha^{s-1}\delta + \dots + \alpha\delta + \delta$$

$$\leq \alpha^{s} \|\nabla F(\tilde{w}_{0})\|^{2} + \delta \frac{1 - \alpha^{s}}{1 - \alpha}$$

$$= \alpha^{s} \|\nabla F(\tilde{w}_{0})\|^{2} + \Delta(1 - \alpha^{s})$$

$$= \alpha^{s} (\|\nabla F(\tilde{w}_{0})\|^{2} - \Delta) + \Delta.$$

$$(33)$$

By adding  $-\Delta$  to both sides, we achieve the desired result.

Based on Theorem 3, we are able to derive the following total complexity for iSARAH in the strongly convex case.

**Corollary 4.** Let  $\eta = \mathcal{O}\left(\frac{1}{L}\right)$ ,  $m = \mathcal{O}(\kappa)$ ,  $b = \mathcal{O}\left(\max\left\{\frac{1}{\epsilon}, \kappa\right\}\right)$  and  $s = \mathcal{O}\left(\log\left(\frac{1}{\epsilon}\right)\right)$  in Theorem 3. Then, the total work complexity to achieve  $\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] \leq \epsilon$  is  $\mathcal{O}\left(\max\left\{\frac{1}{\epsilon}, \kappa\right\}\log\left(\frac{1}{\epsilon}\right)\right)$ .

*Proof.* For example, let  $\eta = \frac{2}{5L}$ ,  $m = 20\kappa - 1$ , and  $b = \max\left\{20\kappa - 10, \frac{20\mathbb{E}\left[\|\nabla f(w_*;\xi)\|^2\right]}{\epsilon}\right\}$ . From (33), we have

$$\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] \le \left(\frac{1}{8} + \frac{1}{4} + \frac{1}{8}\right) \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^2] + \frac{\epsilon}{8}$$

$$\le \frac{1}{2} \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^2] + \frac{\epsilon}{8}$$

$$\le \frac{1}{2^s} \|\nabla F(\tilde{w}_0)\|^2 + \frac{\epsilon}{4}.$$

To guarantee that  $\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] \leq \epsilon$ , it is sufficient to make  $\frac{1}{2^s}\|\nabla F(\tilde{w}_0)\|^2 = \frac{3}{4}\epsilon$  or equivalently  $s = \log\left(\frac{\|\nabla F(\tilde{w}_0)\|^2}{\frac{3}{4}\epsilon}\right)$ . This implies the total complexity to achieve an  $\epsilon$ -accuracy solution is  $(b+m)s = \mathcal{O}\left(\left(\max\left\{\frac{1}{\epsilon},\kappa\right\} + \kappa\right)\log\left(\frac{1}{\epsilon}\right)\right) = \mathcal{O}\left(\max\left\{\frac{1}{\epsilon},\kappa\right\}\log\left(\frac{1}{\epsilon}\right)\right)$ .

#### 3.5.2 General Convex Case

We finally turn to the analysis of the convergence rate of the multiple-loop iSARAH in the general convex case. As mentioned in the introduction, we are able to achieve best sample complexity rates of any stochastic algorithm, to the best of our knowledge, but under an additional reasonably mild assumption. We introduce this assumption below.

**Assumption 4.** Let  $\tilde{w}_0, \dots, \tilde{w}_T$  be the (outer) iterations of Algorithm 1. We assume that there exist M > 0 and N > 0 such that,  $k = 0, \dots, T$ ,

$$F(\tilde{w}_k) - F(w_*) \le M \|\nabla F(\tilde{w}_k)\|^2 + N,$$
 (34)

where  $F(w_*)$  is the optimal value of F.

Let us discuss Assumption 4. First, we note that the assumption only requires to hold for the outer iterations  $\tilde{w}_0,\ldots,\tilde{w}_{\mathcal{T}}$  of Algorithm 1 instead of holding for all  $w\in\mathbb{R}^d$  or for all of the inner iterates. Moreover, this assumption is clearly weaker than the Polyak-Lojasiewicz (PL) condition, which has been studied and discussed in [15, 9, 3] which itself is weaker than strong convexity assumption. Under PL condition, we simply have N=0 in (34) and as we will discuss below we can recover better convergence rate in this case. On the other hand, if PL condition does not hold but if the sequence of iterates  $\{\tilde{w}_k\}$  remains in a set, say  $\mathcal{W}$ , on which the objective function is bounded from above, that is for all  $w\in\mathcal{W}$ ,  $F(w)\leq F_{\max}$  for some finite value  $F_{\max}$ , then Assumption 4 is satisfied with  $N=F_{\max}-F(w_*)$  and M=0, where  $F(w_*)$  is the optimal value of F. In other words, Assumption 4 is a relaxation of the boundedness assumption and the PL condition.

**Theorem 4.** Suppose that Assumptions 1, 3, and 4 hold. Consider iSARAH (Algorithm 1) with the choice of  $\eta$ , m, and b such that

$$\alpha_c = \frac{2M}{\eta(m+1)} + \frac{\eta L}{2 - \eta L} + \frac{8LM - 1}{b(2 - \eta L)} < 1.$$

Then, we have

$$\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] - \Delta_c \le \alpha^s (\|\nabla F(\tilde{w}_0)\|^2 - \Delta_c), \tag{35}$$

where

$$\Delta_c = \frac{\delta_c}{1 - \alpha_c} \text{ and } \delta_c = \frac{2N}{\eta(m+1)} + \frac{8LN}{b(2 - \eta L)} + \frac{4\mathbb{E}\left[\|\nabla f(w_*; \xi)\|^2\right]}{b(2 - \eta L)}.$$

*Proof.* By Lemma 10, with  $\tilde{w} = \tilde{w}_s$  and  $w_0 = \tilde{w}_{s-1}$ , we have

$$\mathbb{E}[\|\nabla F(\tilde{w}_{s})\|^{2}] \leq \frac{2}{\eta(m+1)} \mathbb{E}[F(\tilde{w}_{s-1}) - F(w_{*})] + \frac{\eta L}{2 - \eta L} \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}] + \frac{2}{2 - \eta L} \left(\frac{4L\mathbb{E}[F(\tilde{w}_{s-1}) - F(w_{*})] + 2\mathbb{E}\left[\|\nabla f(w_{*};\xi)\|^{2}\right] - \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}]}{b}\right)$$

$$\stackrel{(34)}{\leq} \left(\frac{2M}{\eta(m+1)} + \frac{\eta L}{2 - \eta L} + \frac{8LM - 1}{b(2 - \eta L)}\right) \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^{2}]$$

$$+ \frac{2N}{\eta(m+1)} + \frac{8LN}{b(2-\eta L)} + \frac{4\mathbb{E}\left[\|\nabla f(w_*;\xi)\|^2\right]}{b(2-\eta L)}$$

$$= \alpha_c \mathbb{E}[\|\nabla F(\tilde{w}_{s-1})\|^2] + \delta_c$$

$$\leq \alpha_c^s (\|\nabla F(\tilde{w}_0)\|^2 - \Delta_c) + \Delta_c.$$

**Remark 1.** From Theorem 4, we also observe that in the case when N=0 and  $\kappa=L\cdot M$ , then  $\Delta_c=\mathcal{O}(\frac{1}{b})$ , which means we can select  $m=\mathcal{O}(\kappa)$  and  $b=\mathcal{O}\left(\max\left\{\frac{1}{\epsilon},\kappa\right\}\right)$  and thus recover the convergence rate for the strongly convex case, simply under Assumption 4 with N=0.

Now choosing the appropriate values for  $\eta$ , m, b and s we can achieve the following complexity result.

**Corollary 5.** Let 
$$\eta = \mathcal{O}\left(\frac{1}{L}\right)$$
,  $m = \mathcal{O}\left(\frac{\max\{M,N\}}{\epsilon}\right)$ ,  $b = \mathcal{O}\left(\frac{\max\{M,N\}}{\epsilon}\right)$  and  $s = \mathcal{O}\left(\log\left(\frac{1}{\epsilon}\right)\right)$  in Theorem 4. Then, the total work complexity to achieve  $\mathbb{E}[\|\nabla F(\tilde{w}_s)\|^2] \leq \epsilon$  is  $(b+m)s = \mathcal{O}\left(\frac{\max\{M,N\}}{\epsilon}\log\left(\frac{1}{\epsilon}\right)\right)$ .

We can observe that, with the help of Assumption 4, iSARAH could achieve the best known complexity among stochastic methods (those which do not have access to exact gradient computation) in the general convex case.

#### 4 Conclusion and Future Work

We have provided the analysis of the inexact version of SARAH, which requires only stochastic gradient information computed on a mini-batch of sufficient size. We analyze the one-loop results (iSARAH-IN) in the general convex and nonconvex cases with only smoothness assumption; and the multiple-loop results (iSARAH) in the strongly convex case and with an additional assumption (Assumption 4) in the general convex case. With this Assumption 4, which we argue is reasonable, iSARAH achieves the best known complexity among stochastic methods. The convergence rate of iSARAH multiple-loop for nonconvex still remains an open question.

## References

- [1] Aaron Defazio, Francis Bach, and Simon Lacoste-Julien. SAGA: A fast incremental gradient method with support for non-strongly convex composite objectives. In *NIPS*, pages 1646–1654, 2014.
- [2] Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *NIPS*, pages 315–323, 2013.
- [3] Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In Paolo Frasconi, Niels Landwehr, Giuseppe Manco, and Jilles Vreeken, editors, *Machine Learning and Knowledge Discovery in Databases*, pages 795– 811, Cham, 2016. Springer International Publishing.
- [4] Jakub Konečný and Peter Richtárik. Semi-stochastic gradient descent methods. *Frontiers in Applied Mathematics and Statistics*, 3:9, 2017.

- [5] Nicolas Le Roux, Mark Schmidt, and Francis Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *NIPS*, pages 2663–2671, 2012.
- [6] Lihua Lei and Michael Jordan. Less than a Single Pass: Stochastically Controlled Stochastic Gradient. In Aarti Singh and Jerry Zhu, editors, *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 148–156, Fort Lauderdale, FL, USA, 20–22 Apr 2017. PMLR.
- [7] Lihua Lei, Cheng Ju, Jianbo Chen, and Michael I Jordan. Non-convex finite-sum optimization via SCSG methods. In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems 30*, pages 2348–2358. Curran Associates, Inc., 2017.
- [8] Yurii Nesterov. *Introductory lectures on convex optimization: a basic course*. Applied optimization. Kluwer Academic Publ., Boston, Dordrecht, London, 2004.
- [9] Yurii Nesterov and Boris T Polyak. Cubic regularization of newton method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.
- [10] Lam Nguyen, Jie Liu, Katya Scheinberg, and Martin Takáč. SARAH: A novel method for machine learning problems using stochastic recursive gradient. *ICML*, pages 2613–2621, 2017.
- [11] Lam Nguyen, Nam Nguyen, Dzung Phan, Jayant Kalagnanam, and Katya Scheinberg. When does stochastic gradient algorithm work well? *arXiv:1801.06159*, 2018.
- [12] Lam Nguyen, Phuong Ha Nguyen, Marten van Dijk, Peter Richtárik, Katya Scheinberg, and Martin Takáč. SGD and Hogwild! convergence without the bounded gradients assumption. *ICML*, 2018.
- [13] Lam M. Nguyen, Jie Liu, Katya Scheinberg, and Martin Takáč. Stochastic recursive gradient algorithm for nonconvex optimization. *CoRR*, abs/1705.07261, 2017.
- [14] Courtney Paquette and Katya Scheinberg. A stochastic line search method with convergence rate analysis. Technical report, Lehigh University, 2018.
- [15] Boris T. Polyak. Some methods of speeding up the convergence of iteration methods. *USSR Computational Mathematics and Mathematical Physics*, 4(5):1–17, 1964.
- [16] Sashank J. Reddi, Ahmed Hefny, Suvrit Sra, Barnabás Póczos, and Alexander J. Smola. Stochastic variance reduction for nonconvex optimization. In *ICML*, pages 314–323, 2016.
- [17] Mark Schmidt, Nicolas Le Roux, and Francis Bach. Minimizing finite sums with the stochastic average gradient. *Mathematical Programming*, 162(1-2):83–112, 2017.