

Topological quantum field theories and quantum gravity.

Ingrid Membrillo

March 20, 2015

Axiomatic formulation of Topological Quantum Field Theory

From the mathematical point of view we expect that all quantum field theories keep

- general features of quantum theory (probabilistic)
- topological framework of spacetime

Both features are embraced by the precise notion of a topological quantum field theory.

M. Atiyah (1988) gave a formal mathematical framework to develop quantum theories in physics.

Axiomatic formulation of Topological Quantum Field Theory

Definition

A topological quantum field theory (TQFT) in dimension $d + 1$ consist of the following data:

Fixed an integer d , consider smooth oriented manifolds M of dimension $d + 1$ (with boundary) and also smooth closed oriented manifolds X of dimension d .

Axiomatic formulation of Topological Quantum Field Theory

Definition

A topological quantum field theory (TQFT) in dimension $d + 1$ consist of the following data:

Fixed an integer d , consider smooth oriented manifolds M of dimension $d + 1$ (with boundary) and also smooth closed oriented manifolds X of dimension d .

- *To each X we associate functorially a **finite-dimensional complex vector space $\mathbf{Z}(X)$** .*

Axiomatic formulation of Topological Quantum Field Theory

Definition

A topological quantum field theory (TQFT) in dimension $d + 1$ consist of the following data:

Fixed an integer d , consider smooth oriented manifolds M of dimension $d + 1$ (with boundary) and also smooth closed oriented manifolds X of dimension d .

- *To each X we associate functorially a **finite-dimensional complex vector space** $\mathbf{Z}(X)$.*
- *To each M , whose boundary is X we associate a **vector** $\mathbf{Z}(M) \in \mathbf{Z}(X)$.*

Axiomatic formulation of Topological Quantum Field Theory

These data are subject to the following properties (**axioms**):

- 1) An orientation preserving diffeomorphism $f : X \rightarrow X'$ induces an isomorphism

$$\mathbf{Z}(f) : \mathbf{Z}(X) \rightarrow \mathbf{Z}(X')$$

and if $g : X' \rightarrow X''$ then

$$\mathbf{Z}(fg) = \mathbf{Z}(f)\mathbf{Z}(g)$$

Axiomatic formulation of Topological Quantum Field Theory

These data are subject to the following properties (**axioms**):

- 1) An orientation preserving diffeomorphism $f : X \rightarrow X'$ induces an isomorphism

$$\mathbf{Z}(f) : \mathbf{Z}(X) \rightarrow \mathbf{Z}(X')$$

and if $g : X' \rightarrow X''$ then

$$\mathbf{Z}(fg) = \mathbf{Z}(f)\mathbf{Z}(g)$$

Axiomatic formulation of Topological Quantum Field Theory

- 2) $\mathbf{Z}(X^*) = \mathbf{Z}(X)^*$ where X^* is X with opposite orientation and $\mathbf{Z}(X)^*$ denotes the dual vector space.

Axiomatic formulation of Topological Quantum Field Theory


- 2) $\mathbf{Z}(X^*) = \mathbf{Z}(X)^*$ where X^* is X with opposite orientation and $\mathbf{Z}(X)^*$ denotes the dual vector space.
- 3) \mathbf{Z} is multiplicative.

Thus, for disjoint unions

$$\mathbf{Z}(X_1 \cup X_2) = \mathbf{Z}(X_1) \otimes \mathbf{Z}(X_2).$$


Axiomatic formulation of Topological Quantum Field Theory

Moreover if $\partial M_1 = X_1 \cup X_3$, $\partial M_2 = X_2 \cup X_3^*$ and $M = M_1 \cup_{X_3} M_2$ is the manifold obtained by identification of the common X_3 – *component*.



Axiomatic formulation of Topological Quantum Field Theory

Moreover if $\partial M_1 = X_1 \cup X_3$, $\partial M_2 = X_2 \cup X_3^*$ and $M = M_1 \cup_{X_3} M_2$ is the manifold obtained by identification of the common X_3 – *component*.



$$Z(\Sigma_1) \otimes Z(\Sigma_3) \otimes Z(\Sigma_3)^* \otimes Z(\Sigma_2) \rightarrow Z(\Sigma_1) \otimes Z(\Sigma_2)$$

Axiomatic formulation of Topological Quantum Field Theory

4a) When X is the empty d -dimensional manifold

$$\mathbf{Z}(\emptyset) = \mathbb{C}. \quad (1)$$

Axiomatic formulation of Topological Quantum Field Theory

4a) When X is the empty d -dimensional manifold

$$\mathbf{Z}(\emptyset) = \mathbb{C}. \quad (1)$$

4b) Similarly, when $M = \emptyset$

$$\mathbf{Z}(\emptyset) = 1 \in \mathbb{C} \quad (2)$$

Axiomatic formulation of Topological Quantum Field Theory

4a) When X is the empty d -dimensional manifold

$$\mathbf{Z}(\emptyset) = \mathbb{C}. \quad (1)$$

4b) Similarly, when $M = \emptyset$

$$\mathbf{Z}(\emptyset) = 1 \in \mathbb{C} \quad (2)$$

Note that when M is a closed d -dimensional manifold so that $\partial M = \emptyset$, then

$$\mathbf{Z}(\mathbf{M}) \in \mathbf{Z}(\emptyset) = \mathbb{C}$$

That means, $\mathbf{Z}(\mathbf{M})$ is a **numerical invariant** of M .

State sum model as a 3D TQFT of gravity

Einstein equations (movement equations) can be derived through minimizing the action S :


$$S = \frac{1}{16\pi} \int R \sqrt{\det(g_{\mu\nu})} d^4x$$

State sum model as a 3D TQFT of gravity

Einstein equations (movement equations) can be derived through minimizing the action S :


$$S = \frac{1}{16\pi} \int R \sqrt{\det(g_{\mu\nu})} d^4x$$

1961 Regge gave a discrete formulation of general relativity using triangulated manifolds.



State sum model as a 3D TQFT of gravity

Regge showed that the curvature of a triangulated n dimensional manifold is encoded in the $(n - 2)$ skeleton.



The discrete form of the action S from which Einstein field equations are derived is given by

$$S_R = \sum |\sigma^i(T)| \varepsilon_i$$

State sum model as a 3D TQFT of gravity

In quantum mechanics the action is replaced by a **partition function Z** :

$$Z = \int e^{iS} D(x)$$


Given a manifold M , $Z(M)$ obtained from the TQFT will have the physical **meaning of the partition function**.

State sum invariants


In 1991 Turaev-Viro developed a 3 dimensional TQFT of gravity called the **state sum model**.

Definition


*Given a 3 dimensional manifold and a triangulation, a **state** is an assignment of an irreducible representation of $SU(2)$ to each edge of the triangulation. This are labeled by a non-half integer parameter j .*



State sum invariants



State sum invariants




Definition

The **weight** of the state is defined by

$$W = \prod_{\text{interior edges}} (-1)^{2j}(2j+1) \prod_{\text{interior triangles}} (-1)^{j_1+j_2+j_3} \prod_{\text{tetrahedra}} \begin{Bmatrix} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{Bmatrix}$$

State sum invariants



Definition

The **weight** of the state is defined by

$$W = \prod_{\text{interior edges}} (-1)^{2j} (2j + 1) \prod_{\text{interior triangles}} (-1)^{j_1 + j_2 + j_3} \prod_{\text{tetrahedra}} \begin{Bmatrix} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{Bmatrix}$$

The **state sum** (partition function) is obtained by summing over all values of the variables, subject to fixed values on the boundary.

$$\mathbf{Z}(M) = \Sigma W$$