

Crystal Symmetry Primer*

Kohei Shinohara[†]

February 26, 2023

Contents

0 Preliminary	3
1 Symmetry operation and space group	4
1.1 Affine group	4
1.2 Euclidean group	6
1.3 Symmetry operation and space group	7
1.4 Working examples from plane groups	7
2 Group theory primer	8
2.1 Definition	8
2.2 Abelian group	8
2.3 Isomorphism, subgroup	8
2.4 Coset decomposition, normal subgroup, factor group	8
2.5 Homomorphism, kernel, image	8
2.6 Conjugation	8
3 Group structure of space groups	9
3.1 Definition of space group	9
3.2 Point group	9
3.3 Vector system	9
3.4 Working examples from plane groups	10
4 Classification of space groups	11
4.1 Transformation matrix, origin shift	11
4.2 Affine space-group type and space-group type	11
4.3 Geometric crystal class	12
4.4 Bravais type of lattice and Bravais class	14
4.5 Arithmetic crystal class	15

*This document is licensed under [CC BY 4.0](#).

[†]<https://lan496.github.io/>

4.6	Lattice system	15
4.7	Crystal system	16
4.8	Crystal family	16
5	Conventions for space groups	18
5.1	Conventional cell	18
5.2	Hermann–Mauguin symbol	18
5.3	Standard settings in ITA	18
5.4	orthorhombic system	19
5.5	Hall symbol	19
6	Magnetic space group	21
6.1	Definition	21
6.2	BNS and OG symbols	21
6.3	Construction type of magnetic space group	21
6.4	Magnetic Hall symbol	21
7	Site symmetry group and normalizer	22
7.1	Action, orbit, stabilizer	22
7.2	Site symmetry group, Wyckoff position, asymmetric unit	22
7.3	Euclidean normalizer	22
7.4	Wyckoff set, equivalent descriptions of crystal structure	22
7.5	Affine normalizer	22
8	Lattice computation	23
8.1	Lattice, choice of basis vectors	23
8.2	Delaunay reduction	23
8.3	Sublattice, Hermite normal form	23
8.4	Smith normal form	23
	References	24

0 Preliminary

This document gives an introduction to crystallography on crystal symmetry and space groups intended for computational materials science. We first consider group theoretic treatments of crystal symmetry in Secs. 1 and 2. Then, we analyze a group structure of space groups in Sec. 3, and the classification of space groups in Sec. 4. There are many conventions to take a representative among “equivalent” objects. We introduce some of the conventions to avoid pitfalls in Sec. 5. The remaining sections are potpourris. In Sec. 6, we consider a group structure of magnetic space groups and conventions to represent them. In Sec. 7, we introduce normalizers of space groups, useful to treat the arbitrariness of transformation. In Sec. 8, we consider standard forms of integer matrices, which can be applied to supercell constructions and generalized regular meshes for Brillouin zone sampling.

There are already many impressive books and lectures covering the topic:

- M. I. Aroyo, editor. *International Tables for Crystallography*, volume A. International Union of Crystallography, December 2016
- M I Aroyo. *Teaching edition of international tables for crystallography - crystallographic symmetry, sixth edition*. IUCr Series. International Tables for Crystallography. John Wiley & Sons, Nashville, TN, 6 edition, May 2021
- Bernd Souvignier. Group theory applied to crystallography. https://www.math.ru.nl/~souvi/krist_09/cryst.pdf
- Ulrich Müller. *Symmetry relationships between crystal structures: applications of crystallographic group theory in crystal chemistry*, volume 18. OUP Oxford, 2013
- Michael Glazer, Gerald Burns, and Alexander N Glazer. *Space groups for solid state scientists*. Elsevier, 2012

In Japanese,

- Yuji Tachikawa, [物理数学 III \(2018\)](#)
- [対称性・群論トレーニングコース](#)
- 野田幸男, 結晶学と構造物性 入門から応用、実践まで (内田老鶴圃, 2017)

If you find typos or errors, please open an [issue](#) or [pull request](#).

1 Symmetry operation and space group

1.1 Affine group

1.1.1 Lattice

Definition 1.1 (lattice). *Lattice* L in \mathbb{R}^n is the set spanned by n independent vectors $\mathbf{a}_1, \dots, \mathbf{a}_n$ as

$$L := \left\{ \sum_{i=1}^n l_i \mathbf{a}_i \mid l_1, \dots, l_n \in \mathbb{Z} \right\}. \quad (1)$$

These vectors $\mathbf{a}_1, \dots, \mathbf{a}_n$ are called *lattice basis* of L .

We introduce the standard inner product in vector space \mathbb{R}^n . Then it is convenient to define the following matrix for calculating distances.

Definition 1.2 (metric tensor). Let $\mathbf{A} := (\mathbf{a}_1, \dots, \mathbf{a}_n)$ be basis vectors of a lattice L . A *metric tensor* of L is

$$\mathbf{G} := \mathbf{A}^\top \mathbf{A}. \quad (2)$$

We remark that metric tensor \mathbf{G} implicitly depends on the choice of basis vectors of L . When basis vectors are changed from \mathbf{A} to $\mathbf{A}\mathbf{P}$, the metric tensor are changed from $\mathbf{G} = \mathbf{A}^\top \mathbf{A}$ to $\mathbf{P}^\top \mathbf{G} \mathbf{P}$.

Be careful basis vectors \mathbf{A} are column-wise whereas programmers often use row-wise basis vectors.

```
import numpy as np
# Row-wise basis vectors
lattice = np.array([
    [ax, ay, az], # a axis
    [bx, by, bz], # b axis
    [cx, cy, cz], # c axis
])
# Fractional coordinates
frac_coords = np.array([
    [p0, q0, r0],
    [p1, q1, r2],
    ...
])
# Cartesian coordinates
cart_coords = np.dot(frac_coords, lattice)
```

1.1.2 Affine mapping

To display components of points, we fix some origin O and basis $\mathbf{a}_1, \dots, \mathbf{a}_n$ in \mathbb{R}^n . We define *affine space* by using O and $\mathbf{a}_1, \dots, \mathbf{a}_n$.

$$\mathbb{A}_n := \left\{ \begin{pmatrix} x_1 \\ \vdots \\ x_n \\ 1 \end{pmatrix} \mid x_1 \cdots, x_n \in \mathbb{R} \right\}. \quad (3)$$

Here, $(0, \dots, 0, 1)^\top \in \mathbb{A}_n$ corresponds to the origin O , and the i th components x_i corresponds to the basis \mathbf{a}_i .

Definition 1.3 (affine group). An *affine mapping* $\begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix}$ ($\mathbf{W} \in \text{GL}(n, \mathbb{R}), \mathbf{w} \in \mathbb{R}^n$) is a mapping that moves a point $\begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix} \in \mathbb{A}_n$ to

$$\begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix} = \begin{pmatrix} \mathbf{W}\mathbf{x} + \mathbf{w} \\ 1 \end{pmatrix} \in \mathbb{A}_n. \quad (4)$$

The matrix \mathbf{W} is called the *linear part* and the vector \mathbf{w} is called the *translation part*.

The *affine group* \mathcal{A}_n is the set of affine mappings

$$\mathcal{A}_n := \left\{ \begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix} \mid \mathbf{W} \in \text{GL}(n, \mathbb{R}), \mathbf{w} \in \mathbb{R}^n \right\}. \quad (5)$$

We often identify a point $\begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix} \in \mathbb{A}_n$ as $\mathbf{x} \in \mathbb{R}^n$ and write an affine mapping $\begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix}$ as (\mathbf{W}, \mathbf{w}) or $\{\mathbf{W} \mid \mathbf{w}\}$. Then the action of the affine mapping is compactly written as

$$(\mathbf{W}, \mathbf{w})\mathbf{x} := \mathbf{W}\mathbf{x} + \mathbf{w} \quad (6)$$

$$\{\mathbf{W} \mid \mathbf{w}\}\mathbf{x} := \mathbf{W}\mathbf{x} + \mathbf{w} \quad (7)$$

The notation for affine mapping in Eq. (6) is called *matrix-column pair*, and the one in Eq. (7) is called *Seitz symbol*¹. The original $(n+1) \times (n+1)$ matrix is also called an *augmented matrix*².

¹Do not write $(\mathbf{W} \mid \mathbf{w})$ or $\{\mathbf{W}, \mathbf{w}\}$!

²The augmented matrix is a 4×4 matrix in three dimensions. Because the fourth column is always constant, the column is often omitted in [computer vision](#).

1.1.3 Combination and inverse of affine mappings

Consider two affine mappings $(\mathbf{W}_1, \mathbf{w}_1)$ and $(\mathbf{W}_2, \mathbf{w}_2)$. Then $(\mathbf{W}_1, \mathbf{w}_1)$ maps \mathbf{x} to \mathbf{x}' and $(\mathbf{W}_2, \mathbf{w}_2)$ maps \mathbf{x}' to \mathbf{x}'' . We define a combination of $(\mathbf{W}_1, \mathbf{w}_1)$ and $(\mathbf{W}_2, \mathbf{w}_2)$ so that it maps \mathbf{x} to \mathbf{x}'' ,

$$\begin{aligned}\mathbf{x}' &= (\mathbf{W}_1, \mathbf{w}_1)\mathbf{x} = \mathbf{W}_1\mathbf{x} + \mathbf{w}_1 \\ \mathbf{x}'' &= (\mathbf{W}_2, \mathbf{w}_2)\mathbf{x}' = \mathbf{W}_2\mathbf{x}' + \mathbf{w}_2 \\ &= \mathbf{W}_2\mathbf{W}_1\mathbf{x} + \mathbf{W}_2\mathbf{w}_1 + \mathbf{w}_2\end{aligned}$$

$$\therefore (\mathbf{W}_2, \mathbf{w}_2)(\mathbf{W}_1, \mathbf{w}_1) := (\mathbf{W}_2\mathbf{W}_1, \mathbf{W}_2\mathbf{w}_1 + \mathbf{w}_2). \quad (8)$$

We define an inverse of $(\mathbf{W}_1, \mathbf{w}_1)$ so that it maps \mathbf{x}' to \mathbf{x} ,

$$\mathbf{x} = \mathbf{W}_1^{-1}\mathbf{x}' - \mathbf{W}_1^{-1}\mathbf{w}_1$$

$$\therefore (\mathbf{W}_1, \mathbf{w}_1)^{-1} := (\mathbf{W}_1^{-1}, -\mathbf{W}_1^{-1}\mathbf{w}_1). \quad (9)$$

These computations can be written as usual matrix operations,

$$\begin{pmatrix} \mathbf{W}_2 & \mathbf{w}_2 \\ \mathbf{0}^\top & 1 \end{pmatrix} \begin{pmatrix} \mathbf{W}_1 & \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{W}_2\mathbf{W}_1 & \mathbf{W}_2\mathbf{w}_1 + \mathbf{w}_2 \\ \mathbf{0}^\top & 1 \end{pmatrix} \quad (10)$$

$$\begin{pmatrix} \mathbf{W}_1 & \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix}^{-1} = \begin{pmatrix} \mathbf{W}_1^{-1} & -\mathbf{W}_1^{-1}\mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix}^{-1}. \quad (11)$$

1.2 Euclidean group

Definition 1.4 (isometry). Let \mathbf{G} be a metric tensor. An affine mapping (\mathbf{W}, \mathbf{w}) is called *isometry* if $\mathbf{W}^\top \mathbf{G} \mathbf{W} = \mathbf{G}$.

We remark that an isometry affine mapping does not change the distance between two points and vice versa.

$$\begin{aligned}\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n, \|\mathbf{A}(\mathbf{W}, \mathbf{w})\mathbf{x} - \mathbf{A}(\mathbf{W}, \mathbf{w})\mathbf{y}\| &= \|\mathbf{A}\mathbf{x} - \mathbf{A}\mathbf{y}\| \\ \iff \forall \mathbf{x} \in \mathbb{R}^n, \|\mathbf{A}\mathbf{W}\mathbf{x}\| &= \|\mathbf{A}\mathbf{x}\| \\ \iff \mathbf{A}\mathbf{W}\mathbf{A}^{-1} &\text{ is orthogonal} \\ \iff \mathbf{W}^\top \mathbf{G} \mathbf{W} &= \mathbf{G}\end{aligned}$$

Definition 1.5 (Euclidean group). The *Euclidean group* \mathcal{E}_n is the set of isometry affine mappings

$$\mathcal{E}_n := \left\{ \begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix} \mid \mathbf{W} \in \text{GL}(n, \mathbb{R}), \mathbf{W}^\top \mathbf{G} \mathbf{W} = \mathbf{G}, \mathbf{w} \in \mathbb{R}^n \right\}. \quad (12)$$

Definition 1.6 (Translation subgroup). The *translation subgroup* of \mathcal{A}_n is the set of affine mappings whose linear parts are identity,

$$\mathcal{T}_n := \left\{ \begin{pmatrix} \mathbf{I}_n & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix} \mid \mathbf{w} \in \mathbb{R}^n \right\}. \quad (13)$$

1.3 Symmetry operation and space group

A *symmetry operation* is an isometry affine mapping between two objects. For example, see Figs 4.1 and 4.2 of Ref. [4] for illustrations of symmetry operations in three dimensions.

Loosely speaking, a *space group* is a set of symmetry operations that preserve a three-dimensional crystal pattern. Similarly, a set of symmetry operations for a two-dimensional crystal pattern is called *plane group*. We consider a more rigorous definition of space groups in Sec. 3.1.

1.3.1 short-hand notation

TODO:

Note: hexagonal axis for $p3$

1.4 Working examples from plane groups

TODO:

Centering: cm

Symmorphic and nonsymmorphic: $p2mm$, $p2mg$, $p2gg$

2 Group theory primer

2.1 Definition

We mention that the affine group \mathcal{A}_n is group: The identity $\begin{pmatrix} I_n & \mathbf{0} \\ \mathbf{0}^\top & 1 \end{pmatrix}$ is in \mathcal{A}_n . Letting $\begin{pmatrix} \mathbf{W}_i & \mathbf{w}_i \\ \mathbf{0}^\top & 1 \end{pmatrix} \in \mathcal{A}_n (i = 1, 2)$, the product of two affine mappings is also affine mapping,

$$\begin{pmatrix} \mathbf{W}_1 & \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix} \begin{pmatrix} \mathbf{W}_2 & \mathbf{w}_2 \\ \mathbf{0}^\top & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{W}_1 \mathbf{W}_2 & \mathbf{W}_1 \mathbf{w}_2 + \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix} \in \mathcal{A}_n. \quad (14)$$

The inverse of an affine mapping is also an affine mapping,

$$\begin{pmatrix} \mathbf{W}_1 & \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix}^{-1} = \begin{pmatrix} \mathbf{W}_1^{-1} & -\mathbf{W}_1^{-1} \mathbf{w}_1 \\ \mathbf{0}^\top & 1 \end{pmatrix} \in \mathcal{A}_n. \quad (15)$$

2.2 Abelian group

translation subgroup

2.3 Isomorphism, subgroup

TODO: Euclidean group is a subgroup of the affine group.

2.4 Coset decomposition, normal subgroup, factor group

TODO: Translation subgroup is a normal subgroup of a space group

\mathbb{Z}_2 in \mathbb{Z}_4

2.5 Homomorphism, kernel, image

point group

2.6 Conjugation

3 Group structure of space groups

3.1 Definition of space group

Let f be a crystal pattern. A space group of f is defined as the stabilizer on \mathcal{E}_n .

Definition 3.1 (space group). The stabilizer of crystal pattern f on \mathcal{E}_n

$$\mathcal{G} := \text{Stab}_{\mathcal{E}_n}(f) = \{g \in \mathcal{E}_n \mid g \cdot f = f\} \quad (16)$$

is called *space group* if $\mathcal{T}(\mathcal{G}) := \mathcal{T}_n \cap \mathcal{G}$ is isomorphic to a n -dimensional lattice. For the space group \mathcal{G} , $\mathcal{T}(\mathcal{G})$ is called *translation subgroup* of \mathcal{G} .

3.2 Point group

Let \mathcal{G} be a space group. The translation subgroup $\mathcal{T}(\mathcal{G})$ is normal subgroup of \mathcal{G} : For any $(\mathbf{W}, \mathbf{w}) \in \mathcal{G}$ and $(\mathbf{I}_n, \mathbf{t}) \in \mathcal{T}(\mathcal{G})$, one has

$$(\mathbf{W}, \mathbf{w})(\mathbf{I}_n, \mathbf{t})(\mathbf{W}, \mathbf{w})^{-1} = (\mathbf{I}_n, \mathbf{W}\mathbf{t}) \in \mathcal{T}(\mathcal{G}). \quad (17)$$

Definition 3.2 (point group). Let \mathcal{G} be a space group. The set of the linear part of \mathcal{G} is called *point group* of \mathcal{G}

$$\mathcal{P} := \{\mathbf{W} \mid (\mathbf{W}, \mathbf{w}) \in \mathcal{G}\}. \quad (18)$$

The point group \mathcal{P} is isomorphic to the factor group $\mathcal{G}/\mathcal{T}(\mathcal{G})$.

We fix some basis vectors $(\mathbf{a}_1, \dots, \mathbf{a}_n)$ of affine space, and act all mappings in \mathcal{P} on the basis vectors. Since mapping in the point group \mathcal{P} is an isometry, it does not change the norm of vectors. Therefore, there are only finitely many acted vectors because the maximum norm of acted vectors is bound by $\max_{i=1, \dots, n} \mathbf{a}_i$.

Theorem 3.3. The point group \mathcal{P} of a space group \mathcal{G} is finite.

When we choose the lattice basis of $\mathcal{T}(\mathcal{G})$ as the basis vectors of the affine spaces, the translation subgroup $\mathcal{T}(\mathcal{G})$ is represented as $\{(\mathbf{I}_n, \mathbf{t}) \mid \mathbf{t} \in \mathbb{Z}^n\}$. And the action of point group \mathcal{P} is described as integer matrices.

3.3 Vector system

In general, a space group is not specified only with a point group and translation subgroup. We need to determine a translation part of each symmetry operation.

Definition 3.4 (vector system). Let \mathcal{P} be a point group of a space group \mathcal{G} . The map $\tau : \mathcal{P} \rightarrow \mathbb{R}^n$ is called a *vector system* if the map satisfies *cocycle condition*:

$$\tau(gh) = g\tau(h) + \tau(g) \pmod{\mathbb{Z}^n} \quad (\forall g, h \in \mathcal{P}). \quad (19)$$


We remark that we require the cocycle condition to have a vector system consistent with the product of two symmetry operations,

$$\begin{pmatrix} g & \tau(g) \\ \mathbf{0}^\top & 1 \end{pmatrix} \begin{pmatrix} h & \tau(h) \\ \mathbf{0}^\top & 1 \end{pmatrix} = \begin{pmatrix} gh & g\tau(h) + \tau(g) \\ \mathbf{0}^\top & 1 \end{pmatrix}$$

TODO: symmorphic and nonsymmorphic space groups

3.4 Working examples from plane groups

4 Classification of space groups



We classify space groups in three dimensions.

4.1 Transformation matrix, origin shift

When we change the origin and basis vectors for the affine space from O and $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$ to \mathbf{p} and $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)\mathbf{P}$, the components of point coordinates and affine mappings are also changed as follows.

The point coordinates \mathbf{x} is transformed to the new coordinates \mathbf{x}'

$$\begin{aligned}\mathbf{x}' &= (\mathbf{P}, \mathbf{p})^{-1} \mathbf{x} \\ &= \mathbf{P}^{-1}(\mathbf{x} - \mathbf{p}).\end{aligned}\tag{20}$$

An affine mapping (\mathbf{W}, \mathbf{w}) is transformed to $(\mathbf{W}', \mathbf{w}')$

$$\begin{aligned}(\mathbf{W}', \mathbf{w}') &= (\mathbf{P}, \mathbf{p})^{-1}(\mathbf{W}, \mathbf{w})(\mathbf{P}, \mathbf{p}) \\ &= \left(\mathbf{P}^{-1} \mathbf{W} \mathbf{P}, \mathbf{P}^{-1}(\mathbf{W} \mathbf{p} + \mathbf{w} - \mathbf{p}) \right).\end{aligned}\tag{21}$$

The metric tensor \mathbf{G} of the lattice spanned the basis vectors of the affine space is transformed to

$$\mathbf{G}' = \mathbf{P}^\top \mathbf{G} \mathbf{P}.\tag{22}$$

4.2 Affine space-group type and space-group type

It is natural to identify two space groups that are transformed into the another by changing coordinate systems.

Definition 4.1 (affine space-group type). Two space groups $\mathcal{G}, \mathcal{G}'$ belong to the same *affine space-group type* if they are conjugate by some affine mapping,

$$\exists(\mathbf{P}, \mathbf{p}) \in \mathcal{A}_3 \quad \text{s.t.} \quad (\mathbf{P}, \mathbf{p})^{-1} \mathcal{G}(\mathbf{P}, \mathbf{p}) = \mathcal{G}'. \quad (23)$$

These space groups are also called *affinely equivalent*.

It is nontrivial fact that affine equivalence completely identifies the isomorphism of space groups.

Theorem 4.2 (Bieberbach). Two space groups are isomorphic if and only if they belong to the same affine type.

In crystallography, a tighter classification of space groups is often used. That is, orientation-preserving affine mapping is only considered in transformations of coordinates systems ³.

Definition 4.3 (space-group type). Two space groups $\mathcal{G}, \mathcal{G}'$ belong to the same (*crystallographic*) *space-group type* if they are conjugate by some orientation-preserving affine mapping,

$$\exists(\mathbf{P}, \mathbf{p}) \in \mathcal{A}_3^+ \quad \text{s.t.} \quad (\mathbf{P}, \mathbf{p})^{-1} \mathcal{G}(\mathbf{P}, \mathbf{p}) = \mathcal{G}', \quad (24)$$

where orientation-preserving affine mapping is an affine mapping whose linear part is positive,

$$\mathcal{A}_n^+ := \left\{ \begin{pmatrix} \mathbf{W} & \mathbf{w} \\ \mathbf{0}^\top & 1 \end{pmatrix} \mid \mathbf{W} \in \text{GL}(n, \mathbb{R}), \det \mathbf{W} > 0, \mathbf{w} \in \mathbb{R}^n \right\}. \quad (25)$$

A pair of affine space-groups types that belong to the different crystallographic space-group types are called *enantiomorphic pair*. For example, there are 11 enantiomorphic pairs in three dimensions as Table. 1.

4.3 Geometric crystal class

We consider classification based on a point group of a space group.

³Some people indicate “space groups” as space-group types in their terminology. I think the clear distinction between space groups and space-group types makes our lives easier.

Table 1: Enantiomorphic pairs in space groups

$P4_1$ (76)	$P4_3$ (78)
$P4_122$ (91)	$P4_322$ (95)
$P4_12_12$ (92)	$P4_32_12$ (96)
$P3_1$ (144)	$P3_2$ (145)
$P3_112$ (151)	$P3_212$ (153)
$P3_121$ (152)	$P3_221$ (154)
$P6_1$ (169)	$P6_5$ (173)
$P6_2$ (170)	$P6_4$ (172)
$P6_122$ (178)	$P6_522$ (179)
$P6_222$ (180)	$P6_422$ (181)
$P4_332$ (212)	$P4_132$ (213)

Table 2: Laue classes in space groups

Laue class	Geometric crystal class
$\bar{1}$	1, $\bar{1}$
$2/m$	2, m , $2/m$
mmm	222, $2mm$, mmm
$\bar{3}$	3, $\bar{3}$
$\bar{3}m$	32, $3m$, $\bar{3}m$
$4/m$	4, $\bar{4}$, $4/m$
$4/mmm$	422, $\bar{4}2m$, $4mm$, $4/mmm$
$6/m$	6, $\bar{6}$, $6/m$
$6/mmm$	622, $\bar{6}2m$, $6mm$, $6/mmm$
$m\bar{3}$	23, $m\bar{3}$
$m\bar{3}m$	432, $\bar{4}32$, $m\bar{3}m$

Definition 4.4 (geometric crystal class). Two subgroups of $GL(3, \mathbb{Z})$, \mathcal{P} and \mathcal{P}' , belong to the same *geometric crystal class* if they are conjugate by some invertible matrix,

$$\exists \mathbf{P} \in GL(3, \mathbb{R}) \quad s.t. \quad \mathbf{P}^{-1} \mathcal{P} \mathbf{P} = \mathcal{P}'. \quad (26)$$

Two space groups \mathcal{G} and \mathcal{G}' with point groups \mathcal{P} and \mathcal{P}' , respectively, belong to the same *geometric crystal class* if \mathcal{P} and \mathcal{P}' belong to the same geometric crystal class.

Two space groups belong to the same *Laue class* if point groups obtained by their point group with inversions belong to the same geometric crystal class (Table 2).

TODO: $3m1$ and $31m$

4.4 Bravais type of lattice and Bravais class

We consider classification based on a translation lattice of a space group.

Definition 4.5 (translation lattice). A *translation lattice* L of a space group \mathcal{G} is a set of translation parts of translation subgroup of \mathcal{G} ,

$$L = \{ \mathbf{t} \mid (\mathbf{I}_3, \mathbf{t}) \in \mathcal{G} \}. \quad (27)$$

Definition 4.6 (Bravais group). A set of isometry mapping that pertains L is called *Bravais group of L* . Letting \mathbf{G} be a metric tensor of L , Bravais group of L is

$$\mathcal{B}(L) := \{ \mathbf{W} \in \text{GL}(3, \mathbb{Z}) \mid \mathbf{W}^\top \mathbf{G} \mathbf{W} = \mathbf{G} \}. \quad (28)$$

We mention that a Bravais group may change as a set by transforming primitive basis.

Definition 4.7 (Bravais type of lattice). Two lattices L and L' belong to the same *Bravais type of lattice* if their Bravais groups are conjugate by some unimodular matrix,

$$\exists \mathbf{P} \in \text{SL}(3, \mathbb{Z}) \quad \text{s.t.} \quad \mathbf{P}^{-1} \mathcal{B}(L) \mathbf{P} = \mathcal{B}(L'). \quad (29)$$

Remarkably, this definition of Bravais types of lattices is independent of choices for primitive basis.

Definition 4.8 (Bravais manifold). Let K be a subgroup of $\text{GL}(3, \mathbb{Z})$. A *space of metric tensors (Bravais manifold) of K* is the space of all metric tensors invariant with K ,

$$\mathbf{M}(K) := \{ \mathbf{G} \in \mathbb{R}_{\text{sym}}^{3 \times 3} \mid \mathbf{W}^\top \mathbf{G} \mathbf{W} = \mathbf{G} \quad (\forall \mathbf{W} \in K) \} \quad (30)$$

Let L and \mathcal{P} be a translation lattice and a point group of a space group \mathcal{G} , respectively. When the dimension of $\mathbf{M}(\mathcal{B}(L))$ is smaller than that of $\mathbf{M}(\mathcal{P})$, the translation lattice L is called to have *spacialized metric*.

TODO: Add example of specialized metric and Euclidean normalizer

Definition 4.9 (Bravais class). Let L be a lattice with a metric tensor \mathbf{G} . A space group \mathcal{G} with a point group \mathcal{P} and a translation lattice L belongs to a *Bravais class corresponding to the Bravais type of L* if $\mathbf{M}(\mathcal{P})$ and $\mathbf{M}(\mathcal{B}(L))$ are conjugate in $\text{SL}(3, \mathbb{Z})$,

$$\exists \mathbf{P} \in \text{SL}(3, \mathbb{Z}) \quad \text{s.t.} \quad \mathbf{P}^{-1} \mathbf{M}(\mathcal{P}) \mathbf{P} = \mathbf{M}(\mathcal{B}(L)). \quad (31)$$

TODO: Note that the definition of Bravais classes is independent of whether a translation lattice is a specialized metric or not.

Definition 4.10 (holohedry). A subgroup \mathcal{P} of $\text{GL}(3, \mathbb{Z})$ is called *holohedry* if there is a lattice L whose Bravais group belongs to the same geometric crystal class of \mathcal{P} .

TODO: limiting case from mP to oS

4.5 Arithmetic crystal class

Definition 4.11 (arithmetic crystal class). Two subgroups of $\text{GL}(3, \mathbb{Z})$, \mathcal{P} and \mathcal{P}' , belong to the same *arithmetic crystal class* if they are conjugate by some unimodular matrix,

$$\exists \mathbf{P} \in \text{SL}(3, \mathbb{R}) \quad s.t. \quad \mathbf{P}^{-1} \mathcal{P} \mathbf{P} = \mathcal{P}'. \quad (32)$$

Two space groups \mathcal{G} and \mathcal{G}' with point groups \mathcal{P} and \mathcal{P}' , respectively, belong to the same *arithmetic crystal class* if \mathcal{P} and \mathcal{P}' belong to the same arithmetic crystal class.

Definition 4.12 (Bravais arithmetic crystal class). The arithmetic crystal class of a space group \mathcal{G} is called a *Bravais arithmetic crystal class* if the point group of \mathcal{G} is the Bravais group of the translation lattice L of \mathcal{G} ,

$$\mathcal{P} = \mathcal{B}(L). \quad (33)$$

Note that Bravais arithmetic crystal classes are classifications for arithmetic crystal classes of space groups, and Bravais classes are classifications for space groups. The definition of Bravais arithmetic crystal classes is compatible with that of Bravais types of lattices (see Table 3).

TODO: Example: cm and pm

4.6 Lattice system

Definition 4.13 (lattice system). Two lattices L and L' belong to the same *lattice system* if their Bravais groups belong to the same geometric crystal class,

$$\exists \mathbf{P} \in \text{SL}(3, \mathbb{Z}) \quad s.t. \quad \mathbf{P}^{-1} \mathcal{B}(L) \mathbf{P} = \mathcal{B}(L'). \quad (34)$$

TODO: definition of lattice system on space groups

Table 4

Table 3: The correspondence of Bravais arithmetic crystal classes and Bravais types of lattices in three dimensions.

Bravais type of lattice	Bravais arithmetic crystal class
aP	$\bar{1}P$ (2)
mP	$2/mP$ (10)
mC	$2/mC$ (12)
oP	$mmmP$ (47)
oS	$mmmC$ (65)
oF	$mmmF$ (69)
oI	$mmmI$ (71)
tP	$4/mmmP$ (123)
tI	$4/mmmI$ (139)
hR	$\bar{3}mR$ (166)
hP	$6/mmmP$ (191)
cP	$m\bar{3}mP$ (221)
cF	$m\bar{3}mF$ (225)
cI	$m\bar{3}mI$ (229)

Table 4: Lattice systems in space groups

Lattice system	Holohedry	Bravais types of lattices
Triclinic	$\bar{1}$	aP
Monoclinic	$2/m$	mP, mS
Orthorhombic	mmm	oP, oS, oF, oI
Tetragonal	$4/mmm$	tP, tI
Rhombohedral	$\bar{3}m$	hR
Hexagonal	$6/mmm$	hP
Cubic	$m\bar{3}m$	cP, cF, cI

4.7 Crystal system

TODO:

Two point groups belong to the same crystal system if and only if the sets of Bravais type of lattices on which these point groups act coincide.

Table 5

4.8 Crystal family

Table 5: Crystal systems in space groups

Crystal system	Geometric crystal classes
Triclinic	$1, \bar{1}$
Monoclinic	$2/m, m, 2$
Orthorhombic	$mmm, mm2, 222$
Tetragonal	$4/mmm, \bar{4}2m, 4mm, 422, 4/m, \bar{4}, 4$
Hexagonal	$6/mmm, \bar{6}2m, 6mm, 622, 6/m, \bar{6}, 6$
Trigonal	$\bar{3}m, 3m, 32, \bar{3}, 3$
Cubic	$m\bar{3}m, \bar{4}3m, 432, m\bar{3}, 23$

Table 6: lattice symmetry directions in three-dimensional space (ITA Table 2.1.3.1)

Lattice system	Primary	Secondary	Tertiary
triclinic	None		
monoclinic (unique axis b)	[010]		
orthorhombic	[100]	[010]	[001]
tetragonal	[001]	$\langle 100 \rangle = [100][010]$	$\langle 1\bar{1}0 \rangle = [110][1\bar{1}0]$
rhombohedral(hexagonal axes)	[001]	$\langle 100 \rangle = [100][010][\bar{1}\bar{1}0]$	
rhombohedral(rhombohedral axes)	[111]	$\langle 1\bar{1}0 \rangle = [1\bar{1}0][01\bar{1}][\bar{1}01]$	
hexagonal	[001]	$\langle 100 \rangle = [100][010][\bar{1}\bar{1}0]$	$\langle 1\bar{1}0 \rangle = [1\bar{1}0][120][\bar{2}\bar{1}0]$
cubic	$\langle 001 \rangle = [100][010][001]$	$\langle 111 \rangle = [111][1\bar{1}\bar{1}][\bar{1}\bar{1}1]$	$\langle 110 \rangle = [1\bar{1}0][110][01\bar{1}][011][\bar{1}01][101]$

Table 7: settings of monoclinic system

unique axis setting	b abc	-b cba	c abc	-c ba\bar{c}	a abc	-a $\bar{a}bc$
transformation to abc	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
Cell choice 1	$C \rightarrow C$	$A \rightarrow C$	$A \rightarrow C$	$B \rightarrow C$	$B \rightarrow C$	$C \rightarrow C$
Cell choice 2	$A \rightarrow A$	$C \rightarrow A$	$B \rightarrow A$	$A \rightarrow A$	$C \rightarrow A$	$B \rightarrow A$
Cell choice 3	$I \rightarrow I$	$I \rightarrow I$	$I \rightarrow I$	$I \rightarrow I$	$I \rightarrow I$	$I \rightarrow I$

5 Conventions for space groups

5.1 Conventional cell

5.2 Hermann–Mauguin symbol

Convention for Hermann-Mauguin symbols is shown in Table 6.

TODO: How to read H-M symbols

5.3 Standard settings in ITA

5.3.1 Monoclinic system(ITA-2.1.3.15)

TODO: Fix unique axes

Axis settings for monoclinic systems are shown in Table 7. Cell choices for monoclinic systems are shown in Table 8 and Fig. 1.

Table 8: transformation of cell choices in monoclinic system

choice	<i>b</i> 1	<i>b</i> 2	<i>b</i> 3
transformation to <i>b</i> 1	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & -1 \end{pmatrix}$
centering	$C \rightarrow C$	$A \rightarrow C$	$I \rightarrow C$




Figure 1: choices of monoclinic system

5.4 orthorhombic system

Axis settings for orthorhombic systems are shown in Table 9.

5.5 Hall symbol

Table 9: settings of orthorhombic system

settings	abc	ba\bar{c}	cab	$\bar{c}ba$	bca	a$\bar{c}b$
transformation to abc	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$
centering	$A \rightarrow A$ $B \rightarrow B$ $C \rightarrow C$	$A \rightarrow B$ $B \rightarrow A$ $C \rightarrow C$	$A \rightarrow C$ $B \rightarrow A$ $C \rightarrow B$	$A \rightarrow C$ $B \rightarrow B$ $C \rightarrow A$	$A \rightarrow B$ $B \rightarrow C$ $C \rightarrow A$	$A \rightarrow A$ $B \rightarrow C$ $C \rightarrow B$

6 Magnetic space group

6.1 Definition

6.2 BNS and OG symbols

6.3 Construction type of magnetic space group

6.4 Magnetic Hall symbol

7 Site symmetry group and normalizer

7.1 Action, orbit, stabilizer

7.2 Site symmetry group, Wyckoff position, asymmetric unit

7.3 Euclidean normalizer

7.4 Wyckoff set, equivalent descriptions of crystal structure

7.5 Affine normalizer

8 Lattice computation

8.1 Lattice, choice of basis vectors

The choice of basis vectors is not unique for a given lattice. A lattice spanned by basis vectors $(\mathbf{a}_1, \dots, \mathbf{a}_n)$ and a lattice spanned by

$$(\mathbf{a}'_1, \dots, \mathbf{a}'_n) := (\mathbf{a}_1, \dots, \mathbf{a}_n) \mathbf{P} \quad (\mathbf{P} \in \mathbb{R}^{n \times n})$$

coincide if and only if $\mathbf{P} \in \text{SL}(n, \mathbb{Z})$, where $\text{SL}(n, \mathbb{Z})$ is the set of $n \times n$ unimodular matrices.

8.2 Delaunay reduction

8.3 Sublattice, Hermite normal form

8.4 Smith normal form

References

- [1] M. I. Aroyo, editor. *International Tables for Crystallography*, volume A. International Union of Crystallography, December 2016.
- [2] M I Aroyo. *Teaching edition of international tables for crystallography - crystallographic symmetry, sixth edition*. IUCr Series. International Tables for Crystallography. John Wiley & Sons, Nashville, TN, 6 edition, May 2021.
- [3] Bernd Souvignier. Group theory applied to crystallography. https://www.math.ru.nl/~souvi/krist_09/cryst.pdf.
- [4] Ulrich Müller. *Symmetry relationships between crystal structures: applications of crystallographic group theory in crystal chemistry*, volume 18. OUP Oxford, 2013.
- [5] Michael Glazer, Gerald Burns, and Alexander N Glazer. *Space groups for solid state scientists*. Elsevier, 2012.