

Contents

1. The canonical and isobaric ensembles via molecular dynamics	1
1.1. Classical non-Hamiltonian statistical mechanics	1
1.2. Matrix exponential	2
1.3. Canonical ensembles	2
1.3.1. Nosé-Hoover chain	2
1.3.2. Conserved quantities of Nosé-Hoover chain equations	2
1.3.3. Microcanonical distribution given by Nosé-Hoover chain equations	3
1.3.4. Integrating the Nosé-Hoover chain equations	3
1.4. The isobaric ensembles	4
1.4.1. Instantaneous stress tensor	4
1.4.2. Isotropic volume fluctuations	5
1.4.2.1. MTK equations for isotropic volume fluctuations	5
1.4.2.2. Conserved quantities of MTK equations	5
1.4.2.3. Microcanonical distribution given by MTK equations	6
1.4.2.4. Integrating the MTK equations for isotropic volume fluctuations	6
1.4.3. Anisotropic cell fluctuations	7
1.4.3.1. MTK equations for anisotropic cell fluctuations	7
1.4.3.2. Conserved quantities of anisotropic MTK equations	9
1.4.3.3. Microcanonical distribution given by anisotropic MTK equations	9
1.4.3.4. Integrating the MTK equations for anisotropic cell fluctuations	9
1.4.3.5. Actions of $e^{iL_1\Delta t}$ and $e^{iL_2\frac{\Delta t}{2}}$	11
1.4.3.6. Action of $e^{iL_{g,1}\Delta t}$	12
1.4.4. Cell fluctuations with mask	12
1.4.4.1. MTK equations	12
1.4.4.2. Conserved quantities	13
1.4.4.3. Integrating the MTK equations	13
Bibliography	13

1. The canonical and isobaric ensembles via molecular dynamics

1.1. Classical non-Hamiltonian statistical mechanics

Consider a dynamical system $\dot{x} = \xi(x)$. We define the compressibility as

$$\kappa(x_t) := \nabla_{x_t} \cdot \dot{x}_t =: \frac{d}{dt} w(x_t). \quad (1)$$

With $\sqrt{g(x)} := e^{-w(x)}$, generalized Liouville's Theorem states that

$$\sqrt{g(x_t)} dx_t = \sqrt{g(x_0)} dx_0. \quad (2)$$

The Liouville operator is defined as

$$i\mathcal{L} := \boldsymbol{\xi}(x) \cdot \nabla_x. \quad (3)$$

Since $\dot{x}_t = iLx_t$, we formally write

$$x_t = e^{it\mathcal{L}}x_0. \quad (4)$$

1.2. Matrix exponential

For square matrices A and B , we define the 2nd order integrator as

$$\begin{aligned} S_2(\lambda) &:= e^{\frac{\lambda}{2}B} e^{\lambda A} e^{\frac{\lambda}{2}B} \\ S_2(\lambda) &= e^{\lambda(A+B)} + O(\lambda^3). \end{aligned} \quad (5)$$

The fourth order integrator [1]

$$\begin{aligned} S_4(\lambda) &:= S_2(x_3\lambda)S_2(x_2\lambda)S_2(x_1\lambda) \\ x_1 = x_3 &:= \frac{1}{2 - 2^{\frac{1}{3}}} \\ x_2 &:= -\frac{2^{\frac{1}{3}}}{2 - 2^{\frac{1}{3}}} \\ S_4(\lambda) &= e^{\lambda(A+B)} + O(\lambda^5). \end{aligned} \quad (6)$$

1.3. Canonical ensembles

1.3.1. Nosé-Hoover chain

Nosé-Hoover chain equations:

$$\begin{aligned} \dot{r}_i &= \frac{p_i}{m_i} \\ \dot{p}_i &= F_i - \frac{p_{\eta_1}}{Q_1} p_i \\ \dot{\eta}_j &= \frac{p_{\eta_j}}{Q_j} \quad (j = 1, \dots, M) \\ \dot{p}_{\eta_1} &= \sum_{i=1}^N \frac{p_i^2}{m_i} - dNkT - \frac{p_{\eta_2}}{Q_2} p_{\eta_1} \\ \dot{p}_{\eta_j} &= \frac{p_{\eta_{j-1}}^2}{Q_{j-1}} - kT - \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \quad (j = 2, \dots, M-1) \\ \dot{p}_{\eta_M} &= \frac{p_{\eta_{M-1}}^2}{Q_{M-1}} - kT \end{aligned} \quad (7)$$

Ref. [2] suggests

$$\begin{aligned} Q_1 &= dNkT\tau^2 \\ Q_j &= kT\tau^2 \quad (j = 2, \dots, M). \end{aligned} \quad (8)$$

1.3.2. Conserved quantities of Nosé-Hoover chain equations

Nosé-Hoover chain equations conserves

$$H' := \mathcal{H}(r^N, p^N) + \sum_{j=1}^M \frac{p_{\eta_j}^2}{2Q_j} + dNkT\eta_1 + kT\eta_c. \quad (9)$$

where $\eta_c := \sum_{j=2}^M \eta_j$. In the absence of external forces ($\sum_i \mathbf{F}_i = 0$), the center-of-mass momentum $\mathbf{P} := \sum_i \mathbf{p}_i$ also has a conserved quantity

$$\mathbf{K} := \mathbf{P}e^{\eta_1}. \quad (10)$$

1.3.3. Microcanonical distribution given by Nosé-Hoover chain equations

The EOM gives the compressibility

$$\kappa(\mathbf{x}) = -dN\dot{\eta}_1 - \dot{\eta}_c. \quad (11)$$

Let us consider the distribution function traced over the extended variables,

$$\begin{aligned} Z_{V,T}(r^N, p^N) &:= \int d\mathbf{p}_\eta^M d\eta^M e^{dN\eta_1 + \eta_c} \delta\left(\mathcal{H}(r^N, p^N) + \sum_{j=1}^M \frac{p_{\eta_j}^2}{2Q_j} + dNkT\eta_1 + kT\eta_c - H'\right) \\ &= \left(\prod_{j=1}^M \sqrt{2\pi kTQ_j}\right) \exp\left(-\frac{1}{kT}(\mathcal{H}(r^N, p^N) - H')\right) \int d\eta_1 \\ &\propto \exp\left(-\frac{1}{kT}\mathcal{H}(r^N, p^N)\right). \end{aligned} \quad (12)$$

This shows that the Nosé-Hoover chain equations generates the canonical ensemble in the physical phase space.

1.3.4. Integrating the Nosé-Hoover chain equations

The Liouville operator

$$\begin{aligned} iL &:= iL_{\text{NHC}} + iL_1 + iL_2 \\ iL_1 &:= \sum_{i=1}^N \frac{\mathbf{p}_i}{m_i} \cdot \frac{\partial}{\partial \mathbf{r}_i} \\ iL_2 &:= \sum_{i=1}^N \mathbf{F}_i \cdot \frac{\partial}{\partial \mathbf{p}_i} \\ iL_{\text{NHC}} &:= -\sum_{i=1}^N \frac{p_{\eta_1}}{Q_1} \mathbf{p}_i \cdot \frac{\partial}{\partial \mathbf{p}_i} + \sum_{j=1}^M \frac{p_{\eta_j}}{Q_j} \frac{\partial}{\partial \eta_j} + \sum_{j=1}^{M-1} \left(G_j - p_{\eta_j} \frac{p_{\eta_{j+1}}}{Q_{j+1}}\right) \frac{\partial}{\partial p_{\eta_j}} + G_M \frac{\partial}{\partial p_{\eta_M}} \\ G_1 &:= \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} - dNkT \\ G_j &:= \frac{p_{\eta_{j-1}}^2}{Q_{j-1}} - kT \quad (j = 2, \dots, M) \end{aligned} \quad (13)$$

$$e^{iL\Delta t} = e^{iL_{\text{NHC}}\frac{\Delta t}{2}} e^{iL_2\frac{\Delta t}{2}} e^{iL_1\Delta t} e^{iL_2\frac{\Delta t}{2}} e^{iL_{\text{NHC}}\frac{\Delta t}{2}} + O(\Delta t^3). \quad (14)$$

$$e^{iL_1\Delta t} \begin{pmatrix} \mathbf{r}_i \\ \mathbf{p}_i \\ \eta_j \\ \mathbf{p}_{\eta_j} \end{pmatrix} = \begin{pmatrix} \mathbf{r}_i + \frac{\mathbf{p}_i}{m_i}\Delta t \\ \mathbf{p}_i \\ \eta_j \\ \mathbf{p}_{\eta_j} \end{pmatrix} \quad (15)$$

$$e^{iL_2\frac{\Delta t}{2}} \begin{pmatrix} \mathbf{r}_i \\ \mathbf{p}_i \\ \eta_j \\ \mathbf{p}_{\eta_j} \end{pmatrix} = \begin{pmatrix} \mathbf{r}_i \\ \mathbf{p}_i + \mathbf{F}_i\frac{\Delta t}{2} \\ \eta_j \\ \mathbf{p}_{\eta_j} \end{pmatrix} \quad (16)$$

$$e^{iL_{\text{NHC}}\frac{\Delta t}{2}} = \left(e^{iL_{\text{NHC}}\frac{\Delta t}{2n}}\right)^n$$

$$e^{iL_{\text{NHC}}\frac{\Delta t}{2n}} = S_4^{\text{NHC}}\left(\frac{\Delta t}{2n}\right) + O\left(\left(\frac{\Delta t}{n}\right)^5\right)$$

$$S_4^{\text{NHC}}\left(\frac{\Delta t}{2n}\right) := \prod_{\alpha=1}^3 S_2^{\text{NHC}}\left(x_\alpha \frac{\Delta t}{2n}\right) \quad \left(\delta_\alpha := x_\alpha \frac{\Delta t}{2n}\right)$$

$$\begin{aligned} S_2^{\text{NHC}}(\delta_\alpha) &:= \exp\left(\frac{\delta_\alpha}{2} G_M \frac{\partial}{\partial \mathbf{p}_{\eta_M}}\right) \\ &\times \prod_{j=M-1}^1 \left(\exp\left(-\frac{\delta_\alpha}{4} \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \frac{\partial}{\partial p_{\eta_j}}\right) \exp\left(\frac{\delta_\alpha}{2} G_j \frac{\partial}{\partial p_{\eta_j}}\right) \exp\left(-\frac{\delta_\alpha}{4} \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \frac{\partial}{\partial p_{\eta_j}}\right) \right) \\ &\times \prod_{i=1}^N \exp\left(-\delta_\alpha \frac{p_{\eta_1}}{Q_1} \mathbf{p}_i \cdot \frac{\partial}{\partial \mathbf{p}_i}\right) \\ &\times \prod_{j=1}^M \exp\left(\delta_\alpha \frac{p_{\eta_j}}{Q_j} \frac{\partial}{\partial \eta_j}\right) \\ &\times \prod_{j=1}^{M-1} \left(\exp\left(-\frac{\delta_\alpha}{4} \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \frac{\partial}{\partial p_{\eta_j}}\right) \exp\left(\frac{\delta_\alpha}{2} G_j \frac{\partial}{\partial p_{\eta_j}}\right) \exp\left(-\frac{\delta_\alpha}{4} \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \frac{\partial}{\partial p_{\eta_j}}\right) \right) \\ &\times \exp\left(\frac{\delta_\alpha}{2} G_M \frac{\partial}{\partial p_{\eta_M}}\right) \end{aligned}$$

Here, we use the following formula to evaluate the actions,

$$\exp\left(cx \frac{\partial}{\partial x}\right) f(x) = f(xe^c). \quad (18)$$

1.4. The isobaric ensembles

Refs. [3]

1.4.1. Instantaneous stress tensor

Let \mathbf{h} be right-handed row-major basis vectors.

Instantaneous stress tensor

$$\mathcal{P}_{\alpha\beta}^{\text{int}} = \frac{1}{\det \mathbf{h}} \sum_{i=1}^N \left[\frac{p_{i\alpha} p_{i\beta}}{m_i} + F_{i\alpha} r_{i\beta} \right] - \frac{1}{\det \mathbf{h}} \sum_{\gamma=1}^3 \frac{\partial U}{\partial h_{\alpha\gamma}} h_{\gamma\beta}. \quad (19)$$

1.4.2. Isotropic volume fluctuations

1.4.2.1. MTK equations for isotropic volume fluctuations

MTK equations:

$$\begin{aligned} \dot{\mathbf{r}}_i &= \frac{\mathbf{p}_i}{m_i} + \frac{p_\varepsilon}{W} \mathbf{r}_i \\ \dot{\mathbf{p}}_i &= \tilde{\mathbf{F}}_i - \left(1 + \frac{d}{N_f} \right) \frac{p_\varepsilon}{W} \mathbf{p}_i - \frac{p_{\eta_1}}{Q_1} \mathbf{p}_i \\ \dot{V} &= \frac{dV}{W} p_\varepsilon \\ \dot{p}_\varepsilon &= dV(\mathcal{P}^{\text{int}} - P) + \frac{d}{N_f} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} - \frac{p_{\xi_1}}{Q'_1} p_\varepsilon \\ \dot{\eta}_j &= \frac{p_{\eta_j}}{Q_j} \\ \dot{\xi}_j &= \frac{p_{\xi_j}}{Q'_j} \\ \dot{p}_{\eta_j} &= G_j - \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \quad (j = 1, \dots, M-1) \\ \dot{p}_{\eta_M} &= G_M \\ \dot{p}_{\xi_j} &= G'_j - \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \quad (j = 1, \dots, M-1) \\ \dot{p}_{\xi_M} &= G'_M \end{aligned} \quad (20)$$

where

$$\begin{aligned} G'_1 &:= \frac{p_\varepsilon^2}{W} - kT \\ G'_j &:= \frac{p_{\xi_{j-1}}^2}{Q'_{j-1}} - kT \quad (j = 2, \dots, M). \end{aligned} \quad (21)$$

Ref. [3] suggests to set

$$\begin{aligned} W &= (N_f + d) kT \tau^2 \\ Q'_1 &= d^2 kT \tau^2 \\ Q'_j &= kT \tau^2 \quad (j = 2, \dots, M), \end{aligned} \quad (22)$$

where τ is a characteristic time scale for barostat.

1.4.2.2. Conserved quantities of MTK equations

The conserved energy

$$H' := \mathcal{H}(r, p) + \frac{p_\varepsilon^2}{2W} + PV + \sum_{j=1}^M \left(\frac{p_{\eta_j}^2}{2Q_j} + \frac{p_{\xi_j}^2}{2Q'_j} + kT\xi_j \right) + N_f kT\eta_1 + kT\eta_c, \quad (23)$$

where $\eta_c := \sum_{j=2}^M \eta_j$.

In the absence of external forces ($\sum_i \mathbf{F}_i = 0$), the center-of-mass momentum $\mathbf{P} := \sum_i \mathbf{p}_i$ also has a conserved quantity

$$\mathbf{K} := \mathbf{P} \exp \left(\left(1 + \frac{d}{N_f} \right) \varepsilon + \eta_1 \right), \quad (24)$$

where $\varepsilon := \frac{1}{d} \log \frac{V}{V_0}$ with some reference volume V_0 .

1.4.2.3. Microcanonical distribution given by MTK equations

Let $Q(N, V, T)$ be a partition function for NVT ensemble. The partition function for isotropic-NPT ensemble is defined as

$$\Delta(N, P, T) := \frac{1}{V_0} \int dV e^{-\beta PV} Q(N, V, T). \quad (25)$$

The EOM gives the compressibility

$$\kappa(\mathbf{x}) = -\frac{d}{dt}(dN\eta_1 + \eta_c + \xi_1 + \xi_c) \quad (26)$$

$$\therefore \sqrt{g(\mathbf{x})} = e^{dN\eta_1 + \eta_c + \xi_1 + \xi_c},$$

where $\xi_c := \sum_{j=2}^M \xi_j$. Because this metric $\sqrt{g(\mathbf{x})}d\mathbf{x}$ does not depends on physical variables, the extended variables are traced out in $\Delta(N, P, T)$.

1.4.2.4. Integrating the MTK equations for isotropic volume fluctuations

$$iL := iL_1 + iL_2 + iL_{\varepsilon,1} + iL_{\varepsilon,2} + iL_{\text{NHC-baro}} + iL_{\text{NHC-thermo}}$$

$$iL_1 := \sum_{i=1}^N \left(\frac{\mathbf{p}_i}{m_i} + \frac{p_\varepsilon}{W} \mathbf{r}_i \right) \cdot \frac{\partial}{\partial \mathbf{r}_i}$$

$$iL_2 := \sum_{i=1}^N \left(\tilde{\mathbf{F}}_i - \left(1 + \frac{d}{N_f} \right) \frac{p_\varepsilon}{W} \mathbf{p}_i \right) \cdot \frac{\partial}{\partial \mathbf{p}_i}$$

$$iL_{\varepsilon,1} := \frac{p_\varepsilon}{W} \frac{\partial}{\partial \varepsilon} \quad (27)$$

$$iL_{\varepsilon,2} := G_\varepsilon \frac{\partial}{\partial p_\varepsilon}$$

$$iL_{\text{NHC-thermo}} := -\sum_{i=1}^N \frac{p_{\eta_1}}{Q_1} \mathbf{p}_i \cdot \frac{\partial}{\partial \mathbf{p}_i} + \sum_{j=1}^M \frac{p_{\eta_j}}{Q_j} \frac{\partial}{\partial \eta_j} + \sum_{j=1}^{M-1} \left(G_j - p_{\eta_j} \frac{p_{\eta_{j+1}}}{Q_{j+1}} \right) \frac{\partial}{\partial p_{\eta_j}} + G_M \frac{\partial}{\partial p_{\eta_M}}$$

$$iL_{\text{NHC-baro}} := -\frac{p_{\xi_1}}{Q'_1} p_\varepsilon \frac{\partial}{\partial p_\varepsilon} + \sum_{j=1}^M \frac{p_{\xi_j}}{Q'_j} \frac{\partial}{\partial \xi_j} + \sum_{j=1}^{M-1} \left(G'_j - p_{\xi_j} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} \right) \frac{\partial}{\partial p_{\xi_j}} + G'_M \frac{\partial}{\partial p_{\xi_M}}$$

with

$$\varepsilon := \frac{1}{d} \ln \frac{V}{V_0}$$

$$\mathcal{P}^{\text{int}} := \frac{1}{d} \text{Tr}[\mathcal{P}^{\text{int}}] = \frac{1}{dV} \sum_{i=1}^N \left(\frac{\mathbf{p}_i^2}{m_i} + \mathbf{r}_i \cdot \mathbf{F}_i \right) \quad (28)$$

$$G_\varepsilon := dV(\mathcal{P}^{\text{int}} - P) + \frac{1}{N} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i}.$$

$$\begin{aligned} e^{iL\Delta t} &= e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}} e^{iL_{\text{NHC-thermo}} \frac{\Delta t}{2}} \\ &\quad \times e^{iL_{\varepsilon,2} \frac{\Delta t}{2}} e^{iL_2 \frac{\Delta t}{2}} \\ &\quad \times e^{iL_{\varepsilon,1} \Delta t} e^{iL_1 \Delta t} \\ &\quad \times e^{iL_2 \frac{\Delta t}{2}} e^{iL_{\varepsilon,2} \frac{\Delta t}{2}} \\ &\quad \times e^{iL_{\text{NHC-thermo}} \frac{\Delta t}{2}} e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}} + O(\Delta t^3). \end{aligned} \quad (29)$$

$$\begin{aligned} e^{iL_1 \Delta t} \mathbf{r}_i &= \exp \left(\Delta t \left(\frac{\mathbf{p}_i}{m_i} + \frac{p_\varepsilon}{W} \mathbf{r}_i \right) \cdot \frac{\partial}{\partial \mathbf{r}_i} \right) \mathbf{r}_i \\ &= \mathbf{r}_i e^{\frac{p_\varepsilon \Delta t}{W}} + \Delta t \frac{\mathbf{p}_i}{m_i} \text{exprel} \left(\frac{p_\varepsilon \Delta t}{W} \right) \end{aligned} \quad (30)$$

$$\begin{aligned} e^{iL_2 \frac{\Delta t}{2}} \mathbf{p}_i &= \exp \left(\frac{\Delta t}{2} \left(\tilde{\mathbf{F}}_i - \left(1 + \frac{d}{N_f} \right) \frac{p_\varepsilon}{W} \mathbf{p}_i \right) \cdot \frac{\partial}{\partial \mathbf{p}_i} \right) \mathbf{p}_i \\ &= \mathbf{p}_i e^{-\frac{\kappa \Delta t}{2W}} + \frac{\Delta t}{2} \tilde{\mathbf{F}}_i \text{exprel} \left(-\frac{\kappa \Delta t}{2W} \right) \end{aligned} \quad (31)$$

$$\kappa := \left(1 + \frac{d}{N_f} \right) p_\varepsilon$$

1.4.3. Anisotropic cell fluctuations

1.4.3.1. MTK equations for anisotropic cell fluctuations

MTK equations:

$$\begin{aligned}
\dot{\mathbf{r}}_i &= \frac{\mathbf{p}_i}{m_i} + \frac{\mathbf{p}_g}{W_g} \mathbf{r}_i \\
\dot{\mathbf{p}}_i &= \tilde{\mathbf{F}}_i - \left(\mathbf{p}_g + \frac{\text{Tr}[\mathbf{p}_g] \mathbf{I}}{N_f} \right) \frac{\mathbf{p}_i}{W_g} - \frac{p_{\eta_1}}{Q_1} \mathbf{p}_i \\
\dot{\mathbf{h}} &= \frac{\mathbf{h} \mathbf{p}_g}{W_g} \\
\dot{\mathbf{p}}_g &= \det[\mathbf{h}] (\mathcal{P}^{\text{int}} - P \mathbf{I}) + \frac{1}{N_f} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} \mathbf{I} - \frac{p_{\xi_1}}{Q'_1} \mathbf{p}_g \\
\dot{\eta}_j &= \frac{p_{\eta_j}}{Q_j} \\
\dot{\xi}_j &= \frac{p_{\xi_j}}{Q'_j} \\
\dot{p}_{\eta_j} &= G_j - \frac{p_{\eta_{j+1}}}{Q_{j+1}} p_{\eta_j} \quad (j = 1, \dots, M-1) \\
\dot{p}_{\eta_M} &= G_M \\
\dot{p}_{\xi_j} &= G'_j - \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \quad (j = 1, \dots, M-1) \\
\dot{p}_{\xi_M} &= G'_M
\end{aligned} \tag{32}$$

where

$$\begin{aligned}
G_1 &:= \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} - N_f kT \\
G_j &:= \frac{p_{\eta_{j-1}}^2}{Q_{j-1}} - kT \quad (j = 2, \dots, M) \\
G'_1 &:= \frac{\text{Tr}[\mathbf{p}_g^\top \mathbf{p}_g]}{W_g} - d^2 kT \\
G'_j &:= \frac{p_{\xi_{j-1}}^2}{Q'_{j-1}} - kT \quad (j = 2, \dots, M)
\end{aligned} \tag{33}$$

Ref. [3] suggests to set

$$\begin{aligned}
W_g &= \frac{N_f + d}{d} kT \tau^2 \\
Q'_1 &= d^2 kT \tau^2 \\
Q'_j &= kT \tau^2 \quad (j = 2, \dots, M),
\end{aligned} \tag{34}$$

where τ is a characteristic time scale for barostat. d^2 in Q'_1 should be substituted with the degree of freedoms in basis vectors.

1.4.3.2. Conserved quantities of anisotropic MTK equations

The conserved energy

$$\begin{aligned}
H' := & \mathcal{H}(r, p) + \frac{\text{Tr}[\mathbf{p}_g^\top \mathbf{p}_g]}{2W_g} + P \det[\mathbf{h}] \\
& + \sum_{j=1}^M \left(\frac{p_{\eta_j}^2}{2Q_j} + \frac{p_{\xi_j}^2}{2Q'_j} \right) + N_f kT \eta_1 + d^2 kT \xi_1 + kT \sum_{j=2}^M (\eta_j + \xi_j)
\end{aligned} \tag{35}$$

In the absence of external forces ($\sum_i \mathbf{F}_i = 0$), the center-of-mass momentum $\mathbf{P} := \sum_i \mathbf{p}_i$ also has a conserved quantity

$$\mathbf{K} := \mathbf{h} \mathbf{P} (\det \mathbf{h})^{1/N_f} e^{\eta_1}. \tag{36}$$

1.4.3.3. Microcanonical distribution given by anisotropic MTK equations

Let $Q(N, V, T)$ be a partition function for NPT ensemble. The partition function for anisotropic-NPT ensemble is defined as

$$\begin{aligned}
\Delta(N, P, T) &:= \frac{1}{V_0} \int dV e^{-\beta PV} \int d\mathbf{h}_0 Q(N, V\mathbf{h}_0, T) \delta(\det \mathbf{h}_0 - 1) \\
&= \frac{1}{V_0} \int dV \int d\mathbf{h} V^{1-d} e^{-\beta PV} Q(N, \mathbf{h}, T) \delta(\det \mathbf{h} - V)
\end{aligned} \tag{37}$$

The EOM gives the compressibility

$$\begin{aligned}
\kappa(\mathbf{x}) &= -\frac{d}{dt} ((1-d) \ln \det \mathbf{h} + dN\eta_1 + \eta_c + d^2\xi_1 + \xi_c) \\
\therefore \sqrt{g(\mathbf{x})} &= (\det \mathbf{h})^{1-d} e^{dN\eta_1 + \eta_c + d^2\xi_1 + \xi_c}.
\end{aligned} \tag{38}$$

Because this metric $\sqrt{g(\mathbf{x})}d\mathbf{x}$ depends on physical variables as same as the integral in $\Delta(N, P, T)$, the extended variables are traced out in $\Delta(N, P, T)$.

1.4.3.4. Integrating the MTK equations for anisotropic cell fluctuations

Ref. [4]

$$\begin{aligned}
iL &:= iL_1 + iL_2 + iL_{g,1} + iL_{g,2} + iL_{\text{NHC-baro}} + iL_{\text{NHC-thermo}} \\
iL_1 &:= \sum_{i=1}^N \left(\frac{\mathbf{p}_i}{m_i} + \frac{\mathbf{p}_g}{W_g} \mathbf{r}_i \right) \cdot \frac{\partial}{\partial \mathbf{r}_i} \\
iL_2 &:= \sum_{i=1}^N \left(\tilde{\mathbf{F}}_i - \left(\mathbf{p}_g + \frac{\text{Tr}[\mathbf{p}_g]}{N_f} \mathbf{I} \right) \frac{\mathbf{p}_i}{W_g} \right) \cdot \frac{\partial}{\partial \mathbf{p}_i} \\
iL_{g,1} &:= \frac{\mathbf{h} \mathbf{p}_g}{W_g} \odot \frac{\partial}{\partial \mathbf{h}} \\
iL_{g,2} &:= \mathbf{G}_g \odot \frac{\partial}{\partial \mathbf{p}_g} \\
iL_{\text{NHC-thermo}} &:= - \sum_{i=1}^N \frac{p_{\eta_i}}{Q_1} \mathbf{p}_i \cdot \frac{\partial}{\partial \mathbf{p}_i} + \sum_{j=1}^M \frac{p_{\eta_j}}{Q_j} \frac{\partial}{\partial \eta_j} + \sum_{j=1}^{M-1} \left(G_j - p_{\eta_j} \frac{p_{\eta_{j+1}}}{Q_{j+1}} \right) \frac{\partial}{\partial p_{\eta_j}} + G_M \frac{\partial}{\partial p_{\eta_M}} \\
iL_{\text{NHC-baro}} &:= - \frac{p_{\xi_1}}{Q'_1} \mathbf{p}_g \odot \frac{\partial}{\partial \mathbf{p}_g} + \sum_{j=1}^M \frac{p_{\xi_j}}{Q'_j} \frac{\partial}{\partial \xi_j} + \sum_{j=1}^{M-1} \left(G'_j - p_{\xi_j} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} \right) \frac{\partial}{\partial p_{\xi_j}} + G'_M \frac{\partial}{\partial p_{\xi_M}}
\end{aligned} \tag{39}$$

with

$$\mathbf{G}_g := \det[\mathbf{h}] (\mathcal{P}^{\text{int}} - P \mathbf{I}) + \frac{1}{N_f} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} \mathbf{I}. \tag{40}$$

$$\begin{aligned}
e^{iL\Delta t} &= e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}} e^{iL_{\text{NHC-thermo}} \frac{\Delta t}{2}} \\
&\times e^{iL_{g,2} \frac{\Delta t}{2}} e^{iL_2 \frac{\Delta t}{2}} \\
&\times e^{iL_{g,1} \Delta t} e^{iL_1 \Delta t} \\
&\times e^{iL_2 \frac{\Delta t}{2}} e^{iL_{g,2} \frac{\Delta t}{2}} \\
&\times e^{iL_{\text{NHC-thermo}} \frac{\Delta t}{2}} e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}} + O(\Delta t^3).
\end{aligned} \tag{41}$$

The actions of $e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}}$ and $e^{iL_{\text{NHC-thermo}} \frac{\Delta t}{2}}$ can be evaluated similarly to the Nosé-Hoover chain equations. The action of $e^{iL_{g,2} \frac{\Delta t}{2}}$ just translates \mathbf{p}_g .

Since \mathbf{p}_g is a symmetric real matrix, we can diagonalize it as

$$\begin{aligned}
\mathbf{p}_g &= \sum_{\mu=1}^3 \lambda_{\mu} \mathbf{u}_{\mu} \mathbf{u}_{\mu}^{\top} \\
\mathbf{U} &:= (\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3) \\
\mathbf{U} \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix} \mathbf{U}^{\top} &= \mathbf{p}_g
\end{aligned} \tag{42}$$

with

$$\begin{aligned}
\lambda_{\mu} &\in \mathbb{R} \\
\mathbf{u}_{\mu}^{\top} \mathbf{u}_{\nu} &= \delta_{\mu\nu}.
\end{aligned} \tag{43}$$

$$e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2}} = \left(e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2n}} \right)^n$$

$$e^{iL_{\text{NHC-baro}} \frac{\Delta t}{2n}} = S_4^{\text{NHC-baro}} \left(\frac{\Delta t}{2n} \right) + O \left(\left(\frac{\Delta t}{n} \right)^5 \right)$$

$$S_4^{\text{NHC-baro}} \left(\frac{\Delta t}{2n} \right) := \prod_{\alpha=1}^3 S_2^{\text{NHC-baro}} \left(x_\alpha \frac{\Delta t}{2n} \right) \quad \left(\delta_\alpha := x_\alpha \frac{\Delta t}{2n} \right)$$

$$\begin{aligned} S_2^{\text{NHC-baro}}(\delta_\alpha) &:= \exp \left(\frac{\delta_\alpha}{2} G'_M \frac{\partial}{\partial p_{\xi_M}} \right) \\ &\times \prod_{j=M-1}^1 \left(\exp \left(-\frac{\delta_\alpha}{4} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \frac{\partial}{\partial p_{\xi_j}} \right) \exp \left(\frac{\delta_\alpha}{2} G'_j \frac{\partial}{\partial p_{\xi_j}} \right) \exp \left(-\frac{\delta_\alpha}{4} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \frac{\partial}{\partial p_{\xi_j}} \right) \right) \\ &\times \exp \left(-\delta_\alpha \frac{p_{\xi_1}}{Q'_1} \mathbf{p}_g \odot \frac{\partial}{\partial \mathbf{p}_g} \right) \\ &\times \prod_{j=1}^M \exp \left(\delta_\alpha \frac{p_{\xi_j}}{Q'_j} \frac{\partial}{\partial \xi_j} \right) \\ &\times \prod_{j=1}^{M-1} \left(\exp \left(-\frac{\delta_\alpha}{4} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \frac{\partial}{\partial p_{\xi_j}} \right) \exp \left(\frac{\delta_\alpha}{2} G'_j \frac{\partial}{\partial p_{\xi_j}} \right) \exp \left(-\frac{\delta_\alpha}{4} \frac{p_{\xi_{j+1}}}{Q'_{j+1}} p_{\xi_j} \frac{\partial}{\partial p_{\xi_j}} \right) \right) \\ &\times \exp \left(\frac{\delta_\alpha}{2} G'_M \frac{\partial}{\partial p_{\xi_M}} \right) \end{aligned}$$

1.4.3.5. Actions of $e^{iL_1 \Delta t}$ and $e^{iL_2 \frac{\Delta t}{2}}$

$$\mathbf{x}_i := \mathbf{U}^\top \mathbf{r}_i$$

$$\begin{aligned} e^{iL_1 \Delta t} \mathbf{r}_i &= \exp \left(\Delta t \left(\frac{\mathbf{p}_i}{m_i} + \frac{1}{W_g} \mathbf{p}_g \mathbf{r}_i \right) \cdot \frac{\partial}{\partial \mathbf{r}_i} \right) \mathbf{r}_i \\ &= \mathbf{U} \exp \left(\Delta t \left(\frac{\mathbf{p}_i}{m_i} + \frac{1}{W_g} \mathbf{p}_g \mathbf{r}_i \right) \cdot \frac{\partial}{\partial \mathbf{r}_i} \right) \mathbf{x}_i \\ &= \mathbf{U} \exp \left(\Delta t \left(\frac{\mathbf{U}^\top \mathbf{p}_i}{m_i} + \frac{1}{W_g} \mathbf{U}^\top \mathbf{p}_g \mathbf{U} \mathbf{x}_i \right) \cdot \frac{\partial}{\partial \mathbf{x}_i} \right) \mathbf{x}_i \\ &= \mathbf{U} \left(\exp \left(\Delta t \left(\frac{[\mathbf{U}^\top \mathbf{p}_i]_\alpha}{m_i} + \frac{1}{W_g} \lambda_\alpha x_{i\alpha} \right) \frac{\partial}{\partial x_{i\alpha}} \right) x_{i\alpha} \right)_{\alpha=1,2,3} \\ &= \mathbf{U} \left(x_{i\alpha} e^{\frac{\lambda_\alpha \Delta t}{W_g}} + \Delta t \frac{[\mathbf{U}^\top \mathbf{p}_i]_\alpha}{m_i} \text{exprel} \left(\frac{\lambda_\alpha \Delta t}{W_g} \right) \right)_{\alpha=1,2,3} \end{aligned} \tag{45}$$

$$\text{exprel}(x) := \frac{e^x - 1}{x}$$

Similarly

$$\mathbf{y}_i := \mathbf{U}^\top \mathbf{p}_i$$

$$\begin{aligned}
e^{iL_2 \frac{\Delta t}{2}} \mathbf{p}_i &= \exp \left(\frac{\Delta t}{2} \left(\tilde{\mathbf{F}}_i - \frac{1}{W_g} \left(\mathbf{p}_g + \frac{\text{Tr}[\mathbf{p}_g]}{N_f} \mathbf{I} \right) \mathbf{p}_i \right) \cdot \frac{\partial}{\partial \mathbf{p}_i} \right) \mathbf{p}_i \\
&= \mathbf{U} \exp \left(\frac{\Delta t}{2} \left(\tilde{\mathbf{F}}_i - \frac{1}{W_g} \left(\mathbf{p}_g + \frac{\text{Tr}[\mathbf{p}_g]}{N_f} \mathbf{I} \right) \mathbf{p}_i \right) \cdot \frac{\partial}{\partial \mathbf{p}_i} \right) \mathbf{y}_i \\
&= \mathbf{U} \exp \left(\frac{\Delta t}{2} \left(\mathbf{U}^\top \tilde{\mathbf{F}}_i - \frac{1}{W_g} \mathbf{U}^\top \left(\mathbf{p}_g + \frac{\text{Tr}[\mathbf{p}_g]}{N_f} \mathbf{I} \right) \mathbf{U} \mathbf{y}_i \right) \cdot \frac{\partial}{\partial \mathbf{y}_i} \right) \mathbf{y}_i \\
&= \mathbf{U} \left(\exp \left(\frac{\Delta t}{2} \left([\mathbf{U}^\top \tilde{\mathbf{F}}_i]_\alpha - \frac{1}{W_g} \left(\lambda_\alpha + \frac{\text{Tr}[\mathbf{p}_g]}{N_f} \right) y_{i\alpha} \right) \frac{\partial}{\partial y_{i\alpha}} \right) y_{i\alpha} \right)_{\alpha=1,2,3} \\
&= \mathbf{U} \left(y_{i\alpha} e^{-\frac{\kappa_\alpha \Delta t}{2W_g}} + \frac{\Delta t}{2} [\mathbf{U}^\top \tilde{\mathbf{F}}_i]_\alpha \exp \left(-\frac{\kappa_\alpha \Delta t}{2W_g} \right) \right)_{\alpha=1,2,3}
\end{aligned} \tag{46}$$

$$\kappa_\alpha := \lambda_\alpha + \frac{\text{Tr}[\mathbf{p}_g]}{N_f}$$

1.4.3.6. Action of $e^{iL_{g,1}\Delta t}$

$$\mathbf{n} := \mathbf{h} \mathbf{U}$$

$$\begin{aligned}
e^{iL_{g,1}\Delta t} \mathbf{h} &= \exp \left(\Delta t \frac{\mathbf{h} \mathbf{p}_g}{W_g} \odot \frac{\partial}{\partial \mathbf{h}} \right) \mathbf{h} \\
&= \exp \left(\frac{\Delta t}{W_g} \text{Tr} \left[(\mathbf{h} \mathbf{p}_g)^\top \frac{\partial}{\partial \mathbf{h}} \right] \right) \mathbf{h} \\
&= \exp \left(\frac{\Delta t}{W_g} \text{Tr} \left[\mathbf{p}_g \mathbf{U} \mathbf{n}^\top \frac{\partial}{\partial \mathbf{n}} \mathbf{U}^\top \right] \right) \mathbf{n} \mathbf{U}^\top \\
&= \left(\prod_{\mu\alpha} \exp \left(\frac{\Delta t \lambda_\alpha}{W_g} n_{\mu\alpha} \frac{\partial}{\partial n_{\mu\alpha}} \right) \mathbf{n} \right) \mathbf{U}^\top \\
&= \left(e^{\frac{\Delta t \lambda_\alpha}{W_g} n_{\mu\alpha}} \right)_{\mu\alpha} \mathbf{U}^\top
\end{aligned} \tag{47}$$

1.4.4. Cell fluctuations with mask

1.4.4.1. MTK equations

Consider the case when the cell fluctuations are allowed only n_c axes of the cell matrix \mathbf{h} . For $c = 1, \dots, n_c$, let p_c be the momentum conjugate to the c th axis of \mathbf{h} .

$$\mathbf{p}_g = \sum_c p_c \mathbf{e}_c \mathbf{e}_c^\top \tag{48}$$

EOM are the same as the anisotropic MTK equations except for

$$\begin{aligned}
\dot{p}_c &= \det[\mathbf{h}] \cdot \mathbf{e}_c^\top (\mathcal{P}^{\text{int}} - P\mathbf{I})\mathbf{e}_c + \frac{1}{N_f} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i} - \frac{p_{\xi_1}}{Q'_1} p_c \quad (c = 1, \dots, n_c) \\
G'_1 &:= \sum_c \frac{p_c^2}{W_g} - n_c kT \\
Q'_1 &= n_c kT \tau^2,
\end{aligned} \tag{49}$$

where τ is a characteristic time scale for barostat.

1.4.4.2. Conserved quantities

The conserved energy

$$\begin{aligned}
H' &:= \mathcal{H}(r, p) + \sum_c \frac{p_c^2}{2W} + P \det[\mathbf{h}] \\
&+ \sum_{j=1}^M \left(\frac{p_{\eta_j}^2}{2Q_j} + \frac{p_{\xi_j}^2}{2Q'_j} \right) + N_f kT \eta_1 + n_c kT \xi_1 + kT \sum_{j=2}^M (\eta_j + \xi_j)
\end{aligned} \tag{50}$$

1.4.4.3. Integrating the MTK equations

$$\begin{aligned}
iL &:= iL_1 + iL_2 + iL_{g,1} + iL_{g,2} + iL_{\text{NHC-baro}} + iL_{\text{NHC-thermo}} \\
iL_{g,2} &:= \sum_c G_c \odot \frac{\partial}{\partial p_c}
\end{aligned} \tag{51}$$

with

$$G_c := \det[\mathbf{h}] \cdot \mathbf{e}_c^\top (\mathcal{P}^{\text{int}} - P\mathbf{I})\mathbf{e}_c + \frac{1}{N_f} \sum_{i=1}^N \frac{\mathbf{p}_i^2}{m_i}. \tag{52}$$

Bibliography

- [1] H. Yoshida, “Construction of higher order symplectic integrators,” *Physics Letters A*, vol. 150, no. 5, pp. 262–268, 1990, doi: [https://doi.org/10.1016/0375-9601\(90\)90092-3](https://doi.org/10.1016/0375-9601(90)90092-3).
- [2] G. J. Martyna, M. L. Klein, and M. Tuckerman, “Nosé-Hoover chains: The canonical ensemble via continuous dynamics,” *The Journal of Chemical Physics*, vol. 97, no. 4, pp. 2635–2643, 1992, doi: 10.1063/1.463940.
- [3] G. J. Martyna, D. J. Tobias, and M. L. Klein, “Constant pressure molecular dynamics algorithms,” *The Journal of Chemical Physics*, vol. 101, no. 5, pp. 4177–4189, 1994, doi: 10.1063/1.467468.
- [4] M. E. Tuckerman, J. Alejandre, R. López-Rendón, A. L. Jochim, and G. J. Martyna, “A Liouville-operator derived measure-preserving integrator for molecular dynamics simulations in the isothermal-isobaric ensemble,” *Journal of Physics A: Mathematical and General*, vol. 39, no. 19, p. 5629, Apr. 2006, doi: 10.1088/0305-4470/39/19/S18.