

The most used equation going forward (by far)

$$\vec{F} = m\vec{a}$$


$$\sum F_x = ma_x \quad \sum F_y = ma_y$$

$$F_{x1} + F_{x2} + \cdots = ma_x$$


$$F_{y1} + F_{y2} + \cdots = ma_y$$

## Question #19


A ball has been tossed straight up. Which is the correct free-body diagram just after the ball has left the hand? Ignore air resistance.




D



C





B




E

## Question #20


A ball, hanging from the ceiling by a string, is pulled back and released. Which is the correct free-body diagram just after its release?




B




C



D





A




E

# Question #21


A car is parked on a hill.  
Which is the correct free-body diagram?




E




C



B





D




A

# Question #21


A car is parked on a hill.  
Which is the correct free-body diagram?




E




C



B





D



A



## Question #22

A car is towed to the right at constant speed. Which is the correct free-body diagram?



## Question #22


A car is towed to the right at constant speed. Which is the correct free-body diagram?



## Question #23

The box is sitting on the floor of an elevator. The elevator is accelerating upward. The magnitude of the normal force on the box is


- a.  $n = 0$ .
- b.  $n = F_g$ .
- c.  $n < F_g$ .
- d. Not enough information to tell.
- e.  $n > F_g$ .






# Speed of a towed car

A 10-kg box is pulled across a rough, level surface by a rope attached to the box. The rope is pulling at a  $20^\circ$  angle above the horizontal with a tension equal to 50 N. The frictional force is equal to 10 N. Starting from rest, how long will it take for this box to travel 10 m?




# Example

You push horizontally on a 50 N block which is sitting on a 30 degree, incline. What is the direction and magnitude of static friction if the block is to remain stationary?




# Box on an incline problem.

A 50 N(5 kg) box is placed on a 30 degree incline and has just begun sliding downward. A 5 N kinetic friction force acts on the box. How far down the incline will the box travel in 3.0 s?



# Gravity: A Force

Attractive, long-ranged force between any two objects.




$$F_{1 \text{ on } 2} = F_{2 \text{ on } 1} = \frac{Gm_1 m_2}{r^2} \quad (\text{Newton's law of gravity})$$

$$G = 6.67 \times 10^{-11} \frac{\text{N m}^2}{\text{kg}^2}$$

Gravitational constant

# Gravity: A force

Will the gravitational force between two people-sized objects be large or small?



$$\vec{F}_G = \vec{F}_{\text{planet on } m} = \left( \frac{GMm}{r^2}, \text{ straight down} \right)$$

$$G = 6.67 \times 10^{-11} \frac{\text{N m}^2}{\text{kg}^2}$$


$$M_E = 5.98 \times 10^{24} \text{ kg}$$

$$R_E = 6.37 \times 10^6 \text{ m}$$

# Gravity: A force

Will the gravitational force between two people-sized objects be large or small?

When will the gravitational force be large?



$$\vec{F}_G = \vec{F}_{\text{planet on } m} = \left( \frac{GMm}{r^2}, \text{ straight down} \right)$$

$$G = 6.67 \times 10^{-11} \frac{\text{N m}^2}{\text{kg}^2}$$


$$M_E = 5.98 \times 10^{24} \text{ kg}$$

$$R_E = 6.37 \times 10^6 \text{ m}$$

# Gravity: A force

Will the gravitational force between two people-sized objects be large or small?

When will the gravitational force be large?



$$\vec{F}_G = \vec{F}_{\text{planet on } m} = \left( \frac{GMm}{r^2}, \text{ straight down} \right) = (mg, \text{ straight down})$$

$$G = 6.67 \times 10^{-11} \frac{\text{N m}^2}{\text{kg}^2}$$

$$M_E = 5.98 \times 10^{24} \text{ kg}$$

$$R_E = 6.37 \times 10^6 \text{ m}$$