

# **An assessment of the Ecological Conservation Redline: Unlocking priority areas for conservation**

Wenze Yue<sup>a</sup>, Bi'ou Feng<sup>a</sup>, Qiushi Zhou<sup>a</sup>, Ronghua Xu<sup>a</sup>, Mengmeng Li<sup>a,b\*</sup>

<sup>a</sup> Department of Land Management, Zhejiang University, Hangzhou 310029, P.R. China

<sup>b</sup> Institute for Environmental Studies, VU University Amsterdam, 1081HV, Amsterdam, the Netherlands

\* **Correspondence to:** Mengmeng Li ([mengmeng.li@zju.edu.cn](mailto:mengmeng.li@zju.edu.cn))

**Abstract:** Protected areas (PAs) are established to conserve wildlife habitats and biodiversity. To this end, the Chinese central government has initiated a pioneering environmental planning and management policy known as the Ecological Conservation Redline (ECR). Whilst ecological benefits of ECR policy have been extensively evaluated, spatially explicit assessment of the ECR remains understudies. Here, we propose an element-function-structure framework for assessing the concurrence of ECR areas and ecological conservation hotspots to further outline priority areas for conservation. Results show that 67% of existing PAs are protected by ECR areas, while that of ecological corridors is only 11%. Regional variation in the ECR representativeness can be leveraged by deliberately protecting tailored ecological conservation hotspots in specific locations and stepping stones in ecological corridors. This study highlights the substantial space for ecological management to achieve the goals of ECR policy, and discourses on the co-production of knowledge from researchers and policy makers.

**Keyword:** Protected areas; biodiversity conservation; natural resource management; ecological security; assessment framework

## 25    **1. Introduction**

26    Originally, terrestrial Protected Areas (PAs) are established worldwide to constrain destructive human  
27    activities and safeguard endangered species in perpetuity (Folharini et al., 2021; Kroner et al., 2019;  
28    Ribeiro et al., 2021). Yet, evidence suggests that PAs are confronted with increasing alterations from  
29    human activities that characterize the Anthropocene (Ellis & Ramankutty, 2008; Geldmann et al., 2014),  
30    despite the rapid expansion of PA coverage over the past decades (Buchadas et al., 2022). At a global scale,  
31    only 42% of PAs are free of any detectable human disturbance, whereas approximately 33% are  
32    extensively modified (Jones et al., 2018). For example, in the United States, national parks, wilderness  
33    areas, community conserved areas, and nature reserves are declining in their ability to conserve  
34    biodiversity due to human disturbance (Radeloff et al., 2010). Furthermore, the established species-  
35    oriented indicators have played a key role in assessing the outcome of PAs (Gray et al., 2016), but they  
36    may fall short in the context of increasingly divergent goals of conservation. For instance, Maxwell et al.  
37    (2020) state that PAs must effectively contribute to meeting global goals, ranging from preventing specie  
38    extinctions to retaining the most-intact ecosystems.

39    Consistent with the inclusive goals of conservation efforts, the central government of China has initiated  
40    a pioneering environmental planning and management policy known as the Ecological Conservation  
41    Redline (ECR, see Gao (2019) and Schmidt-Traub et al. (2020)). The ECR policy, which aims to protect  
42    more than one-quarter of China's territory, is currently one of the nation's highest priorities for policy  
43    implementation (Zhang et al., 2022). Basically, ECR boundaries outline regions with privileged ecological  
44    functions by assimilating the principles of ecosystem service, ecological sensitivity, and biodiversity  
45    priorities into management practices (Bai et al., 2021; Gao et al., 2020). Ideally, the establishment of ECR  
46    can benefit habitat quality and maintain habitat connectivity (Luo et al., 2020). Therefore, the ECR vision  
47    resembles the concept of ecological networks in terms of integrity and connectivity, but it is more  
48    conducive to safeguarding ecological security in terms the willingness for political action (Ma et al., 2022;  
49    Montoya et al., 2006).

50    The joint legislation regarding the ECR policy by multiple central government departments<sup>1</sup> is intended  
51    to play a critical role in ecological conservation, and assessments of management practices have been  
52    extensively conducted in such context (Gao et al., 2020). These assessments help better reconcile

---

<sup>1</sup> [http://gi.mnr.gov.cn/202208/t20220819\\_2756940.html](http://gi.mnr.gov.cn/202208/t20220819_2756940.html)


conservation goals with local economic development under the premise that ECR areas have been optimally identified (Busch & Amarjargal, 2022). To the best of our knowledge, however, few studies have assessed the rationality of the issued ECR coverage. Given that no clear planning framework has been set to maximize the representativeness and effectiveness of conservation goals, government and local authorities are facing the dilemma of ECR conflicts with regional socio-economic development, despite the fact that they inherently prefer socio-economic development in most cases. This issue has largely led to spatial mismatch, isolation, and severe human interference inside or around the ECR areas. For example, Liang et al. (2022) find that, in Jiangsu province of China, a large proportion of built-up land is located in ECR areas, in addition that ECR boundaries do not always match local landforms, which is prone to absurd patterns of regional development due to land-use conflict (de Jong et al., 2021).

As such, the objective of this study is to facilitate a framework for a comprehensive science-based assessment of ECR, and possibly to unlock priority areas for future optimization of ECR delineation. To achieve this, ecological conservation hotspots are firstly identified based on ecosystem element, function, and structure. These ecological conservation hotspots are further used to delineate the ecological representativeness of ECR coverage at both provincial and county scales. Finally, priority areas for conservation are identified by synthesizing ecological function zones and ecological stepping stones. We implement this approach for the case of Zhejiang province, which plays a critical role in China's ecological protection, as it is the place where the ECR policy originates.

## **2. Materials and methods**

In this study, we propose a stepwise method to assess the effectiveness of ECR coverage (Fig. 1). First, ecological conservation hotspots are identified accounting for ecological element, function, and structure. Specifically, ecological element is depicted by integrating the wide range of existing PAs. Ecological Functional Zones (EFZs) maximize the dominant ecological functions and serve the functional objective of ECR patches, which are connected by ecological corridors that serve the holistic ECR system objective. Second, the entire retention of these three ecological conservation hotspots in ECR areas is investigated (i.e., ecological representativeness) for each of the three components. Third, we compare the variation in ECR representativeness among Major Function-oriented Zones, the blueprint for the future development and protection pattern of China's territory (Fan, 2015), in order to better reflect practical issues across different counties. In addition, management characteristics of ECR are discussed from the perspective of regional competition for land. Finally, we identify priority areas for conservation, to illustrate the gap

83 between management practice and policy goals, and to inform sound policy-making for regional  
84 optimization of ECR coverage.




85  
86 **Figure 1. Workflow for the assessment of ECR coverage.** The three columns in gray represent the  
87 element-function-structure that altogether depict ecological conservation hotspots.

## 88 2.1. Study area

89 Zhejiang province is located in the southeast coast of China (Fig. 2). This province covers an estimate of  
90 105,000 km<sup>2</sup>, of which forest land and cultivated land account for 62% and 23%, respectively (Table S1).  
91 Owing to its abundant vegetation and valuable biological resources, Zhejiang province performs essential  
92 ecological functions at both regional and national scales. Notably, in 2000, a local flagship program  
93 analogous to the current ECR policy was first launched in the Anji County of Zhejiang province (Gao,  
94 2019, also see Fig. 2).

95 In addition, Zhejiang is a densely populated province in the south wing of the Yangtze River Delta and it

96 is now experiencing rapid urbanization (Wang et al., 2017). It governs 11 cities and 89 counties, with a  
 97 total GDP and per capita GDP ranking among the top five in China. Currently, the terrestrial ECR areas  
 98 in Zhejiang province cover an area of 24,852 km<sup>2</sup>, more than 23% of the land (Table S2).



99

100 **Figure 2. Location and land use patterns of Zhejiang province.** Land use data is derived from the  
 101 *National Land Cover Datasets (NLCDs, see Liu et al. (2014))* for the year 2018. The topographic basemap  
 102 is publicly available at ESRI ArcGIS®.

## 103 2.2. Dataset

104 Various datasets are used in this study (Table S3). These datasets can be categorized into 4 classes: 1)  
 105 Land use. Land use data with a spatial resolution of 30 m are obtained from the Resource and  
 106 Environmental Science and Data Center (<https://www.resdc.cn/>), and was reclassified into six categories:  
 107 cultivated land, forest land, grassland, open water, constructed land, and barren land. 2) Terrain. Digital

elevation model (DEM) with a spatial resolution of 30 m is downloaded from the Geospatial Data Cloud (<http://www.gscloud.cn/>). 3) Biophysical elements. Normalized Difference Vegetation Index (NDVI) and Net Primary Productivity (NPP) are derived from Landsat-8 satellite imaginaries; Soil properties are provided by the World Soil Database, which includes soil depth, organic matter content, soil composition and others; Temperature and precipitation are derived from the observation data of 42 meteorological stations covering Zhejiang Province, and precipitation and potential evapotranspiration in 1995–2018 are interpolated using the Kriging method. 4) Socioeconomic factors. Road Network is obtained from the OpenStreetMap (OSM, <https://www.openstreetmap.org/>); ECR boundaries, major function-oriented zones, and existing PAs are acquired from the local government (<https://www.zj.gov.cn/>).

Pre-processing operations such as geo-referencing and merging are implemented on ArcGIS 10.5 platform. All data layers are uniformly projected to the spatial reference of WGS\_1984\_UTM\_Zone\_50N, and the grid size is resampled to 100 m × 100 m.

### **2.3. Determination of ecological conservation hotspots**

In this section, we elaborate the methods used to extract existing PAs, EFZs, as well as ecological corridors, which are the three most relevant components for ECR delineation (i.e., element, function, and structure, respectively).

#### **2.3.1. Protected Areas (PAs)**

Existing spectrum of PAs in China consists of national parks, nature reserves and parks (Miller-Rushing et al., 2017), with a decreasing sequence of ecological importance and management intensity accordingly. We compile the scheme of PAs, and manually merge PAs regardless of their relative importance in the ArcGIS platform (<https://www.esri.com/>).

#### **2.3.2. Important Ecological Functional Zones (EFZs)**

Ecological functions can be portrayed as the ecosystem services (ESs) and ecological vulnerability (Peng et al., 2019). According to Zhejiang's ecological conditions and expert knowledge, we select and assess four types of ESs and one indicator for ecological vulnerability with the InVEST model, NPP index, and the Revised Universal Soil Loss Equation (RUSLE). For more technical details, please see the Supplementary Method. These four ESs subsequently are normalized using Min-Max value method, and summed with equal weights. Lastly, we extract 20% of areas with the largest values as EFZs (Bai et al.,



2018).

### 2.3.3. Potential ecological corridors

Regions protected by ECR areas are not isolated space for many large species that are largely dependent on habitat outside the ECR, and corridors linking them together are therefore crucial elements for animal movement (Dai et al., 2021; Ghoddousi et al., 2022). Literally, an ecological corridor that connects multiple sources is a narrow strip of vegetation that is distinct from other nearby ecological elements. The graph-based approach and circuit theory have been widely used to identify potential corridors (Huang et al., 2020; Peng et al., 2018). Following the circuit theory, landscape can be regarded as a conductive surface and the probability in the biological migration conforms to Ohm's law:

$$I = U / R \quad (1)$$

where  $R$  characterizes the resistance of the conductive,  $U$  represents the voltage across a given node, and  $I$  is the current through the node. In this study, the regional ecological network is regarded as a circuit, and the flow is considered as the current between ecological sources.

As for the application of circuit theory, the key is to determine ecological sources and resistance surfaces. Ecological sources are the origin of species dispersal and provide important ecosystem services as well as take great structural importance (Dai et al., 2021). In light of this, ecological sources are identified as these core areas with high ecological functions. Specifically, the research objects of the Morphological Spatial Pattern Analysis (MSPA) are divided into two categories: foreground (forest land, grassland, open water) and background (cultivated land, constructed land, barren land) to distinguish the core area (Lin et al., 2021). Consequently, the value of ecological functions is categorized into one out of the five levels following the natural break method, with the top 20% bin considered as ecological function zones (Peng et al., 2018).

Meanwhile, resistance value indicates the difficulty for species to disperse, which are determined by natural conditions and human activities. According to the characteristics of the study area and existing researches (Wang et al., 2021), resistance value can be represented by land use/cover, elevation, slope, distance from road, and distance from constructed land (Table S4).

## 2.4. Ecological representativeness of ECR coverage

We analyze the overall pattern of these different types of ecological hotspots (i.e., ecological element,

function, and structure), the proportion of ecological conservation hotspots (PPH) within its area and the percentage of overlap with hotspots (PEH), to measure ecological representativeness (Bai et al., 2021). PPH is obtained by calculating the percentage of area of all hotspots that are protected in Zhejiang province, to assay the extent of the protection of ecological hotspots. PEH is obtained by calculating the percentage of ECR areas that concur with ecological conservation hotspots in each of the 89 counties, interpreting the representativeness of the current ECR. Thus, PPH reveals the overall protection of all ecological conservation hotspots, and PEH represent the percentage of ECR areas covers hotspots relative to non-hotspot areas (Bai et al., 2021). To further examine the regional variation of ECR representativeness in Zhejiang province, we distinguish four clusters according to the major function-oriented zones, determined by natural conditions and human activities therein (Fan et al., 2012).

## 2.5. Determination of conservation priorities

Implementing official adjustments of the issued ECR coverage may be unrealistic in the short term. Hence, we determine the ecological optimization target through the overall analysis with regards to EFZ and ecological stepping stones.

### 2.5.1. Importance zoning of EFZ

After dissolving the core areas of the established PAs in ECR areas, conservation priority of non-overlapping EFZ is graded into one out of the four levels according to the summed level of ecological functions and landscape connectivity of their ecological sources.

Moreover, we use graph-based connectivity indicators to characterize the overall connectivity in a landscape and the importance of each patch for connectivity (Saura & Pascual-Hortal, 2007). Specifically, Integral Index of Connectivity (*IIC*) and Probability of Connectivity (*PC*) are chosen to calculate the relative importance (*dI*) of landscape connectivity by using Conefor 2.6 software. The main calculation formula is as follows:

$$IIC = \frac{\sum_{i=1}^n \sum_{j=1}^n \frac{a_i * a_j}{1 + n l_{ij}}}{A_L^2} \quad (2)$$

$$PC = \frac{\sum_{i=1}^n \sum_{j=1}^n P_{ij}^* a_i * a_j}{A_L^2} \quad (3)$$

$$dI = 0.5 * dIIC + 0.5 * dPC \quad (4)$$



where  $n$  is the total number of patches;  $a_i$  and  $b_j$  are areas of patches  $i$  and  $j$  respectively;  $nl_{ij}$  is the number of connections between patches  $i$  and  $j$ ;  $A_L$  is the area of the entire landscape; and  $P_{ij}^*$  represents the maximum probability that patch  $i$  and  $j$  are connected.


### 2.5.2. Ecological stepping stones

Ecological corridors play a crucial role in biological dispersal and species persistence (Rocha et al., 2021), while linear construction projects such as roads and railways threaten the protection of ecological corridors (Marcantonio et al., 2013). In that case, however, stepping stones in corridors can still provide essential ecological linkage (Rocha et al., 2021; Saura et al., 2014). In this study, we use the Pinchpoint Mapper to identify ecological stepping stones following the circuit theory. According to the maximum migration distance of terrestrial animals and plants (Rouget et al., 2006), the width of potential ecological corridors is determined to be 1.2 km, and the cost-weighted density (CWD) of corridors is graded into five classes using the natural break method. We assign the areas that fall into the highest density bin as the essential locations for ecological flow. Considering the exclusivity of ecological elements and integrity (Saura et al., 2014), we eventually define ecological patches with an area  $> 10 \text{ km}^2$  as stepping stones, of which the total current value can further reflect their relative importance.

## 3. Results

### 3.1. Ecological representativeness of ECR coverage

Overall, ecological conservation hotspots cover approximately 31% of the total terrestrial area of Zhejiang Province (Fig. 3), and this proportion is moderately higher than the national target of ECR policy (25%). Specifically, PAs, which cover a terrestrial area of  $10110 \text{ km}^2$ , are mostly distributed in Hangzhou City and Wenzhou City (Fig. 3a and Table S5). Moreover, 84% of these PAs are established to restrict human activity in forestry areas (Table S5). In contrast, the total area of EFZ hotspots reaches nearly  $25,000 \text{ km}^2$  (Fig. 3b and Table S6), which is also dominated by forestry (Table S6). Ecological corridors, however, only cover a small area of  $8297 \text{ km}^2$  (Fig. 3c), with an average length of 56 km. The vast majority of land in ecological corridors is currently used for forestry and agricultural production, accounting for 47% and 36%, respectively (Table S7).




216

217 **Figure 3. Ecological conservation hotspots. (a) Protected Areas; (b) Ecological Function Zones; and (c)**  
218 **Ecological Corridors. To further explore the enhanced visualization of these results, please check the**  
219 **interactive map viewer at <https://landbigdata.github.io/ECR/>**

220 The PPH, or the proportion of ecological conservation hotspots protected by ECR areas, for all of Zhejiang  
221 is only 48%, suggesting an enormous space for spatial optimization. Yet, there is a large variation in terms  
222 of PPH among the three dimensions (i.e., ecological element, function, and structure). In the dimension  
223 of ecological elements (i.e., PAs), PPH is roughly 67%. Local governments reserve space for tourism and  
224 construction development, and thus ECR protection is prone to leave gaps in PAs (Fig. 4a), especially in  
225 nature parks (e.g., in the northeast of Wenzhou city). In the dimension of ecological functions, 48% of the  
226 entire EFZ is protected by ECR areas (Fig. 4b). However, ECR areas in the south and northwest of  
227 Zhejiang province cannot sufficiently protect EFZ, and these in central Zhejiang province protect a large  
228 proportion of land outside the EFZ. In terms of ecological structure (i.e., ecological corridors), however,  
229 PPH is only 11%, highlighting the remarkable insufficiency of ecological corridors protected by ECR  
230 areas.


231



**Figure 4. Concurrence of ECR and ecological conservation hotspots. (a) ECR and PAs, and (b) ECR and EFZ. For enhanced visualization of these results, please also explore the interactive map at <https://landbigdata.github.io/ECR/>**

### 3.2. Regional variation in ECR delineation

As the spatial congruity between ECR areas and ecological conservation hotspots differs across space, we compare the representativeness (PEH) among different counties, particularly with regards to four types of major function-oriented zones (i.e., urban development areas, priority development areas, main agricultural areas, and key eco-function areas). Results show that the proportion of non-ecological land in ECR areas increases substantially with the increasing priority for socio-economic activities in the 89 counties, manifested as the low ecological representativeness of ECR coverage (Fig. 5a-b).



244

245 **Figure 5. Regional variations in the ecological representativeness of ECR coverage.** (a) division of  
246 major function-oriented zones; (b) bivariate correlation between ECR ratio and percentage of ecological  
247 land; (c) distribution of the ecological representativeness of three hotspots (PEH) for each county, with  
248 color indicating the categories.


Among the four types of major function-oriented zones, the urban development areas exhibit the highest representativeness in terms of PAs (Fig. 5c). As the main area of economic activities, local authorities tend to strictly protect the existing PAs in the progress of socioeconomic development to meet local ecological needs. However, given the frequent disturbance of human activities, ecological corridors are largely pre-occupied for residence and food production. In particular, many counties like Lucheng county are in such a dilemma with 5% of construction land located in ECR areas (Tables S8 and S9). Priority development areas are dominated by counties in the northeast of Zhejiang, where people act more multifarious and concentrated, and thus ecological functions are generally under the median (Fig 5c), which can be attributed to the scarcity of ecological land. Notably, the ratio of non-ecological land roughly increases with increase of ECR coverage. For example, in Cixi County and Haining county, nearly 60% of the ECR areas are in non-ecological regions (Tables S8 and S9).

Overall, ECR areas in the main agricultural production zones are less representative in terms of ecological function and structure (i.e., EFZ and ecological corridors, respectively). There is quite some non-ecological land that is protected by ECR areas (Table S9). For example, 63% of total land in Pinghu County is used for cultivation, whereas 57% of its total cultivated land is found in ECR areas (Tables S8 and S9). By contrast, almost all counties in the key eco-function areas effectively represent ecological functions. The proportion of ecological land in these counties is over 80% (Tables S8 and S9), and thus land-use competition is less intensive in ECR areas.

### 3.3. Priority areas for conservation

To increase the ecological representativeness of ECR coverage at the provincial level, counties in the southwest of Zhejiang province can act as the leverage point for balancing ecological functions and landscape connectivity (Fig. 6a). Thus, these counties are responsible for the most protection, but are still plagued by inadequate EFZ protection despite the highest ECR ratios. Counties in the western mountains and coastal hills have high ecological functional importance and good landscape connectivity ( $dI$ -level = 3), which are potential areas for ECR protection. Notably, counties in the west of Zhejiang province already have large proportions of ecological conservation hotspots protected by the ECR areas, such as Lin'an County, Tonglu County, and Kecheng County. However, counties in coastal hills show relative low representativeness of ecological hotspots, such as Xianju County, Qingtian County, and Jinyun County. Moreover, the coverage of ECR areas in the urban development areas, such as Yuyao County and Yongjia County, have only marginal contributions to the overall ecological security ( $dI$ -level = 1/2).





279

280 **Figure 6. Priority areas for conservation (a) and ecological stepping stones at ecological corridors (b).**  
281 *To facilitate sound policy-making for local government and decision-makers, both maps can be explored*  
282 *interactively in the map viewer at <https://landbigdata.github.io/ECR/>*

283 Several counties assume the important responsibility of maintaining landscape connectivity, although they  
284 may provide minimal ecosystem functions (Fig. 6b). Based on the circuit theory, we identify 59 stepping  
285 stones with an average size of 2.5 km<sup>2</sup> and a total area of 149 km<sup>2</sup>, which accounts for 38% in size of  
286 potential ecological corridors. The hills and basins in the geographical center of Zhejiang province, such  
287 as Keqiao County, Zhuji County, and Jiande County are hotspots of ecological stepping stones (Fig. 6b  
288 and Table S10). Similarly, a large proportion of land in Ou Hai County, Haining County, and Shangyu  
289 County is along river banks that act as ecological stepping stones (see the interactive map). However,  
290 many stepping stones in these counties have been occupied by non-ecological land (Table S10), especially  
291 in Jiaxing City, where the vast majority of these stepping stones are used for agricultural production (Table  
292 S11).

## 293 4. Discussion

### 294 4.1. Conservation policies in China

295 To resolve sustainability crisis, the central government of China has enacted a wide range of nature  
296 conservation policies in the past few decades, including Three-north Shelter Forest Program, Natural

Forest Conservation Program, and Grain for Green Program, among others (Bryan et al., 2018). In particular, these programs have largely contributed to forest restoration and biodiversity conservation in China (Chen et al., 2019; Deng et al., 2014; Viña et al., 2016). For example, China plays a leading role during the greening process on Earth (25%) with forests contributing to 42% of greening in China (Chen et al., 2019).

Moreover, the Chinese government have also launched a number of programs to address biodiversity issues, particularly, Wildlife Conservation and Nature Protection Program, by which PAs are established all over the country (Bryan et al., 2018). In order to foster the effectiveness of PAs, which now covers ~18% of the country's land, the Chinese government recently established a new scheme of PAs that consists of national parks, nature reserves and parks (Miller-Rushing et al., 2017; Wei et al., 2020). In addition, a series of efforts has been carried out to warrant the maintenance of PAs, including establishment of conservation monitoring networks, synthesis of species catalogues, and legislation of conservation priority areas (Ma et al., 2017; Wei et al., 2020). These efforts altogether have substantially contributed to the nation-wide strategy of ecological civilization (Wei et al., 2020).

#### **4.2. Significance of this study and implications for ECR Policy**

ECR policy is supposed to be distinguished from other conservation policies in China with regard to the integrity of ecological protection, strict requirements for spatial accuracy, and overall-process management (Gao et al., 2020). To the best of our knowledge, this study is a unique exploration in the assessment of ECR coverage, and can therefore benefit the conservation community in the following aspects.

While other studies have explored approaches to the assessment of ECR coverage (Hu et al., 2020; Zuo & Gao, 2021), this study broadens the scope for ecological assessments by incorporating multiple goals for conservation. Ecological benefits, among social outcome and nature-human interactions, have always been a key indicator in examining the management effectiveness of PAs (Ghoddousi et al., 2022). Prior studies, on the one hand, overwhelmingly investigate ecological vulnerability and habitat quality in PAs to illustrate protection effectiveness (Ke et al., 2017; Ribeiro et al., 2021; Xu et al., 2017), and on the other hand, analyze social and conservation outcomes of PAs (Oldekop et al., 2016; Porter-Bolland et al., 2012). These assessments are frequently carried out within a given area. However, ECR is the bottom line (threshold) of ecological security, which is irreplaceable space for ecosystem conservation. We prioritize



potential ecological areas by assessing the ecological importance of the entire area.

Although ECR areas have been formally delineated across China (Gao et al., 2020), local governments and authorities may fail to achieve their overall targets for conservation without an effective operationalization at an appropriate scale. The lack of definite standards still causes a large uncertainty on how to define and implement ECR. However, we downscaled targets and analysis to counties, which are the operators of ECR delineation. In addition, the characteristics of counties with different functions are clustered, thus enabling local authorities to make informed decisions and optimization. Moreover, unlike the common research on a macro-scale (Barber et al., 2014; Edgar et al., 2014) or micro-scale (Jones et al., 2020; Liu et al., 2001), this study supplements the discussion from an intermediate perspective.

A preferable pathway to ECR implementation is site-specific management at the county scale as it is the scale at which environmental planning and management policies are primarily targeted, in the meanwhile, this can account for spatial heterogeneity therein. Specifically, we find that the proportion of ecological conservation hotspots protected by ECR is nearly 70% in many counties in the southwest of Zhejiang province. Yet, this proportion for many counties in the northeast of Zhejiang province is less than 5%. With equity as one consideration, we suggest that ECR coverage in the southwest counties should be extended to the highest priority regions identified in Fig. 6a, while ECR coverage in counties elsewhere could be extended to less prioritized regions. Moreover, we suggest to play close attention to landscape connectivity in northeast counties by protecting ecological corridors, and particularly stepping stones, as results show that only 11% of ecological corridors are protected by ECR areas. Finally, the core and non-core areas of ecological conservation need to be distinguished, and hierarchical regulation must be applied. Given the complexity of ecological security, establishing buffer zones outside ECR areas and coordinate all ecological land through hierarchical supervision are necessary to maintain ecosystem integrity.

### **4.3. Embracing the multifaceted goals of conservation**

Globally, there is a growing recognition that government and local authorities should enact policies that are capable to mitigate trade-offs between biodiversity protection and other aspects of human well-being including cultural values, social relations, as well as public health (de Omena & Hanazaki, 2022; McKinnon et al., 2016). Similarly, the original goals of the ECR policy in China include protecting species from extinction, maintaining and restoring wildlife habitats, enhancing ecosystem services, and protecting biological diversity (Gao et al., 2020; Wei et al., 2020). However, government and local authorities

increasingly acknowledge that Indigenous community and cultural landscape need to be accounted in the practice of ECR delineation. According to the latest protocol on the implementation of ECR policy, Indigenous people are authorized to build houses, raise livestock, and cultivate land in ECR areas, that is, ECR areas are not exclusively for biodiversity conservation. Our assessment demonstrates that a large proportion of ECR areas in non-hotspot regions (in or nearby urbanized areas), which we speculate that this is because that the goal of ECR delineation to conserve natural habitat and biodiversity is compromised to people's demand for aesthetic enjoyment and perception (O'Connor et al., 2021).

Conservation targets are challenging to be clearly set as multiple conservation goals are often closely intertwined (Díaz et al., 2020). In particular, there are often trade-offs and synergies in terms of conservation outcomes for nature or for people (Carranza et al., 2020; O'Connor et al., 2021). For instance, studies investigating multiple outcomes of conservation policy suggest synergies between habitat quality and species richness (Zhou et al., 2021), but may also demonstrate potential trade-offs between habitat quality and economic prosperity (Jones et al., 2020). As such, the evaluation of conservation effectiveness should be critical, and preferably outcome-oriented (Ghoddousi et al., 2022; Maxwell et al., 2020; Rodrigues & Cazalis, 2020). To better implement of ECR, government and local authorities need to improve the transparency and promote public participation of ECR delineation to bring multiple targets of conservation to the table, ensuring that ECR policy can spontaneously contribute to the conservation outcomes for nature and for people. Moreover, open-access monitoring network is required to safeguard the implementation of ECR policy in a real-time mode (Mi et al., 2021; Wang et al., 2021), as this allows for prompt feedbacks from governments, local communities and stakeholders, among many others.

#### **4.4. Limitations**

We acknowledge that our assessment has uncertainties and limitations. Conceptually, our assessment focuses on the ecological importance of ECR coverage without social dimensions. As a policy tool, researchers, government, residents, and other stakeholders are supposed to work together on the delimitation of ECR (Bai et al., 2018). However, their negotiation may affect the marginal distribution of local patches, causing ambiguous effects on the identification of regional ecological conservation hotspots. Therefore, the proposed framework is more helpful in intuitively judging the ecological representativeness of ECR coverage. Methodologically, although the InVEST model and circuit theory are widely adopted in the construction of ecological networks (Cai et al., 2020; Luo & Wu, 2021), theoretical consensus on the determination of key parameters and thresholds, such as the width of ecological corridors, are still not

reached. However, exploratory analysis suggests that variation in thresholds has only marginal effects on the representativeness of ECR coverage. Lastly, our assessment does not include dynamic factors. The ecosystem is in a dynamic state under the interaction of biology and environment, and the ecological data of a single year may be accidental. Thus, carrying out more meaningful prioritization and clearer policy guidance becomes challenging. In the following research, long-term data will be synthesized to increase the reliability and robustness of the assessment.

## 5. Conclusions

In this study, we propose a comprehensive framework of “element-function-structure” to assess the of coverage of ECR areas in Zhejiang province, China. First, the InVEST model, MSPA analysis, and circuit theory are employed to identify three types of ecological conservation hotspots, and these hotspots are further used to assess the ecological representativeness of ECR coverage. Moreover, regional variation of ECR coverage among different counties is analyzed with regard to major function-oriented zones, the official zoning scheme to coordinate future development and protection pattern. Lastly, priority areas for conservation are outlined to leverage ecological benefit. This framework provides a new perspective and feasible approach to ensure that key components for ecological security are prioritized, contributing to the optimization of ECR coverage. The conclusions are as follows:

- ECR areas and ecological conservation hotspots often do not coincide. The retention rate of existing PAs in ECR areas is nearly 67%, followed by ecological function areas, which is approximately 48%. However, the existing ECR coverage is remarkably insufficient to protect potential ecological structure, namely, ecological corridors.
- The ecological representativeness of ERC areas corresponds to key function zones. In key ecological function areas, for example, ECR areas account for a relative large proportion of ecological lands.
- The priority areas for conservation suggest that regions in the southwest of Zhejiang province can be the leverage point to increase ecological benefits by maintaining ecological functions. While counties elsewhere with less ecological function can also contribute to ecological conservation, particularly by maintaining landscape connectivity via stepping stones.
- We argue that assessment of ECR coverage needs to account for multiple demands from conservation, ranging from biodiversity conservation to human well-being.

**Data Availability Statement.** Datasets that support findings of this article are available at <https://landbigdata.github.io/ECR/>

**Declaration of Competing Interest.** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Reference

- Bai, Y., Fang, Z., & Hughes, A. C. (2021). Ecological redlines provide a mechanism to maximize conservation gains in Mainland Southeast Asia. *One Earth*, 4(10), 1491-1504. doi:10.1016/j.oneear.2021.09.010
- Bai, Y., Wong, C. P., Jiang, B., Hughes, A. C., Wang, M., & Wang, Q. (2018). Developing China's Ecological Redline Policy using ecosystem services assessments for land use planning. *Nature Communications*, 9(1), 3034. doi:10.1038/s41467-018-05306-1
- Barber, C. P., Cochrane, M. A., Souza, C. M., & Laurance, W. F. (2014). Roads, deforestation, and the mitigating effect of protected areas in the Amazon. *Biological Conservation*, 177, 203-209. doi:10.1016/j.biocon.2014.07.004
- Bryan, B. A., Gao, L., Ye, Y., Sun, X., Connor, J. D., Crossman, N. D., . . . Hou, X. (2018). China's response to a national land-system sustainability emergency. *Nature*, 559(7713), 193-204. doi:10.1038/s41586-018-0280-2
- Buchadas, A., Qin, S., Meyfroidt, P., & Kuemmerle, T. (2022). Conservation frontiers: understanding the geographic expansion of conservation. *Journal of Land Use Science*, 1-14. doi:10.1080/1747423x.2021.2018516
- Busch, J., & Amarjargal, O. (2022). 100 global bright spots of green growth: Co-occurrence of nighttime light gain and forest gain, 1990–2015. *Global Environmental Change*, 75. doi:10.1016/j.gloenvcha.2022.102556
- Cai, A., Wang, J., MacLachlan, I., & Zhu, L. (2020). Modeling the trade-offs between urban development and ecological process based on landscape multi-functionality and regional ecological networks. *Journal of Environmental Planning and Management*, 63(13), 2357-2379. doi:10.1080/09640568.2020.1723507
- Carranza, D. M., Varas-Belemmi, K., De Veer, D., Iglesias-Müller, C., Coral-Santacruz, D., Méndez, F. A., . . . Gaymer, C. F. (2020). Socio-environmental conflicts: An underestimated threat to biodiversity conservation in Chile. *Environmental Science & Policy*, 110, 46-59. doi:10.1016/j.envsci.2020.04.006
- Chen, C., Park, T., Wang, X., Piao, S., Xu, B., Chaturvedi, R. K., . . . Myneni, R. B. (2019). China and India lead in greening of the world through land-use management. *Nature Sustainability*, 2, 122-129. doi:10.1038/s41893-019-0220-7
- Dai, L., Liu, Y., & Luo, X. (2021). Integrating the MCR and DOI models to construct an ecological security network for the urban agglomeration around Poyang Lake, China. *Science of the Total Environment*, 754, 141868. doi:10.1016/j.scitotenv.2020.141868
- de Jong, L., De Bruin, S., Knoop, J., & van Vliet, J. (2021). Understanding land-use change conflict: a systematic review of case studies. *Journal of Land Use Science*, 16(3), 223-239. doi:10.1080/1747423x.2021.1933226
- de Omena, M. T. R. N., & Hanazaki, N. (2022). How do Brazilian National Park managers evaluate the

- relationship between conservation and public use? *Environmental Science & Policy*, 131, 1-9.  
doi:10.1016/j.envsci.2022.01.009
- Deng, L., Liu, G. B., & Shangguan, Z. P. (2014). Land-use conversion and changing soil carbon stocks in  
China's 'Grain-for-Green' Program: a synthesis. *Global Change Biology*, 20(11), 3544-3556.  
doi:10.1111/gcb.12508
- Díaz, S., Zafra-Calvo, N., Purvis, A., Verburg, P. H., Obura, D., Leadley, P., . . . Zanne, A. E. (2020). Set  
ambitious goals for biodiversity and sustainability. *Science*, 370(6515), 411-413.  
doi:10.1126/science.abe1530
- Edgar, G. J., Stuart-Smith, R. D., Willis, T. J., Kininmonth, S., Baker, S. C., Banks, S., . . . Thomson, R. J.  
(2014). Global conservation outcomes depend on marine protected areas with five key features.  
*Nature*, 506(7487), 216-220. doi:10.1038/nature13022
- Ellis, E. C., & Ramankutty, N. (2008). Putting people in the map: anthropogenic biomes of the world.  
*Frontiers in Ecology and the Environment*, 6(8), 439-447. doi:10.1890/070062
- Fan, J. (2015). Draft of major function oriented zoning of China. *Acta Geographica Sinica*, 70(2), 186-  
201.
- Fan, J., Sun, W., Zhou, K., & Chen, D. (2012). Major Function Oriented Zone: New method of spatial  
regulation for reshaping regional development pattern in China. *Chinese Geographical Science*,  
22(2), 196-209. doi:10.1007/s11769-012-0528-y
- Folharini, S. d. O., Melo, S. N. d., & Cameron, S. R. (2021). Effect of protected areas on forest crimes in  
Brazil. *Journal of Environmental Planning and Management*, 65(2), 272-287.  
doi:10.1080/09640568.2021.1881885
- Gao, J. (2019). How China will protect one-quarter of its land. *Nature*, 569(7755), 457-458.  
doi:10.1038/d41586-019-01563-2
- Gao, J., Wang, Y., Zou, C., Xu, D., Lin, N., Wang, L., & Zhang, K. (2020). China's ecological conservation  
redline: A solution for future nature conservation. *Ambio*, 49(9), 1519-1529. doi:10.1007/s13280-  
019-01307-6
- Gao, J., Zou, C., Zhang, K., Xu, M., & Wang, Y. (2020). The establishment of Chinese ecological  
conservation redline and insights into improving international protected areas. *Journal of  
Environmental Management*, 264, 110505. doi:10.1016/j.jenvman.2020.110505
- Geldmann, J., Joppa, L. N., & Burgess, N. D. (2014). Mapping change in human pressure globally on land  
and within protected areas. *Conservation Biology*, 28(6), 1604-1616. doi:10.1111/cobi.12332
- Ghoddousi, A., Loos, J., & Kuemmerle, T. (2022). An Outcome-Oriented, Social-Ecological Framework  
for Assessing Protected Area Effectiveness. *BioScience*, 72(2), 201-212.  
doi:10.1093/biosci/biab114
- Gray, C. L., Hill, S. L., Newbold, T., Hudson, L. N., Borger, L., Contu, S., . . . Scharlemann, J. P. (2016).  
Local biodiversity is higher inside than outside terrestrial protected areas worldwide. *Nature  
Communications*, 7, 12306. doi:10.1038/ncomms12306
- Hu, T., Peng, J., Liu, Y., Wu, J., Li, W., & Zhou, B. (2020). Evidence of green space sparing to ecosystem  
service improvement in urban regions: A case study of China's Ecological Red Line policy.  
*Journal of Cleaner Production*, 251. doi:10.1016/j.jclepro.2019.119678
- Huang, J., Hu, Y., & Zheng, F. (2020). Research on recognition and protection of ecological security  
patterns based on circuit theory: a case study of Jinan City. *Environmental Science and Pollution  
Research*, 27(11), 12414-12427. doi:10.1007/s11356-020-07764-x
- Jones, K. R., Venter, O., Fuller, R. A., Allan, J. R., Maxwell, S. L., Negret, P. J., & Watson, J. E. M. (2018).  
One-third of global protected land is under intense human pressure. *Science*, 360(6390), 788-791.  
doi:10.1126/science.aap9565

- 501 Jones, N., Graziano, M., & Dimitrakopoulos, P. G. (2020). Social impacts of European Protected Areas  
502 and policy recommendations. *Environmental Science & Policy*, 112, 134-140.  
503 doi:10.1016/j.envsci.2020.06.004
- 504 Jones, N., Malesios, C., Kantartzis, A., & Dimitrakopoulos, P. G. (2020). The role of location and social  
505 impacts of protected areas on subjective wellbeing. *Environmental Research Letters*, 15(11).  
506 doi:10.1088/1748-9326/abb96e
- 507 Ke, X., Gui, S., Huang, H., Zhang, H., Wang, C., & Guo, W. (2017). Ecological risk assessment and source  
508 identification for heavy metals in surface sediment from the Liaohe River protected area, China.  
509 *Chemosphere*, 175, 473-481. doi:10.1016/j.chemosphere.2017.02.029
- 510 Kroner, R. E. G., Qin, S., Cook, C. N., Krithivasan, R., Pack, S. M., Bonilla, O. D., . . . Mascia, M. B.  
511 (2019). The uncertain future of protected lands and waters. *Science*, 364(6443), 881-886.  
512 doi:10.1126/science.aau5525
- 513 Liang, X., Jin, X., He, J., Wang, X., Xu, C., Qiao, G., . . . Zhou, Y. (2022). Impacts of land management  
514 practice strategy on regional ecosystems: Enlightenment from ecological redline adjustment in  
515 Jiangsu, China. *Land Use Policy*, 119. doi:10.1016/j.landusepol.2022.106137
- 516 Lin, J., Huang, C., Wen, Y., & Liu, X. (2021). An assessment framework for improving protected areas  
517 based on morphological spatial pattern analysis and graph-based indicators. *Ecological Indicators*,  
518 130. doi:10.1016/j.ecolind.2021.108138
- 519 Liu, J., Kuang, W., Zhang, Z., Xu, X., Qin, Y., Ning, J., . . . Chi, W. (2014). Spatiotemporal characteristics,  
520 patterns, and causes of land-use changes in China since the late 1980s. *Journal of Geographical*  
521 *Sciences*, 24(2), 195-210. doi:10.1007/s11442-014-1082-6
- 522 Liu, J., Linderman, M., Ouyang, Z., An, L., Yang, J., & Zhang, H. (2001). Ecological degradation in  
523 protected areas: the case of Wolong Nature Reserve for giant pandas. *Science*, 292(5514), 98-101.  
524 doi:10.1126/science.1058104
- 525 Luo, Y., & Wu, J. (2021). Linking the minimum spanning tree and edge betweenness to understand arterial  
526 corridors in an ecological network. *Landscape Ecology*, 36(5), 1549-1565. doi:10.1007/s10980-  
527 021-01201-1
- 528 Luo, Y., Wu, J., Wang, X., Wang, Z., & Zhao, Y. (2020). Can policy maintain habitat connectivity under  
529 landscape fragmentation? A case study of Shenzhen, China. *Science of the Total Environment*, 715,  
530 136829. doi:10.1016/j.scitotenv.2020.136829
- 531 Ma, K., Shen, X., Grumbine, R. E., & Corlett, R. (2017). China's biodiversity conservation research in  
532 progress. *Biological Conservation*, 210, 1-2. doi:10.1016/j.biocon.2017.05.029
- 533 Ma, S., Xue, M., & Ji, S. (2022). An improved emergy ecological footprint method for ecological security  
534 assessment and quantitative analysis of influencing factors: a case study of Zhejiang Province.  
535 *Journal of Environmental Planning and Management*, 1-25. doi:10.1080/09640568.2022.2088339
- 536 Marcantonio, M., Rocchini, D., Geri, F., Bacaro, G., & Amici, V. (2013). Biodiversity, roads, & landscape  
537 fragmentation: Two Mediterranean cases. *Applied Geography*, 42, 63-72.  
538 doi:10.1016/j.apgeog.2013.05.001
- 539 Maxwell, S. L., Cazalis, V., Dudley, N., Hoffmann, M., Rodrigues, A. S. L., Stolton, S., . . . Watson, J. E.  
540 M. (2020). Area-based conservation in the twenty-first century. *Nature*, 586(7828), 217-227.  
541 doi:10.1038/s41586-020-2773-z
- 542 McKinnon, M. C., Cheng, S. H., Dupre, S., Edmond, J., Garside, R., Glew, L., . . . Woodhouse, E. (2016).  
543 What are the effects of nature conservation on human well-being? A systematic map of empirical  
544 evidence from developing countries. *Environmental Evidence*, 5(1). doi:10.1186/s13750-016-  
545 0058-7
- 546 Mi, X., Feng, G., Hu, Y., Zhang, J., Chen, L., Corlett, R. T., . . . Ma, K. (2021). The global significance of

- 547 biodiversity science in China: an overview. *National Science Review*, 8(7), nwab032.  
548 doi:10.1093/nsr/nwab032
- 549 Miller-Rushing, A. J., Primack, R. B., Ma, K., & Zhou, Z.-Q. (2017). A Chinese approach to protected  
550 areas: A case study comparison with the United States. *Biological Conservation*, 210, 101-112.  
551 doi:10.1016/j.biocon.2016.05.022
- 552 Montoya, J. M., Pimm, S. L., & Solé, R. V. (2006). Ecological networks and their fragility. *Nature*,  
553 442(7100), 259-264. doi:10.1038/nature04927
- 554 O'Connor, L. M. J., Pollock, L. J., Renaud, J., Verhagen, W., Verburg, P. H., Lavorel, S., . . . Thuiller, W.  
555 (2021). Balancing conservation priorities for nature and for people in Europe. *Science*, 372(6544),  
556 856-860. doi:10.1126/science.abc4896
- 557 Oldekop, J. A., Holmes, G., Harris, W. E., & Evans, K. L. (2016). A global assessment of the social and  
558 conservation outcomes of protected areas. *Conservation Biology*, 30(1), 133-141.  
559 doi:10.1111/cobi.12568
- 560 Peng, J., Wang, A., Luo, L., Liu, Y., Li, H., Hu, Y., . . . Wu, J. (2019). Spatial identification of conservation  
561 priority areas for urban ecological land: An approach based on water ecosystem services. *Land*  
562 *Degradation & Development*, 30(6), 683-694. doi:10.1002/ldr.3257
- 563 Peng, J., Yang, Y., Liu, Y., Hu, Y., Du, Y., Meersmans, J., & Qiu, S. (2018). Linking ecosystem services  
564 and circuit theory to identify ecological security patterns. *Science of the Total Environment*, 644,  
565 781-790. doi:10.1016/j.scitotenv.2018.06.292
- 566 Porter-Bolland, L., Ellis, E. A., Guariguata, M. R., Ruiz-Mallén, I., Negrete-Yankelevich, S., & Reyes-  
567 García, V. (2012). Community managed forests and forest protected areas: An assessment of their  
568 conservation effectiveness across the tropics. *Forest Ecology and Management*, 268, 6-17.  
569 doi:10.1016/j.foreco.2011.05.034
- 570 Radeloff, V. C., Stewart, S. I., Hawbaker, T. J., Gimmi, U., Pidgeon, A. M., Flather, C. H., . . . Helmers,  
571 D. P. (2010). Housing growth in and near United States protected areas limits their conservation  
572 value. *Proceedings of the National Academy of Sciences*, 107(2), 940-945.  
573 doi:10.1073/pnas.0911131107
- 574 Ribeiro, S., Moreira, L. F. B., Overbeck, G. E., & Maltchik, L. (2021). Protected Areas of the Pampa  
575 biome presented land use incompatible with conservation purposes. *Journal of Land Use Science*,  
576 16(3), 260-272. doi:10.1080/1747423x.2021.1934134
- 577 Rocha, É. G. d., Brigatti, E., Niebuhr, B. B., Ribeiro, M. C., & Vieira, M. V. (2021). Dispersal movement  
578 through fragmented landscapes: the role of stepping stones and perceptual range. *Landscape*  
579 *Ecology*, 36(11), 3249-3267. doi:10.1007/s10980-021-01310-x
- 580 Rodrigues, A. S. L., & Cazalis, V. (2020). The multifaceted challenge of evaluating protected area  
581 effectiveness. *Nature Communications*, 11(1), 5147. doi:10.1038/s41467-020-18989-2
- 582 Rouget, M., Cowling, R. M., Lombard, A. T., Knight, A. T., & Kerley, G. I. (2006). Designing large-scale  
583 conservation corridors for pattern and process. *Conservation Biology*, 20(2), 549-561.  
584 doi:10.1111/j.1523-1739.2006.00297.x
- 585 Saura, S., Bodin, Ö., Fortin, M.-J., & Frair, J. (2014). Stepping stones are crucial for species' long-distance  
586 dispersal and range expansion through habitat networks. *Journal of applied ecology*, 51(1), 171-  
587 182. doi:10.1111/1365-2664.12179
- 588 Saura, S., & Pascual-Hortal, L. (2007). A new habitat availability index to integrate connectivity in  
589 landscape conservation planning: Comparison with existing indices and application to a case study.  
590 *Landscape and Urban Planning*, 83(2-3), 91-103. doi:10.1016/j.landurbplan.2007.03.005
- 591 Schmidt-Traub, G., Locke, H., Gao, J., Ouyang, Z., Adams, J., Li, L., . . . Wei, F. (2020). Integrating  
592 climate, biodiversity, and sustainable land-use strategies: innovations from China. *National*



*Science Review*, 8(7). doi:10.1093/nsr/nwaa139

- Viña, A., McConnell, W. J., Yang, H., Xu, Z., & Liu, J. (2016). Effects of conservation policy on China's forest recovery. *Science advances*, 2(3), e1500965. doi:10.1126/sciadv.1500965
- Wang, M., Jiang, B., Alatalo, J. M., Bai, Y., Wang, Q., Tan, J., . . . Su, J. (2021). Improved ecological monitoring for urban ecosystem protection in China. *Ecological Indicators*, 120. doi:10.1016/j.ecolind.2020.106950
- Wang, T., Li, H., & Huang, Y. (2021). The complex ecological network's resilience of the Wuhan metropolitan area. *Ecological Indicators*, 130. doi:10.1016/j.ecolind.2021.108101
- Wang, Y., Jin, C., Lu, M., & Lu, Y. (2017). Assessing the suitability of regional human settlements environment from a different preferences perspective: A case study of Zhejiang Province, China. *Habitat International*, 70, 1-12. doi:10.1016/j.habitatint.2017.09.010
- Wei, F., Cui, S., Liu, N., Chang, J., Ping, X., Ma, T., . . . Locke, H. (2020). Ecological civilization: China's effort to build a shared future for all life on Earth. *National Science Review*, 8(7). doi:10.1093/nsr/nwaa279
- Xu, W., Xiao, Y., Zhang, J., Yang, W., Zhang, L., Hull, V., . . . Ouyang, Z. (2017). Strengthening protected areas for biodiversity and ecosystem services in China. *Proceedings of the National Academy of Sciences*, 114(7), 1601-1606. doi:10.1073/pnas.1620503114
- Zhang, K., Zou, C., Lin, N., Qiu, J., Pei, W., Yang, Y., . . . Zhang, Z. (2022). The Ecological Conservation Redline program: A new model for improving China's protected area network. *Environmental Science & Policy*, 131, 10-13. doi:10.1016/j.envsci.2022.01.012
- Zhou, M., Deng, J., Lin, Y., Zhang, L., He, S., & Yang, W. (2021). Evaluating combined effects of socio-economic development and ecological conservation policies on sediment retention service in the Qiantang River Basin, China. *Journal of Cleaner Production*, 286. doi:10.1016/j.jclepro.2020.124961
- Zuo, L., & Gao, J. (2021). Investigating the compounding effects of environmental factors on ecosystem services relationships for Ecological Conservation Red Line areas. *Land Degradation & Development*, 32(16), 4609-4623. doi:10.1002/ldr.4059