

Wigner Resolution from Action Capacity

Brian S. Mulloy[✉]

Independent Researcher, Corktown, Detroit, Michigan, USA*

(Dated: 2026-06-11)

Phase-space formulations of quantum mechanics usually ignore that resolution is bounded below by Planck's quantum of action. In the Wigner formulation, the justification is that even though pointwise resolution forces the quasi-probability negative, those values carry physical structure. Zurek located that structure at scales set by the state's action capacity. We identify those scales as the principal axes of a family of de Gosson quantum blobs. The blob family deforms at constant action $\hbar/2$ and is circumscribed by the state's elliptical Heisenberg cell of action capacity $\pi \Delta x \Delta p$. Convolution of the Wigner function to the scale of each $\hbar/2$ blob restores resolution to Planck's quantum of action and renders it non-negative. Integrating over the family reaches the resolution $\pi \hbar^2 / (\Delta x \Delta p)$, the symplectic polar dual of the Heisenberg cell. The resulting portrait is everywhere non-negative and preserves the structure of the Wigner function. This resolution carries no free parameter and sharpens as the action capacity grows.

The Wigner function represents a quantum state in phase space [1, 2]. It is defined pointwise, below Planck's quantum of action, and Hudson's theorem forces it negative there [3]. Its negativity is a resource for quantum advantage [4, 5] and a measure of nonclassicality [6], while a non-negative portrait admits a probability interpretation [7–10] and supports classical sampling [11]. Here a family of $\hbar/2$ quantum blobs, circumscribed by the state's action capacity $\pi \Delta x \Delta p$, resolves the Wigner function to $\pi \hbar^2 / (\Delta x \Delta p)$, the resolution symplectic polar duality sets on any non-negative portrait.




FIG. 1. Heisenberg cell, principal quantum blobs, and quorum cell. The Heisenberg cell is the state's covariance ellipsoid with action capacity $A = \pi \Delta x \Delta p \geq \hbar/2$. The quantum blob family (Fig. 2) has two members at the principal axes, $a_{\theta=0} = \pi \Delta x \delta p = \hbar/2$ and $a_{\pi/2} = \pi \delta x \Delta p = \hbar/2$. Zurek's relations $\delta p \sim \hbar/\Delta x$ and $\delta x \sim \hbar/\Delta p$ locate the semi-axis scales from the state's action capacity. The quorum cell $\tilde{a}$ is an ellipse with semi-axes $\delta x, \delta p$ and action $\tilde{a} = \pi \delta x \delta p$. A circumscribes $a_{\theta=0}$ and $a_{\pi/2}$, which circumscribe $\tilde{a}$.




FIG. 2. Quadrature family of quantum blobs and quorum cell. The Heisenberg cell A circumscribes the quantum blob family $\{a_{\theta}\}$, which circumscribes the quorum cell $\tilde{a}$. Across θ the family deforms at constant action $a_{\theta} = \hbar/2$. At the principal angles $\theta = 0$ and $\theta = \pi/2$, the family recovers $a_{\theta=0}$ and $a_{\pi/2}$ from Fig. 1.

action [12]. Heisenberg's uncertainty principle $\sigma_x \sigma_p \geq \hbar/2$ [13–15] fixes the limit. Geometrically, de Gosson gave this action region the form of the state's covariance ellipsoid, with symplectic capacity $\pi \Delta x \Delta p$ [16] and semi-axes $\Delta x = \sqrt{2} \sigma_x$ and $\Delta p = \sqrt{2} \sigma_p$. This is the Heisenberg cell of action capacity (Figs. 1, 2),

$$A = \pi \Delta x \Delta p \geq \hbar/2. \quad (1)$$

The Wigner function carries structure at small scales. Zurek's relations $\delta p \sim \hbar/\Delta x$ and $\delta x \sim \hbar/\Delta p$ locate those scales from the state's action capacity [17]. We identify them as the principal axes of a family of de Gosson quantum blobs (Fig. 1),

$$a_{\theta=0} = \pi \Delta x \delta p = \hbar/2, \quad a_{\pi/2} = \pi \delta x \Delta p = \hbar/2. \quad (2)$$

Planck divided phase space into elementary regions of




FIG. 3. **Harmonic oscillator.** Rows correspond to Fock states $n = 0, 1, 2, 20$ with action capacity $A/(h/2) = 2n + 1$. Column 1: Wigner function $W(x, p)$ (negativity in blue) overlaid with the Heisenberg cell A and quorum cell $\tilde{a}$. Centered at the covariance origin, $\tilde{a}$ marks the scale of the finest structure (the Zurek scale) of the bare, unconvolved Wigner function. Column 2: Cross-section $W(x, 0)$ exhibiting negative dips. Column 3: Quantum kernel $K_{\pi/2}$ overlaid with A and the quantum blob $a_{\pi/2}$ (action $h/2$, restoring resolution up to Planck's quantum of action). Column 4: Convolved cross-section $\tilde{W}_{\pi/2}(x, 0)$ at quadrature angle $\theta = \pi/2$, which is non-negative. Column 5: The portrait $\tilde{W}(x, p)$, integrated over the blob family across all angles, overlaid with A and $\tilde{a}$ (symplectic polar duals). It is everywhere non-negative, with positive lobes matching the count and position of those in W , resolved at scale $\tilde{a}$. For the vacuum ($n = 0$), A and $\tilde{a}$ coincide as a Gaussian blob at $A = h/2$. As n increases, A grows and $\tilde{a}$ reciprocally shrinks ($\tilde{a} A = (h/2)^2$). At $n = 20$, the rings condense toward a classical energy trajectory, leaving $\tilde{a}$ as a fine central cell. Axes are in the oscillator scales $x_0 = \sqrt{\hbar/m\omega}$ and $p_0 = \sqrt{\hbar m\omega}$.

The capacity $\Delta x, \Delta p$ runs along the major axes, the resolution $\delta x, \delta p$ along the minor.

Between these principal members the quantum blob a_θ deforms across $\theta \in [0, \pi)$ at constant action $h/2$ (Fig. 2), the preimage of the rigid ellipse $\hat{a}_\theta$ under the scaling $X = x/\Delta x$, $P = p/\Delta p$,

$$a_\theta = \{(x, p) : (X, P) \in \hat{a}_\theta\}. \quad (3)$$

The rotated quadratures $X_\theta = X \cos \theta + P \sin \theta$ and $P_\theta = -X \sin \theta + P \cos \theta$ are the homodyne quadratures of optical tomography at phase θ . In the scaled coordinates $\hat{a}_\theta$ is the rotation by θ of a single ellipse,

$$\hat{a}_\theta = \left\{ (X, P) : X_\theta^2 + \frac{A^2}{(\pi\hbar)^2} P_\theta^2 \leq 1 \right\}. \quad (4)$$

The Heisenberg cell A circumscribes each blob a_θ .

The Wigner function of the state is

$$W(x, p) = \frac{1}{\pi\hbar} \int \psi^*(x+s) \psi(x-s) e^{2ips/\hbar} ds. \quad (5)$$

The Wigner function of each quantum blob a_θ is the centered Gaussian whose $h/2$ ellipse is a_θ ,

$$K_\theta(x, p) = \frac{1}{\pi\hbar} \exp \left[-X_\theta^2 - \frac{A^2}{(\pi\hbar)^2} P_\theta^2 \right]. \quad (6)$$

The convolution of W with each quantum kernel K_θ is

$$\tilde{W}_\theta(x, p) = (W * K_\theta)(x, p). \quad (7)$$

Convolving W with the kernel K_θ of a single quantum blob a_θ lifts it up to Planck's quantum of action and




FIG. 4. **Squeezed vacuum, eigenstates, and the Kerr crescent.** Rows correspond to the squeezed vacuum ($r = 0.5$), the Morse eigenstate ($n = 8$), the double-well eigenstate ($n = 5$), and the Kerr crescent, a coherent state sheared at constant photon number. Columns are as in Fig. 3. The Kerr state is shown in the principal frame of its covariance, where the cell overlays are axis-aligned. The portrait $\widetilde{W}$ (Column 5) remains strictly non-negative across all four states, resolved at the scale of $\tilde{a}$.

yields a non-negative function [8, 9, 18]. The convolution $\widetilde{W}_\theta$ is non-negative everywhere.

Zurek also located the scale $a \sim \hbar \times (\hbar/A)$ [17, 19] of the chessboard structure in the compass state (Fig. 5, column 1). We identify its rotationally invariant form as the quorum cell $\tilde{a}$. Named after the quorum of observables in optical tomography [20], $\tilde{a}$ is the ellipse with semi-axes $\delta x, \delta p$ (Figs. 1, 2) and action

$$\tilde{a} = \pi \delta x \delta p = \frac{\pi \hbar^2}{\Delta x \Delta p}. \quad (8)$$

$\tilde{a}$ is the scale of the state's finest structure in the bare Wigner function (Figs. 3, 4, 5, column 1). The quorum cell $\tilde{a}$ is the symplectic polar dual of the Heisenberg cell A [21, 22]. The duality is reflexive, so A and $\tilde{a}$ each fix the other, and antimonotone, so $\tilde{a}$ shrinks as A grows. Each blob a_θ is circumscribed by A and circumscribes $\tilde{a}$, nesting as $A \supseteq a_\theta \supseteq \tilde{a}$ (Fig. 2). A quorum of these a_θ blobs, each respecting $\hbar/2$, reaches $\tilde{a}$ together. Resolu-

tion and capacity are reciprocal at the quantum of action, $\tilde{a} A = (\hbar/2)^2$.

Integrating $\widetilde{W}_\theta$ over the family gives the phase-space portrait of the state at the resolution $\tilde{a}$,

$$\widetilde{W}(x, p) = \frac{1}{\pi} \int_0^\pi \widetilde{W}_\theta(x, p) d\theta. \quad (9)$$

$\widetilde{W}_\theta \geq 0$ for every θ , so $\widetilde{W} \geq 0$ everywhere. Tomographic reconstruction inverts marginals pointwise to recover W [23–25]. Over the same θ , convolving W with the quantum kernel family $\{K_\theta\}$ builds the non-negative portrait $\widetilde{W}$.

A quantum blob a_θ too large for A to circumscribe, or too small to circumscribe $\tilde{a}$, is a blob of another state and portrays that state, not this one [21, 26]. Because every blob a_θ of this state circumscribes the quorum cell $\tilde{a}$, and a non-negative portrait $\widetilde{W}$ smooths W by a Gaussian of action at least $\hbar/2$ [8, 9], $\tilde{a}$ is the finest resolution




FIG. 5. **Cat states.** Rows correspond to the 2-cat, 3-cat, 4-cat, and Zurek compass states. Columns are as in Fig. 3. As before, the portrait $\tilde{W}$ (Column 5) is everywhere non-negative, with positive lobes matching the count and location of those in W , resolved at scale $\tilde{a}$. The 4-cat and Zurek compass states differ by a $\pi/4$ rotation in phase space.

any non-negative portrait attains, and the family $\{a_\theta\}$ reaches it across $\theta \in [0, \pi)$.

The principal frame, where the Heisenberg cell A is axis-aligned, costs no generality. The Wigner function transforms covariantly under the symplectic group [18], so the map that diagonalizes a state's covariance carries it to this frame at preserved action. The Kerr crescent, a sheared coherent state, is resolved this way (Fig. 4). In one degree of freedom, A fixes the family uniquely. In higher dimensions, action $h/2$ no longer singles out a blob [26], symplectic capacity and volume diverge, and the higher-dimensional case remains for future work.

A fixed vacuum-width Gaussian [7] matches the state only at the vacuum $A \sim h/2$ and for larger states erases structure at resolution $\tilde{a}$. Methods with a free parameter [27–29] can resolve more finely, the scale chosen as a parameter. The resolution here carries no free parameter, fixed by the action capacity A alone.

Figures 3, 4, and 5 show numerical results across twelve states. The cross spectral phase between W and $\tilde{W}$ vanishes, so the portrait holds every feature in place, and $\tilde{W}$

is non-negative to machine precision [30].

The uncertainty product $\sigma_x \sigma_p$ grows unbounded, the micro-to-macro transition Heisenberg [31] and Bohr [32] identified as requiring special care. The quantum blob does not. As A grows, the family $\{a_\theta\}$ deforms at fixed action $h/2$. Squeezing reaches the same floor by external apparatus [33], while here the deformation is inherent to the state. A Schrödinger cat with capacity $\Delta x \sim 10$ cm and $\Delta p \sim 1$ kg·m/s has positional resolution $\delta x = \hbar/\Delta p \sim 10^{-34}$ m, vastly smaller than the radioactive atom that triggers its apparatus [34]. The correspondence principle follows from $A \rightarrow \infty$, the physical limit underlying the heuristic $h \rightarrow 0$.

The scale called sub-Planck is the minor axis δ of an $h/2$ blob, narrowing as the conjugate capacity Δ grows, the scale Zurek tied to a state's sensitivity [17, 19] and the scale to which $\tilde{W}$ resolves. The negative Wigner function W and its non-negative portrait $\tilde{W}$ have their structure at a single scale $\tilde{a}$, set by the state's action capacity A . Zurek scales and de Gosson quantum blobs give the res-

olution. Heisenberg's product gives the capacity. As the action capacity grows to $\pi \Delta x \Delta p \geq h/2$ the resolution sharpens to $\pi \hbar^2 / (\Delta x \Delta p) \leq h/2$. We read this structure as the geometry of phase space itself, which de Gosson takes as prior to the uncertainty principle [35].

The author thanks M. de Gosson for helpful comments. This research received no external funding. Numerical computations were performed with QuTiP [36] and SciPy [37]. The code reproducing the numerical results, together with Supplemental Material mapping this Letter onto Zurek and de Gosson, is openly available [30].

* bmulloy@umich.edu

- [1] E. Wigner, On the quantum correction for thermodynamic equilibrium, *Physical Review* **40**, 749 (1932).
- [2] M. Hillery, R. F. O'Connell, M. O. Scully, and E. P. Wigner, Distribution functions in physics: Fundamentals, *Physics Reports* **106**, 121 (1984).
- [3] R. L. Hudson, When is the Wigner quasi-probability density non-negative?, *Reports on Mathematical Physics* **6**, 249 (1974).
- [4] V. Veitch, C. Ferrie, D. Gross, and J. Emerson, Negative quasi-probability as a resource for quantum computation, *New Journal of Physics* **14**, 113011 (2012).
- [5] A. Mari and J. Eisert, Positive Wigner functions render classical simulation of quantum computation efficient, *Physical Review Letters* **109**, 230503 (2012).
- [6] A. Kenfack and K. Życzkowski, Negativity of the Wigner function as an indicator of non-classicality, *Journal of Optics B: Quantum and Semiclassical Optics* **6**, 396 (2004).
- [7] K. Husimi, Some formal properties of the density matrix, *Proc. Phys.-Math. Soc. Jpn., 3rd Ser.* **22**, 264 (1940).
- [8] N. D. Cartwright, A non-negative Wigner-type distribution, *Physica A: Statistical Mechanics and its Applications* **83**, 210 (1976).
- [9] M. de Gosson, Cellules quantiques symplectiques et fonctions de Husimi–Wigner, *Bulletin des Sciences Mathématiques* **129**, 211 (2005).
- [10] D. Dragoman, Phase space formulation of quantum mechanics. Insight into the measurement problem, *Phys. Scr.* **72**, 290 (2005).
- [11] A. Polkovnikov, Phase space representation of quantum dynamics, *Annals of Physics* **325**, 1790 (2010).
- [12] M. Planck, *Vorlesungen über die Theorie der Wärmestrahlung* (Johann Ambrosius Barth, Leipzig, 1906) english translation: M. Masius, *The Theory of Heat Radiation* (Blakiston, Philadelphia, 1914).
- [13] E. H. Kennard, Zur Quantenmechanik einfacher Bewegungstypen, *Zeitschrift für Physik* **44**, 326 (1927).
- [14] H. P. Robertson, The uncertainty principle, *Physical Review* **34**, 163 (1929).
- [15] E. Schrödinger, Zum Heisenbergschen Unschärfeprinzip, *Sitzungsber. Preuss. Akad. Wiss., Phys.-Math. Kl.*, 296 (1930).
- [16] M. A. de Gosson, The symplectic camel and the uncertainty principle: The tip of an iceberg?, *Foundations of Physics* **39**, 194 (2009).
- [17] W. H. Zurek, Sub-Planck structure in phase space and its relevance for quantum decoherence, *Nature* **412**, 712 (2001).
- [18] E. Cordero, M. de Gosson, M. Dörfler, and F. Nicola, On the symplectic covariance and interferences of time-frequency distributions, *SIAM Journal on Mathematical Analysis* **50**, 2178 (2018).
- [19] W. H. Zurek, Decoherence, einselection, and the quantum origins of the classical, *Reviews of Modern Physics* **75**, 715 (2003).
- [20] G. M. D'Ariano, L. Maccone, and M. G. A. Paris, Quorum of observables for universal quantum estimation, *Journal of Physics A: Mathematical and General* **34**, 93 (2001).
- [21] M. de Gosson and C. de Gosson, Symplectic polar duality, quantum blobs, and generalized Gaussians, *Symmetry* **14**, 1890 (2022).
- [22] M. A. de Gosson, Polar duality and the reconstruction of quantum covariance matrices from partial data, *Journal of Physics A: Mathematical and Theoretical* **57**, 205303 (2024).
- [23] K. Vogel and H. Risken, Determination of quasiprobability distributions in terms of probability distributions for the rotated quadrature phase, *Physical Review A* **40**, 2847 (1989).
- [24] A. I. Lvovsky and M. G. Raymer, Continuous-variable optical quantum-state tomography, *Reviews of Modern Physics* **81**, 299 (2009).
- [25] M. A. de Gosson, Symplectic Radon transform and the metaplectic representation, *Entropy* **24**, 761 (2022).
- [26] M. A. de Gosson, Quantum blobs, *Foundations of Physics* **43**, 440 (2013).
- [27] K. Wódkiewicz, Operational approach to phase-space measurements in quantum mechanics, *Phys. Rev. Lett.* **52**, 1064 (1984).
- [28] D. M. Appleby, Generalized Husimi functions: Analyticity and information content, *J. Mod. Opt.* **46**, 825 (1999).
- [29] K. E. Cahill and R. J. Glauber, Density operators and quasiprobability distributions, *Physical Review* **177**, 1882 (1969).
- [30] B. S. Mulloy, Code, figures, and supplemental material for “Wigner Resolution from Action Capacity” (2026), the supplemental material is archived in this deposit, under the same DOI.
- [31] W. Heisenberg, The physical content of quantum kinematics and mechanics, in *Quantum Theory and Measurement*, edited by J. A. Wheeler and W. H. Zurek (Princeton University Press, 1983) pp. 62–84.
- [32] N. Bohr, The quantum postulate and the recent development of atomic theory, *Nature* **121**, 580 (1928).
- [33] D. F. Walls, Squeezed states of light, *Nature* **306**, 141 (1983).
- [34] E. Schrödinger, The present situation in quantum mechanics, *Proceedings of the American Philosophical Society* **124**, 323 (1980), translated by J. D. Trimmer from *Naturwissenschaften* **23** (1935) 807–812, 823–828, 844–849.
- [35] M. A. de Gosson, On quantum indeterminacy and the uncertainty principle (2026), arXiv:2605.01103 [quant-ph].
- [36] N. Lambert *et al.*, QuTiP 5: The quantum toolbox in Python, *Phys. Rep.* **1153**, 1 (2025).
- [37] P. Virtanen *et al.*, SciPy 1.0: Fundamental algorithms for scientific computing in Python, *Nat. Methods* **17**, 261 (2020).