

M1C03 Lecture 23

Functions

Jeremy Lane

Nov 8, 2021

Announcement(s)

- ① Test 1 results are out
- ② Quiz due Friday
- ③ Assignment 4 posted

Definition and examples of functions

Reference: Notes on functions (Avenue), Lakins Chapter 5

A *function* consists of three things:

- a set X called the *domain*,
- a set Y called the *codomain*, and
- a *correspondence* (or rule, or formula) that assigns to every element of the domain a unique element of the codomain.

Let

$$\mathcal{L} = \{“a”, “b”, \dots, “z”\}, \quad \mathcal{U} = \{“A”, “B”, \dots, “Z”\}, \quad \mathcal{A} = \mathcal{L} \cup \mathcal{U}.$$

Let *Cap* be the correspondence that converts a letter to upper case.

Let *Low* be the correspondence that converts a letter to lower case.

Consider the set of binary sequences of length 4,

$$B_4 = \left\{ \begin{array}{cccccccc} 0000, & 1000, & 0100, & 0010, & 0001, & 1100, & 1010, & 1001, \\ 0101, & 0011, & 0110, & 1110, & 1101, & 1011, & 0111, & 1111 \end{array} \right\}.$$

- (bit flip) $n: B_4 \rightarrow B_4$ flips each digit in the sequence.
 - (right shift) $r: B_4 \rightarrow B_4$ shifts every digit to the right. The leftmost digit becomes 0 and the rightmost digit disappears.
 - (left shift) $l: B_4 \rightarrow B_4$ shifts every digit to the left. The rightmost digit becomes 0 and the leftmost digit disappears.
-
- $t: B_4 \rightarrow B_3$ removes the leftmost digit.
 - $a: B_3 \rightarrow B_4$ appends 0 as the leftmost digit.

Consider the correspondences:

- $f(x) = \sqrt{x}$ (where \sqrt{x} means the non-negative square root of x), and
- $g(x) = x^2$.

Example

Let $\mathcal{C} = \{\text{"black"}, \text{"red"}, \text{"blue"}, \text{"yellow"}, \text{"white"}, \text{"turquoise"}\}$. Let \mathcal{S} be the set of students in the classroom.

Let s be the correspondence that assigns a colour in \mathcal{C} to students in the classroom that are wearing clothes of that colour.

Does this define a function $s: \mathcal{C} \rightarrow \mathcal{S}$? Why or why not?

Example

Let P be the set of polynomials of any degree in a variable x .

What are some examples of functions whose domain and codomain involves P ?

Example

Let $M_{m \times n}(\mathbb{R})$ be the set of $m \times n$ matrices (a $m \times n$ square array of numbers).

What are some examples of functions whose domain and codomain involves $M_{m \times n}(\mathbb{R})$?

Example

What are some examples of functions in real-life?