

Applying Google Earth Engine to Wildfire Disturbance Detection in the State of Alaska

Forrest M. Hoffman^{1,2}, Zachary L. Langford²,
Jitendra Kumar^{1,2}, Steve Norman³, and William W. Hargrove³

¹Oak Ridge National Laboratory, ²University of Tennessee Knoxville, and
³USDA Forest Service Eastern Forest Environmental Threat Assessment Center


2018 US-IALE Annual Meeting
Chicago, Illinois, USA
April 11, 2018

CLIMATE CHANGE
SCIENCE INSTITUTE
OAK RIDGE NATIONAL LABORATORY




Motivations

- ▶ Can we estimate wildfire area extents based on the 2004 wildfire season?
- ▶ Can we find the most important attributes from MODIS sensors and daily meteorological variables using Google Earth Engine?
- ▶ Can we locate fires located under 1000 acres? (<1000 acres is cutoff from Monitoring Trends in Burn Severity (MTBS) dataset)




Numerous forest fires (outlined in red) were burning in the Yukon Flats region of east-central Alaska in mid-June 2004 captured by MODIS sensor. (Image Source: NASA)




Monitoring Trends in Burn Severity (MTBS) dataset for Alaska from 2000-2015, with 2004 having the most area (km^2) burned.

2004 Wildfire Season

- ▶ One of the warmest and driest summer on record.
- ▶ Most lightning strikes recorded during summer.
- ▶ Wildland fires burned the largest acreage in recorded Alaska history.




Departure from average temperature across Alaska for every year since 1949. (Image Source: Alaska Climate Research Center)



Number of lightnings strikes (6538) in Alaska from June 5 – 19 2004. The grand total was over 147,642 strikes [2].


Study Region

- ▶ Study region is based on climate-division boundaries in Interior Alaska [1].
 - ▶ Wildfires are persistent in this region due to climate patterns, for example, the length of the growing season has increased 45% over the last century [3].
-
- ▶ Monitoring Trends in Burn Severity (MTBS) dataset was used for training MODIS and Daymet variables.
 - ▶ Binary classification with 0 representing no wildfire and 1 representing a wildfire.
- 
- The map displays a satellite view of a region in Interior Alaska, characterized by its rugged terrain and diverse land cover. A large black polygon outlines the study area. Numerous red spots of varying sizes are scattered across the map, representing wildfire locations. The legend on the right side provides a key for both land cover types and wildfire extent. Land cover categories include Open Water, Perennial Ice/Snow, Developed, Open Space, Developed, Low Intensity, Developed, Medium Intensity, Developed, High Intensity, Barren Land (Rock/Sand/Clay), Unconsolidated Shore, Deciduous Forest, Evergreen Forest, Mixed Forest, Dwarf Scrub, Shrub/Scrub, Grassland/Herbaceous, Sedge/Herbaceous, Lichens, Moss, Pasture/Hay, Cultivated Crops, Woody Wetlands, and Emergent Herbaceous Wetlands. The scale bar at the bottom indicates distances up to 400 km, and a north arrow is present in the bottom right corner.


Geospatial Datasets

We used Google Earth Engine (GEE) for processing images and building models. Two types of datasets were looked into:

- ▶ MODIS datasets of MOD09A1 (Surface Reflectance 8-Day L3 Global 500m) and MOD11A2 (Land Surface Temperature and Emissivity 8-Day L3 Global 1km) were analyzed [7, 8].
- ▶ Daymet, which is a collection of algorithms designed to interpolate and extrapolate from daily meteorological observations to produce gridded estimates of daily weather parameters [6].



Daymet V3 average annual minimum temperature for 1980 and 2015 for a subset of the Daymet domain in Alaska and western Canada.




MOD09A1 RGB composite from June 17, 2004.

Geospatial Datasets

Description	Resolution	Variable
GMTED2010	225 m	elevation (m)
	225 m	slope (%)
MOD09A1	500 m at 8 days	NDVI
	500 m at 8 days	EVI
	500 m at 8 days	SAVI
	500 m at 8 days	Bands 1–7 (459–2155 nm)
MOD11A2	1 km at 8 days	Daytime LST (Kelvin)
Daymet	1 km at daily	Daylight period (seconds)
	1 km at daily	Precipitation (mm)
	1 km at daily	Snow water equivalent (kg/m^2)
	1 km at daily	Maximum temperature ($^{\circ}\text{C}$)
	1 km at daily	Minimum temperature ($^{\circ}\text{C}$)
	1 km at daily	Shortwave Radiation (W/m^2)
	1 km at daily	Vapor Pressure (Pa)

Daymet and MODIS products were processed from early-April through late-October in 2004.

Data Workflow



- ▶ Perform image processing methods for classification.
- ▶ Build models with MODIS (288), MODIS/Daymet (456), and Daymet (168) variables and the MTBS dataset.
- ▶ Right Figure: Google Earth Engine interactive development environment [5].






Image Processing

- ▶ Increased resolution to 500 m for all datasets, GEE performs nearest neighbor resampling.
- ▶ Linear interpolation for missing values.
- ▶ Savitzky-Golay filter was applied to smooth out noise [4].
- ▶ Daymet variables were merged into 8-day averages.




Example image processing workflow applied to a large wildfire, which occurred on June 13, 2004.



Fire severity for the Boundary fire based on Landsat 7.
(Source: USGS and US Forest Service)


Random Forest

- ▶ Random Forest: estimator that fits a number of decision tree classifiers on various sub-samples of the dataset and use averaging to improve the predictive accuracy and control over-fitting.
- ▶ We split up dataset based on 100×100 pixel tiles, with 163 tiles used for training and 73 tiles used for testing.




Results: MODIS

Validation Metrics



- Precision, recall, and F1-score for non-wildfire class ($n=493710$) was 0.99, 1.00, and 1.00, respectively.
- Precision, recall, and F1-score for wildfire class ($n=16878$) was 0.93, 0.78, and 0.85, respectively.


Feature Importance



- EVI, NDVI, and MODIS Red band contributed the most using the Gini feature importance metric.
- Elevation, slope, SAVI, and SWIR1 contributed the least.


Results: Daymet

Validation Metrics



- ▶ Precision, recall, and F1-score for non-wildfire class ($n=493710$) was 0.99, 1.00, and 0.99, respectively.
- ▶ Precision, recall, and F1-score for wildfire class ($n=16878$) was 0.62, 0.53, and 0.57, respectively.


Feature Importance



- ▶ Most variables contributed equally for feature importance.
- ▶ Snow water equivalent, minimum temperature, and precipitation were the highest scoring features.


Results: MODIS/Daymet

Validation Metrics




- Precision, recall, and F1-score for non-wildfire class ($n=493710$) was 0.99, 1.00, and 1.00, respectively.
- Precision, recall, and F1-score for wildfire class ($n=16878$) was 0.93, 0.78, and 0.85, respectively.

Feature Importance




- Daymet variables (minimum temperature, maximum temperature, shortwave radiation) contributed most for Daymet/MODIS classification.
- MODIS Green, Red, and SAVI variables contributed the most.

Results: Test (Bonanza Creek Wildfire)



Fire severity for the Bonanza Creek wildfire based on Landsat 7. (Source: USGS and US Forest Service)



Conclusions

- ▶ MODIS (NIR, SWIR), indices (EVI, NDVI, SAVI), and Daymet variables (minimum temperature, maximum temperature, snow water equivalent, shortwave radiation) are the most important drivers for this study.
- ▶ Random Forest provides a good approach for understanding feature importance.
- ▶ Google Earth Engine provides a great platform for processing and analyzing datasets.
- ▶ Future Work:
 - ▶ Does varying responses in different years provide a better classification?
 - ▶ Investigate class imbalance problem when subsetting data into training and testing.
 - ▶ Does including daily meteorological variables for entire year provide better results?

Acknowledgements



U.S. DEPARTMENT OF
ENERGY

Office of Science

This research was sponsored by the Climate and Environmental Sciences Division (CESD) of the Biological and Environmental Research (BER) Program in the US Department of Energy Office of Science and the US Department of Agriculture Forest Service, and the National Science Foundation (AGS-1048890). This research used resources of the Oak Ridge Leadership Computing Facility (OLCF) at Oak Ridge National Laboratory (ORNL), which is managed by UT-Battelle, LLC, for the US Department of Energy under Contract No. DE-AC05-00OR22725.

References I

-  Peter A. Bieniek, Uma S. Bhatt, Richard L. Thoman, Heather Angeloff, James Partain, John Papineau, Frederick Fritsch, Eric Holloway, John E. Walsh, Christopher Daly, Martha Shulski, Gary Hufford, David F. Hill, Stavros Calos, and Rudiger Gens.
Climate divisions for alaska based on objective methods.
Journal of Applied Meteorology and Climatology,
51(7):1276–1289, 2012.

References II

-  F. Stuart Chapin, Sarah F. Trainor, Orville Huntington, Amy L. Lovecraft, Erika Zavaleta, David C. Natcher, A. David McGuire, Joanna L. Nelson, Lily Ray, Monika Calef, Nancy Fresco, Henry Huntington, T. Scott Rupp, La'ona DeWilde, and Rosamond L. Naylor.
Increasing wildfire in alaska's boreal forest: Pathways to potential solutions of a wicked problem.
BioScience, 58(6):531–540, 2008.
-  F. Stuart Chapin III, Sarah F. Trainor, Patricia Cochran, Henry Huntington, Carl J. Markon, Molly McCommon, A. David McGuire, and Mark Serreze.
Alaska.
Technical report, Washington, D.C., 2014.
Report.

References III

-  Jin Chen, Per. Jnsson, Masayuki Tamura, Zhihui Gu, Bunkei Matsushita, and Lars Eklundh.
A simple method for reconstructing a high-quality ndvi time-series data set based on the savitzkygolay filter.
Remote Sensing of Environment, 91(3):332 – 344, 2004.
-  Noel Gorelick, Matt Hancher, Mike Dixon, Simon Ilyushchenko, David Thau, and Rebecca Moore.
Google earth engine: Planetary-scale geospatial analysis for everyone.
Remote Sensing of Environment, 202:18 – 27, 2017.
Big Remotely Sensed Data: tools, applications and experiences.

References IV

-  P.E. Thornton, M.M. Thornton, B.W. Mayer, Y. Wei,
R. Devarakonda, R.S. Vose, and R.B. Cook.
Daymet: Daily surface weather data on a 1-km grid for north
america, version 3, 2017.
-  Eric Vermote.
Mod09a1 modis/terra surface reflectance 8-day l3 global 500m
sin grid v006, 2015.
-  Z. Wan, S. Hook, and G. Hulley.
Mod11a2 modis/terra land surface temperature/emissivity
8-day l3 global 1km sin grid v006, 2015.