

# Regress adaptation index on a feature

YC, LL, LG

December 12, 2022

We observe data points  $(a_i, b_i, x_i)$ , for  $i = 1, \dots, n$ .  $a_i$  is the control response,  $b_i$  is the test response,  $x_i$  is a feature extracted from the input image presented (e.g. contrast, orientation, etc.). For simplicity, we assume  $x_i$  is univariate, even though the theory can be extended to the multivariate situation. We are interested in estimating the effect of  $x$  on the adaptation index. In the noiseless setting, it is just finding a functional relationship

vector output from ANN

$$\frac{b_i}{a_i} = f(x_i).$$

If we restrict ourselves to linear function,  $f(x) = x \cdot \beta + c$  and  $\beta$  can be estimated via linear regression of  $\frac{b_i}{a_i}$  on  $x_i$ . In the presence of noise, the situation is a bit more complicated. There are two natural estimators.

1. **Estimator 1: take ratio then regress.** We calculate the ratio  $\frac{b_i}{a_i}$  and the regress  $\frac{b_i}{a_i}$  on  $x_i$  to find  $\beta$ .

$$\hat{\beta}_{1, -} = \arg \min_{\beta, \beta_0} \sum_{i=1}^n \left( \frac{b_i}{a_i} - x_i \cdot \beta - \beta_0 \right)^2.$$

2. **Estimator 2: regress  $b_i$ .** Regress  $b_i$  on  $a_i \cdot x_i$  and  $a_i$ , get the coefficient in front  $a_i \cdot x_i$  as an estimator of  $\beta$ .

$$\hat{\beta}_{2, -, -} = \arg \min_{\beta, \gamma, \gamma_0} \sum_{i=1}^n (b_i - (a_i x_i) \cdot \beta - a_i \gamma - \gamma_0)^2$$

or predict R1 from stim feature via ANN

3. **Estimator 3: two-stage estimator.** First, regress  $a_i$  on  $x_i$  to obtain  $\alpha, \alpha_0$ . Then regress  $b_i$  on  $((\alpha x_i + \alpha_0)x_i)$  and  $a_i$  as follows to obtain  $\beta$ :

$$\hat{\beta}_{3, -, -} = \arg \min_{\beta, \gamma, \gamma_0} \sum_{i=1}^n (b_i - ((\alpha x_i + \alpha_0)x_i) \cdot \beta - a_i \gamma - \gamma_0)^2$$

The goal of this write-up is show that the third estimator is less sensitive to noise in  $a_i$ , through a short math derivation and some simulations.

We make two main assumptions.

**Assumption 1.** For each  $i \in \{1, \dots, n\}$ ,  $(a_i, b_i)$  is independent and identically distributed (i.i.d.).

INITIAL

**Assumption 2.** For each  $i \in \{1, \dots, n\}$ ,  $\mu_{A,i}$  is generated

assume R1 depends on stim

assume adaptation depends on stim

INITIAL

$$\mu_{A,i} = x_i \cdot \alpha^* + \alpha_0 + \epsilon_{\alpha,i}.$$

$\theta_i$  is generated as

$$\theta_i = x_i \cdot \beta^* + \beta_0^* + \epsilon_{\theta,i},$$

where  $\epsilon_{\alpha,i}, \epsilon_{\theta,i}$  are small Gaussian noise  $\mathcal{N}(0, \sigma_\theta^2)$ . Given  $\theta_i$ ,  $(a_i, b_i)$  is a random sample drawn from the joint Gaussian distribution

$$\begin{bmatrix} A \\ B \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \mu_{A,i} \\ \theta_i \mu_{A,i} \end{bmatrix}, \begin{bmatrix} \sigma_A^2 & 0 \\ 0 & \sigma_B^2 \end{bmatrix} \right),$$

with parameters  $\mu_{A,i}, \theta_i, \sigma_A^2$  and  $\sigma_B^2$ . In other words,  $a_i = \mu_{A,i} + \epsilon_A, b_i = \theta_i \mu_{A,i} + \epsilon_B$ ,  $(\epsilon_A, \epsilon_B)$  has a joint Gaussian distribution.

The above assumptions can be quite restrictive in practice, they are open for discussion.

Note that unlike in the previous draft, we made  $\rho$  the correlation between the noise in  $A$  and  $B$  be zero here!!! It is an assumption that makes the third estimator much better, but it is not clear whether can assume this in practice. It would be nice to test.

INITIAL

## 1 The distribution of residuals for true $\beta^*$ in two estimators

Under the above assumptions, for the first estimator, if we plugin the true  $\beta^*$ , we have

$$\begin{aligned} \frac{b_i}{a_i} - x_i \cdot \beta^* - \beta_0^* - 0 &= \frac{\theta_i \mu_A + \epsilon_B}{\mu_A + \epsilon_A} - x_i \beta^* - \beta_0^* - 0 \\ &= \frac{\theta_i(\mu_A + \epsilon_A) - \theta_i \epsilon_A + \epsilon_B}{\mu_A + \epsilon_A} - x_i \beta^* - \beta_0^* - 0 \\ &= \epsilon_{\theta,i} + \frac{-\theta_i \epsilon_A + \epsilon_B}{\mu_A + \epsilon_A} \\ &= \epsilon_{\theta,i} + \underbrace{\frac{-(x_i \beta^* + \epsilon_{\theta,i}) \epsilon_A + \epsilon_B}{\mu_A + \epsilon_A}}_{E_1} x_i \beta^* \end{aligned}$$

**The main concern** is that the division by  $\mu_A + \epsilon_A$  might cause the residual to have high variance. Also because of that division, the best estimator might be biased for estimating  $\beta^*$ .

For the second estimator, if we plugin the true  $\beta^*$ , we have

$$\begin{aligned} b_i - (a_i x_i) \cdot \beta^* - a_i \beta_0^* + 0 &= (\theta_i \mu_A + \epsilon_B) - ((\mu_A + \epsilon_A) x_i) \beta^* - (\mu_A + \epsilon_A) \beta_0^* \\ &= (x_i \beta^* + \beta_0^* + \epsilon_{\theta,i}) \mu_A + \epsilon_B - (\mu_A + \epsilon_A) (x_i \beta^* + \beta_0^*) \\ &= \underbrace{\mu_A \epsilon_{\theta,i} + \epsilon_B + (x_i \beta^* + \beta_0^*) \epsilon_A}_{E_2}. \end{aligned}$$

The residual for true  $\beta^*$  in the second estimator should have smaller variance than that in the previous one, especially in the case where  $\mu_A + \epsilon_A$  can be close to 0. **The main concern** here is that the error  $E_2$  is correlated with  $a_i x_i$  and  $a_i$ .

In the third estimator, the two-stage estimator, the first-stage of estimator decorrelates

## 2 The distribution of estimated $\beta$ 's.

In multivariate linear regression, for response  $\mathbf{y} \in \mathbb{R}^n$  and design matrix  $\mathbf{X} \in \mathbb{R}^{n \times d}$ , the least squares solution has a generic form

$$\hat{\beta} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}.$$

Consider the design matrix  $\begin{bmatrix} x_1 & 1 \\ \vdots & \vdots \\ x_n & 1 \end{bmatrix}$  as fixed and the condition number is not too bad, then

the variance of  $\hat{\beta}_1$  is approximately proportional to the variance of  $E_1$ . We are skipping the exact calculation of the variance, which is feasible but too tedious.

Similarly, if the condition number of the design matrix  $\begin{bmatrix} a_1 x_1 & a_1 & 1 \\ \vdots & \vdots & \vdots \\ a_n x_n & a_n & 1 \end{bmatrix}$  is not too bad,

the variance of  $\hat{\beta}_2$  is approximately proportional to the variance of  $E_2$ .

## 3 Simulations

In this section, we carry out several simulations under the two assumptions 1 and 2 to support/visualize the claims in the previous sections.

### 3.1 Case 1

We simulate  $N = 10000$  data points of  $(a_i, b_i, x_i)$  i.i.d. from Assumption 2 with parameter choices

$$\begin{aligned} x_i &\sim \text{Uniform}[0.4, 1.2] \\ \beta^* &= 1 \\ \alpha^* &= 2 \\ \mu_{A,i} &= \beta^* * x_i + 0.01\mathcal{N}(0, 1) \\ \theta_i &= \alpha^* * x_i + 0.01\mathcal{N}(0, 1) \\ \rho &= 0 \\ \sigma_A &= \sigma_B = 0.3 \end{aligned}$$

We use the  $\beta$  estimate  $\pm 1.96$  standard error as 95% confidence interval (CI), and obtain

$$\begin{aligned} \hat{\beta}_1 &= 0.9041, \quad 95\%CI = [0.829, 0.979] \\ \hat{\beta}_2 &= 1.6683, \quad 95\%CI = [1.640, 1.696] \\ \hat{\beta}_3 &= 0.9989, \quad 95\%CI = [0.985, 1.013] \end{aligned}$$

### Obervations:

- $\hat{\beta}_2$  is clearly wrong, because the residual is correlated with the covariates as we explained above.

- $\hat{\beta}_1$  is OK but the CI is slightly off, because the actual tail is much heavier than Gaussian tail. The problem will become more serious in the next experiment.
- $\hat{\beta}_3$  is good.

### 3.2 Case 2

We increase the variance. We simulate  $N = 10000$  data points of  $(a_i, b_i, x_i)$  i.i.d. from Assumption 2 with parameter choices

$$\begin{aligned}
x_i &\sim \text{Uniform}[0.4, 1.2] \\
\beta^* &= 1 \\
\alpha^* &= 2 \\
\mu_{A,i} &= \beta^* \cdot x_i + 0.01\mathcal{N}(0, 1) \\
\theta_i &= \alpha^* \cdot x_i + 0.01\mathcal{N}(0, 1) \\
\rho &= 0 \\
\sigma_A = \sigma_B &= \textcolor{red}{0.6}
\end{aligned}$$

We use the  $\beta$  estimate  $\pm 1.96$  standard error as 95% confidence interval (CI), and obtain

$$\begin{aligned}
\hat{\beta}_1 &= 1.2201, \quad 95\%CI = [0.692, 1.748] \\
\hat{\beta}_2 &= 1.5791, \quad 95\%CI = [1.541, 1.617] \\
\hat{\beta}_3 &= 0.9982, \quad 95\%CI = [0.978, 1.018]
\end{aligned}$$

#### Observations:

- Again,  $\hat{\beta}_2$  is clearly wrong, same reason as above.
- Now  $\hat{\beta}_1$  has very large standard error, because the error distribution is not Gaussian (because of division by 0) so the error variance is large.
- $\hat{\beta}_3$  is good.

## 4 Takeaways

If the two assumptions are satisfied, clearly Estimator 3 (the two-stage estimator) is a better estimator for estimating  $\beta$ . However, one needs to be careful about the assumptions and check whether these assumptions are reasonable in real data.

1.  $b_i, a_i$  are all approximately Gaussian distributed.
2. There are no correlation in variance for  $b_i$  and  $a_i$ : they are correlated because their mean is related. [it could be a strong assumption](#)
3. Both relationships are approximately linear:

$$\mu_{A,i} = x_i \cdot \alpha^* + \alpha_0 + \epsilon_{\alpha,i}.$$

$$\theta_i = x_i \cdot \beta^* + \beta_0^* + \epsilon_{\theta,i},$$

If a nonlinear model is known to be better, this part may be extended to nonlinear models.