

Halloween_Candy_lab: Spooky Bioinformatics!

Monika Ramos

Exploratory Analysis of Halloween Candy

```
candy_file <- "candy-data.csv"
```

```
candy = read.csv(candy_file, row.names=1)  
head(candy)
```

	chocolate	fruity	caramel	peanutyalmondy	nougat	crispedricewafer
100 Grand	1	0	1	0	0	1
3 Musketeers	1	0	0	0	1	0
One dime	0	0	0	0	0	0
One quarter	0	0	0	0	0	0
Air Heads	0	1	0	0	0	0
Almond Joy	1	0	0	1	0	0

	hard	bar	pluribus	sugarpercent	pricepercent	winpercent
100 Grand	0	1	0	0.732	0.860	66.97173
3 Musketeers	0	1	0	0.604	0.511	67.60294
One dime	0	0	0	0.011	0.116	32.26109
One quarter	0	0	0	0.011	0.511	46.11650
Air Heads	0	0	0	0.906	0.511	52.34146
Almond Joy	0	1	0	0.465	0.767	50.34755

Q1. How many different candy types are in this dataset?

```
dim(candy)
```

```
[1] 85 12
```

There are 85 candy types and 12 candy groups/measurements in the columns.

Q2. How many fruity candy types are in the dataset?

```
sum(candy$fruity)
```

```
[1] 38
```

There are 38 fruity candy types.

2. What is your favorite candy? One of the most interesting variables in the dataset is winpercent. For a given candy this value is the percentage of people who prefer this candy over another randomly chosen candy from the dataset (what 538 term a matchup). Higher values indicate a more popular candy.

```
candy["Twix", ]$winpercent
```

```
[1] 81.64291
```

Q3. What is your favorite candy in the dataset and what is its winpercent value?

```
candy["100 Grand",]$winpercent
```

```
[1] 66.97173
```

My favorite candy is 100 grand bars because it is a hybrid of crunch bars and twix! Its winpercent value is 66.9%.

Q4. What is the winpercent value for "Kit Kat"?

```
candy["Kit Kat",]$winpercent
```

```
[1] 76.7686
```

Kit Kat winpercent value is 76%.

Q5. What is the winpercent value for "Tootsie Roll Snack Bars"?

```
candy["Tootsie Roll Snack Bars",]$winpercent
```

[1] 49.6535

This one is slightly lower than others. Its winpercent is 49.65%

There is a useful `skim()` function in the `skimr` package that can help give you a quick overview of a given dataset. Let's install this package and try it on our candy data.

```
library(skimr)
skim(candy)
```

Table 1: Data summary

Name	candy
Number of rows	85
Number of columns	12
Column type frequency:	
numeric	12
Group variables	None

Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
chocolate	0	1	0.44	0.50	0.00	0.00	0.00	1.00	1.00	
fruity	0	1	0.45	0.50	0.00	0.00	0.00	1.00	1.00	
caramel	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
peanutyalmondy	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
nougat	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
crispedricewafer	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
hard	0	1	0.18	0.38	0.00	0.00	0.00	0.00	1.00	
bar	0	1	0.25	0.43	0.00	0.00	0.00	0.00	1.00	
pluribus	0	1	0.52	0.50	0.00	0.00	1.00	1.00	1.00	
sugarpercent	0	1	0.48	0.28	0.01	0.22	0.47	0.73	0.99	
pricepercent	0	1	0.47	0.29	0.01	0.26	0.47	0.65	0.98	
winpercent	0	1	50.32	14.71	22.45	39.14	47.83	59.86	84.18	

Q6. Is there any variable/column that looks to be on a different scale to the majority of the other columns in the dataset? The histogram is the only column when viewing with `skim()` function that is a character value rather than integers or doubles with the other columns. Some


columns such as chocolate are either 1 or 0 and nothing in between. Mean appears to have the widest range of values.

Q7. What do you think a zero and one represent for the candy\$chocolate column?

I think it is similar to logical values where 1 is yes/TRUE and 0 is No/False. This means is the candy considered a chocolate type of candy? If yes, 1 and If no, 0.

Q8. Plot a histogram of winpercent values

```
hist(candy$winpercent)
```



Q9. Is the distribution of winpercent values symmetrical? The distribution is not symmetrical. In a symmetrical distribution, mean = median = mode and in this case the mean doesn't **exactly** equal the median, though it is close.

Q10. Is the center of the distribution above or below 50%?

```
summary(candy$winpercent)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
22.45	39.14	47.83	50.32	59.86	84.18

The center of distribution is roughly at 50% given that the median is 47.83 and the mean is 50.32. With this information, i would say it is just below 50%.

Q11. On average is chocolate candy higher or lower ranked than fruit candy?

```
mean(candy$winpercent[as.logical(candy$chocolate)])
```

```
[1] 60.92153
```

```
mean(candy$winpercent[as.logical(candy$fruity)])
```

```
[1] 44.11974
```

On average, it appears that chocolate candy is ranked higher than fruity candy.

Q12. Is this difference statistically significant?

```
t.test(x = (candy$winpercent[as.logical(candy$chocolate)]), y=(candy$winpercent[as.logical(candy$fruity)]))
```

Welch Two Sample t-test

```
data: (candy$winpercent[as.logical(candy$chocolate)]) and (candy$winpercent[as.logical(candy$fruity)])
t = 6.2582, df = 68.882, p-value = 2.871e-08
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 11.44563 22.15795
sample estimates:
mean of x mean of y
 60.92153  44.11974
```

The difference is statistically significant with a p-value < 0.05 .

3. Overall Candy Rankings use the base R `order()` function together with `head()` to sort the whole dataset by winpercent

Q13. What are the five least liked candy types in this set?

```
library(tidyverse)
```

```
-- Attaching packages ----- tidyverse 1.3.2 --
v ggplot2 3.3.6      v purrr   0.3.5
v tibble  3.1.8      v dplyr   1.0.10
v tidyr   1.2.1      v stringr 1.4.1
v readr   2.1.3      v forcats 0.5.2
-- Conflicts ----- tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()     masks stats::lag()
```

```
candy_sorted <- arrange(.data = candy, winpercent)
head(candy_sorted)
```

	chocolate	fruity	caramel	peanutyalmondy	nougat
Nik L Nip	0	1	0	0	0
Boston Baked Beans	0	0	0	1	0
Chiclets	0	1	0	0	0
Super Bubble	0	1	0	0	0
Jawbusters	0	1	0	0	0
Root Beer Barrels	0	0	0	0	0

	crispedricewafer	hard	bar	pluribus	sugarpercent	pricepercent
Nik L Nip	0	0	0	1	0.197	0.976
Boston Baked Beans	0	0	0	1	0.313	0.511
Chiclets	0	0	0	1	0.046	0.325
Super Bubble	0	0	0	0	0.162	0.116
Jawbusters	0	1	0	1	0.093	0.511
Root Beer Barrels	0	1	0	1	0.732	0.069

	winpercent
Nik L Nip	22.44534
Boston Baked Beans	23.41782
Chiclets	24.52499
Super Bubble	27.30386
Jawbusters	28.12744
Root Beer Barrels	29.70369

The five least liked candies are Nik L Nip, Boston Baked Beans, Chiclets, Super Bubble and Jawbusters. Ouch, jawbusters are hard to chew!

Q14. What are the top 5 all time favorite candy types out of this set?


```
tail(candy_sorted)
```

	chocolate	fruity	caramel	peanut	almond	nougat
Reese's pieces	1	0	0		1	0
Snickers	1	0	1		1	1
Kit Kat	1	0	0		0	0
Twix	1	0	1		0	0
Reese's Miniatures	1	0	0		1	0
Reese's Peanut Butter cup	1	0	0		1	0
	crisped rice wafer	hard bar	pluribus	sugar	percent	
Reese's pieces		0	0	0	1	0.406
Snickers		0	0	1	0	0.546
Kit Kat		1	0	1	0	0.313
Twix		1	0	1	0	0.546
Reese's Miniatures		0	0	0	0	0.034
Reese's Peanut Butter cup		0	0	0	0	0.720
	price	percent	win	percent		
Reese's pieces	0.651	73.43499				
Snickers	0.651	76.67378				
Kit Kat	0.511	76.76860				
Twix	0.906	81.64291				
Reese's Miniatures	0.279	81.86626				
Reese's Peanut Butter cup	0.651	84.18029				

The top 5 all time favorite candies are Reese's Peanut butter cup, Reese's Miniatures, Twix, Kit Kat, and Snickers.


Q15. Make a first barplot of candy ranking based on winpercent values.

```
library(ggplot2)
ggplot(candy) +
  aes(winpercent, (rownames(candy))) + geom_col()
```



Q16. This is quite ugly, use the `reorder()` function to get the bars sorted by winpercent?


```
ggplot(candy) +  
  aes(winpercent, reorder(rownames(candy), winpercent)) + geom_col()
```

Time to add some useful color. setup a color vector.

```
my_cols=rep("black", nrow(candy))
my_cols[as.logical(candy$chocolate)] = "chocolate"
my_cols[as.logical(candy$bar)] = "brown"
my_cols[as.logical(candy$fruity)] = "pink"

ggplot(candy) +
  aes(winpercent, reorder(rownames(candy),winpercent)) +
  geom_col(fill=my_cols)
```



Q17. What is the worst ranked chocolate candy? The worst ranked is Sixlets.

Q18. What is the best ranked fruity candy?


The best fruity is starburst.

4. Taking a look at pricepercent

```
library(ggrepel)

# How about a plot of price vs win
ggplot(candy) +
  aes(winpercent, pricepercent, label=rownames(candy)) +
  geom_point(col=my_cols) +
  geom_text_repel(col=my_cols, size=3.3, max.overlaps = 5)
```

Warning: ggrepel: 65 unlabeled data points (too many overlaps). Consider increasing max.overlaps



Q19. Which candy type is the highest ranked in terms of winpercent for the least money - i.e. offers the most bang for your buck?

Reeses miniatures

Q20. What are the top 5 most expensive candy types in the dataset and of these which is the least popular?

```
ord <- order(candy$pricepercent, decreasing = TRUE)
head( candy[ord,c(11,12)], n=5 )
```

	pricepercent	winpercent
Nik L Nip	0.976	22.44534
Nestle Smarties	0.976	37.88719
Ring pop	0.965	35.29076
Hershey's Krackel	0.918	62.28448
Hershey's Milk Chocolate	0.918	56.49050


Nik L Nip.

5 Exploring the correlation structure

```
library(corrplot)
```

corrplot 0.92 loaded

```
cij <- cor(candy)
corrplot(cij)
```



Q22. Examining this plot what two variables are anti-correlated (i.e. have minus values)? Chocolate and Fruit are anti-correlated Q23. Similarly, what two variables are most positively correlated? chocolate and bar. also chocolate and winpercent...

6. Principal Component Analysis

```
pca <- prcomp(candy, scale. = T)
summary(pca)
```


Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	2.0788	1.1378	1.1092	1.07533	0.9518	0.81923	0.81530
Proportion of Variance	0.3601	0.1079	0.1025	0.09636	0.0755	0.05593	0.05539
Cumulative Proportion	0.3601	0.4680	0.5705	0.66688	0.7424	0.79830	0.85369


	PC8	PC9	PC10	PC11	PC12
Standard deviation	0.7811	0.6745	0.6625	0.6411	0.6211
Proportion of Variance	0.05539	0.04961	0.04961	0.04961	0.04961
Cumulative Proportion	0.90899	0.95860	1.00821	1.05782	1.10743

Standard deviation	0.74530	0.67824	0.62349	0.43974	0.39760
Proportion of Variance	0.04629	0.03833	0.03239	0.01611	0.01317
Cumulative Proportion	0.89998	0.93832	0.97071	0.98683	1.00000

```
plot(pca$x[,1:2])
```




```
plot(pca$x[,1:2], col=my_cols, pch=16)
```



```
# Make a new data-frame with our PCA results and candy data
my_data <- cbind(candy, pca$x[,1:3])
```

```
p <- ggplot(my_data) +
  aes(x=PC1, y=PC2,
      size=winpercent/100,
      text=rownames(my_data),
      label=rownames(my_data)) +
  geom_point(col=my_cols)
```

p




```
library(ggrepel)
```

```
p + geom_text_repel(size=3.3, col=my_cols, max.overlaps = 7) +
  theme(legend.position = "none") +
  labs(title="Halloween Candy PCA Space",
       subtitle="Colored by type: chocolate bar (dark brown), chocolate other (light brown)",
       caption="Data from 538")
```

Warning: ggrepel: 60 unlabeled data points (too many overlaps). Consider increasing max.overlaps

Halloween Candy PCA Space

Colored by type: chocolate bar (dark brown), chocolate other (light brown),




Data from 538

```
#library(plotly)
#ggplotly(p)
```

finish by taking a quick look at PCA our loadings. Do these make sense to you? Notice the opposite effects of chocolate and fruity and the similar effects of chocolate and bar (i.e. we already know they are correlated).

```
par(mar=c(8,4,2,2))
barplot(pca$rotation[,1], las=2, ylab="PC1 Contribution")
```



Q24. What original variables are picked up strongly by PC1 in the positive direction? Do these make sense to you?

Fruity, hard and pluribus. Yes it makes sense because chocolate is usually individually packaged. Fruity is usually many like sour patch kids and are harder than chocolate since they are hard sugar.