

黑洞

December 19, 2016

1 Black holes in the nuclei of galaxies

Dead stars with masses greater than $3M_{\odot}$ must be black holes,

The only properties which isolated black holes can possess are mass, angular momentum and electric charge.

For a non-rotating black hole, a *Schwarzschild black hole*, there is a spherical surface about the black hole from which electromagnetic radiation suffers an infinite gravitational redshift, as observed from outside this surface. This surface of infinite redshift has radius

$$r_g = \frac{2GM}{c^2} = 3 \left(\frac{M}{M_{\odot}} \right) \text{ km} \quad (1)$$

and is known as the *Schwarzschild radius*. Radiation with frequency ν_0 emitted at radius r from the black hole suffers a gravitational redshift, so that the frequency of the radiation as observed at an infinite distance from the black hole ν_{∞} is

$$\nu_{\infty} = \nu_0 \left(1 - \frac{2GM}{rc^2} \right)^{1/2} = \nu_0 \left(1 - \frac{r_g}{r} \right)^{1/2} \quad (2)$$

There is a last stable circular orbit about a Schwarzschild black hole at radius $r = 3r_g$.

Within this radius, test particles spiral inevitably into the black hole, contributing to its mass and angular momentum. As will be shown below, the **speed of a test particle on the last stable circular orbit** of a Schwarzschild black hole is $v_\phi = c/2$.

In the case of black holes with **finite angular momentum J** , the *Kerr black holes*, the **surface of infinite redshift** occurs at radius

$$r_\infty = \frac{GM}{c^2} + \left[\left(\frac{GM}{c^2} \right)^2 - \left(\frac{J}{Mc} \right)^2 \right]^{1/2} \quad (3)$$

There is a **maximum angular momentum** which a rotating black hole can possess, $J_{\max} = GM^2/c$. The radius of the surface of infinite redshift for a maximally rotating black hole then occurs at $r_\infty = GM/c^2 = r_g/2$, that is, half the Schwarzschild radius of a non-rotating black hole.

There is a last stable orbit about a Kerr black hole, but now test particles can orbit in either the **corotating or counter-rotating directions with respect to the angular momentum axis of the black hole**. For a maximally rotating Kerr black hole, the last stable circular orbit for corotating test particles coincides with r_∞ , that is $r = GM/c^2$, one sixth of the corresponding radius for a non-rotating, Schwarzschild black hole.

The **binding energies of particles on the last stable orbit** for Schwarzschild and maximally rotating Kerr black holes, relative to their rest mass energies, are

$$\text{Schwarzschild} \quad \left[1 - \sqrt{\frac{8}{9}} \right] \quad \text{Kerr} \quad \left[1 - \sqrt{\frac{1}{3}} \right] \quad (4)$$

corresponding to 5.72% and 42.3% of their rest mass energies, respectively. A fraction of the rotational energy of a rotating black hole can also be made available to the external

Universe. In terms of the rest-mass energy of the black hole, this fraction is

$$1 - 2^{-1/2} \left\{ 1 + [1 - (J/J_{\max})^2]^{1/2} \right\}^{1/2} \quad (5)$$

amounting to 29% for a maximally rotating Kerr black hole, $J = J_{\max}$.

2 引力波