

The Effects of Crowding on Metabolic Activity in *Hemigrapsus oregonensis*

Celeste Castaneda-Lopez, Mia Doan, Julia Huber,
Seila Lai, and Lily Peterson


Background

Shoreline Armoring

- 25% of Puget Sound coastline
- 92% of Seattle coastline
- (WA Association of Land Trusts, 2018)

Coastal Squeeze (Pontee, N., 2013)

- Rising sea levels → migration of habitat
- Shoreline armoring reduces intertidal habitat availability (Long et al., 2011)
 - Seawalls
 - Rocks
 - Sandbags



Background

Physical crowding, aggression & metabolic activity

- Higher density increases physical force and defensive displays in hermit crabs (Hazlett, 1968).
- Repeated exposure to intruder conspecifics elevated baseline heart rate in the european shore crab (Rovero et al, 2000).
- Respiration increases during and after agonistic interactions in swimming crabs (Smith & Taylor, 1993).
- Longer fights → greater oxygen demand (Smith & Taylor, 1993).

Water Chemistry, crowding & metabolic activity

- Conspecific chemical cues change the duration and intensity of agonistic interactions in swimming crabs (Smith et al, 1995).
- Shore crab chemical cues can signal mating, predators, and dominance (Hardege, 2002; Trussell, 2003).
- Crowding-related changes in water chemistry increase metabolic stress in shrimp, raising respiration to maintain homeostasis and reducing energy available for growth (Nga et al, 2005).




Fig 1: The mean durations of interactions between *N. puher* exposed to seawater (control) or to water conditioned by females and males. Error bars represent 95% confidence intervals).





Fig 2: Number of agonistic encounters as a function of density in hermit crabs



3

Fig. Example of scaphognathite rate changes of *N. puher* during an agonistic interaction and recovery from it. The interaction started at time 0 and lasted 38 min (from Smith & Taylor, 1993).

Background

Hairy Shore Crab (*Hemigrapsus oregonensis*)

- Model species for intertidal crab studies
- Range: Resurrection Bay to Alaska to Baja California (Visser et al. 2004)
- Habitats: Common intertidal areas like mudflats, bays, estuaries, seagrass beds, open beaches (Visser et al. 2004)

Behavioral Traits & Density

- Typically remain in or near their individual burrows and display aggressive behavior toward other crabs during encounters, especially when foraging outside burrows (Visser et al. 2004)
- Reported densities (Jensen et al. 2002):
 - Avg. 472 crabs/m² under boulders (Puget Sound, WA)
 - Max 624 crabs/m² under oyster shells (Bodega Bay, CA)



Research Questions

How does crowding influence the metabolic activity of *H. oregonensis*?

How does altered water chemistry from crowded conditions influence the metabolic activity of *H. oregonensis*?



Physical Crowding Hypotheses

Null Hypothesis (H_0): Physical crowding does not significantly affect the metabolic activity of *H. oregonensis*.

Alternative Hypothesis (H_1): Physical crowding significantly affects the metabolic activity of *H. oregonensis*.

Alternative Hypothesis (H_{1a}): Physical crowding significantly increases the metabolic activity of *H. oregonensis*.

Alternative Hypothesis (H_{1b}): Physical crowding significantly decreases the metabolic activity of *H. oregonensis*.

Water Chemistry and Crowding Hypotheses

Null Hypothesis (H_0): Altered water chemistry from crowded conditions does not significantly affect the metabolic activity of *H. oregonensis*.


Alternative Hypothesis (H_1): Altered water chemistry from crowded conditions significantly affects the metabolic activity of *H. oregonensis*.

Alternative Hypothesis (H_{1a}): Altered water chemistry from crowded conditions significantly increases the metabolic activity of *H. oregonensis*.

Alternative Hypothesis (H_{1b}): Altered water chemistry from crowded conditions significantly decreases the metabolic activity of *H. oregonensis*.

Experimental Design

- Subject:
 - 10 crabs per tank
- Treatment Groups:
 - Non Crowded condition: Crabs housed in a larger tank (212 cm^2).
 - Crowded condition: Crabs housed in a smaller tank (150 cm^2) to simulate crowding stress.
- Setup:
 - Week 1: 10 crabs are assigned to either crowded or non crowded conditions with fresh water.
 - Week 2 & 3: Randomly selected crabs from each treatment are used. Water from the previous week's crowded treatment is transferred to the non crowded tank of the following week.
- Measurement:
 - Resazurin
 - Time it takes for dye to turn pink and fluorescence using fluorometer
 - Mass of crabs before and after treatment
 - Conditions of water (pH, salinity) before and after treatment



* Three crabs from each tank, along with their corresponding tank water, will be removed for respirometry measurements.

Expected Results




Fig 4: Average fluorescence of resofurin in each treatment group, indicating aerobic respiration per crab.

References

- Nga, B. T., Lürling, M., Peeters, E. T. H. M., Roijackers, R., Scheffer, M., & Nghia, T. T. (2005). Chemical and physical effects of crowding on growth and survival of *Penaeus monodon* Fabricius post-larvae. *Aquaculture*, 246(1), 455–465. <https://doi.org/10.1016/j.aquaculture.2005.02.026>
- Hazlett, B. A. (1968). Effects of Crowding on the Agonistic Behavior of the Hermit Crab *Pagurus Bernhardus*. *Ecology (Durham)*, 49(3), 573–575. <https://doi.org/10.2307/1934131>
- Rovero, F., Hughes, R. N., Whiteley, N. M., & Chelazzi, G. (2000). Estimating the energetic cost of fighting in shore crabs by noninvasive monitoring of heartbeat rate. *Animal Behaviour*, 59(4), 705–713. <https://doi.org/10.1006/anbe.1999.1353>
- Smith, I. P., & Taylor, A. C. (1993). The energetic cost of agonistic behaviour in the velvet swimming crab, *Necora* (= *Liocarcinus*) *puber* (L.). *Animal Behaviour*, 45(2), 375–391. <https://doi.org/10.1006/anbe.1993.1042>
- Huntingford, F. A., Taylor, A. C., Smith, I. P., & Thorpe, K. E. (1995). Behavioural and physiological studies of aggression in swimming crabs. *Journal of Experimental Marine Biology and Ecology*, 193(1), 21–39. [https://doi.org/10.1016/0022-0981\(95\)00108-5](https://doi.org/10.1016/0022-0981(95)00108-5)
- HARDEGE, J. D., JENNINGS, A., HAYDEN, D., MÜLLER, C. T., PASCOE, D., BENTLEY, M. G., & CLARE, A. S. (2002). Novel behavioural assay and partial purification of a female-derived sex pheromone in *Carcinus maenas*. *Marine Ecology. Progress Series (Halstenbek)*, 244, 179–189. <https://doi.org/10.3354/meps244179>
- Trussell, G. C., Ewanchuk, P. J., & Bertness, M. D. (2003). Trait-Mediated Effects in Rocky Intertidal Food Chains: Predator Risk Cues Alter Prey Feeding Rates. *Ecology (Durham)*, 84(3), 629–640. [https://doi.org/10.1890/0012-9658\(2003\)084\[0629:tmeiri\]2.0.co;2](https://doi.org/10.1890/0012-9658(2003)084[0629:tmeiri]2.0.co;2)

References

- JENSEN, G. C., MCDONALD, P. S., & ARMSTRONG, D. A. (2002). East meets west: competitive interactions between green crab *Carcinus maenas*, and native and introduced shore crab *Hemigrapsus* spp. *Marine Ecology. Progress Series (Halstenbek)*, 225, 251–262. <https://doi.org/10.3354/meps225251>
- Visser, E. P., McDonald, P. S., & Armstrong, D. A. (2004). The Impact of Yellow Shore Crabs, *Hemigrapsus oregonensis*, on Early Benthic Phase Dungeness Crabs, *Cancer magister*, in Intertidal Oyster Shell Mitigation Habitat. *Estuaries*, 27(4), 699–715. <https://doi.org/10.1007/BF02907654>
- Boonzaier, K. (2018, May 9). *Beaches rebound in the sound*. Washington Association of Land Trusts. <https://walandtrusts.org/news/beaches-rebound-sound/>
- Long, W. C., Grow, J. N., Majoris, J. E., & Hines, A. H. (2011). Effects of anthropogenic shoreline hardening and invasion by phragmites australis on habitat quality for juvenile blue crabs (*Callinectes sapidus*). *Journal of Experimental Marine Biology and Ecology*, 409(1–2), 215–222. <https://doi.org/10.1016/j.jembe.2011.08.024>
- Pontee, N. (2013). Defining coastal squeeze: A discussion. *Ocean & Coastal Management*, 84, 204–207. <https://doi.org/10.1016/j.ocecoaman.2013.07.010>