



The Waisman Laboratory
for Brain Imaging and Behavior



University of Wisconsin
SCHOOL OF MEDICINE
AND PUBLIC HEALTH

Lecture 1 Simplicial homology and persistent homology




Moo K. Chung

Department of Biostatistics and Medical Informatics
Waisman Laboratory for Brain Imaging and Behavior
University of Wisconsin-Madison

www.stat.wisc.edu/~mchung

Euler characteristic

Topologically equivalent
Deformation-invariant




Sphere

$$\begin{aligned} \text{EC} &= N - E + F \\ &= 8 - 12 + 6 \\ &= 2 \end{aligned}$$

Solid ball

$$\begin{aligned} \text{EC} &= N - E + F - V \\ &= 8 - 12 + 6 - 1 \\ &= 1 \end{aligned}$$


Computing Euler characteristic by parts




Cover an object
with polyhedrons

Incorrect computation


$$\begin{aligned} EC &= N - E + F \\ &= 8 - 8 + 1 \\ &= 1 \end{aligned}$$




$$\begin{aligned} EC &= N - E + F \\ &= (8+4) - (8+6) + (1+1) \\ &= 0 \end{aligned}$$



$$\begin{aligned} EC &= N - E + F \\ &= 8 - (8+2) + (1+1) \\ &= 0 \end{aligned}$$



$$\begin{aligned} EC &= N - E + F \\ &= (8+2) - (8+4) + (1+1) \\ &= 0 \end{aligned}$$





Euler characteristic of a surface mesh from SurfStat

$N - E + F = 2$ for a surface topologically equivalent to a sphere. For each triangle, there are three edges. Since two adjacent triangles share the same edge, the total number of edges is $E = 3F/2$. Hence, we have $F=2N - 4$ for a closed surface.



By checking the **Euler characteristic** of the binary volume of a mandible, holes in the binary volume can be detected. This process is necessary to make the mandible binary volume to be topologically equivalent to a solid sphere.

Topology correction in segmentation



Hole & handles
corrected using
Euler characteristic

Geometry to Topology

Gauss–Bonnet theorem The integral of the Gauss curvature on a 2D Riemannian manifold is $2\pi\chi(\mathcal{M})$, where $\chi(\mathcal{M})$ is the Euler characteristic of \mathcal{M} .

Keith Worsley's random field theory


Random field, stochastic process

$$P\left(\sup_{x \in \mathbb{M}} T(x) > h\right)$$

$$A_h = \{x \in \mathbb{M} : T(x) > h\}$$

$$P\left(\sup_{x \in \mathbb{M}} T(x) > h\right) = \mathbb{E}\chi(A_h)$$

$$\chi(A_h) = \sum_j (-1)^j \beta_j(A_h)$$



Milnor 1963 Morse theory

Adler, 1994 The geometry of random fields

Worsley et al., 1996 Human Brain Mapping

Matlab toolbox

PH-STAT

Statistical Inference on Persistent Homology

<https://github.com/laplcebeltrami/PH-STAT>

Topological Data Analysis

- Branch of data science that uses topology
- Study properties of data that remain invariant under continuous transformations
- Identify underlying patterns using topology

Persistent Homology

- Captures the topological features of data across different scales
- More persistent topological features → signal
- Less persistent topological features → noise
- Multiscale approach

References

Gunnar Carlsson 2009, A User's Guide to Topological Data Analysis



Herbert Edelsbrunner and John L. Harer
Computational Topology: An Introduction 2010,
American Mathematical Society

Chung et al. 2020 Review: Toplogical distance and losses in brain networks arXiv:2102.08623

n -simplex


The basic building block of persistent homology
The smallest convex set containing $n+1$ points

$$\sum_{i=0}^n x_i = 1, x_i \geq 0$$




Simplicial complex

A simplicial complex is a set composed of points, line segments, triangles, and their n-dimensional counterparts.



Simplicial complex




Not simplicial complex


Ex. surface mesh, graphs (including hypergraphs), networks

k-skeleton


A simplicial complex consisting of points and line segments only → graphs & network




Cubical complex (2D or 3D images)




Left central gyrus



4-neighbor connectivity




Lung blood vessel



6-neighbor connectivity

Betti numbers β_i

of i-dimensional
holes/loops




$\beta_0 = \# \text{ of connected components} = 3$

$\beta_1 = \# \text{ of cycles} = 1$

Euler characteristic: $\chi = 3 - 1 = 2$

Betti numbers β_i # of i-dimensional holes/loops



$\beta_0 = \# \text{ of connected components} = 3$
 $\beta_1 = \# \text{ of 1D holes} = 1$
 $\beta_2 = \# \text{ of 2D cavities} = 0$

Betti-number representation:
 $(3, 1, 0, 0, \dots)$

Euler characteristic:

$$\chi = \beta_0 - \beta_1 = 2$$

$\beta_0 = 1, \beta_1 = 2, \beta_2 = 1$
 $(1, 2, 1, 0, 0, \dots)$

Filtration



$$\mathcal{G}_1 \subset \mathcal{G}_2 \subset \mathcal{G}_3 \subset \dots$$

Sequence of nested objects or vector spaces



Extract persistent homological features

Persistent diagram, barcodes

Morse Filtration


Most useful in functional and
time series data

Morse theory for functional data

$$Y = \mu + \epsilon$$

Chung et al., 2009 *Information Processing in Medical Imaging (IPMI)* 5636:386-397.


Unknown signal μ is assumed to be a Morse function: all critical values are unique.




Sublevel set
 $R(y) = \mu^{-1}(-\infty, y]$

Number of connected components $\#R(y)$





Critical values capture the pattern of signal changes



$$f(t) = e(t)$$



$$f(t) = t + e(t)$$



Morse filtration


Consider a sublevel set

$$R(y) = \mu^{-1}(-\infty, y]$$

For critical values

$$b < c$$

$$R(b) \subset R(c)$$




of connected components

$$\#R(b) = \#R(c) - 1$$

Persistence Diagram (PD)

$O(n \log n)$



Pair the time of death with the time of the closest earlier birth.
Birth I is paired to infinity or ignored.

Pairing Brackets

((((0)(0)(0 ((0 0))) 00000((((0))))

((((0)(0)(0 ((0 0))) 00000((((0))))

((((0)(0)(0 ((0 0))) 00000((((0))))


((((0)(0)(0 ((0 0))) 00000((((0))))

((((0)(0)(0 ((0 0))) 00000((((0))))

((((0)(0)(0 ((0 0))) 00000((((0))))

((((0)(0)(0 ((0 0))) 00000((((0))))

Persistence Diagram (PD)







Pair the time of death with the time of the closest earlier birth

Chung et al., 2009
Information Processing
in Medical Imaging
(IPMI) 5636:386-397.

Surface Data




T. Kim⁴

mathematics
behavior
JSA



*First persistent
homology study
applied to medical
imaging*



Persistent homology on cortical manifolds



Rips filtrations

1-skeleton of point cloud data



$\varepsilon = 70\text{mm}$



Recovering underlying topology

1-skeleton of point cloud data



$\epsilon = 70\text{mm}$



Better approach: perform kernel smoothing and then Morse filtration




Persistence Diagram (PD) of a protein molecule

Rips filtration on distance between 8000 atoms







Extremely slow computation → Simply use graph filtration

Bottleneck distance




*For definition, simply
read the assigned reading PDF*





What is wrong with the arbitrary thresholding?

Edge weight ρ_{ij} between node i and j


→ Connectivity matrix $\rho = (\rho_{ij})$




Threshold at 0.5



Threshold at 0.7



Single threshold often suboptimal → TDA-based network analysis



Graph filtrations

Baseline filtration for brain networks first introduced in

Lee et al. 2011 MICCAI 302-309

Lee et al. 2012 IEEE Transactions on Medical Imaging 31:2267-2277

Rips filtration

vs.

graph filtration


Metric space

$$\mathcal{X} = (V, w) \quad w = (w_{ij})$$

Node set Metric

$$w_{ik} < w_{ij} + w_{jk}$$

Rips complex = Simplicial complex



Rips filtration

$$\mathcal{X}_{\epsilon_0} \subset \mathcal{X}_{\epsilon_1} \subset \mathcal{X}_{\epsilon_2} \subset \dots$$

for increased radius


$$\epsilon_0 < \epsilon_1 < \epsilon_2 < \dots$$

Weighted graph

$$\mathcal{X} = (V, w) \quad w = (w_{ij})$$

Node set Edge weight

Binary graph: 1-skeleton




Graph filtration

$$\mathcal{X}_{\epsilon_0} \supset \mathcal{X}_{\epsilon_1} \supset \mathcal{X}_{\epsilon_2} \supset \dots$$

for increased edge weights


$$\epsilon_0 < \epsilon_1 < \epsilon_2 < \dots$$

Graph filtration=single linkage clustering



Graph filtrations on resting-state fMRI

MZ-twins



0.1

0.2

0.3

0.4

0.5

DZ-twins



0.1

0.2

0.3

0.4

0.5


Graph filtration (filtration on 1-skeleton)

Rips filtration is computationally expensive:

For n -nodes, $O(n^{3k+3})$ for the k -th Betti number.



For 1-skeleton, graph filtration is $O(n \log n)$ for both 0-th and 1-st Betti number.

Graph filtration on directed graphs



Building persistent homology on directed graphs
is not trivial and important → [Research project](#)

How to compute the number of cycles in big network data?



Radeon Vega64 eGPU

How many cycles in the network?

Betti numbers


Monotonicity of Betti-0 plot

Monotonicity of β_0


The **deletion** of edge **increases** the the number of connected components by at most 1.

β_0 increases by 0 or 1.


Case 1



Case 2



0-th Betti plot on PET correlation network



24 attention deficit hyperactivity disorder (ADHD) children
26 autism spectrum disorder (ASD) children
11 pediatric control subjects

Monotonicity of Betti-1 plot

Monotonicity of β_1 :

The **deletion** of edge (in the filtration download) decreases the the number of **cycles** by at most 1.
 β_1 decreases by 0 or 1.

Euler characteristic
for l -skeleton:


$$\chi = \beta_0 - \beta_1 = p - q$$

\uparrow \uparrow
nodes edges

$$\beta_1 = \beta_0 - p + q$$

\uparrow \uparrow \uparrow \uparrow
-l, 0 0, +l fixed -l

How Betti numbers change over downward graph filtration



$$\beta_0 = 2$$


$$\beta_0 - \beta_1 = 1$$

$$\beta_1 = 1$$

$$p = 4$$

$$p - q = 1$$

$$q = 3$$



$$\beta_0 = 3$$

$$\beta_0 - \beta_1 = 3$$

$$\beta_1 = 0$$

$$p = 4$$

$$p - q = 3$$

$$q = 1$$

Computation of Betti-plots in practice

Computation of β_0 : Many existing algorithms. Can use a built-in function in MATLAB.

```
[beta_0, S] = graphconncomp(adj)
```


Computation of β_1 : As a function of β_0

$$\beta_1 = \beta_0 - p + q$$

```
q=sum(sum(adj))/2;  
beta_1 = beta_0 - p + q;
```


This is not efficient. Need an incremental algorithm that updates as we delete one edge at a time.

Betti-plots in 116 nodes network




0-cyles

1-cyles



year 2011






year 2019




Birth and death decomposition

Persistence = Life time (death – birth) of a feature

Edges destroy cycles




Edges create components






Theorem Birth & death sets partition the edge set

E_1 Edges destroy cycles



E_0 Edges create components






$$\#(E_1) = 1 + \frac{|V|(|V| - 3)}{2}$$

$$\#(E) = \frac{|V|(|V| - 1)}{2}$$

$$\#(E_0) = |V| - 1$$

Maximum
spanning
tree



$O(|E| \log |V|)$

Hodge Laplacian

Multiplicity of eigenvalues


Multiplicity m of eigenvalue λ :

$$L\mathbf{v}_1 = \lambda\mathbf{v}_1, \dots, L\mathbf{v}_m = \lambda\mathbf{v}_m$$

Theorem: The multiplicity of eigenvalue $\lambda_0 = 0$ is the number of connected components: **0-th Betti number** β_0 .

Equivalently $\beta_0 = p - \text{rank}(L)$

Multiplicity of eigenvalues



>> [F, D] = eig(L)

F =

| | | | | |
|---------|---------|---------|---------|---------|
| 0 | -0.5774 | -0.7071 | 0 | 0.4082 |
| 0 | -0.5774 | 0.0000 | 0 | -0.8165 |
| 0 | -0.5774 | 0.7071 | 0 | 0.4082 |
| -0.7071 | 0 | 0 | -0.7071 | 0 |
| -0.7071 | 0 | 0 | 0.7071 | 0 |

$$L = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

D =

| | | | | |
|---|--------|--------|--------|--------|
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0.0000 | 0 | 0 | 0 |
| 0 | 0 | 1.0000 | 0 | 0 |
| 0 | 0 | 0 | 2.0000 | 0 |
| 0 | 0 | 0 | 0 | 3.0000 |


$$\text{rank}(L) = 3$$

$$\beta_0 = 2$$

Theorem: $\beta_0 = p - \text{rank}(L)$

Proof. $\text{rank}(L) = \text{rank}(MM^\top) = \text{rank}(M^\top)$

$$\ker L = \{\sigma : L\sigma = 0\} = \{\sigma : M^\top \sigma = 0\}$$




$$\sigma = [a, a, b, c, d]^\top$$

$$\begin{aligned}\text{rank}(\ker L) &= 4 \\ \text{rank}(L) &= 1\end{aligned}$$

Graph with 2 disconnected components

$$\ker L = \{\sigma : L\sigma = 0\} = \{\sigma : M^\top \sigma = 0\}$$



$$\sigma = [a, a, a, b, b]^\top$$

$$\text{rank}(\ker L) = \beta_0 = 2$$

$$\text{rank}(L) = p - 2$$

Rank-nullity theorem for graph Laplacian

$$\ker L = \{\sigma : L\sigma = 0\} = \{\sigma : M^\top \sigma = 0\}$$

$$img L = \{L\sigma : \sigma \in \mathbb{R}^p\}$$

$$\text{rank}(img L) + \text{rank}(\ker L) = p$$


$$\text{rank}(L) = n - \beta_0$$

How to find the connected components?

$$\ker L = \{\sigma : L\sigma = 0\} = \{\sigma : M^\top \sigma = 0\}$$

The kernel space is spanned by the eigenvectors corresponding to zero eigenvalue


```
>> [F, D] = eig(L)
```



Boundary operators ∂_k

∂_k Removes the filled-in interior of k -simplexes


$$\partial_k : C_k \rightarrow C_{k-1}$$



$$\partial_2$$

$$\partial_2 \sigma = [v_1, v_2] + [v_2, v_3] + [v_3, v_1]$$

$$\partial_1 \partial_2 \sigma = 0$$



$$\partial_1 \partial_2 \sigma = v_2 - v_1 + v_3 - v_2 + v_1 - v_3 = 0$$

Theorem. $\partial_{k-1} \partial_k \sigma = 0$

Example. Boundary of boundary of a filled-in tetrahedron = edges

→ Does the sum of edges vanish?





Image of boundary operator

$$img \ \partial_{k+1} = \{\partial_{k+1}\sigma : \sigma \in C_{k+1}\}$$

collection of k -boundaries

$$\partial_2\sigma = [v_1, v_2] + [v_2, v_3] + [v_3, v_1] \text{ is I-boundary}$$



Kernel of boundary operator

$$\ker \partial_k = \{\sigma \in C_k : \partial_k \sigma = 0\}$$

collection of k -cycles

$$\partial_1 \partial_2 \sigma = v_2 - v_1 + v_3 - v_2 + v_1 - v_3 = 0$$

$\partial_2 \sigma = [v_1, v_2] + [v_2, v_3] + [v_3, v_1]$ is 1-cycle




Image of boundary operator

From theorem $\partial_k \partial_{k+1} \sigma = 0$
boundary $\partial_{k+1} \sigma$ is always a cycle.

$$\ker \partial_k \supset \text{img} \partial_{k+1}$$

Set of cycles

Set of boundaries

Quotient space:

$$H_k = \ker \partial_k / \text{img} \partial_{k+1}$$

Total number of algebraically
independent cycles
(# of basis).

$$\beta_k = \text{rank}(\ker \partial_k) - \text{rank}(\text{img} \partial_{k+1})$$

Boundary matrix = incidence matrix ∂_k

(i,j) -th entry = 1 if $\tau_i \subset \sigma_j$

Sign depends on the orientation of τ_i


of $(k-l)$ -dimensional simplices τ_i

of k -dimensional simplices σ_j

| | | | | | | | |
|---|---|---|-----|-----|---|---|----|
| | 1 | 0 | 1 | ... | 1 | 0 | 1 |
| • | | | | | | • | |
| • | | | | | | • | |
| • | | | | | | • | |
| | 1 | 1 | | | 0 | 0 | |
| 0 | 1 | 0 | ... | | 0 | 1 | -1 |


∂_k

Boundary matrix ∂_0



$$\partial_0 = \begin{matrix} & v_1 & v_2 & v_3 & v_4 & v_5 \\ 0 & (& 0 & 0 & 0 & 0 & 0 \end{matrix}$$


Boundary matrix ∂_1



$$\partial_1 \partial_2 \sigma = v_2 - v_1 + v_3 - v_2 + v_1 - v_3 = 0$$

$$\partial_1 = \begin{pmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \end{pmatrix} \left(\begin{array}{ccccc} \sigma & e_{12} & e_{23} & e_{31} & e_{24} & e_{41} & e_{45} \\ & -1 & 0 & 1 & 0 & 1 & 0 \\ & 1 & -1 & 0 & -1 & 0 & 0 \\ & 0 & 1 & -1 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 1 & -1 & -1 \\ & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right)$$


Boundary matrix ∂_2



$$\partial_2 \sigma = [v_1, v_2] + [v_2, v_3] + [v_3, v_1]$$


$$\partial_2 = \begin{pmatrix} \sigma & e_{12} & e_{23} & e_{31} \\ & e_{24} & e_{41} & e_{45} \end{pmatrix}$$

Rank nullity theorem for boundary matrix



$$\beta_k = \text{rank}(\ker \partial_k) - \text{rank}(\partial_{k+1})$$

Computing Betti numbers



$$S(\partial_1) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad S(\partial_2) = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\beta_0 = \text{rank}(\ker \partial_0) - \text{rank}(\partial_1) = 5-4$$

$$\beta_1 = \text{rank}(\ker \partial_1) - \text{rank}(\partial_2) = 2-1$$

k-th Hodge Laplacian

$$\Delta_k = \partial_{k+1} \partial_{k+1}^\top + \partial_k^\top \partial_k$$

0th Hodge Laplacian
Graph Laplacian

$$\Delta_0 = \partial_1 \partial_1^\top$$

of nodes
of nodes

~~$$\Delta_1 = \partial_2 \partial_2^\top + \partial_1^\top \partial_1$$~~

1st Hodge Laplacian for graphs:

$$\Delta_1 = \partial_1^\top \partial_1$$

of edges
of edges

k-th Hodge Laplacian

$$\Delta_k = \partial_{k+1} \partial_{k+1}^\top + \partial_k^\top \partial_k$$

0th Hodge Laplacian
Graph Laplacian

$$\Delta_0 = \partial_1 \partial_1^\top$$

of nodes

of nodes

1st Hodge Laplacian


$$\Delta_1 = \partial_2 \partial_2^\top + \partial_1^\top \partial_1$$

of edges


of edges

of edges

of edges



Eigenvectors of Hodge Laplacian of zero eigenvalue



$$Diag(\lambda_0, \dots, \lambda_5) = \begin{pmatrix} 0.0000 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.8299 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2.6889 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3.0000 & 0 & 0 \\ 0 & 0 & 0 & 0 & 4.0000 & 0 \\ 0 & 0 & 0 & 0 & 0 & 4.4812 \end{pmatrix}$$



$$(f_0, \dots, f_5) = \begin{pmatrix} 0.4082 & -0.0000 & 0.0000 & 0.5774 & 0.7071 & -0.0000 \\ -0.2041 & 0.1993 & -0.5765 & 0.5774 & -0.3536 & 0.3578 \\ -0.2041 & -0.1993 & 0.5765 & 0.5774 & -0.3536 & -0.3578 \\ 0.6124 & -0.4325 & 0.1793 & -0.0000 & -0.3536 & 0.5299 \\ 0.6124 & 0.4325 & -0.1793 & -0.0000 & -0.3536 & -0.5299 \\ 0.0000 & -0.7392 & -0.5207 & 0.0000 & 0.0000 & -0.4271 \end{pmatrix}$$

1st Hodge Laplacian for graphs

1st Hodge Laplacian for 2-simplex

$$\Delta_1 = \partial_2 \partial_2^\top + \partial_1^\top \partial_1$$

$$\Delta_1 = \partial_1^\top \partial_1$$



vector representation of l-cycles

$$cycle = \sum_{i < j} a_{ij} e_{ij}$$

| | β_1 | |
|------------|-----------|------|
| # of edges | 2.3 | 1.39 |
| . | ... | . |
| . | . | . |
| . | . | . |
| . | . | . |
| -1.4 | ... | 0 |

Linearly independent columns

Gaussian elimination →

| | β_1 | |
|------------|-----------|------|
| # of edges | 1 | 0... |
| 0 | 0 | . |
| 0 | 1 | 0 |
| 0 | 1 | . |
| 1. | ... | 0 |
| | | 1 |

More meaningful representation

Five biggest cycle differences (male – female) in HCP

