

AdvStDaAn, Worksheet, Week 1

Micheal Lappert

31.03.2022

Contents

Exercise 1	1
Exercise 1.a)	2
Exercise 1.b)	3
Exercise 1.c)	6
Exercise 1.d)	6
Exercise 2	7
Exercise 2.a)	8
STILL TO DO!!	9

Exercise 1

```
path <- file.path('Datasets', 'Softdrink.dat')
df <- read.table(path, header=TRUE)

summary(df)
```

Dataset loading and sanity check:

```
##      Time      volume      distance      location
## Min.   : 8.00   Min.   : 2.00   Min.   : 10.8   Length:25
## 1st Qu.:13.75   1st Qu.: 4.00   1st Qu.: 45.0   Class :character
## Median :18.11   Median : 7.00   Median : 99.0   Mode  :character
## Mean   :22.38   Mean    : 8.76   Mean    :122.8
## 3rd Qu.:21.50   3rd Qu.:10.00   3rd Qu.:181.5
## Max.   :79.24   Max.    :30.00   Max.    :438.0
```

```
head(df)
```

```
##      Time volume distance location
## 1 16.68      7      168 San Diego
## 2 11.50      3       66 San Diego
```

```
## 3 12.03      3      102 San Diego
## 4 14.88      4       24 San Diego
## 5 13.75      6       45 San Diego
## 6 18.11      7       99 San Diego
```

```
tail(df)
```

```
##      Time volume distance      location
## 20 35.10     17    231.0      Austin
## 21 17.90     10     42.0      Austin
## 22 52.32     26    243.0      Austin
## 23 18.75      9    135.0      Austin
## 24 19.83      8    190.5 Minneapolis
## 25 10.75      4     45.0 Minneapolis
```

Data looks just fine.

Exercise 1.a)


```
mod1.1 <- lm(Time ~ volume, data = df)
summary(mod1.1)
```

```
##
## Call:
## lm(formula = Time ~ volume, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -7.5811 -1.8739 -0.3493  2.1807 10.6342
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    3.321      1.371   2.422  0.0237 *
## volume         2.176      0.124  17.546 8.22e-15 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 4.181 on 23 degrees of freedom
## Multiple R-squared:  0.9305, Adjusted R-squared:  0.9275
## F-statistic: 307.8 on 1 and 23 DF,  p-value: 8.22e-15
```


The model looks fine: - Volume is significant on the 5% niveau and the R-squared has a score of 0.93.

We have to do a residual and sensitivity analysis with stochastic simulation to investigate the correctness of the model.

```
plot(mod1.1)
```



```
plot.lmSim(mod1.1, SEED = 1)
```



Interpretation:

1. Tukey-Anscombe plot: Shows outlier with index $i=9$ which affects the smoother. In the simulation it is visible that the original curve is extreme.
=> The expectation of the residuals cannot be constant.
2. Q-Q plot: In the lower as well as in the higher part of the plot some points differ from the straight line. Most of them are within the stochastic fluctuation except $i=9$.
=> The assumption of normal distributed residuals seems violated.
3. Scale-location plot: Shows a clear upwards trend. In the simulation it is visible that the original curve is extreme.
=> The variance of the residuals is not constant.
4. Residuals vs. Leverage: Observations $i = 9$ & 22 have Cooke's Distance > 1 and are therefore too influential. Both observations have additionally too much leverage.
=> Residuals are not normally distributed.

CONCLUSION: The fit is not satisfactory. Trying transformations of response and explanatory variable. Since the nonconstant variance seems to be the most severe problem, log-transformations might help.

Exercise 1.b)

```
df$lVolume <- log(df$volume)
df$lTime <- log(df$Time)

head(df)
```

Tukey's first-aid transformations:

```
##      Time volume distance location lVolume    lTime
## 1 16.68      7      168 San Diego 1.945910 2.814210
## 2 11.50      3       66 San Diego 1.098612 2.442347
## 3 12.03      3      102 San Diego 1.098612 2.487404
## 4 14.88      4       24 San Diego 1.386294 2.700018
```


```
## 5 13.75      6      45 San Diego 1.791759 2.621039
## 6 18.11      7      99 San Diego 1.945910 2.896464
```

Model with transformed variables:


```
mod1.2 <- lm(lTime ~ lVolume, data = df)
summary(mod1.2)
```

```
##
## Call:
## lm(formula = lTime ~ lVolume, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.33794 -0.11068 -0.01232  0.12385  0.36222
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  1.50560    0.10897   13.82 1.26e-12 ***
## lVolume      0.74575    0.05317   14.03 9.25e-13 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.1738 on 23 degrees of freedom
## Multiple R-squared:  0.8953, Adjusted R-squared:  0.8908
## F-statistic: 196.7 on 1 and 23 DF,  p-value: 9.252e-13
```

```
plot(mod1.2)
```



```
plot.lmSim(mod1.2, SEED = 1)
```



Interpretation:

1. Tukey-Anscombe plot: The smoother shows a somewhat strange banana form with the low in the middle which is outside the stochastic fluctuation.
=> The assumption of constant expectation is therefore violated.

2. Q-Q plot: The data scatters nicely around the straight line and seems to be within the stochastic fluctuation.
=> The assumption of Gaussian distributed errors seems not violated.
3. Scale-location plot: The smoother shows a slightly decreasing trend but seems to be ok and lies within the stochastic fluctuation.
=> There is no evidence against the assumption of constant variance of the residuals.
4. Residuals vs. Leverage: All observations have Cook's Distance <1 and therewith no too influential points.
=> No too influential (dangerous) observations

CONCLUSION: The model does still not fit adequately the data, although it is much better than the one before.

An alternative transformation for volume could be the square-root transformation. So let's try out:

```
df$sVolume <- sqrt(df$volume)
```


```
mod1.3 <- lm(Time ~ sVolume, data = df)
summary(mod1.3)
```

```
##
## Call:
## lm(formula = Time ~ sVolume, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.817 -3.266 -1.284  2.509 18.212
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  -17.788      3.510  -5.067 3.95e-05 ***
## sVolume       14.390      1.186  12.133 1.77e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 5.83 on 23 degrees of freedom
## Multiple R-squared:  0.8649, Adjusted R-squared:  0.859
## F-statistic: 147.2 on 1 and 23 DF,  p-value: 1.775e-11
```


The R^2 is with 0.8649 higher than before.

Residual and Sensitivity Analysis:

```
plot(mod1.3)
```



```
plot.lmSim(mod1.3, SEED = 1)
```



Interpretation:

1. Tukey-Anscombe plot: The smoother shows still a somewhat strange banana form with the low in the middle but like that it is inside the stochastic fluctuation.
=> The assumption of constant expectation is not violated.
2. Q-Q plot: The data scatters nicely around the straight line (except $i=9$) and seems to be within the stochastic fluctuation.
=> The assumption of Gaussian distributed errors seems not violated.
3. Scale-location plot: The smoother looks ok and lies within the stochastic fluctuation.
=> There is no evidence against the assumption of constant variance of the residuals.
4. Residuals vs. Leverage: All observations have Cook's Distance < 1 and therewith no too influential points.
=> No too influential (dangerous) observations

CONCLUSION: The model does still fit adequately the data.

Exercise 1.c)

The fitte model in 1.b) is:

$$Time_i = \exp(\beta_0) + \exp(\beta_1) * \sqrt{volume_i} + \exp(E_i)$$

with

$$\mu = 0$$

$$\sigma = \sigma$$

Exercise 1.d)

Extending the model adequately with the second explanatory variable 'distance':


```
df$lDistance <- log(df$distance)
mod1.4 <- lm(lTime ~ sVolume + lDistance, data = df)
summary(mod1.4)
```

```
##
## Call:
## lm(formula = lTime ~ sVolume + lDistance, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.17012 -0.09203 -0.01658  0.07866  0.26425
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.14704    0.14300   8.021 5.65e-08 ***
## sVolume       0.41553    0.03649  11.389 1.08e-10 ***
```


```
## lDistance    0.14401    0.04234    3.401    0.00256 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.1296 on 22 degrees of freedom
## Multiple R-squared:  0.9443, Adjusted R-squared:  0.9393
## F-statistic: 186.6 on 2 and 22 DF,  p-value: 1.587e-14
```

The R^2 increases to 0.9443 which is a really good fit. Lets check the model:

```
plot(mod1.4)
```



```
plot.lmSim(mod1.4, SEED = 1)
```



Interpretation:

1. Tukey-Anscombe plot: The smoother shows a really nice almost straight line which is within the stochastic fluctuation.
=> The assumption of constant expectation is not violated.
2. Q-Q plot: The data scatters nicely around the straight line and seems to be within the stochastic fluctuation.
=> The assumption of Gaussian distributed errors seems not violated.
3. Scale-location plot: The smoother shows a almost straight line and is within the stochastic fluctuation.
=> There is no evidence against the assumption of constant variance of the residuals.
4. Residuals vs. Leverage: All observations have Cook's Distance < 1 and therewith no too influential points. Some are leverage point with leverage $> 2 * 3/25 = 0.24$ (25 examples in dataset)
=> No too influential (dangerous) observations

CONCLUSION: The model does fit adequately the data.

Exercise 2

```
path <- file.path('Datasets', 'Windmill.dat')
df <- read.table(path, header = TRUE)

summary(df)
```

Loading and Checking the data

```
##      velocity      DC.output
## Min.   : 5.482   Min.    :0.123
## 1st Qu.: 8.838   1st Qu.:1.144
## Median :13.424   Median :1.800
## Mean   :13.720   Mean    :1.610
## 3rd Qu.:18.235   3rd Qu.:2.166
## Max.   :22.822   Max.    :2.386
```

```
dim(df)
```

```
## [1] 25  2
```

```
head(df)
```

```
##      velocity DC.output
## 1 11.187073    1.582
## 2 13.424487    1.822
## 3  7.607209    1.057
## 4  6.041019    0.500
## 5 22.374145    2.236
## 6 21.702921    2.386
```

```
tail(df)
```

```
##      velocity DC.output
## 20 12.193909    1.501
## 21 20.360472    2.303
## 22 22.821628    2.310
## 23  9.173400    1.194
## 24  8.837787    1.144
## 25  5.481666    0.123
```

Exercise 2.a)

Start with fitting an ordinary regression model:

```
mod2.1 <- lm(DC.output ~ velocity, data = df)
summary(mod2.1)
```


```
##
## Call:
## lm(formula = DC.output ~ velocity, data = df)
##
```




```
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.59869 -0.14099  0.06059  0.17262  0.32184
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  0.130875   0.125989   1.039   0.31
## velocity    0.107780   0.008514  12.659 7.55e-12 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2361 on 23 degrees of freedom
## Multiple R-squared:  0.8745, Adjusted R-squared:  0.869
## F-statistic: 160.3 on 1 and 23 DF,  p-value: 7.546e-12
```

The model seems to fit not too bad. Lets check this:

```
plot(mod2.1)
```



```
plot.lmSim(mod2.1)
```



STILL TO DO!!

Interpretation:

1. Tukey-Anscombe plot: The smoother shows a really nice almost straight line which is within the stochastic fluctuation.
=> The assumption of constant expectation is not violated.
2. Q-Q plot: The data scatters nicely around the straight line and seems to be within the stochastic fluctuation.
=> The assumption of Gaussian distributed errors seems not violated.
3. Scale-location plot: The smoother shows a almost straight line and is within the stochastic fluctuation.
=> There is no evidence against the assumption of constant variance of the residuals.
4. Residuals vs. Leverage: All observations have Cook's Distance < 1 and therewith no too influential points. Some are leverage point with leverage $> 2 * 3/25 = 0.24$ (25 examples in dataset)
=> No too influential (dangerous) observations

CONCLUSION: The model does fit adequately the data.