



# **Marine Ecological Modelling Global Climate Change**

**Ecological niches and geographic distributions**

Jorge Assis, PhD // theMarineDataScientist, jmassis@ualg.pt  
2020, Centre of Marine Sciences, University of Algarve



# Macroecology

It's a big-picture statistical approach to the study of ecology (not a separate field or sub discipline; coined ~30 years ago).

Focuses on **patterns and processes** operating **at large spatial and temporal scales** and ignores localized and fine-scaled details.

Aims to **uncover the relationship between** complex ecological systems (**organisms, populations, species**) **and their environment**, and to characterise and explain statistical patterns of abundance, distribution and diversity.

e.g.,

*How global climate change may affect marine biodiversity.*



# Macroecology

**Multiple comparisons** across numerous locations, communities, populations and species to **scale up** results and hypotheses from local **to global processes**.

**In opposition to reductionalist experimental approaches**, which have difficulty in generating predictive theory.



## Niche concept is central in macroecology


**Distributional limits are shaped by** constraints on **dispersal** (i.e., the process by which species expand ranges, including both the **movement of individuals** and their **successful establishment**);

With no dispersal limits, every species could potentially be distributed everywhere, and spatial patterns of distribution and diversity would be absent or random.



**Well defined biogeographic patterns of diversity and distribution.**

Following **classification systems based on species composition** the definition of **biogeographic regions** like the **Marine Ecoregions of the World**; nested system of 12 realms, 62 provinces, and 232 ecoregions. A standard for conservation planning units.







# Niche concept is central in macroecology

Hutchinson (1987; now-classic reference), defined the niche as **the hypervolume defined by the environmental dimensions within which a species can survive and reproduce**. The niche as a property of a species, not the environment.

**Fundamental niche:** the response of species to the environment in absence of biotic interactions (competition, predation, facilitation);

**Realized niche:** the environmental dimensions in which a species can survive and reproduce, including the effects of biotic interactions (realised niche smaller than the fundamental niche due to negative interspecific interactions).






Hypervolume of environmental dimensions where species can survive and reproduce (fundamental niche).


e.g.,

A Mediterranean coral with the niche defined by thermal conditions, wave intensity and salinity.




**Movement** also determines whether a species is present in suitable habitats: **a species can be absent from a suitable habitat due to dispersal limitations.** Also, populations may go locally extinct in response to stochasticity and dispersal will determine how fast the empty habitats will be recolonised.





## Movement

**Younger species** (evolutionary speaking) **occupy less of their potential range extents than older species**: species ranges likely determined by time for dispersal, rather than ecological limits on dispersal (i.e., potentially species can get to suitable habitats, but it takes time).




## Which abiotic factors set range limits?










A variety of abiotic factors may set the range limits of species to create biogeographic patterns. These may depend on the part of that species' range that is being considered.

e.g.,

Poleward limits set by tolerance to extreme minimum temperatures, low latitude limits set by maximum temperatures and western limits set by limiting nutrient conditions.



# Mediterranean coral

Predictor (unit)	Oc. range	LP model	LP literature	Relative contribution(%)
Temperature min. (°C)	11.4 - 16.0	<11.5	<12.0 <sup>1,2</sup>	
Temperature max. (°C)	14.1 - 24.4	>25.5	>25.0 <sup>1,3,4</sup>	
Slope (degree)	3.2e <sup>-2</sup> - 22.2	<0.1	steep <sup>1</sup>	
Silicate max. (μmol/L)	1.6 - 19.9	>20.1		
Productivity min. (gC/m <sup>3</sup> /day)	7.5e <sup>-6</sup> - 3.3e <sup>-3</sup>	<4.0e <sup>-6</sup>		
Phosphate min. (μmol/L)	1.6e <sup>-4</sup> - 0.5	<2.2e <sup>-5</sup>	< 0.08 <sup>5</sup>	
Phosphate max. (μmol/L)	2.4e <sup>-2</sup> - 0.77	>0.6		
Nitrate min. (μmol/L)	1.0e <sup>-6</sup> - 5.2	<7.2e <sup>-7</sup>	< 2.0 <sup>5</sup>	
Nitrate max. (μmol/L)	2.5e <sup>-3</sup> - 10.9	>5.4		



## Layer

Temperature

Salinity

Sea ice concentration

Sea ice thickness

Current velocity

Nitrate

Phosphate

Silicate

Dissolved molecular oxygen

Dissolved iron

Chlorophyll

Phytoplankton\*

Primary productivity\*

Light at the bottom

## Abiotic factors setting range limits?

*Non-estuarine cold-temperate fish (N Atlantic)*


*Non-estuarine tropical fish (Equator)*

*Cold-temperate macroalgae (N Atlantic)*

*Warm-temperate seagrass (Mediterranean)*

*Cold-temperate coral (Mediterranean)*




(...)



## Species response functions

Whittaker (1956, 1960, 1967) developed key ideas to analyse the **relationship between the distribution of species and the values of environmental gradients**. It has been hypothesized that species responses are bell-shaped (Gaussian), **equally spaced and of equal amplitude, with their width restricted by competition**.





# Species response functions

Yet, **unimodal responses are the most common**, typically, **positive or negative relationship**. Skewed response curves are expected (physiological stress limiting occurrence at the “harsh”). Bimodal or multimodal response curves have also been hypothesized to result from competition.

Liebig’s law of the minimum suggests that **the true response of species to one factor can only be detected when all other factors are non-limiting**.





# Modelling with abiotic conditions

**Ecophysiological knowledge should guide the modeller** to characterize potential interactions among factors in determining species distributions.

## **My main questions**

Which environmental variables drive the distribution of my model species?

Which response I expect for each environmental variable?

## **My main hypotheses based on ecophysiological knowledge**

(...)




# Niche conservatism

Species range limits are not simply set by unsuitable abiotic and biotic conditions at their range margins, but also by the failure of organisms to adapt to unsuitable conditions.


Niche conservatism is the tendency of species to retain ancestral ecological traits; the idea that **species physiological tolerance limits will remain similar over time.**

(proposed and tested 20 years ago; Peterson et al., 1999)



## Niche conservatism

For a given environmental space, the clade originates as a single ancestral species. In time, there is a niche shift to use an alternate environment. At the present time, there are four species using the ES 1 and two using ES 2. Niche conservatism in species 1-4 limits the individuals of using ES 2.




## Niche conservatism

**Species may adapt** to different conditions over time, but biogeographic patterns show **well defined physiological tolerances at range margins**.

Without such limits, every species could be everywhere, and again there would be few non-random biogeographic patterns.

**Communities become saturated with ecologically similar species over evolutionary time** (e.g. given tens of millions of years): not possible for “new species” to invade the ecological space - competitive exclusion.



## Niche conservatism

An extensive review shows evidence for ecological niche characteristics being highly conserved over short-to-moderate time spans (i.e., from individual life spans up to tens or hundreds of thousands of years).



## Niche conservatism

A clear evidence for niche conservatism is how ongoing warming is shifting the distribution of species globally, particularly at low latitude range edges.

Without niche conservatism, species would persist locally while climate conditions shift.